

CPT and Lorentz Tests with Muons

Robert Bluhm,¹ V. Alan Kostelecký,² and Charles D. Lane²

¹*Physics Department, Colby College, Waterville, Maine 04901*

²*Physics Department, Indiana University, Bloomington, Indiana 47405*

(Received 30 July 1999)

Precision experiments with muons are sensitive to Planck-scale *CPT* and Lorentz violation that is undetectable in other tests. Existing data on the muonium ground-state hyperfine structure and on the muon anomalous magnetic moment could be analyzed to provide dimensionless figures of merit for *CPT* and Lorentz violation at the levels of 4×10^{-21} and 10^{-23} .

PACS numbers: 11.30.Er, 11.30.Cp, 13.40.Em, 14.60.Ef

The minimal standard model of particle physics is *CPT* and Lorentz invariant. However, spontaneous breaking of these symmetries may occur in a more fundamental theory incorporating gravity [1,2]. Minuscule low-energy signals of *CPT* and Lorentz breaking could then emerge in experiments sensitive to effects suppressed by the ratio of a low-energy scale to the Planck scale. At presently attainable energies, the resulting effects would be described by a general standard-model extension [3] that allows for *CPT* and Lorentz violation but otherwise maintains conventional properties of quantum field theory, including gauge invariance, renormalizability, and energy conservation.

In the present work, we study the sensitivity of different muon experiments to *CPT* and Lorentz violation. Planck-scale sensitivity to possible effects is known to be attainable in certain experiments without muons. These include, for example, tests with neutral-meson oscillations [4,5], searches for cosmic birefringence [3,6,7], clock-comparison experiments [8,9], comparisons of particles and antiparticles in Penning traps [10,11], spectroscopic comparisons of hydrogen and antihydrogen [12], measurements of the baryon asymmetry [13], and observations of high-energy cosmic rays [14]. However, in the context of the standard-model extension, dominant effects in the muon sector would be disjoint from those in any of the above experiments because the latter involve only photons, hadrons, and electrons. Moreover, if the size of *CPT* and Lorentz violation scales with mass, high-precision experiments with muons would represent a particularly promising approach to detecting lepton-sector effects from the Planck scale.

The standard *CPT* test involving muons compares the g factors for μ^- and μ^+ , with a bound [15,16] given by the figure of merit

$$r_g^\mu \equiv |g_{\mu^+} - g_{\mu^-}|/g_{av} \lesssim 10^{-8}. \quad (1)$$

We show here that data from experiments normally not associated with *CPT* or Lorentz tests, including muonium microwave spectroscopy [17] and $g - 2$ experiments on μ^+ alone [18], can indeed provide Planck-scale sensitivity to *CPT* and Lorentz violation.

For the experiments considered here, it suffices to consider a quantum-electrodynamics limit of the standard-

model extension incorporating only muons, electrons, and photons. Other terms in the full standard-model extension would be irrelevant or lead only to subdominant effects. In natural units with $\hbar = c = 1$, the Lorentz-violating Lagrangian terms of interest are

$$\begin{aligned} \mathcal{L} = & -a_{\kappa AB} \bar{l}_A \gamma^\kappa l_B - b_{\kappa AB} \bar{l}_A \gamma_5 \gamma^\kappa l_B \\ & - \frac{1}{2} H_{\kappa\lambda AB} \bar{l}_A \sigma^{\kappa\lambda} l_B + \frac{1}{2} i c_{\kappa\lambda AB} \bar{l}_A \gamma^\kappa \overleftrightarrow{D}^\lambda l_B \\ & + \frac{1}{2} i d_{\kappa\lambda AB} \bar{l}_A \gamma_5 \gamma^\kappa \overleftrightarrow{D}^\lambda l_B. \end{aligned} \quad (2)$$

Here, the lepton fields are denoted by l_A with $A = 1, 2$ corresponding to e^- , μ^- , respectively, and $iD_\lambda \equiv i\partial_\lambda - qA_\lambda$ with charge $q = -|e|$. To avoid confusion with four-vector indices, the symbol μ is reserved in this Letter solely as a label for the muon.

The terms associated with the parameters $a_{\kappa AB}$, $b_{\kappa AB}$ are *CPT* odd, while the others are *CPT* even. All the parameters in Eq. (2) are assumed small, and they all are Hermitian 2×2 matrices in flavor space. For example,

$$b_\kappa = \begin{pmatrix} b_\kappa^e & b_\kappa^{e\mu} \\ b_\kappa^{\mu e} & b_\kappa^\mu \end{pmatrix}, \quad (3)$$

where b_κ^e , b_κ^μ are associated with terms preserving lepton number while the others are associated with terms violating it. Since the usual standard model conserves lepton number, leading-order rates for processes that violate lepton number in the standard-model extension must be quadratic in the flavor-nondiagonal parameters $b_\kappa^{e\mu}$, etc. In contrast, processes violating Lorentz symmetry but preserving lepton number can depend linearly on flavor-diagonal parameters b_κ^e , b_κ^μ , etc. This means that experimental bounds from processes preserving lepton number are typically many orders of magnitude sharper than bounds involving lepton-number violation.

Consider first spectroscopic studies of muonium M , which is a μ^+e^- bound state. In experiments at RAL and LANL, precisions of about 20 ppb have been attained both for the $1S-2S$ transition [19] and for the ground-state Zeeman hyperfine transitions [17]. However, we restrict attention here to the latter because the hyperfine transition frequencies are much smaller than the $1S-2S$ ones, which implies better absolute energy resolution and corresponding sensitivity to *CPT* and Lorentz violation [20].

The four hyperfine ground states of M can be labeled 1, 2, 3, 4 in order of decreasing energy. The Zeeman hyperfine transitions ν_{12} , ν_{34} have been measured in a 1.7 T magnetic field [21] with a precision of about 40 Hz (~ 20 ppb), and the hyperfine interval has been extracted.

Since electromagnetic transitions in M conserve lepton number, dominant effects in the standard-model extension arise from flavor-diagonal terms in Eq. (2). For the case of an antimuon μ^+ , the modified Dirac equation is

$$(i\gamma^\lambda D_\lambda - m_\mu + a_\lambda^\mu \gamma^\lambda - b_\lambda^\mu \gamma_5 \gamma^\lambda + \frac{1}{2} H_{\kappa\lambda}^\mu \sigma^{\kappa\lambda} + ic_{\kappa\lambda}^\mu \gamma^\kappa D^\lambda + id_{\kappa\lambda}^\mu \gamma_5 \gamma^\kappa D^\lambda) \psi = 0, \quad (4)$$

where ψ is a four-component μ^+ field of mass m_μ . A similar equation exists for the e^- , containing parameters a_λ^e , b_λ^e , $H_{\kappa\lambda}^e$, $c_{\kappa\lambda}^e$, $d_{\kappa\lambda}^e$. The associated Hamiltonians are found using established procedures [11]. The Coulomb potential in M is $A^\lambda = (|e|/4\pi r, \vec{0})$.

The leading-order Lorentz-violating energy shifts in M can be obtained from these Hamiltonians using perturbation theory and relativistic two-fermion techniques [22]. For the four Zeeman hyperfine levels in a 1.7 T magnetic field, we thereby can determine the corresponding shifts $\delta\nu_{12}$, $\delta\nu_{34}$ in the frequencies ν_{12} , ν_{34} . We find

$$\delta\nu_{12} \approx -\delta\nu_{34} \approx -\tilde{b}_3^\mu/\pi, \quad (5)$$

where $\tilde{b}_3^\mu \equiv b_3^\mu + d_{30}^\mu m_\mu + H_{12}^\mu$. Although in a weak or zero field [23] the results would depend on a combination of both muon and electron parameters for Lorentz violation, only the muon parameters appear in Eq. (5) because in a 1.7 T field the relevant transitions essentially involve pure muon-spin flips. Note that subleading-order Lorentz-violating effects are further suppressed by powers of α or $\mu_\mu B/m_\mu \approx 5 \times 10^{-15}$ and can therefore be neglected.

Since the laboratory frame rotates with the Earth, and since the frequency shifts (5) depend on spatial components of the parameters for CPT and Lorentz violation, the frequencies ν_{12} , ν_{34} oscillate about a mean value with frequency equal to the Earth's sidereal frequency $\Omega \approx 2\pi/(23 \text{ h } 56 \text{ m})$. Note that no signal of this type emerges at any perturbative order in the usual standard model without Lorentz violation. Also, the anticorrelation of the variations of $\delta\nu_{12}$ and $\delta\nu_{34}$ could help exclude environmental systematic effects in analyzing real data.

The result (5) could directly be used to place a bound on CPT and Lorentz violation in the laboratory frame.

However, for purposes of comparison among experiments it is much more useful to work with quantities defined with respect to a nonrotating frame. A suitable choice of basis $\{\hat{X}, \hat{Y}, \hat{Z}\}$ for a nonrotating frame is standard celestial equatorial axes, with the \hat{Z} direction oriented along the Earth's rotational north pole [9]. Then, the laboratory-frame quantity \tilde{b}_3^μ can be written as

$$\tilde{b}_3^\mu = \tilde{b}_Z^\mu \cos\chi + (\tilde{b}_X^\mu \cos\Omega t + \tilde{b}_Y^\mu \sin\Omega t) \sin\chi, \quad (6)$$

where the nonrotating-frame quantity \tilde{b}_J^μ with $J = X, Y, Z$ is defined by $\tilde{b}_J^\mu \equiv b_J^\mu + m_\mu d_{J0}^\mu + \frac{1}{2} \epsilon_{JKL} H_{KL}^\mu$, and where χ is the angle between \hat{Z} and the quantization axis defined by the laboratory magnetic field.

Suppose, for definiteness, that a reanalysis of the data in Ref. [17] using time stamps on the frequency measurements places a bound of 100 Hz on the amplitude of sidereal variations $\delta\nu_{12}$. In terms of nonrotating-frame components, this corresponds to the constraint

$$|\sin\chi| \sqrt{(\tilde{b}_X^\mu)^2 + (\tilde{b}_Y^\mu)^2} \lesssim 2 \times 10^{-22} \text{ GeV}. \quad (7)$$

An appropriate dimensionless figure of merit for this result is the ratio $(r_{\text{hf}}^\mu)_{\text{sidereal}}$ of the amplitude of energy variations to the relativistic energy of M . The bound (7) gives

$$(r_{\text{hf}}^\mu)_{\text{sidereal}} \approx 2\pi |\delta\nu_{12}|/m_\mu \approx 2\pi |\delta\nu_{34}|/m_\mu \lesssim 4 \times 10^{-21}, \quad (8)$$

which is comparable to the dimensionless ratio of the μ^+ mass to the Planck scale M_P , $m_\mu/M_P \approx 10^{-21}$.

We consider next measurements of the muon anomalous magnetic moment [15,16,18]. The most recent experiment [18] measures the angular anomaly frequency ω_a , which is the difference between the spin-precession frequency ω_s and the cyclotron frequency ω_c . This BNL experiment uses relativistic polarized μ^+ moving in a constant 1.45 T magnetic field. The μ^+ have momentum $p \approx 3.09$ GeV and "magic" $\gamma \approx 29.3$, which eliminates the dependence of ω_a on the electric field. Positrons from the decay $\mu^+ \rightarrow e^+ + \nu_e + \bar{\nu}_\mu$ are detected and their decay spectrum is fitted to a specified time function. The anomaly frequency ω_a , which in conventional theory is proportional to $(g-2)/2$, is measured to about 10 ppm. An accuracy below 1 ppm is expected in the near future.

In the standard-model extension, the relativistic Hamiltonian for a μ^+ with an anomalous magnetic moment in a magnetic field \vec{B} is

$$\begin{aligned} \hat{H} = & \gamma^0 \vec{\gamma} \cdot \vec{\pi} + m_\mu (1 - c_{00}^\mu) \gamma^0 + \frac{1}{2} (g-2) \mu_\mu \gamma^0 \vec{\Sigma} \cdot \vec{B} - a_0^\mu - (c_{0j}^\mu + c_{j0}^\mu) \pi^j - [b_0^\mu + (d_{0j}^\mu + d_{j0}^\mu) \pi^j] \gamma_5 \\ & - [a_j^\mu + (c_{jk}^\mu + c_{00}^\mu \delta_{jk}) \pi^k] \gamma^0 \gamma^j - i H_{0j}^\mu \gamma^j - [b_j^\mu + (d_{jk}^\mu + d_{00}^\mu \delta_{jk}) \pi^k] \gamma_5 \gamma^0 \gamma^j + [\frac{1}{2} \epsilon_{jkl} H_{kl}^\mu + m d_{j0}^\mu] \gamma_5 \gamma^j, \end{aligned} \quad (9)$$

where $\Sigma^j = \gamma_5 \gamma^0 \gamma^j$, μ_μ is the muon magneton, and $\vec{\pi} = \vec{p} - q\vec{A}$, with $q = +|e|$ for μ^+ . This Hamiltonian contains no terms that provide leading-order corrections to the g factors for μ^+ or μ^- . Instead, the dominant sensi-

tivity to CPT violation results from the sensitivity to small frequency shifts associated with the spin precession. The conventional figure of merit r_g^μ therefore is zero at leading

order despite the presence of explicit CPT violation, which means alternative figures of merit are needed [11].

A Foldy-Wouthuysen transformation [24] can be used to convert the Hamiltonian \hat{H} to another Hamiltonian \hat{H}' in which the 2×2 off-diagonal blocks contain only first-order terms in the magnetic field \vec{B} [25] and in the parameters for CPT and Lorentz violation. We find $\hat{H}' = \exp(\gamma^0 \gamma_5 \phi) \hat{H} \exp(-\gamma^0 \gamma_5 \phi)$, with $\tan 2\phi = |\vec{\Sigma} \cdot \vec{\pi}|/m_\mu$ and $|\vec{\Sigma} \cdot \vec{\pi}|^2 = \vec{\pi}^2 - q\vec{\Sigma} \cdot \vec{B}$. The off-diagonal blocks in \hat{H}' are irrelevant at leading order since here they produce effects that are at least quadratic in small parameters.

The upper-left 2×2 block of \hat{H}' is the relevant relativistic Hamiltonian for the μ^+ in the laboratory frame [26]. It has the form

$$\hat{H}' = \mathcal{E}_0 + \mathcal{E}_1 + \frac{1}{2} \vec{\sigma} \cdot (\vec{\omega}_{s,0} + f_1 \vec{\beta} + \vec{f}_2), \quad (10)$$

where $\mathcal{E}_0 = \gamma m$ and $\gamma = (1 - \beta^2)^{-1/2}$ with three-velocity $\vec{\beta}$. The term \mathcal{E}_1 contains irrelevant spin-independent corrections. The quantity $\vec{\omega}_{s,0} = (g - 2 + 2/\gamma) \mu_\mu \vec{B}$ is the usual spin-precession frequency. The term $f_1 \vec{\beta}$ is proportional to $\vec{\beta}$, and its contributions average to zero since the detectors in the $(g - 2)$ experiments are spread around the ring and their data are summed. The term \vec{f}_2 depends on the parameters for CPT and Lorentz violation and partially on $\vec{\beta}$, but again only the $\hat{\beta}$ -independent terms are relevant here.

The spin-precession frequency ω_s is calculated as $\frac{d\vec{\sigma}}{dt} = i[\hat{H}', \vec{\sigma}] = \vec{\omega}_s \times \vec{\sigma}$. Since the detectors are in the \hat{x} - \hat{y} plane in the laboratory frame, only the vertical component ω_s is measured. Substituting for \hat{H}' and keeping only the velocity-independent terms along the \hat{z} direction gives for μ^+ the result $\omega_s \approx \omega_{s,0} + 2\check{b}_3^\mu$, where $\check{b}_3^\mu \equiv b_3^\mu/\gamma + m_\mu d_{30}^\mu + H_{12}^\mu$. Note that \check{b}_3^μ reduces to \check{b}_3^μ in the nonrelativistic limit [27].

The cyclotron frequency ω_c is obtained from $[\hat{H}', \vec{r}] = \vec{\pi}/\mathcal{E}_0$, which contains a term $\vec{\omega}_c \times \vec{r}$. However, no leading-order corrections to the usual cyclotron frequency appear: $\omega_c \approx \omega_{c,0} = 2\mu_\mu B/\gamma$. Subleading-order terms do in fact contribute but are of lower order than those in ω_s and therefore can be ignored.

Combining the above results and converting to the nonrotating frame as in Eq. (6), we find the correction to the μ^+ anomaly frequency $\omega_a = \omega_s - \omega_c$ due to CPT and Lorentz violation is

$$\delta\omega_a^{\mu^+} \approx 2\check{b}_Z^\mu \cos\chi + 2(\check{b}_X^\mu \cos\Omega t + \check{b}_Y^\mu \sin\Omega t) \sin\chi, \quad (11)$$

where χ is now the colatitude of the experiment. The corresponding expression $\delta\omega_a^{\mu^-}$ for μ^- is obtained by the substitution $b_J^\mu \rightarrow -b_J^\mu$ in the expressions for $\check{b}_X^\mu, \check{b}_Y^\mu, \check{b}_Z^\mu$.

These results suggest two interesting types of experimental signal. The first involves the difference $\Delta\omega_a^\mu \equiv \delta\omega_a^{\mu^+} - \delta\omega_a^{\mu^-}$, which is $\Delta\omega_a^\mu \approx 4b_3^\mu/\gamma$ in the laboratory frame [28]. It is impractical to measure $g - 2$ for

both μ^+ and μ^- simultaneously, so instead one can directly consider the time-averaged difference $\overline{\Delta\omega_a^\mu}$. In the nonrotating frame,

$$\overline{\Delta\omega_a^\mu} \approx \frac{4}{\gamma} b_Z^\mu \cos\chi. \quad (12)$$

An appropriate figure of merit $r_{\Delta\omega_a^\mu}^\mu$ here is the relative energy difference between μ^+ and μ^- caused by their different spin precessions:

$$r_{\Delta\omega_a^\mu}^\mu \approx \overline{\Delta\omega_a^\mu}/m_\mu. \quad (13)$$

The CERN $g - 2$ experiments compared average μ^+ and μ^- anomaly frequencies, finding [15] $\overline{\Delta\omega_a^\mu}/2\pi \approx 5 \pm 3$ Hz. This gives a value of $r_{\Delta\omega_a^\mu}^\mu$ on the order of 2×10^{-22} , corresponding to $b_Z^\mu \approx (2 \pm 1) \times 10^{-22}$ GeV. A subsequent measurement at BNL [18] provides a μ^+ result within 1 standard deviation of the CERN μ^- result. If the BNL experiment eventually limits the frequency difference to 1 ppm, it would provide a sensitivity at the level of $r_{\Delta\omega_a^\mu}^\mu \lesssim 10^{-23}$, corresponding to $b_Z^\mu \lesssim 10^{-23}$ GeV.

The second interesting type of experimental signal involves sidereal variations in the anomaly frequency. It can be studied using μ^+ alone, in which case time stamps on frequency measurements would permit a bound on sidereal variations of $\omega_a^{\mu^+}$. An appropriate figure of merit $(r_{\omega_a^\mu}^\mu)_{\text{sidereal}}$ is the relative size of the amplitude of energy variations compared to the total energy. Assuming a precision of 1 ppm, we estimate an attainable bound of

$$(r_{\omega_a^\mu}^\mu)_{\text{sidereal}} \approx |\delta\omega_a^{\mu^+}|/m_\mu \lesssim 10^{-23}. \quad (14)$$

The associated bound on parameters in the nonrotating frame is

$$|\sin\chi| \sqrt{(\check{b}_X^\mu)^2 + (\check{b}_Y^\mu)^2} \lesssim 5 \times 10^{-25} \text{ GeV}, \quad (15)$$

which again represents sensitivity to the Planck scale. Note that this test involves different sensitivity to CPT violation than the previous one: the two figures of merit $r_{\Delta\omega_a^\mu}^\mu, (r_{\omega_a^\mu}^\mu)_{\text{sidereal}}$ depend on independent components of parameters for CPT and Lorentz violation.

In addition to effects in flavor-diagonal processes, off-diagonal terms of the type in Eq. (3) arising in the standard-model extension allow Lorentz-violating contributions to flavor-changing processes. For example, precision searches have been performed for the radiative muon decay $\mu \rightarrow e\gamma$, which has a branching ratio below 5×10^{-11} [29]. This decay has previously been analyzed using a CPT - and rotation-invariant model with Lorentz and lepton-number violation that involves terms equivalent (up to field renormalizations) to those of the form $c_{00}^{e\mu}$ and $d_{00}^{e\mu}$ in Eq. (2) [14]. The results of this analysis indicate that combinations of the dimensionless parameters $c_{00}^{e\mu}$ and $d_{00}^{e\mu}$ are bounded at the level of about 10^{-12} by rest-frame muon decays or by muon lifetime measurements in the CERN $g - 2$ experiments, and at the level of about 10^{-19} by constraints from horizontal

air showers of cosmic-ray muons. As expected from the discussion following Eq. (3), these bounds are several orders of magnitude weaker than those from lepton-number preserving processes. An extension of this analysis to include all types of term in Eq. (2) would provide the best existing bounds on the flavor-nondiagonal parameters in the electron-muon sector of the standard-model extension. Useful constraints on these parameters could also be extracted from other future experiments. These include the proposed tests for muon-electron conversion [30], which have an estimated sensitivity to the process $\mu^- + N \rightarrow e^- + N$ of 2×10^{-17} , and the various precision tests that might be envisaged at a future muon collider.

We thank D. Hertzog, K. Jungmann, and B. L. Roberts for discussion. This work is supported in part by the U.S. D.O.E. under Grant No. DE-FG02-91ER40661 and by the N.S.F. under Grant No. PHY-9801869.

-
- [1] V. A. Kostelecký and S. Samuel, Phys. Rev. D **39**, 683 (1989); **40**, 1886 (1989); Phys. Rev. Lett. **63**, 224 (1989); **66**, 1811 (1991); V. A. Kostelecký and R. Potting, Nucl. Phys. **B359**, 545 (1991); Phys. Lett. B **381**, 89 (1996).
- [2] For recent discussions of experimental and theoretical approaches to testing *CPT* and Lorentz symmetry, see, for example, *CPT and Lorentz Symmetry*, edited by V. A. Kostelecký (World Scientific, Singapore, 1999).
- [3] D. Colladay and V. A. Kostelecký, Phys. Rev. D **55**, 6760 (1997); **58**, 116002 (1998).
- [4] B. Schwingerheuer *et al.*, Phys. Rev. Lett. **74**, 4376 (1995); L. K. Gibbons *et al.*, Phys. Rev. D **55**, 6625 (1997); OPAL Collaboration, R. Ackerstaff *et al.*, Z. Phys. C **76**, 401 (1997); DELPHI Collaboration, M. Feindt *et al.*, Report No. DELPHI 97-98 CONF 80, 1997.
- [5] V. A. Kostelecký and R. Potting, in *Gamma Ray-Neutrino Cosmology and Planck Scale Physics*, edited by D. B. Cline (World Scientific, Singapore, 1993) (hep-th/9211116); Phys. Rev. D **51**, 3923 (1995); D. Colladay and V. A. Kostelecký, Phys. Lett. B **344**, 259 (1995); Phys. Rev. D **52**, 6224 (1995); V. A. Kostelecký and R. Van Kooten, Phys. Rev. D **54**, 5585 (1996); V. A. Kostelecký, Phys. Rev. Lett. **80**, 1818 (1998); Phys. Rev. D **61**, 016002 (2000).
- [6] S. M. Carroll, G. B. Field, and R. Jackiw, Phys. Rev. D **41**, 1231 (1990).
- [7] R. Jackiw and V. A. Kostelecký, Phys. Rev. Lett. **82**, 3572 (1999); M. Pérez-Victoria, Phys. Rev. Lett. **83**, 2518 (1999); J. M. Chung, Phys. Lett. B **461**, 138 (1999).
- [8] V. W. Hughes, H. G. Robinson, and V. Beltran-Lopez, Phys. Rev. Lett. **4**, 342 (1960); R. W. P. Drever, Philos. Mag. **6**, 683 (1961); J. D. Prestage *et al.*, Phys. Rev. Lett. **54**, 2387 (1985); S. K. Lamoreaux *et al.*, Phys. Rev. A **39**, 1082 (1989); T. E. Chupp *et al.*, Phys. Rev. Lett. **63**, 1541 (1989); C. J. Berglund *et al.*, Phys. Rev. Lett. **75**, 1879 (1995).
- [9] V. A. Kostelecký and C. D. Lane, Phys. Rev. D **60**, 116010 (1999); J. Math. Phys. (N.Y.) **40**, 6245 (1999).
- [10] R. S. Van Dyck, Jr., P. B. Schwinberg, and H. G. Dehmelt, Phys. Rev. D **34**, 722 (1986); Phys. Rev. Lett. **59**, 26 (1987); G. Gabrielse *et al.*, Phys. Rev. Lett. **82**, 3198 (1999); R. Mittleman *et al.*, in Ref. [2]; Phys. Rev. Lett. **83**, 2116 (1999); H. G. Dehmelt *et al.*, Phys. Rev. Lett. **83**, 4694 (1999).
- [11] R. Bluhm, V. A. Kostelecký, and N. Russell, Phys. Rev. Lett. **79**, 1432 (1997); Phys. Rev. D **57**, 3932 (1998).
- [12] R. Bluhm, V. A. Kostelecký, and N. Russell, Phys. Rev. Lett. **82**, 2254 (1999).
- [13] O. Bertolami *et al.*, Phys. Lett. B **395**, 178 (1997).
- [14] S. Coleman and S. L. Glashow, Phys. Rev. D **59**, 116008 (1999).
- [15] J. Bailey *et al.*, Nucl. Phys. **B150**, 1 (1979).
- [16] F. J. M. Farley and E. Picasso, in *Quantum Electrodynamics*, edited by T. Kinoshita (World Scientific, Singapore, 1990).
- [17] W. Liu *et al.*, Phys. Rev. Lett. **82**, 711 (1999).
- [18] R. M. Carey *et al.*, Phys. Rev. Lett. **82**, 1632 (1999).
- [19] F. E. Maas *et al.*, Phys. Lett. A **187**, 247 (1994); V. Meyer *et al.*, hep-ex/9907013.
- [20] In hydrogen and antihydrogen, a 1S-2S frequency resolution of about 1 mHz might be attainable in principle. However, leading-order sensitivity to *CPT* and Lorentz violation requires magnetic mixing of the 1S and 2S spin states, which introduces experimental difficulties associated with Zeeman field broadening [12]. Similar issues would be relevant for experiments measuring the 1S-2S transition in muonium.
- [21] In these and $g - 2$ experiments, the magnetic field is determined in terms of the proton NMR frequency. In principle, this could introduce additional Lorentz- and *CPT*-violating effects involving the proton. However, clock-comparison experiments [8,9] suggest any such effects are too small to affect the muon tests considered here.
- [22] See, for example, G. Breit, Phys. Rev. **34**, 553 (1929).
- [23] Even in a zero field, the hyperfine interval is split.
- [24] L. Foldy and S. Wouthuysen, Phys. Rev. **78**, 29 (1950).
- [25] H. Mendlowitz and K. M. Case, Phys. Rev. **97**, 33 (1955).
- [26] An expression for the full Hamiltonian \hat{H}' in the nonrelativistic limit is presented in Ref. [9].
- [27] Intuition about this result can be obtained by considering an instantaneous observer Lorentz boost from the muon rest frame to the laboratory. The frequency ω_s^μ scales as γ , b_3^μ is unaffected since it is perpendicular to the boost, and d_{30}^μ and H_{12}^μ each scale as γ . The net effect produces the stated γ dependence in the definition of \check{b}_3^μ .
- [28] Note that $\Delta\omega_a^\mu$ involves only the *CPT*-violating parameter b_3^μ instead of the combination \check{b}_3^μ . The *CPT*-preserving parameters d_{30}^μ and H_{12}^μ cancel in particle-antiparticle comparisons.
- [29] R. D. Bolton *et al.*, Phys. Rev. D **38**, 2077 (1988).
- [30] M. Bachman *et al.*, Brookhaven National Laboratory Proposal, 1997.