

Charged-Particle Multiplicity near Midrapidity in Central Au + Au Collisions at $\sqrt{s_{NN}} = 56$ and 130 GeV

B. B. Back,¹ M. D. Baker,² D. S. Barton,² S. Basilev,⁵ B. D. Bates,⁵ R. Baum,⁹ R. R. Betts,^{1,7} A. Białas,⁴ R. Bindel,⁸
 W. Bogucki,³ A. Budzanowski,³ W. Busza,⁵ A. Carroll,² M. Ceglia,² Y.-H. Chang,⁶ A. E. Chen,⁶ T. Coghen,³
 C. Conner,⁷ W. Czyż,⁴ B. Dąbrowski,³ M. P. Decowski,⁵ M. Despet,³ P. Fita,⁵ J. Fitch,⁵ M. Friedl,⁵ K. Gałuszka,³
 R. Ganz,⁷ E. Garcia,⁸ N. George,¹ J. Godlewski,³ C. Gomes,⁵ E. Griesmayer,⁵ K. Gulbrandsen,⁵ S. Gushue,²
 J. Halik,³ C. Halliwell,⁷ P. Haridas,⁵ A. Hayes,¹⁰ G. A. Heintzelman,² C. Henderson,⁵ R. Hollis,⁷ R. Hołyński,³
 B. Holzman,⁷ E. Johnson,¹⁰ J. Kane,⁵ J. Katzy,^{5,7} W. Kita,³ J. Kotuła,³ H. Kraner,² W. Kucewicz,⁷ P. Kulinič,⁵
 C. Law,⁵ M. Lemler,³ J. Ligocki,³ W. T. Lin,⁶ S. Manly,^{10,11} D. McLeod,⁷ J. Michałowski,³ A. Mignerey,⁸
 J. Mülmenstädt,⁵ M. Neal,⁵ R. Nouicer,⁷ A. Olszewski,^{2,3} R. Pak,² I. C. Park,¹⁰ M. Patel,⁵ H. Pernegger,⁵ M. Plesko,⁵
 C. Reed,⁵ L. P. Remsberg,² M. Reuter,⁷ C. Roland,⁵ G. Roland,⁵ D. Ross,⁵ L. Rosenberg,⁵ J. Ryan,⁵ A. Sanzgiri,¹¹
 P. Sarin,⁵ P. Sawicki,³ J. Scaduto,² J. Shea,⁸ J. Sinacore,² W. Skulski,¹⁰ S. G. Steadman,⁵ G. S. F. Stephans,⁵
 P. Steinberg,² A. Strączek,³ M. Stodulski,³ M. Stręk,³ Z. Stopa,³ A. Sukhanov,² K. Surowiecka,⁵ J.-L. Tang,⁶
 R. Teng,¹⁰ A. Trzupek,³ C. Vale,⁵ G. J. van Nieuwenhuizen,⁵ R. Verdier,⁵ B. Wadsworth,⁵ F. L. H. Wolfs,¹⁰
 B. Wosiek,³ K. Woźniak,³ A. H. Wuosmaa,¹ B. Wysłouch,⁵ K. Zalewski,⁴ and P. Żychowski³

(PHOBOS Collaboration)

¹Physics Division, Argonne National Laboratory, Argonne, Illinois 60439-4843

²Chemistry and C-A Departments, Brookhaven National Laboratory, Upton, New York 11973-5000

³Institute of Nuclear Physics, Kraków, Poland

⁴Department of Physics, Jagellonian University, Kraków, Poland

⁵Laboratory for Nuclear Science, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139-4307

⁶Department of Physics, National Central University, Chung-Li, Taiwan

⁷Department of Physics, University of Illinois at Chicago, Chicago, Illinois 60607-7059

⁸Department of Chemistry and Biochemistry, University of Maryland, College Park, Maryland 20742

⁹Department of Physics, University of Maryland, College Park, Maryland 20742

¹⁰Department of Physics and Astronomy, University of Rochester, Rochester, New York 14627

¹¹Department of Physics, Yale University, New Haven, Connecticut 06520

(Received 19 July 2000)

We present the first measurement of pseudorapidity densities of primary charged particles near midrapidity in Au + Au collisions at $\sqrt{s_{NN}} = 56$ and 130 GeV. For the most central collisions, we find the charged-particle pseudorapidity density to be $dN/d\eta|_{|\eta|<1} = 408 \pm 12(\text{stat}) \pm 30(\text{syst})$ at 56 GeV and $555 \pm 12(\text{stat}) \pm 35(\text{syst})$ at 130 GeV, values that are higher than any previously observed in nuclear collisions. Compared to proton-antiproton collisions, our data show an increase in the pseudorapidity density per participant by more than 40% at the higher energy.

PACS numbers: 25.75.-q

In June 2000, the Relativistic Heavy-Ion Collider (RHIC) at Brookhaven National Laboratory delivered the first collisions between Au nuclei at the highest center of mass energies achieved in the laboratory to date. In this paper we present data taken with the PHOBOS detector during the first collider run at energies of $\sqrt{s_{NN}} = 56$ and 130 GeV. The ultimate goal of our work is to understand the behavior of strongly interacting matter at conditions of extreme density and temperature. Quantum chromodynamics (QCD), the fundamental theory of strong interactions, predicts that for sufficiently high-energy density a new state of matter will be formed, the so-called quark-gluon plasma [1]. The measurements shown here represent the first step toward the development of a full picture of the dynamical evolution of nucleus-nucleus collisions at RHIC energies.

Studying the dependence of charged-particle densities on energy and system size provides information on the interplay between hard parton-parton scattering processes, which can be calculated using perturbative QCD, and soft processes, which are treated by phenomenological models that describe the nonperturbative sector of QCD. Predictions for multiparticle production in high-energy heavy-ion collisions, obtained from a variety of models, typically vary by up to a factor of 2 [2].

In this Letter we report data for the most central Au + Au collisions detected in our apparatus. We have determined the energy dependence of the density of primary charged particles emitted near 90° to the beam axis, characterized by the pseudorapidity density $dN/d\eta|_{|\eta|<1}$, where $\eta = -\ln \tan(\theta/2)$ and θ is the polar angle from the beam axis. These data provide the first means to constrain

models of heavy-ion collisions at RHIC energies. They will allow the extraction of basic information about the initial conditions in these collisions, in particular the energy density, and thus form an essential element for the proper prediction or description of other observables.

The PHOBOS detector employs silicon pad detectors to perform tracking, vertex detection, and multiplicity measurements. Details of the setup and the layout of the silicon sensors can be found elsewhere [3,4]. For the initial running period of the accelerator only a small fraction of the full setup was installed. It included the first six layers of the silicon spectrometer (SPEC), part of the two-layer silicon vertex detector (VTX), and one ladder of the large acceptance Octagon multiplicity detector (see Fig. 1). In total, the installed sensors had 20 000 readout channels, of which less than 2% were nonfunctional. The detector setup also included two sets of 16 scintillator counters (“paddle counters”) located at distances of -3.21 m (PN) and 3.21 m (PP) from the nominal interaction point along the beam (z) axis. These counters subtended pseudorapidities between $3 < |\eta| < 4.5$. They served as the primary event trigger and were used for event selection. Two zero-degree calorimeters (ZDCP, ZDCN) at $z = \pm 18.5$ m provided additional information for event selection by measuring the energy deposited by spectator neutrons.

Monte Carlo (MC) simulations of the detector performance were based on the HIJING event generator [5] and the GEANT 3.21 simulation package, folding in the signal response for scintillator counters and silicon sensors.

In this analysis, we used the data from the SPEC subdetector to reconstruct tracks and to obtain the position, z_{VTX} , of the collision vertex. The SPEC and VTX subdetectors were then used to count the multiplicity of charged particles near $\eta = 0$. Their acceptance, covering $-1 < \eta < 2.5$, is shown in the inset of Fig. 1. The strong dependence on z_{VTX} allows us to check our understanding of acceptance and backgrounds by testing the stability of our results as a function of z_{VTX} .

The readout trigger selected events based on the coincidence of at least one hit in both PP and PN within a




FIG. 1. Detector setup for the initial running period. The inset shows the η acceptance of the SPEC and VTX subdetectors as a function of primary vertex position.

time window of 38 ns. This condition selected most collision events as well as various backgrounds. Off-line event selection was achieved by requiring the paddle time difference to be less than 8 ns, corresponding to a maximum displacement of roughly ± 120 cm relative to $z = 0$. Additional background rejection can be achieved by requiring the ZDC time difference to be less than 20 ns. Since the ZDCs detect only spectator neutrons, they are slightly inefficient for the most central events. Thus, we accepted events with good ZDC timing or high multiplicity in the paddles or both. Double beam backgrounds were studied by using experimental runs where the beams did not collide. We found that the overall rate was less than 1% of all events and that there was no background misidentified as central events, which are characterized by large signals in both paddle counters.

To select the most central events we have used an estimator based on the mean of gain-normalized analog-to-digital converter values in the 16 scintillator counters in PP and PN. Primary charged particles each leave approximately 1.8 MeV in the scintillator. Slow secondary particles that traverse the counters at large angles may deposit larger amounts of energy, and thus mimic a larger multiplicity. To reduce this effect, for each event we discarded the four scintillators in each set of paddle counters with the largest signals. The average of the pulse heights of the remaining 12 scintillators, PP_{12} and PN_{12} , was then calculated for each set. Finally, we used the paddle mean $\frac{1}{2}(PP_{12} + PN_{12})$ as an observable proportional to the number of particles hitting the paddle counters.

Assuming that the number of particles hitting the paddle counters increases monotonically with an increasing number of participants, a cut on the large paddle mean selects the events with the largest number of participants, in both the data and the simulation. This has been confirmed with the ZDCs for our more central events. We observe a good anticorrelation between the paddle mean and the total energy deposited in the ZDCs. For this analysis central events were selected by choosing the 6% of events with the largest paddle mean (see Fig. 2). We obtained a sample of 382 events for $\sqrt{s_{NN}} = 56$ GeV and 724 events for $\sqrt{s_{NN}} = 130$ GeV.

We estimated the average number of participants $\langle N_{\text{part}} \rangle$ chosen by the 6% cut using simulations [5]. MC studies showed that up to 10% of the inelastic cross section fails to produce enough particles to satisfy our trigger conditions. We also studied the distribution of PP_{12} after a 6% cut on PN_{12} and vice versa. This showed the effect of additional fluctuations introduced by slow secondary particles. Considering both of these effects, we estimate that the systematic uncertainty in $\langle N_{\text{part}} \rangle$ is 5%. Applying the 6% centrality cut to MC events we deduce that the number of participants in the data is $\langle N_{\text{part}} \rangle = 330 \pm 4(\text{stat})$ for $\sqrt{s_{NN}} = 56$ GeV and $343 \pm 4(\text{stat})$ for 130 GeV. The rms of the distribution of N_{part} for the MC events is 27 at 56 GeV and 25 at 130 GeV.




FIG. 2. Paddle signal distribution for data events at $\sqrt{s_{NN}} = 130$ GeV.

The analysis proceeded as follows: Using the information from the SPEC sensor planes a road-following tracking algorithm reconstructed straight line tracks passing through at least four layers of the detector and having a χ^2 probability of better than 1% for a nonvertex constrained straight line fit. MC studies show that in this step the tracking efficiency is better than 90%, with less than 10% ghost tracks. For central events, on average 13 tracks were reconstructed at $\sqrt{s_{NN}} = 56$ GeV and 18 tracks at 130 GeV.

The position of the primary collision vertex was determined as the point of closest approach for the found tracks. The distribution of collision vertices, which define the beam orbit, was found to be very stable within each data set, while an offset of 0.7 mm between the two beam energies was observed. To optimize the stability of the vertex finding algorithm, which is essential for the multiplicity determination, we performed a second track finding analysis of the data by confining the vertex position to within 3 mm distance to the previously determined beam orbit in the transverse direction and within $-25 < z_{\text{vtx}} < 15$ cm longitudinally. For central events in this region, MC simulations show that the vertex finding algorithm is more than 99% efficient. As a cross-check we compared the pointing accuracy of the found tracks to the fitted vertex for data and MC and found a most probable value of 1 mm in both cases.

Within the vertex fiducial volume we found a total of 103 central events for the low-energy and 151 central events in the high-energy data set. For the selected events, the multiplicity of charged particles was determined using the SPEC and VTX subdetectors independently. The measurement was done by counting tracklets, which are two hit combinations in consecutive layers of the SPEC or VTX subdetector consistent with a track originating at the primary vertex. For every hit the coordinates in η and ϕ were calculated relative to the fitted vertex position, where ϕ is the azimuthal angle in radians. Then we determined for each hit in the first layer the closest hit in the second layer. Finally, hit combinations with a distance

of $\sqrt{\delta\eta^2 + \delta\phi^2} < 0.015$ in the SPEC subdetector and $|\delta\eta| < 0.1$ in the VTX subdetector were retained as tracklets. To obtain a constraint on possible random channel-by-channel inefficiencies in the sensors and readout chain, we compared the multiplicities of single hits in VTX and SPEC, two-hit tracklets in VTX and SPEC and tracks in SPEC. A detection efficiency of less than 100% would enter in different powers into the different measurements. We find no evidence for inefficiencies beyond the $< 2\%$ of dead channels identified in our standard calibration procedure.

MC studies show that the number of tracklets per event near $\eta = 0$ is approximately proportional to the number of primary charged particles per event. Here we define as primary particles all charged hadrons produced in the collision, including the products of strong and electromagnetic decays, but excluding feed-down products from weak decays and hadrons produced in secondary interactions.

The pseudorapidity density of primary charged particles, $dN/d\eta|_{|\eta|<1}$, was obtained from the observed distribution of tracklets, $dN/d\eta$ (tracklets), using the following procedure: HIJING events were propagated through the GEANT simulation of our setup. The simulated data were then processed with the identical reconstruction chain as the real data, including the event selection based on the paddle counter signals. For each simulated event we determined the number of tracklets, $N_{\text{tracklets}}$, found in the region $|\eta| < 1$ and the number of primary charged particles $N_{\text{primaries}}$ in the same pseudorapidity region. We then determined the vertex dependent proportionality factor $\alpha(z_{\text{vtx}})$ by calculating $\langle N_{\text{tracklets}}/N_{\text{primaries}} \rangle$ as a function of z_{vtx} . For VTX we found $\alpha(z_{\text{vtx}}) \approx 0.1$ and for SPEC $\alpha(z_{\text{vtx}}) \approx 0.06$. The dominant contribution comes from the well known geometrical acceptance of the subdetectors, with all remaining corrections combined contributing only 10% of the value of α . Corrections for inefficiencies and combinatorial background are less than 10%.

MC studies showed that less than 5% of the particles emitted into the angular acceptance of the SPEC and VTX detectors stop in the material of the beam pipe or the first detector layer. The multiplicities we report are corrected for this missing fraction based on the HIJING momentum distributions.

We further studied the contamination of the tracklet distribution by feed-down products from weak decays of strange particles. Because of the proximity of our detectors to the beam line and the good pointing accuracy in the tracklet reconstruction, the contribution was found to be small ($< 4\%$). Again, the multiplicities reported here are corrected based on the HIJING distributions.

Figure 3 shows a direct comparison of the SPEC (left) and VTX (right) tracklet $dN/d\eta$ distributions for data (symbols) and MC events (solid lines), normalized per event. Scaling factors of 1.15 for $\sqrt{s_{NN}} = 56$ GeV and 0.98 for $\sqrt{s_{NN}} = 130$ GeV were applied to the MC distribution to match the integrals to the data. The shape of the




FIG. 3. Tracklet pseudorapidity density in the detector acceptance per event for data at $\sqrt{s_{NN}} = 56$ (circles) and 130 GeV (triangles) for SPEC (left) and VTX (right), in comparison to scaled HIJING simulations (solid lines).

distribution agrees well between simulation and data, confirming our understanding of detector geometry and tracklet efficiency. Based on the comparison of data and MC tracklet distributions, the comparison of results from the SPEC and VTX tracklet analysis and the comparison of tracklet and single hit multiplicities, we estimate the overall systematic uncertainty to be less than 8%. The largest single source of systematic uncertainty is the correction for combinatorial noise in the SPEC subdetector.

As a further cross-check we repeated the analysis procedure using the VENUS [6] event generator and by using $\alpha(z_{VTX})$ obtained from running the event generators at various different energies. For all MC runs, α agreed to within $\pm 2.5\%$, demonstrating the robustness of the tracklet counting procedure.

As a result of the procedure we obtain a primary charged-particle density of $dN/d\eta|_{|\eta|<1} = 408 \pm 12(\text{stat}) \pm 30(\text{syst})$ for $\sqrt{s_{NN}} = 56$ GeV and $555 \pm 12(\text{stat}) \pm 35(\text{syst})$ at $\sqrt{s_{NN}} = 130$ GeV. From the simulation of the paddle counter selection we obtain for the mean number of participating nucleons $\langle N_{\text{part}} \rangle = 330 \pm 4(\text{stat})_{-15}^{+10}(\text{syst})$ for $\sqrt{s_{NN}} = 56$ GeV and $343 \pm 4(\text{stat})_{-14}^{+7}(\text{syst})$ for $\sqrt{s_{NN}} = 130$ GeV.

Normalizing per participant pair, we deduce $dN/d\eta|_{|\eta|<1}/0.5\langle N_{\text{part}} \rangle = 2.47 \pm 0.1(\text{stat}) \pm 0.25(\text{syst})$ and $3.24 \pm 0.1(\text{stat}) \pm 0.25(\text{syst})$, respectively. Taking the strong correlation between the systematic errors at the two energies into account, we observe an increase in the charged-particle density per participant by a factor of $1.31 \pm 0.04(\text{stat}) \pm 0.05(\text{syst})$ from 56 to 130 GeV.

In Fig. 4 we show the normalized yield per participant obtained for Au + Au collisions, proton-antiproton ($p\bar{p}$) collisions [7], and central Pb + Pb collisions at the CERN SPS [8]. The $dN/d\eta$ value for the Pb + Pb data was obtained by numerically integrating the momentum distributions shown in [8].

Several important features of the data emerge: First, the central Au + Au collisions show a significantly larger




FIG. 4. Measured pseudorapidity density normalized per participant pair for central Au + Au collisions. Systematic errors are shown as shaded area. Data are compared with $p\bar{p}$ data and Pb + Pb data from the CERN SPS. Also shown are results of a HIJING simulation (with a line to guide the eye) and a parametrization of the $p\bar{p}$ data [7].

charged-particle density per participant than, for example, nonsingle diffractive (NSD) $p\bar{p}$ collisions at comparable energies. This rules out simple superposition models such as the wounded nucleon model [9] and is compatible with predictions of models like HIJING that include particle production via hard-scattering processes.

Second, the observed increase by 31% from 56 to 130 GeV in central Au + Au collisions is significantly steeper than the increase shown by a $p\bar{p}$ parametrization (see Fig. 4) for the same energy interval [7]. Finally, comparing our data to those obtained at the CERN SPS for Pb + Pb collisions at $\sqrt{s_{NN}} = 17.8$ GeV, we find a 70% higher particle density per participant near $\eta = 0$ at $\sqrt{s_{NN}} = 130$ GeV. General arguments (cf. Bjorken's estimate [10]) suggest that this increase should correspond to a similar increase in the maximal energy density achieved in the collision.

We acknowledge the generous support of the entire RHIC project personnel and the C-A and Chemistry Departments at BNL. We thank Fermilab and CERN for help in silicon detector assembly. We thank the MIT School of Science and LNS for financial support. This work was partially supported by U.S. DoE Grants No. DE-AC02-98CH10886, No. DE-FG02-93ER40802, No. DE-FC02-94ER40818, No. DE-FG02-94ER40865, No. DE-FG02-99ER41099, No. W-31-109-ENG-38, and NSF Grants No. 9603486, No. 9722606, and No. 0072204. The Polish groups were partially supported by KBN Grant No. 2 P03B 04916. The NCU group was partially supported by NSC of Taiwan under Contract No. NSC 89-2112-M-008-024.

- [1] See, for example, J. P. Blaizot, Nucl. Phys. **A661**, 3 (1999).
- [2] *Proceedings of the 14th Quark Matter Conference* [Nucl. Phys. **A661** (1999)].
- [3] B. Back *et al.*, Nucl. Phys. **A661**, 690 (1999).

-
- [4] H. Pernegger *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **419**, 549 (1998).
- [5] X.-N. Wang and M. Gyulassy, Phys. Rev. D **44**, 3501 (1991). We used HIJING v1.35 (April 1998) with standard parameter settings.
- [6] K. Werner, Phys. Rep. **232**, 87 (1993).
- [7] F. Abe *et al.*, Phys. Rev. D **41**, 2330 (1990).
- [8] J. Bächler *et al.*, Nucl. Phys. **A661**, 45 (1999).
- [9] A. Białas, B. Bleszyński, and W. Czyż, Nucl. Phys. **B111**, 461 (1976).
- [10] J.D. Bjorken, Phys. Rev. D **27**, 140 (1983).