

## Intermediate Mass Strangelets are Positively Charged

Jes Madsen

*Institute of Physics and Astronomy, University of Aarhus, DK-8000 Århus C, Denmark*

(Received 18 August 2000)

For a limited range of parameters, stable strange quark matter may be negatively charged in bulk due to one gluon exchange interactions. However, the reduction in strange quark occupation in the surface layer, which is responsible for surface tension, more than compensates this for intermediate mass strangelets, which therefore always have positive quark charge (e.g., for a baryon number between  $10^2$  and  $10^{18}$  assuming  $\alpha_S = 0.9$ ). While details are sensitive to the choice of renormalization, the general conclusion is not. This rules out a scenario where negatively charged strangelets produced in ultrarelativistic heavy ion colliders might grow indefinitely with potentially disastrous consequences.

PACS numbers: 12.38.Mh, 12.39.Ba, 24.85.+p, 25.75.-q

There has recently been some concern about the possibility that a negatively charged strangelet formed in ultrarelativistic heavy ion collisions might grow by absorbing nuclei to, in principle, swallow the Earth. Two investigations have described a number of reasons, theoretical as well as experimental, why such a scenario is exceedingly unlikely [1].

Many unlikely coincidences are required for a dangerous situation to develop. Positively charged quark matter repels ordinary nuclei, so at the very least the small strangelets (baryon number  $A \ll 4 \times 10^2$ , where  $4 \times 10^2$  is the total number of nucleons involved in a collision of two gold or lead nuclei) hypothetically formed in ultrarelativistic heavy ion collisions must have negative quark charge. And for growth to proceed, not only must the ultimate state of stable bulk strange quark matter be negatively charged, but so must the path of intermediate mass strangelets all the way from low  $A$  to  $A \rightarrow \infty$ .

The present study demonstrates that even in the unlikely case where bulk strange quark matter is both stable and has negative quark charge, and where a long-lived, negatively charged strangelet is formed in a collision, such a strangelet *cannot* grow significantly by absorbing nuclei. The reason is that the quark charge of intermediate mass strangelets near the lowest energy state is always positive regardless of the bulk charge. Thus the potentially dangerous strangelet growth path is blocked by Coulomb repulsion.

Following the original suggestions of Bodmer [2] and Witten [3], Farhi and Jaffe [4] presented a detailed study of strange quark matter properties in bulk, as well as finite size strangelets. Concerning the issue of the electric charge of bulk strange matter they showed that the quark charge of stable strange matter is always positive for a small strong interaction constant,  $\alpha_S$  [5]. This is easy to understand physically, because of the fortuitous cancellation of charge for a gas with equal numbers of up, down, and strange quarks. For finite strange quark mass and up and down quark masses close to zero, the number of  $s$  quarks will be reduced, and the net quark charge will be positive, even though the actual numbers are small, with a

typical charge-to-baryon number ratio of  $10^{-3}$ – $10^{-7}$ . One gluon exchange interactions are repulsive for massless up and down quarks but attractive for massive, nonrelativistic quarks. Thus, for sufficiently high  $\alpha_S$  Farhi and Jaffe showed that there is a part of parameter space where this effect increases the abundance of  $s$  quarks enough to give the system a total negative quark charge. Farhi and Jaffe noted that details were quite sensitive to the choice of renormalization point and speculated that it might not be stable with respect to including higher orders in  $\alpha_S$ . But negative quark charge for bulk strange quark matter seems a valid, albeit unlikely, possibility.

The energy and quark content of finite size lumps of strange quark matter (sometimes called strangelets, at least in the low- $A$  regime) are modified relative to that of bulk strange quark matter. These finite size effects (surface tension and curvature energy) generally destabilize strangelets [4,6–10], but they also change the overall charge of the system, and as shown below they always lead to a wide range of intermediate mass strangelets with a positive rather than a negative charge. Physically the main effect of the finite size is a depletion of  $s$  quarks in the surface layer, which removes negative charge from the system relative to the bulk solution. The underlying mechanism is a simple consequence of quantum mechanics. Whereas a relativistic, low-mass quark may have nonzero density at the system boundary, the wave function (and therefore the density) must be zero in the nonrelativistic limit of a very massive quark. The massive  $s$  quarks are therefore suppressed more than  $u$  and  $d$  near the surface [11].

It will be shown below that finite size effects increase the total charge per baryon by an amount  $\Delta Z/A \approx 0.3A^{-1/3}$ . This number is to be compared with a negative charge per baryon (numerically) less than  $10^{-6}$  for stable bulk quark matter with  $\alpha_S = 0.9$ , and  $10^{-3}$  for  $\alpha_S = 1.2$ . Thus the total charge of strangelets becomes positive for  $A < 10^{18}$  and  $A < 10^7$ , respectively.

In the ideal Fermi-gas approximation the energy of a system composed of quark flavors  $i$  is given by

$$E = \sum_i (\Omega_i + N_i \mu_i) + BV. \quad (1)$$

Here  $\Omega_i$ ,  $N_i$ , and  $\mu_i$  denote thermodynamic potentials, total number of quarks, and chemical potentials, respectively.  $B$  is the bag constant;  $V$  is the bag volume. (For simplicity, all thermodynamical equations are written without explicit electron/positron terms. Such terms were included in the numerical calculations.)

In the multiple reflection expansion framework of Balian and Bloch [12], the thermodynamical quantities can be derived from a density of states of the form  $\frac{dN_i}{dk} = 6\left\{\frac{k^2 V}{2\pi^2} + f_S\left(\frac{m_i}{k}\right)kS + f_C\left(\frac{m_i}{k}\right)C + \dots\right\}$ , where area  $S = 4\pi R^2$  and curvature  $C = 8\pi R$  for a sphere.

The volume term is universal, whereas the functions  $f_S$  and  $f_C$  depend on the boundary conditions. The number of quarks of flavor  $i$  is  $N_i = \int_0^{k_{Fi}} (dN_i/dk) dk = n_{i,V}V + n_{i,S}S + n_{i,C}C$ , with Fermi momentum  $k_{Fi} = \mu_i(1 - \lambda_i^2)^{1/2}$ ;  $\lambda_i \equiv m_i/\mu_i$ . The corresponding thermodynamical potentials are  $\Omega_i = \Omega_{i,V}V + \Omega_{i,S}S + \Omega_{i,C}C$ , where  $\partial\Omega_i/\partial\mu_i = -N_i$  and  $\partial\Omega_{i,j}/\partial\mu_i = -n_{i,j}$ . Minimizing the total energy at fixed  $N_i$  gives the energy for a spherical quark lump as  $E = \sum_i N_i \mu_i + (1/3)\Omega_{i,S}S + (2/3)\Omega_{i,C}C$ .

In the following I assume MIT bag-model boundary conditions [13], but the main conclusion depends only on the fact that  $f_S$  is negative.

The volume terms including first order  $\alpha_S$  corrections are given by

$$\begin{aligned} \Omega_{i,V} = & -\frac{\mu_i^4}{4\pi^2} \left( (1 - \lambda_i^2)^{1/2} \left(1 - \frac{5}{2} \lambda_i^2\right) + \frac{3}{2} \lambda_i^4 \ln \frac{1 + (1 - \lambda_i^2)^{1/2}}{\lambda_i} \right. \\ & - \frac{2\alpha_S}{\pi} \left\{ 3\left\{ (1 - \lambda_i^2)^{1/2} - \lambda_i^2 \ln[1 + (1 - \lambda_i^2)^{1/2}] \right\}^2 - 2(1 - \lambda_i^2)^2 - 3\lambda_i^4 \ln^2 \lambda_i + 6 \ln\left(\frac{\sigma}{\mu_i}\right) \right. \\ & \left. \left. \times \left[ \lambda_i^2(1 - \lambda_i^2)^{1/2} - \lambda_i^4 \ln\left(\frac{1 + (1 - \lambda_i^2)^{1/2}}{\lambda_i}\right) \right] \right\} \right), \quad (2) \end{aligned}$$

and  $n_{i,V} = -\partial\Omega_{i,V}/\partial\mu_i$ . The expression for  $\Omega_{i,V}$  is taken from [4] apart from the correction of a sign misprint. Farhi and Jaffe discuss at length the choice of renormalization scale,  $\sigma$ . In accordance with their work I have picked a value of  $\sigma = 313$  MeV in the following. Details of the

numerical results are sensitive to this choice, especially at high  $\alpha_S$ , but calculations for other choices of  $\sigma$  assure that the conclusions are invariant.

The surface contribution follows from  $f_S(m/k) = -[1 - (2/\pi)\tan^{-1}(k/m)]/8\pi$  [6]:

$$\begin{aligned} \Omega_{i,S} = & \frac{3}{4\pi} \mu_i^3 \left[ \frac{(1 - \lambda_i^2)}{6} - \frac{\lambda_i^2(1 - \lambda_i)}{3} \right. \\ & \left. - \frac{1}{3\pi} \left( \tan^{-1} \left[ \frac{(1 - \lambda_i^2)^{1/2}}{\lambda_i} \right] - 2\lambda_i(1 - \lambda_i^2)^{1/2} + \lambda_i^3 \ln \left[ \frac{1 + (1 - \lambda_i^2)^{1/2}}{\lambda_i} \right] \right) \right]; \quad (3) \end{aligned}$$

$$n_{i,S} = -\frac{3}{4\pi} \mu_i^2 \left[ \frac{(1 - \lambda_i^2)}{2} - \frac{1}{\pi} \left( \tan^{-1} \left[ \frac{(1 - \lambda_i^2)^{1/2}}{\lambda_i} \right] - \lambda_i(1 - \lambda_i^2)^{1/2} \right) \right]. \quad (4)$$

For massless quarks  $\Omega_{i,S} = n_{i,S} = 0$ . For massive quarks  $\Omega_{i,S}$  (which is always positive, except for  $\lambda_i = 0$  or 1) has the properties of a surface tension. The corresponding change in quark number per unit area,  $n_{i,S}$ , is always negative, approaching 0 for  $\lambda_i \rightarrow 0$  (massless quarks) or 1 (infinitely massive quarks, i.e., no  $i$  quarks present).

Curvature terms ( $\Omega_{i,C}$  and  $n_{i,C}$ ) are decisive for understanding the average physical properties of small  $A < 100$  strangelets but are not very important for the results at higher  $A$ . They were self-consistently included in the numerical calculations, using the expressions from [10]. Coulomb energies are negligible because of the low charge-to-mass ratio.

Figures 1–4 show results of applying the above equations to strange quark matter at  $\alpha_S = 0.9$ , varying  $B$

and  $m_s$ , and assuming massless  $u$  and  $d$  quarks. Figure 1 shows contour curves for equal energy per baryon of bulk strange quark matter in chemical equilibrium, i.e., with  $\mu_d = \mu_s$  maintained mainly by  $u + d \leftrightarrow s + u$ , and  $\mu_d = \mu_u + \mu_e$ , normally maintained by reactions like  $d \rightarrow u + e^- + \bar{\nu}_e$ . Typical quark chemical potentials are 300 MeV, whereas  $\mu_e$  is a few MeV. A special situation occurs when  $-m_e < \mu_e < m_e$ , since here the weak reactions involving electrons or positrons are blocked, and the total quark charge of the system is zero with neither electrons nor positrons present. For  $\mu_e > m_e$  local charge neutrality is maintained by a small fraction of electrons, whereas positrons take over for  $\mu_e < -m_e$ . Dotted curves in Fig. 1 are contours for a fixed number of electrons per baryon (positrons counted as a negative number of




FIG. 1. Solid curves going from upper left to lower right show constant energy per baryon of bulk strange quark matter in MeV as a function of  $B^{1/4}$  and  $m_s$  for  $\alpha_S = 0.9$ .  $B$  is bounded from below by requiring that  $E/A > 930$  MeV for up-down quark matter ( $m_s \rightarrow \infty$ ). Dotted curves give the total number of electrons per baryon (equal to the net quark charge per baryon). Solid curves labeled 0.51, 0, and  $-0.51$  indicate limits where the electron chemical potential equals  $m_e$ , 0, and  $-m_e$ .

electrons), which equals the net quark charge per baryon,  $Z/A$ . A characteristic property of strange quark matter is a very low  $Z/A$ , except for very high  $m_s$ , where  $s$  quarks are energetically unfavorable. For  $-m_e < \mu_e < m_e$ ,  $Z/A$  becomes 0, but for low  $\alpha_S$  this happens only for unrealistically low  $m_s$ , and negative bulk charge does not occur at all. For high  $\alpha_S$ , the regime of zero charge propagates to intermediate  $m_s$  values, and a region of negative bulk  $Z/A$  shows up. For  $\alpha_S = 0.9$  the bulk  $Z/A$  is still numerically very small, reaching only values of  $-10^{-6}$ , whereas  $Z/A \approx -10^{-3}$  may be reached for a high  $\alpha_S = 1.2$ . It is these regimes of negative (or to a lesser extent zero) charge which could in principle lead to disastrous consequences if a negative strangelet were formed in ultrarelativistic heavy ion collisions and began to grow.

However, the inclusion of finite size effects rules out such a disaster scenario, because they reduce the number of negatively charged  $s$  quarks in the surface layer sufficiently to render the total  $Z/A$  positive, as illustrated in Figs. 2–4.

Quantitatively, stable strangelets are electrically positive for radii from a few fm up to  $10^6$  fm ( $10^2 < A < 10^{18}$ ) for  $\alpha_S = 0.9$ , and for  $10^2 < A < 10^7$  for  $\alpha_S = 1.2$ . This




FIG. 2. As Fig. 1, but for a strangelet with radius  $10^6$  fm, corresponding to  $A \approx 10^{18}$ . Negative quark charge is now limited to the small region in the middle right part of the diagram. For systems smaller than this, finite size effects make the total quark charge positive for all stable strangelets ( $E/A < 930$  MeV).

is seen from the detailed numerical calculations but can be understood qualitatively from the expression for  $n_{s,S}$ , which is the main contributor to the change in charge for  $A > 10^2$ . Thus  $\Delta Z \approx -(1/3)n_{s,S}S \approx 0.18(\mu_s R)^2$  (the number assuming maximum depletion of  $s$  quarks, but the actual depletion is close to this for a wide range of  $\lambda_s$ ). With  $A \approx V\mu_d^3/\pi^2$  and inserting typical numbers for chemical potentials, this leads to

$$\Delta Z/A \approx 0.3A^{-1/3} \approx 0.3R_{\text{fm}}^{-1}, \quad (5)$$

in close agreement with the results of the full numerical calculations.

The numerical results depend on the choice of renormalization, but qualitatively similar results are found for other choices of  $\sigma$ , including the original [14] (but physically questionable [4]) choice of  $\sigma = m_s$ . A perturbative expansion to first order in  $\alpha_S$  is unreliable at high  $\alpha_S$ . In particular, the scheme used here breaks down at  $\alpha_S = \pi/2$ , where light quark pressures become negative, but in practice results for even somewhat lower  $\alpha_S$  should be treated with caution. One might also worry that the finite size terms being known only to zeroth order could be inconsistent, but, since these terms are perturbations on the bulk solution itself, this omission should matter only at order  $\alpha_S^2$ .




FIG. 3. As Fig. 2, but for  $R = 1000$  fm ( $A \approx 10^9$ ). The total quark charge is positive throughout the diagram, typically with  $Z/A \approx 5 \times 10^{-4}$ .

But in spite of these reservations the general qualitative result demonstrated here in quantitative detail to first order in  $\alpha_S$  within the MIT bag model, namely, that surface depletion of  $s$  quarks removes sufficient negative charge to make strangelets positive below some high  $A$  value, seems quite robust. It depends only on the fact that  $f_s$  is negative, which leads to a negative  $n_{s,S}$ , and a positive surface tension. This would seem a natural condition in any model for stable strangelets [11].

Thus the conclusion is that even within the limited range of parameters where bulk strange quark matter is stable and negatively charged, intermediate mass strangelets have positive quark charge because of surface depletion of  $s$  quarks. Even if very small strangelets might still be created with negative charge in ultrarelativistic heavy ion collisions due perhaps to stabilizing shell effects [15] (such effects would play no practical role at high  $A$ ), they would not be able to bridge the positively charged regime of intermediate  $A$  on their way to the putative state of bulk, negatively charged strange quark matter. This rules out the disaster scenarios discussed recently.

I thank Bob Jaffe and Jack Sandweiss for comments. This work was supported in part by the Theoretical Astrophysics Center under the Danish National Research Foundation.




FIG. 4. As Fig. 2, but for  $R = 3$  fm ( $A \approx 30$ ).  $Z/A$  is positive almost everywhere, but a region of negative strangelet charge appears for high  $m_s$ , caused mainly by the increasingly important curvature contributions. Notice the significant increase in energy per baryon typical of small strangelets because of surface and curvature energy contributions. These small strangelets are at most metastable.

- [1] R. L. Jaffe, W. Busza, J. Sandweiss, and F. Wilczek, *Rev. Mod. Phys.* (to be published); A. Dar, A. De Rujula, and U. Heinz, *Phys. Lett. B* **470**, 142 (1999).
- [2] A. R. Bodmer, *Phys. Rev. D* **4**, 1601 (1971).
- [3] E. Witten, *Phys. Rev. D* **30**, 272 (1984).
- [4] E. Farhi and R. L. Jaffe, *Phys. Rev. D* **30**, 2379 (1984).
- [5] The matter is kept locally charge neutral by electrons (positrons). Finite size strangelets are surrounded by a thin atmosphere of electrons (positrons), below which the quark "core" has a net positive (negative) charge.
- [6] M. S. Berger and R. L. Jaffe, *Phys. Rev. C* **35**, 213 (1987); **44**, 566(E) (1991).
- [7] J. Madsen, *Phys. Rev. Lett.* **70**, 391 (1993).
- [8] E. P. Gilson and R. L. Jaffe, *Phys. Rev. Lett.* **71**, 332 (1993).
- [9] J. Madsen, *Phys. Rev. D* **47**, 5156 (1993).
- [10] J. Madsen, *Phys. Rev. D* **50**, 3328 (1994).
- [11] Similar effects explain the positive surface energy in other quantum systems, such as nuclei.
- [12] R. Balian and C. Bloch, *Ann. Phys. (N.Y.)* **60**, 401 (1970).
- [13] T. A. DeGrand, R. L. Jaffe, K. Johnson, and J. Kiskis, *Phys. Rev. D* **12**, 2060 (1975).
- [14] B. A. Freedman and L. D. McLerran, *Phys. Rev. D* **17**, 1109 (1978); V. Baluni, *Phys. Rev. D* **17**, 2092 (1978).
- [15] See, e.g., [8]; J. Schaffner-Bielich, C. Greiner, A. Diener, and H. Stöcker, *Phys. Rev. C* **55**, 3038 (1997).