

Resolving the Antibaryon-Production Puzzle in High-Energy Heavy-Ion Collisions

Ralf Rapp and Edward V. Shuryak

Department of Physics and Astronomy, State University of New York, Stony Brook, New York 11794-3800
(Received 14 September 2000)

We argue that the observed antiproton production in heavy-ion collisions at CERN-SPS energies can be understood if (contrary to most sequential scattering approaches) the backward direction in the process $p\bar{p} \leftrightarrow \bar{n}\pi$ (with $\bar{n} = 5-6$) is consistently accounted for within a thermal framework. Employing the standard picture of subsequent chemical and thermal freezeout, which induces an oversaturation of pion number with associated chemical potentials of $\mu_\pi \approx 60-80$ MeV, enhances the backward reaction substantially. The resulting rates turn out to be large enough to maintain an antiproton abundance at thermal freezeout in accordance with the measured \bar{p}/p ratio in Pb(158A GeV) + Pb collisions.

DOI: 10.1103/PhysRevLett.86.2980

PACS numbers: 25.75.-q

During the past decade remarkable progress has been made in the understanding of the dynamics of strong interactions probed through (ultra-)relativistic heavy-ion collisions. Although the main challenge of an unambiguous identification of the quantum chromodynamic (QCD) phase transition to the quark-gluon plasma (QGP) persists, we have greatly advanced our knowledge on properties of highly excited hadronic matter close to the expected phase boundary. A variety of collective phenomena has been observed, indicating that the produced systems have indeed reached macroscopically large sizes, justifying the use of equilibrium techniques such as thermodynamics and hydrodynamics.

One of the important results that will be used below is that the final-state hadron abundances, including antibaryons, can be rather accurately characterized by the so-called *chemical freezeout* stage [1] with a common temperature T_{ch} and baryon chemical potential μ_B^{ch} , the specific values depending on collision energy [note that a precise description of all hadron species requires corrections to an ideal gas ensemble, e.g., excluded (eigen-)volumes [1,2] to mimic short-range repulsions, or “strangeness suppression” factors [2]. Such corrections are not important for our subsequent analysis and will be neglected. For a contrasting view of hadron production in heavy-ion collisions, see, e.g., Ref. [3]].

At CERN-SPS energies, the chemical freezeout is clearly distinct from the *thermal* one (with an associated temperature $T_{\text{th}} < T_{\text{ch}}$), from where the particles stream freely to the detectors. This follows from the kinetics of hadronic reactions [4,5], i.e., a significant difference between elastic and inelastic collision rates at low relative energies, and has been confirmed by experimental evidence (see, e.g., Ref. [6]). In central Pb + Pb collisions, e.g., a nucleon at midrapidity is elastically rescattered on average about 10–15 times, but less than once inelastically [7]. Also, as shown in [5], most of the collective flow effects at SPS are generated *in between* the two freezeouts, and their observation leaves no doubt about the existence of such an intermediate stage.

Another consequence is that abundances of secondary mesons (pions, kaons, etc.) are not subject to significant changes when the system evolves from T_{ch} to T_{th} . Again, this is conceivable as the rates for number changing reactions are too small to maintain chemical equilibrium on the time scales of the hadronic fireball lifetime, $t \approx 10$ fm/c. Since, on the other hand, elastic scattering (e.g., $\pi\pi \rightarrow \rho \rightarrow \pi\pi$) is still effective, thermal equilibrium is approximately maintained. In statistical mechanics language such a scenario entails additional conservation laws, which can be implemented via effective (pion-, kaon-, etc.) chemical potentials to guarantee fixed particle numbers. For pions (kaons) typical values of $\mu_\pi = 60-80$ MeV ($\mu_K = 100-130$ MeV) are reached at $T_{\text{th}} = 110-120$ MeV (see [4,5,8]). The ensuing “oversaturation” of the pion phase space plays a key role in what follows.

The situation for antibaryons is different from mesonic secondaries as the pertinent *inelastic* (or annihilation) cross section is *not* small. At the relevant (thermal) energies in the comoving frame of collective expansion, $\sqrt{s} = 2(m_N + E_N^{\text{th}}) \approx 2.3$ GeV, one has $\sigma_{p\bar{p} \rightarrow n\pi} \approx 50$ mb. Taking an average baryon density of $\rho_B = 0.75\rho_0$ ($\rho_0 = 0.16$ fm $^{-3}$) around $T = 150$ MeV in the course of the hadronic evolution and a typical (anti-)proton velocity of $v_{\text{th}} = p/E_{\text{tot}} = 0.56c$ [from $E_p^{\text{th}} = (3/2)kT \approx 225$ MeV], we obtain, for the chemical equilibration time scale,

$$\tau_{\text{ch}} = \frac{1}{\sigma_{\text{ann}} \rho_B v_{\text{th}}} \approx 3 \text{ fm}/c. \quad (1)$$

This is well below the fireball lifetime in the pure hadronic phase of $\tau_{\text{had}} \approx 7$ fm/c [5,8], cf. Fig. 1; in other words, only a fraction of $\exp(-\tau_{\text{had}}/\tau_{\text{ch}}) \approx 10\%$ does *not* rescatter towards thermal freezeout. Thus, in spite of the good agreement of the final \bar{p} yields with the standard chemical freezeout predictions, antibaryon production ought to be reconsidered.

Naively one might expect most of the antiprotons to be annihilated, and various transport calculations (e.g., a




FIG. 1. Chemical relaxation time (solid line) for antiprotons in central Pb + Pb collisions at SPS energies employing an annihilation cross section of $\sigma_{\text{ann}} = (40-50)$ mb and a baryon-density profile as obtained within an isentropically expanding fireball model [8]. The dash-dotted curve indicates the remaining hadronic fireball lifetime until thermal freezeout.

relativistic cascade [9] and ultrarelativistic quantum molecular dynamics [10]) have indeed been unable to account for the measured number, falling short by significant factors. Consequently, speculations have been raised that the puzzle could be resolved by certain in-medium effects leading to either a reduction of the annihilation rate [9] or an enhanced production [10].

A generic problem with transport/cascade simulations is that, although multibody resonance decays are included, the inverse reactions are not. As is well known, this violates detailed balance and, in principle, prevents the simulations from reaching the proper thermodynamic limit [11]. The discussions are typically focused on three-body reactions; in the present context it is even 5- or 6-pion collisions which are most relevant for producing baryon-antibaryon pairs. A widespread belief is that those reactions have insignificant rates under the conditions probed in the hadronic stages of heavy-ion collisions. However, as we will argue below, this is not the case; inverse reactions have to be addressed and are, in fact, capable of explaining the observed antiproton yield (although the \bar{p} enhancement in nucleus-nucleus over p - p collisions thus loses its proposed direct relation to QGP formation, the latter is by no means excluded by the subsequent arguments).

The expression for the thermal reaction rate for the process $p\bar{p} \leftrightarrow n\pi$ can be written as (see also Ref. [12])

$$\mathcal{R}_{\text{th}} = \int d^3\tilde{k}_p d^3\tilde{k}_{\bar{p}} d^3\tilde{k}_{\pi_1} \dots d^3\tilde{k}_{\pi_n} \delta^{(4)}(K_{\text{tot}}) |\mathcal{M}_n|^2 \times \left\{ z_p z_{\bar{p}} \exp\left[-\frac{E_p + E_{\bar{p}}}{T}\right] - z_{\pi}^n \exp\left[-\sum_{i=1}^n \frac{\omega_{\pi,i}}{T}\right] \right\}, \quad (2)$$

where \mathcal{M}_n denotes the invariant scattering matrix element (which, of course, is identical for the backward and forward directions due to time-reversal invariance of strong interactions), $d^3\tilde{k}_x = d^3k_x/(2\pi)^3$ are the phase space integrations, and $K_{\text{tot}} = k_p + k_{\bar{p}} - k_{\pi_1} - \dots - k_{\pi_n}$ is the total four-momentum; $z_x = e^{\mu_x/T}$ are the fugacities of particle species x (in Boltzmann approximation), and n is the number of pions produced in a $\bar{p}p$ annihilation at a given energy (to be discussed in more detail below). For a nonvanishing net nucleon number and in chemical equilibrium, one has $\mu_N > 0$, $\mu_{\bar{N}} = -\mu_N$, and $\mu_{\pi} = 0$. Under SPS conditions, typical values at chemical freezeout are [1] $(T_{\text{ch}}, \mu_N^{\text{ch}}) \simeq (170, 260)$ MeV, which results in an antiproton-to-proton ratio of

$$\frac{\bar{p}}{p} \propto \frac{\exp[-(E_{\bar{p}} + \mu_N)/T]}{\exp[-(E_p - \mu_N)/T]} = \exp[-2\mu_N/T] = 4.7\%, \quad (3)$$

consistent with the experimentally measured value of $(5.5 \pm 1)\%$ [13,14]. Note also that at chemical freezeout $z_p z_{\bar{p}} = 1$, $z_{\pi} = 1$, and the forward and backward rates in Eq. (2) are simply equal (the Boltzmann factors in both terms contain the same total energy). This implies, e.g., that cascade simulations ignoring the backreaction at comparable pion densities ($\rho_{\pi} \simeq 0.2-0.25 \text{ fm}^{-3}$) cannot give a proper account of the antiproton production.

The evolution of the system towards thermal freezeout can be constructed using a thermal fireball model which, based on isentropic expansion, leads to $(T_{\text{th}}, \mu_N^{\text{th}}) \simeq (120, 415)$ MeV. At first sight, this gives a \bar{p}/p ratio of $\exp[-2\mu_N/T] \simeq 0.1\%$, a factor of ~ 50 below the experimental result. However, this estimate is lacking an important ingredient. A correct statistical treatment including pion-number conservation forces us to introduce individual chemical potentials (or fugacities) for antiprotons/protons and pions. Insisting on equilibrium for the reaction in question, the following relation holds:

$$z_{\bar{p}} = (z_p)^{-1} \langle z_{\pi}^n \rangle. \quad (4)$$

Since antibaryons constitute only a small fraction of the produced secondaries, they have an insignificant impact on the evolution of nucleon- and pion-chemical potentials. Therefore, Eq. (4) provides an estimate of the antiproton fugacity at thermal freezeout. The crucial point here is that pion oversaturation will generate a strong enhancement of the backreaction $n\pi \rightarrow p\bar{p}$.

For a more quantitative assessment it is important to have a rather accurate determination of the pion multiplicity distributions in $p\bar{p}$ annihilation. A nice discussion of its systematics was given in Ref. [15], which we will follow here to incorporate the empirical knowledge. For $\bar{p}p$ annihilation at rest ($\sqrt{s} = 2m_N$), the data can be represented by a least-squares fit to a Gaussian probability distribution (normalized to one),

$$P(n) = \frac{1}{\sqrt{2\pi}\sigma} \exp[-(n - \langle n \rangle)^2 / 2\sigma^2], \quad (5)$$

with a mean $\langle n \rangle = 5.02$ and width $\sigma = 0.90$ (see also Ref. [16]). At larger cm energies, both the average number and the width increase. For the former, linear [17] and logarithmic [18] dependencies have been proposed,

$$\begin{aligned} \langle n \rangle &= c_1 + c_2 s^{0.5}, \\ \langle n \rangle &= \tilde{c}_1 + \tilde{c}_2 \ln(s) \end{aligned} \quad (6)$$

[s in (GeV^2)], in which both reproduce the measured multiplicities up to at least $\sqrt{s} = 5$ GeV using the parameters $c_1 = 2.6 \pm 0.5$, $c_2 = (1.3 \pm 0.2) \text{ GeV}^{-1}$, and $\tilde{c}_1 = 2.65$, $\tilde{c}_2 = 1.78$, respectively. For the latter, the energy dependence of the width has also been given as [18]

$$\sigma^2 = 0.174 \langle n \rangle s^{0.2}. \quad (7)$$

For our application in a thermal environment at $T = 150$ MeV (implying $\sqrt{s} = 2.33$ GeV) we fix $\langle n \rangle = 5.65$ [in accordance with Eq. (6)] together with a 10% increase in σ [as suggested by Eq. (7)] to extract discrete weights $w_n = P(n; \langle n \rangle, \sigma)$. The averaged pion-fugacity enhancement factor then follows as

$$\langle z_\pi^n \rangle = \sum_{n=2}^{n_{\max}} w_n \exp[n\mu_\pi/T], \quad (8)$$

where $n_{\max} = 9$ for any practical purpose. Inserting now the thermal freezeout values $T_{\text{th}} = 120$ MeV and $\mu_\pi^{\text{th}} \simeq 65$ MeV (as arising in a thermal fireball model [8]) yields $\langle z_\pi^n \rangle = 25$. This entails a large enhancement of the antiproton-to-proton ratio, from 0.1% to 2.5%. In fact, owing to the high power of the pion fugacity, slightly larger chemical potentials of $\mu_\pi = 75$ –80 MeV result in an enhancement factor of 42–54, rendering the pertinent \bar{p}/p ratio in line with the observed (chemical freezeout) value (cf. Fig. 2).

Such slightly increased values for the pion chemical potential close to thermal freezeout can indeed be easily argued for. Within the thermal fireball model of Ref. [8], elastic $\pi N \rightarrow B$ scattering (B : baryonic resonances up to $m_B \simeq 1.7$ GeV) was assumed to be frequent enough to maintain (relative) chemical equilibrium for the occupation of the excited baryonic states. However, with typical corresponding cross sections of $\sigma_{\pi N \rightarrow B} \simeq 15$ –30 mb [19], this might not be fully justified anymore for the last few fm/c prior to thermal freezeout. Consequently, a larger fraction of the pion number resides in explicit pionic degrees of freedom rather than in excited resonances, which translates into an effectively larger μ_π .

Let us finally comment on implications of our observations for relativistic heavy-ion collider (RHIC). Close to the expected chemical freezeout the pion density is very similar to SPS conditions. Thus the rate of producing antiprotons through multipion annihilation per unit time and volume is essentially the same in both cases. The crucial




FIG. 2. Antiproton-to-proton ratio as a function of (decreasing) temperature in an isentropically expanding fireball. The dashed curve represents the naive ratio, $\exp[-2\mu_N/T]$, whereas the solid curves are for finite pion chemical potentials, indicating uncertainties as discussed in the text. The experimental data point is from Ref. [13].

difference is, however, that the *total* density of antiprotons is much larger around midrapidity at RHIC due to substantially smaller baryon chemical potentials. More quantitatively, by using typical thermal model estimates [20] with $\rho_B^{\text{tot}} \simeq 0.2\rho_0$ shortly after chemical freezeout (further reduced thereafter), one obtains $\tau_{\text{ch}}^{\text{RHIC}} \simeq 11$ fm/c. With the lifetime of the hadronic phase at RHIC being comparable to that at SPS energies, chemical equilibrium in the $p\bar{p} \leftrightarrow n\pi$ reaction cannot be maintained until thermal freezeout (also, the emerging pion oversaturation is less pronounced in a baryon-poor regime). The observed antiprotons at RHIC should therefore mostly originate from earlier stages, corresponding to the standard hadrochemical freezeout in the vicinity of the phase boundary. Nevertheless, our time scale estimate indicates that, even under RHIC conditions, antibaryon annihilation will be partially compensated by the inverse reactions.

To summarize, we have analyzed the \bar{p}/p ratio at SPS energies employing a thermal approach. So far this observable has been difficult to understand within, e.g., transport models which only included the annihilation channel, causing doubts whether the latter is actually active or whether unconventional mechanisms for enhanced production need to be invoked. We have shown, however, that the “puzzle” can be resolved in a rather standard statistical-mechanics framework upon inclusion of the inverse process of multipion scattering into $\bar{p}p$ pairs, which can be supported until thermal freezeout. Our main ingredient was that effective pion-number conservation generates pion oversaturation at the later stages of a heavy-ion collision, as described by the buildup of appreciable pion chemical potentials. Raised to a large power ($n \sim 6$) the corresponding pion fugacities sustain a high antibaryon fraction, thus counterbalancing the loss from $B\bar{B}$ annihilation. This mechanism also

complies with the measured centrality dependence being essentially constant, as to be expected from a hadrochemistry varying little with impact parameter (for sufficiently peripheral collisions the applicability of thermal model analyses, of course, ceases and \bar{p} production, normalized to the number of participant nucleons, approaches its value in p - p collisions, which lies about 30% below the one in central nucleus-nucleus reactions [14]).

Finally we should note again that our findings are not contradictory to an earlier QGP formation. Rather, they give further support to the equilibrium concept of subsequent chemical and thermal freezeout stages, which has already proven successful in the explanation of a variety of observables, such as hadron abundances, collective flow, Hanbury-Brown–Twiss radii, and electromagnetic radiation.

This work was supported by the U.S. Department of Energy under Grant No. DE-FG02-88ER40388.

-
- [1] P. Braun-Munzinger, J. Stachel, J.P. Wessels, and N. Xu, Phys. Lett. B **365**, 1 (1996); P. Braun-Munzinger, I. Heppel, and J. Stachel, Phys. Lett. B **465**, 15 (1999).
[2] G.D. Yen and M.I. Gorenstein, Phys. Rev. C **59**, 2788 (1999).

- [3] J. Rafelski and J. Letessier, nucl-th/9903018.
[4] E. Shuryak, Phys. Lett. B **207**, 345 (1988); Phys. Rev. D **42**, 1764 (1990); P. Gerber, H. Leutwyler, and J.L. Goity, Phys. Lett. B **246**, 513 (1990); H. Bebie, P. Gerber, J.L. Goity, and H. Leutwyler, Nucl. Phys. **B378**, 95 (1992).
[5] C. M. Hung and E. Shuryak, Phys. Rev. C **57**, 1891 (1998).
[6] R. Stock, Nucl. Phys. **A661**, 282 (1999).
[7] H. Sorge, Phys. Rev. C **52**, 3291 (1995).
[8] R. Rapp and J. Wambach, Eur. Phys. J. A **6**, 415 (1999).
[9] Y. Pang, D.E. Kahana, and S.H. Kahana, Phys. Rev. Lett. **78**, 3418 (1997).
[10] M. Bleicher, M. Belkacem, S.A. Bass, S. Soff, and H. Stöcker, Phys. Lett. B **485**, 133 (2000).
[11] M. Belkacem *et al.*, Phys. Rev. C **58**, 1727 (1998).
[12] P. Koch, B. Müller, and J. Rafelski, Phys. Rep. **142**, 167 (1986).
[13] NA44 Collaboration, M. Kaneta *et al.*, Nucl. Phys. **A638**, 419 (1998).
[14] NA49 Collaboration, F. Sikler *et al.*, Nucl. Phys. **A661**, 45 (1999).
[15] C.B. Dover, T. Gutsche, M. Maruyama, and A. Faessler, Prog. Part. Nucl. Phys. **29**, 87 (1992).
[16] W. Blümel and U. Heinz, Z. Phys. C **67**, 281 (1995).
[17] P. Gregory *et al.*, Nucl. Phys. **B102**, 189 (1976).
[18] R. Stenbacka *et al.*, Nuovo Cimento A **51**, 63 (1979).
[19] Particle Data Group, D.E. Groom *et al.*, Eur. Phys. J. C **15**, 1 (2000).
[20] S. Bass *et al.*, Nucl. Phys. **A661**, 205c (1999).