

Charged-Particle Pseudorapidity Density Distributions from Au + Au Collisions at $\sqrt{s_{NN}} = 130$ GeV

B. B. Back,¹ M. D. Baker,² D. S. Barton,² R. R. Betts,³ R. Bindel,⁴ A. Budzanowski,⁵ W. Busza,⁶ A. Carroll,²
M. P. Decowski,⁶ E. Garcia,⁴ N. George,¹ K. Gulbrandsen,⁶ S. Gushue,² C. Halliwell,³ G. A. Heintzelman,²
C. Henderson,⁶ D. J. Hofman,³ R. Hołyński,⁵ B. Holzman,^{2,3} E. Johnson,⁷ J. L. Kane,⁶ J. Katzy,⁶ N. Khan,⁷
W. Kucewicz,³ P. Kulinich,⁶ W. T. Lin,⁸ S. Manly,⁷ D. McLeod,³ J. Michałowski,⁵ A. C. Mignerey,⁴ J. Mülmenstädt,⁶
R. Nouicer,³ A. Olszewski,^{2,5} R. Pak,² I. C. Park,⁷ H. Pernegger,⁶ C. Reed,⁶ L. P. Remsberg,² M. Reuter,³ C. Roland,⁶
G. Roland,⁶ L. Rosenberg,⁶ P. Sarin,⁶ P. Sawicki,⁵ W. Skulski,⁷ S. G. Steadman,⁶ P. Steinberg,² G. S. F. Stephans,⁶
M. Stodulski,⁵ A. Sukhanov,² J.-L. Tang,⁸ R. Teng,⁷ A. Trzupek,⁵ C. Vale,⁶ G. J. van Nieuwenhuizen,⁶ R. Verrier,⁶
B. Wadsworth,⁶ F. L. H. Wolfs,⁷ B. Wosiek,⁵ K. Woźniak,⁵ A. H. Wuosmaa,¹ and B. Wysłouch⁶

(PHOBOS Collaboration)

¹*Physics Division, Argonne National Laboratory, Argonne, Illinois 60439-4843*

²*Chemistry and C-A Departments, Brookhaven National Laboratory, Upton, New York 11973-5000*

³*Department of Physics, University of Illinois at Chicago, Chicago, Illinois 60607-7059*

⁴*Department of Chemistry and Biochemistry, University of Maryland, College Park, Maryland 20742*

⁵*Institute of Nuclear Physics, Kraków, Poland*

⁶*Laboratory for Nuclear Science, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139-4307*

⁷*Department of Physics and Astronomy, University of Rochester, Rochester, New York 14627*

⁸*Department of Physics, National Central University, Chung-Li, Taiwan*

(Received 6 June 2001; published 21 August 2001)

The charged-particle pseudorapidity density $dN_{ch}/d\eta$ has been measured for Au + Au collisions at $\sqrt{s_{NN}} = 130$ GeV at RHIC, using the PHOBOS apparatus. The total number of charged particles produced for the 3% most-central Au + Au collisions for $|\eta| \leq 5.4$ is found to be 4200 ± 470 . The evolution of $dN_{ch}/d\eta$ with centrality is discussed, and compared to model calculations and to data from proton-induced collisions. The data show an enhancement in charged-particle production at midrapidity, while in the fragmentation regions, the results are consistent with expectations from pp and pA scattering.

DOI: 10.1103/PhysRevLett.87.102303

PACS numbers: 25.75.-q

Nucleus-nucleus collisions at highly relativistic energies from the Relativistic Heavy Ion Collider (RHIC) present a new opportunity to study strongly interacting matter under conditions of very high temperature and energy density. Data from central Au + Au collisions at energies of $\sqrt{s_{NN}} = 56$ and 130 GeV [1] have shown an increase in midrapidity particle production over that observed in Pb + Pb collisions at the CERN SPS [2]. In addition, the charged-particle yields, when scaled by the number of participating nucleons (N_{part}), exceed the values observed in proton-proton collisions at similar $\sqrt{s_{NN}}$. These results have been used to infer an energy density at the highest value of $\sqrt{s_{NN}}$ that is considerably larger than was achieved at CERN, and which is well within the region where new physics is to be expected.

The distribution of $dN_{ch}/d\eta$ over the entire range of pseudorapidity η , where $\eta = -\ln[\tan(\theta/2)]$ and θ is the angle with respect to the beam, represents a time integral of particle production throughout the collision, and reflects all effects that contribute to the production of charged particles. In addition to the influence of both hard and soft production processes, $dN_{ch}/d\eta$ is sensitive to nuclear effects in the initial parton distributions, as well as from hadronic reinteractions in the final state.

The data for $dN_{ch}/d\eta$ at midrapidity ($|\eta| < 1$) for the most central Au + Au collisions at $\sqrt{s_{NN}} = 56$ and 130 GeV [1] are in reasonable agreement with the predictions of a number of models including HIJING [3], a saturation model (EKRT) [4], and purely hadronic models (e.g., LUCIFER [5,6]). The centrality dependence of $dN_{ch}/d\eta$ at midrapidity has also been measured [7,8]. These latter results suggest some deviation from both the HIJING and EKRT model predictions, although they are in broad agreement with the results of calculations by Kharzeev and Nardi [9].

An extension of the measurements of $dN_{ch}/d\eta$ data beyond midrapidity, for a range of impact parameters, is necessary to further constrain models. It is also of interest to determine whether the observed scaling of the charged-particle yield with N_{part} at midrapidity is modified at large values of η where, in proton-nucleus (pA) collisions [10–12], rescattering, stopping, and target fragmentation influence the shape of the $dN_{ch}/d\eta$ distributions. In order to address these questions, we have used the PHOBOS apparatus to measure the charged-particle pseudorapidity density $dN_{ch}/d\eta$ from Au + Au collisions at $\sqrt{s_{NN}} = 130$ GeV over the range $|\eta| < 5.4$.

The PHOBOS experiment at RHIC largely consists of several arrays of silicon pad detectors. The details of the experimental arrangement are described in Ref. [13]. The procedures used for event selection, the determination of the collision-vertex position, event centrality, and the estimation of N_{part} , have been described in Refs. [1,8]. The specific elements of the experiment used in the current measurement and the analysis procedures leading to $dN_{ch}/d\eta$ are described below.

The data samples included in the current analysis were taken at a collision energy of $\sqrt{s_{NN}} = 130$ GeV. The collision vertices were confined to a region within ± 10 cm of the nominal beam crossing and center of the apparatus ($z = 0$). At midrapidity, charged particles were detected, and their energy deposition measured, with an octagonal array of pad detectors approximately 1 m long (the ‘‘octagon’’) that surrounds the thin-walled Be beam pipe. The octagon subtends the full azimuthal range, except for regions where sensors that would intercept the acceptance for the tracking spectrometers and vertex finding detectors are removed. For collision vertices within $|z| < 10$ cm, the pseudorapidity coverage of the octagon is complete for $|\eta| \leq 3.2$. Six rings of silicon pad detectors, placed at distances of $|z| \approx 1, 2,$ and 5 m, detected particles in pseudorapidity ranges of $3 \leq |\eta| \leq 4$, $4 \leq |\eta| \leq 4.7$, and $4.7 \leq |\eta| \leq 5.4$, respectively. Thus, for collisions within the 20 cm long region in the center of the experiment, there are no significant gaps in the η coverage from one subdetector to the next. The total numbers of pads in the octagon and rings are 11 040 and 3072, respectively.

Two complementary methods were used to analyze the pseudorapidity distribution data. The first method (‘‘hit counting’’) uses the segmentation of the multiplicity detector. After merging of signals in neighboring pads, in cases where a particle travels through more than a single pad, the deposited energy was corrected for the angle of incidence, so that all tracks originating from the collision vertex possess a common average value of the deposited energy ($\Delta E \approx 80$ keV). Pads containing more than 75% of this value were counted as occupied. This requirement largely suppresses hits from background and from secondary particles not originating from the primary collision vertex. Then, for a given value of η and bin i in collision centrality, the number of hit pads $N(\eta, i)$ was corrected for the effects of multiple occupancy, where more than one particle travels through a given pad, as well as for contributions from the remaining secondary particles, absorption in the beam pipe, and weak decays of primary particles. The probability $P(N)$ of N particles passing through a given pad was assumed to be Poisson distributed. The mean occupancy $\mu(\eta, i)$ was then determined from the ratio of occupied to unoccupied pads in a range of η for each centrality bin. As a check, the occupancy was also determined from the energy-deposition spectra, where a fitting procedure was used to determine the relative contribution to the total energy deposition of one or more particles. Maximum

values of approximately 1.6 particles per hit pad were obtained for the most central collisions at midrapidity.

To account for effects or biases not treated in the above analysis procedures, a final correction was deduced from GEANT [14] simulations of the detector response using events from the HIJING [3], RQMD [15], and VENUS [16] event generators. The ratio between the simulated, occupancy-corrected $dN_{ch}/d\eta$ distributions and the known ‘‘truth’’ distributions formed this last set of corrections, which ranged up to 15% in the octagon and up to 50% in the rings. These background correction factors, dependent upon both η and centrality bin, were applied to the occupancy-corrected data, yielding $dN_{ch}/d\eta$. The final results obtained using both occupancy determination methods were in good agreement.

In the second (‘‘analog’’) method, the pseudorapidity distribution was extracted directly from the energy deposition $\Delta E(\eta)$ in the multiplicity detectors, which was transformed into $dN_{ch}/d\eta$ using the results of Monte Carlo simulations. The average energy per track (ΔE_{tr}) was determined as a function of η using particles from HIJING events, passed through the GEANT simulation of the detector. The fraction of primary particles $f_{\text{prim}}(\eta)$ was determined from the same simulations. Then, $dN_{ch}/d\eta = \frac{\Delta E(\eta) \times f_{\text{prim}}(\eta)}{\Delta E_{\text{tr}}(\eta) \Delta \eta}$. Although the two methods are significantly different, they yield results that differ generally by $\leq 5\%$ throughout the range in η , well within the systematic uncertainties (see below), which are approximately 10%.

The systematic uncertainty in the occupancy correction for the hit-counting analysis was obtained by comparing the results from the full analysis chain using Poisson-derived occupancy corrections with those derived from the measured ΔE spectra. The average deviations are less than 3%, yielding a partial systematic error of $\approx 3\%$ at midrapidity. The systematic uncertainties from the Monte Carlo simulations have been estimated by using different assumptions in the GEANT simulation, as well as different event generators, including RQMD [15] and VENUS [16]. The variations observed in the derived background corrections are between 4% and 8%, suggesting a total systematic uncertainty of approximately 10%. The systematic errors in the analog analysis arise from uncertainties in the Monte Carlo simulations, and in the absolute energy calibrations of the silicon-pad detectors. The latter are approximately 5%, yielding a total systematic error for the analog method of approximately 10%, similar to that for the hit-counting method.

Our final results are presented in Figs. 1(a)–1(f), which show the error-weighted average values of $dN_{ch}/d\eta$ from the two procedures for six different centrality bins. The error bars represent a convolution of the estimated systematic errors in the different analyses. The different centrality bins are denoted by the corresponding fraction of the estimated total cross section σ_{tot} based on the observed fraction $\sigma_{\text{obs}} = (0.97 \pm 0.03)\sigma_{\text{tot}}$, as well as by the deduced average number of participant nucleons ($\langle N_{\text{part}} \rangle$). The




FIG. 1. Charged-particle pseudorapidity density $dN_{ch}/d\eta$ from $\sqrt{s_{NN}} = 130$ GeV Au + Au collisions, for different centrality bins, as defined by different fractions of the total cross section. The error bars reflect primarily the overall systematic uncertainties. The average number of participants $\langle N_{part} \rangle$ for each bin is also indicated. For fractions of the cross section $>45\%$ the systematic uncertainties in the $\langle N_{part} \rangle$ determination are still under study and no value is quoted.

present analyses give values of $dN_{ch}/d\eta|_{|\eta|<1}$ that are in good agreement from the independent analysis presented in [8]. For example, for the 6% most central collisions, we find $dN_{ch}/d\eta|_{|\eta|\leq 1} = 547 \pm 55$, compared to 580 ± 25 from Ref. [8].

The integral over $|\eta| < 5.4$ of the distributions, N_{ch}^{tot} , plotted in Fig. 2(a) as a function of centrality, is a direct measure of the total entropy produced in the collisions. Predictions of its magnitude have varied by as much as a factor of 2 [17]. With increasing $\langle N_{part} \rangle$, the observed values of N_{ch}^{tot} change smoothly from 910 ± 100 for the 40%–45% centrality bin to 4200 ± 470 for the 3% most central collisions. Per participant pair, these numbers correspond, respectively, to 21.8 ± 2.6 and 23.7 ± 2.7 , compared with the pp and $p\bar{p}$ nondiffractive total charged particle multiplicity of 18.5 ± 0.7 [18]. The uncertainties contain a contribution in quadrature from the systematic errors in $\langle N_{part} \rangle$ which range from approximately 3% to 5% [8]. The predictions of the HIJING model reproduce the general trend of the N_{ch}^{tot} centrality dependence, but systematically underpredict the observed values by 10%–15%.

The shapes of the pseudorapidity distributions evolve gradually with increasing centrality, as shown in Figs. 1(a)–1(f). For all centrality bins, there is a plateau region between $-2 < \eta < 2$, followed by a rapid dropoff towards larger pseudorapidities. A more detailed study of the centrality dependence of the shape is given in




FIG. 2. (a) Total number of charged particles detected within the range $-5.4 < \eta < 5.4$. The symbols are the measured data, and the line represents the prediction of the code HIJING. The error bars reflect systematic uncertainties. (b)–(f) Centrality dependence of $dN_{ch}/d\eta$ for different ranges of $|\eta|$. The filled symbols are the measured data, and the solid curves are the HIJING predictions. The open diamonds and triangles refer to the predictions of the AMPT and LUCIFER models, respectively, for the 6% most central collisions. The open circles represent the values from pp and $p\bar{p}$ collisions.

Figs. 2(b)–2(f) and Fig. 3(a). In Figs. 2(b)–2(f), the N_{part} dependence of $dN_{ch}/d\eta$ normalized per participant pair $\langle N_{part}/2 \rangle$ is plotted for five pseudorapidity bins ranging from $|\eta| < 1$ to $5 < |\eta| < 5.4$. Also plotted are data from pp [19] and $p\bar{p}$ [20] collisions, scaled as described below, as open circles, and predictions from the HIJING model as solid lines. The statistical uncertainties are small and the systematic uncertainties are comparable to those described above. Figure 3(a) shows the pseudorapidity distribution for peripheral (35%–45%) and central (0%–6%) Au + Au collisions, scaled by the respective $\langle N_{part}/2 \rangle$.

For all pseudorapidity bins in Figs. 2(b)–2(f), the data evolve smoothly from the most peripheral to the most central collisions. As seen in Figs. 2(b), 2(c), and 3(a), central collisions yield a 10%–15% higher $dN_{ch}/d\eta$ per participant in the plateau region, compared to peripheral events. This difference between central and peripheral collisions changes character between $|\eta| = 3$ and 4, beyond which a higher yield per participant is found in peripheral collisions. In the highest 1.5 units of pseudorapidity, as is observed in pA collisions [10–12], the scaled charged-particle density actually falls with N_{part} , being reduced by nearly a factor of 2 for $|\eta| > 5$. Qualitatively, the changes in the distributions from peripheral to central collisions in




FIG. 3. (a) $dN_{ch}/d\eta/(\langle N_{part} \rangle/2)$ for central (0%–6%, open symbols) and peripheral (35%–45%, filled symbols) collisions. The error bars have been suppressed for clarity. (b) Comparison of the measured pseudorapidity distribution for the 0%–6% centrality bin with the HIJING (solid curve) and AMPT (dashed curve) models. The grey band represents data from pp and $p\bar{p}$ collisions interpolated to $\sqrt{s_{NN}} = 130$ GeV.

the fragmentation regions are similar to those observed in pA collisions [10–12].

Finally, Fig. 3(b) shows a comparison between $(dN_{ch}/d\eta)/\langle N_{part} \rangle/2$ from 0%–6% central Au + Au collisions (data points) and scaled data from pp and $p\bar{p}$ collisions (grey band). The $pp/p\bar{p}$ distribution was obtained by scaling the measured $dN_{ch}/d\eta$ distributions from pp collisions at $\sqrt{s} = 53$ GeV and $p\bar{p}$ collisions at $\sqrt{s} = 200$ and 546 GeV [19,20] horizontally by $y_{max}(130 \text{ GeV})/y_{max}(\sqrt{s})$, where $y_{max}(\sqrt{s}) = \ln(\sqrt{s}/m_p)$, and vertically using the parametrization of $dN/d\eta|_{|\eta|<1}$ from [20]. [This empirical scaling is chosen such that the extrapolation of three sets of pp or $p\bar{p}$ data are confined within the band shown in Fig. 3(b).] The grey band gives an estimate of the uncertainty of this extrapolation procedure.

The scaled $dN_{ch}/d\eta$ is found to be higher in central Au + Au collisions than in $pp/p\bar{p}$ over the full pseudorapidity range out to $|\eta| = 4$, with the largest excess observed in the central plateau region. This is in contrast with the HIJING prediction (solid curve), which shows an excess in Au + Au collisions only for $|\eta| \leq 2-3$. A possible origin of this qualitative difference is suggested by the AMPT model of Zhang *et al.* [21]. In this model, the initial state parton distribution is obtained as in HIJING, but is followed by a parton cascade [22], string fragmentation, and hadronic rescattering using a relativistic transport model [23]. This model reproduces the excess in particle production at higher pseudorapidities seen in Au + Au collisions relative to $pp/p\bar{p}$. Furthermore, predictions of $dN_{ch}/d\eta$ in a completely hadronic framework (e.g., LUCIFER) [6] are very similar to those of the AMPT calculations, and are also in good agreement with the data for the most cen-

tral collisions. These observations suggest that effects in the hadronic phase, e.g., rescattering, should be taken into account to provide a full description of the data. A comparison of the predicted centrality dependence of the full distributions from these models with the data would also be of interest. Further insight will be gained from future RHIC data, which should include more precise reference data from pp collisions, as well as nucleus-nucleus data at different collision energies. These results will provide a basis for separating the effects of new phenomena from conventional hadronic physics.

This work was supported in part by U.S. DoE Grants No. DE-AC02-98CH10886, No. DE-FG02-93ER-404802, No. DE-FC02-94ER40818, No. DE-FG02-94ER40865, No. DE-FG02-99ER41099, No. W-31-109-ENG-38, and NSF Grants No. 9603486, No. 9722606, and No. 0072204. The Polish groups were partially supported by KBN Grant No. 2P03B 04916. The NCU group was partially supported by the NSC of Taiwan under Contract No. NSC 89-2112-M-008-024.

- [1] B. B. Back *et al.*, Phys. Rev. Lett. **85**, 3100 (2000).
- [2] J. Bälcher *et al.*, Nucl. Phys. **A661**, 45 (1999).
- [3] X. N. Wang and M. Gyulassy, Phys. Rev. D **44**, 3501 (1991).
- [4] K. J. Eskola *et al.*, Nucl. Phys. **B570**, 379 (2000).
- [5] D. E. Kahana and S. H. Kahana, Phys. Rev. C **59**, 1651 (1999).
- [6] D. E. Kahana and S. H. Kahana, Phys. Rev. C **63**, 031901(R) (2001).
- [7] PHENIX Collaboration, K. Adcox *et al.*, Phys. Rev. Lett. **86**, 3500 (2001).
- [8] B. B. Back *et al.*, e-print nucl-ex/0105011.
- [9] D. Kharzeev and M. Nardi, Phys. Lett. B **507**, 121 (2001).
- [10] W. Busza, Acta. Phys. Pol. B **8**, 333 (1978).
- [11] C. De Marzo *et al.*, Phys. Rev. D **26**, 1019 (1982).
- [12] D. S. Barton *et al.*, Phys. Rev. D **27**, 2580 (1983).
- [13] B. B. Back *et al.*, Nucl. Phys. **A661**, 690 (1999).
- [14] GEANT Detector Description and Simulation Tool v. 3.21, CERN.
- [15] H. Sorge, Phys. Rev. C **52**, 3291 (1995).
- [16] K. Werner, Phys. Rep. **232**, 87 (1993).
- [17] S. A. Bass *et al.*, Nucl. Phys. **A661**, 205 (1999).
- [18] A. Wroblewski, in *Proceedings of the XIV International Symposium on Multiparticle Dynamics, Lake Tahoe, 1983* (World Scientific, Singapore, 1984), p. 579.
- [19] G. J. Alner *et al.*, Z. Phys. C **33**, 1 (1986).
- [20] F. Abe *et al.*, Phys. Rev. D **41**, 2330 (1990).
- [21] B. Zhang *et al.*, Phys. Rev. C **61**, 067901 (2000).
- [22] B. Zhang, Comput. Phys. Commun. **109**, 193 (1998).
- [23] B. A. Li and C. M. Ko, Phys. Rev. C **52**, 2037 (1995).