

Measurement of G_{E_p}/G_{M_p} in $\vec{e}p \rightarrow e\vec{p}$ to $Q^2 = 5.6 \text{ GeV}^2$

O. Gayou,^{1,7} K. A. Aniol,⁹ T. Averett,¹ F. Benmokhtar,² W. Bertozzi,²⁴ L. Bimbot,²⁶ E. J. Brash,⁴ J. R. Calarco,²⁵ C. Cavata,²⁷ Z. Chai,²⁴ C.-C. Chang,²³ T. Chang,¹⁵ J.-P. Chen,⁵ E. Chudakov,⁵ R. De Leo,¹⁶ S. Dieterich,² R. Endres,² M. B. Epstein,⁹ S. Escoffier,²⁷ K. G. Fissum,²² H. Fonvieille,⁷ S. Frullani,¹⁸ J. Gao,⁸ F. Garibaldi,¹⁸ S. Gilad,²⁴ R. Gilman,^{2,5} A. Glamazdin,²¹ C. Glashauser,² J. Gomez,⁵ V. Gorbenko,²¹ J.-O. Hansen,⁵ D. W. Higinbotham,^{24,*} G. M. Huber,⁴ M. Iodice,¹⁷ C. W. de Jager,⁵ X. Jiang,² M. K. Jones,⁵ J. J. Kelly,²³ M. Khandaker,³ A. Kozlov,⁴ K. M. Kramer,¹ G. Kumbartzki,² J. J. LeRose,⁵ D. Lhuillier,²⁷ R. A. Lindgren,²⁶ N. Liyanage,⁵ G. J. Lolos,⁴ D. J. Margaziotis,⁹ F. Marie,²⁷ P. Markowitz,¹² K. McCormick,²⁰ R. Michaels,⁵ B. D. Milbrath,¹¹ S. K. Nanda,⁵ D. Neyret,²⁷ Z. Papandreou,⁴ L. Pentchev,^{1,†} C. F. Perdrisat,¹ N. M. Piskunov,¹⁹ V. Punjabi,³ T. Pussieux,²⁷ G. Quéméner,¹⁴ R. D. Ransome,² B. A. Raue,¹² R. Roché,¹³ M. Rvachev,²⁴ A. Saha,⁵ C. Salgado,³ S. Širca,²⁴ I. Sitnik,¹⁹ S. Strauch,^{2,‡} L. Todor,¹⁰ E. Tomasi-Gustafsson,²⁷ G. M. Urciuoli,¹⁸ H. Voskanyan,²⁹ K. Wijesooriya,⁶ B. B. Wojtsekhowski,⁵ X. Zheng,²⁴ and L. Zhu²⁴

(Jefferson Lab Hall A Collaboration)

¹College of William and Mary, Williamsburg, Virginia 23187

²Rutgers, The State University of New Jersey, Piscataway, New Jersey 08855

³Norfolk State University, Norfolk, Virginia 23504

⁴University of Regina, Regina, Saskatchewan, Canada S4S 0A2

⁵Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606

⁶Argonne National Laboratory, Argonne, Illinois 60439

⁷Université Blaise Pascal/CNRS-IN2P3, F-63177 Aubière, France

⁸California Institute of Technology, Pasadena, California 91125

⁹California State University, Los Angeles, California 90032

¹⁰Carnegie Mellon University, Pittsburgh, Pennsylvania 15213

¹¹Eastern Kentucky University, Richmond, Kentucky 40475

¹²Florida International University, Miami, Florida 33199

¹³Florida State University, Tallahassee, Florida 32306

¹⁴Institut des Sciences Nucléaires, CNRS-IN2P3, F-38026 Grenoble, France

¹⁵University of Illinois, Urbana-Champaign, Illinois 61801

¹⁶INFN, Sezione di Bari and University of Bari, 70126 Bari, Italy

¹⁷INFN, Sezione di Roma-III, 00146 Roma, Italy

¹⁸INFN, Sezione Sanità and Istituto Superiore di Sanità, 00161 Rome, Italy

¹⁹JINR-LHE, 141980 Dubna, Moscow Region, Russian Federation

²⁰Kent State University, Kent, Ohio 44242

²¹Kharkov Institute of Physics and Technology, Kharkov 61108, Ukraine

²²University of Lund, P.O. Box 118, S-221 00 Lund, Sweden

²³University of Maryland, College Park, Maryland 20742

²⁴Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

²⁵University of New Hampshire, Durham, New Hampshire 03824

²⁶Institut de Physique Nucléaire, F-91406 Orsay, France

²⁷DAPNIA/SPhN CEA/Saclay, F-91191 Gif-sur-Yvette, France

²⁸University of Virginia, Charlottesville, Virginia 22901

²⁹Yerevan Physics Institute, Yerevan 375036, Armenia

(Received 15 November 2001; published 12 February 2002)

The ratio of the electric and magnetic form factors of the proton G_{E_p}/G_{M_p} , which is an image of its charge and magnetization distributions, was measured at the Thomas Jefferson National Accelerator Facility (JLab) using the recoil polarization technique. The ratio of the form factors is directly proportional to the ratio of the transverse to longitudinal components of the polarization of the recoil proton in the elastic $\vec{e}p \rightarrow e\vec{p}$ reaction. The new data presented span the range $3.5 < Q^2 < 5.6 \text{ GeV}^2$ and are well described by a linear Q^2 fit. Also, the ratio $\sqrt{Q^2} F_{2p}/F_{1p}$ reaches a constant value above $Q^2 = 2 \text{ GeV}^2$.

DOI: 10.1103/PhysRevLett.88.092301

PACS numbers: 25.30.Bf, 13.40.Gp, 14.20.Dh, 24.85.+p

The nucleon electromagnetic form factors are a key ingredient to describe its internal structure and eventually understand the strong interaction. Experimental values for the proton have been obtained over the last 50 years

via electron-proton scattering, often using the Rosenbluth separation technique [1]. They show that the magnetic form factor, G_{M_p} , follows approximately a dipole form factor $G_D = [1 + Q^2/0.71(\text{GeV}^2)]^{-2}$, where Q^2 is the

four-momentum transfer squared [2–8]. However, measuring the charge form factor G_{E_p} by Rosenbluth separation becomes difficult for $Q^2 > 1 \text{ GeV}^2$, because the charge scattering contributes only little to the differential cross section. Extending the measurement of the form factors to larger Q^2 is important, for example, to test the perturbative QCD (pQCD) scaling predictions for the Dirac and Pauli form factors F_{1_p} and F_{2_p} [9]. The recoil polarization method, proposed in the 1970s [10], has been established as the most effective available technique for measuring the ratio G_{E_p}/G_{M_p} at large Q^2 [11–14]. The results of Ref. [11] showed a surprising, roughly linear, decrease of this ratio as a function of Q^2 up to 3.5 GeV^2 . In a nonrelativistic approach, this faster decrease of G_{E_p} can be interpreted as confinement of the charge distribution in the Breit frame to a larger region of space than the magnetism distribution.

In the one-photon exchange approximation for elastic ep scattering, a longitudinally polarized electron beam transfers its polarization to the recoil proton with two nonzero components, P_t , perpendicular to, and P_ℓ , parallel to, the proton momentum in the scattering plane. P_t and P_ℓ are proportional to $G_{E_p}G_{M_p}$ and $G_{M_p}^2$, respectively, so that the ratio of the form factors follows directly from the simultaneous measurements of these two polarization components [10]:

$$\frac{G_{E_p}}{G_{M_p}} = -\frac{P_t}{P_\ell} \frac{(E_e + E_{e'})}{2m} \tan \frac{\theta_e}{2}. \quad (1)$$

Here m is the proton mass, θ_e is the lab scattering angle, and E_e and $E_{e'}$ are the incident and scattered energies of the electron.

We present the results of new measurements of the ratio $\mu_p G_{E_p}/G_{M_p}$, where μ_p is the magnetic moment of the proton, up to $Q^2 = 5.6 \text{ GeV}^2$ performed in Hall A at Jefferson Lab. A polarized electron beam from the Continuous Electron Beam Accelerator was scattered on a 15-cm-long circulating liquid hydrogen target. A strained GaAs crystal excited by circularly polarized laser light produced the polarized electron beam, with an average current of $40 \mu\text{A}$. A typical longitudinal beam polarization at the target of ~ 0.70 was measured with both a Møller polarimeter [15] (with an uncertainty of $\sim 3\%$) and a Compton polarimeter [16] (with an uncertainty of $\sim 1.4\%$ [17]). The helicity of the beam was flipped pseudorandomly at 30 Hz.

Recoil protons were detected in the left high resolution spectrometer (HRS) [18]. The HRS has a central bend angle of 45° and accepts a maximum central momentum of $4 \text{ GeV}/c$ with a 6.5 msr acceptance; it has a $\pm 5\%$ momentum acceptance and a $< 2 \times 10^{-4}$ momentum resolution. Two vertical drift chambers located at the focal plane, along with the knowledge of the optics of the three quadrupoles and the dipole of the HRS, allow measurements of the proton position at the target with an absolute

accuracy of 0.2 mm and of the angle of its trajectory with an absolute accuracy of 0.7 mrad in the dispersive, and 0.3 mrad in the transverse, direction. As the data acquisition was triggered by a single proton in the HRS, we also detected the scattered electron in order to isolate elastic ep scattering events and reject the significant background in the spectrometer, mostly from pion electroproduction. The polarization transfer in this reaction can be different in magnitude and sign from the polarization transfer in elastic scattering.

For the measurement at $Q^2 = 3.5 \text{ GeV}^2$, the electron was detected in the second (right) HRS, and the trigger was a coincidence between an electron and a proton, as described in Ref. [11]. For the measurements at higher Q^2 , at a fixed beam energy of 4.6 GeV , the electron was scattered at a larger angle than the proton and thus defined the rate of the reaction. To maximize the number of elastic events selected, the electron was detected in a calorimeter with a large solid angle. The $1.35 \times 2.55 \text{ m}^2$ calorimeter was assembled with blocks of lead glass with a cross-sectional area of $15 \times 15 \text{ cm}^2$ each, in 9 columns and 17 rows. The use of lead glass eliminates the pion background, which does not produce Čerenkov light in this material. At each Q^2 , the calorimeter was located at a distance from the target where the electron solid angle matched the proton HRS acceptance according to the Jacobian of the reaction. This distance ranges from 9 m at $Q^2 = 5.6 \text{ GeV}^2$ to 17 m at $Q^2 = 4.0 \text{ GeV}^2$. The trigger was defined by a proton in the HRS, signaled by a coincidence of two planes of scintillators in the focal plane. For each single proton event in the left HRS, the energy and time information from the calorimeter was read out for all blocks, and elastic events were selected by applying software cuts to the calorimeter data. Our analysis showed that the calorimeter registered an energy signal in about ten blocks for each trigger. A 10-ns time cut was applied to eliminate signals that were not in coincidence with the proton. To select the elastic ep reaction from events surviving the time window, a cut was made on the difference between the measured angle of the particle in the calorimeter and the predicted angle from the proton angle and momentum assuming ep elastic kinematics. The background remaining after these two cuts was evaluated by extrapolating the distribution of events that were not correlated in position to the elastic region; it represents less than 2% of the accepted events and is taken into account in the polarization analysis by measuring the polarization of the rejected events. It corresponds to a correction of at most 3% on the ratio $\mu_p G_{E_p}/G_{M_p}$. The uncertainty on this correction was included in the final systematic error. About 5% of elastic events were rejected, because of missing lead-glass blocks in the calorimeter and discriminator threshold.

The recoil proton polarization was measured by the focal plane polarimeter (FPP) located behind the focal plane of the left HRS [19]. The FPP determines the two polarization components perpendicular to the momentum,

P_t^{fpp} and P_n^{fpp} , by measuring asymmetries in the azimuthal angular distribution after scattering the proton in an analyzer. To improve the figure of merit, the usual graphite analyzer was replaced by polyethylene, 60 cm thick at $Q^2 = 3.5 \text{ GeV}^2$ and 100 cm thick for the other kinematics. The angular distribution is measured by detecting the trajectory of the proton in two sets of two straw chambers, one before and one after the scattering in the analyzer; the distribution is given by

$$N(\vartheta, \varphi) = N_0(\vartheta) \{1 + [A_y(\vartheta)P_t^{\text{fpp}} + a_{\text{in}}] \sin \varphi - [A_y(\vartheta)P_n^{\text{fpp}} + b_{\text{in}}] \cos \varphi\}, \quad (2)$$

where $N_0(\vartheta)$ is the number of protons scattered in the polarimeter to a polar angle ϑ , φ is the azimuthal angle after scattering, and $A_y(\vartheta)$ is the analyzing power; a_{in} and b_{in} are instrumental asymmetries. Such a distribution was measured for the two states of the electron beam helicity, positive and negative. The difference in the beam polarization for these two helicity states was compatible with zero at the 0.3% level [17]. The difference between these two distributions $N^+/N_0^+ - N^-/N_0^-$ cancels the instrumental asymmetries to first order. It also gives us access to the transferred, helicity-dependent polarization, which is the quantity of interest. The induced, helicity-independent polarization is zero in the case of elastic scattering from the proton. Figure 1 shows this difference distribution, fitted (solid line) with a cosine function $C \cos(\varphi + \delta)$, where the amplitude C is $\sqrt{(P_n^{\text{fpp}})^2 + (P_t^{\text{fpp}})^2}$ and the phase shift δ is such that $\tan \delta = P_t^{\text{fpp}}/P_n^{\text{fpp}}$. Since P_t^{fpp} is related to the interference term $G_{E_p}G_{M_p}$, this phase shift is a measure of G_{E_p} . The dashed line represents what the distribution would look like if $\mu_p G_{E_p}/G_{M_p} = 1$. The vertical lines at $\varphi = 90^\circ$ and $\varphi = 270^\circ$ emphasize the phase shift δ .

The proton spin precesses through the magnetic fields of the HRS. The polarization vector at the analyzer of the FPP, \mathbf{P}^{fpp} , is related to the polarization vector at the target, \mathbf{P} , by the spin transfer matrix \mathbf{S} : $\mathbf{P}^{\text{fpp}} = \mathbf{S} \times \mathbf{P}$. Because protons with different angles and interaction points at the




FIG. 1. Difference distribution for positive and negative electron beam helicity, for $Q^2 = 5.6 \text{ GeV}^2$. See text for details.

target see different magnetic fields in the HRS, the matrix elements S_{ij} must be calculated for each event from the reconstructed target coordinates. The matrix elements were determined using a model of the HRS based on optics studies and using the differential algebra-based code COSY [20].

The polarization components $hA_y P_t$ and $hA_y P_\ell$ are obtained by maximizing the likelihood function [21] $L(P_t, P_\ell)$ defined as

$$L(P_t, P_\ell) = \prod_{i=1}^{N_p} \{1 \pm A_y(\vartheta_i) (S_{tt,i} h P_t + S_{t\ell,i} h P_\ell) \sin \varphi_i \mp A_y(\vartheta_i) (S_{nt,i} h P_t + S_{n\ell,i} h P_\ell) \cos \varphi_i\}, \quad (3)$$

where the product runs over all events, N_p , \pm stands for the sign of the beam helicity, and h is the beam polarization. The analyzing power and beam helicity eventually cancel in forming the ratio $hA_y P_t/hA_y P_\ell$.

The new results for the ratio $\mu_p G_{E_p}/G_{M_p}$ are presented in Fig. 2, with statistical error bars, together with the data of Ref. [11]. The systematic errors are represented by the bands at the top. The new data are tabulated in Table I, with their statistical and systematic errors. The main sources of systematic errors are related to the spin precession. Those can be divided into three parts. Our analysis shows that the major part is the error associated with the uncertainty in the total bending angle in the nondispersive plane of the spectrometer, due to misalignment of the magnetic elements of the spectrometer. A careful study of this misalignment has been done recently in Hall A [27], reducing the systematic error compared to Ref. [11] at $Q^2 = 3.5 \text{ GeV}^2$ by a factor of 6. The other sources of error in the precession are related to uncertainties in the




FIG. 2. The ratio $\mu_p G_{E_p}/G_{M_p}$ from this experiment and Jones *et al.* (Ref. [11]), compared with theoretical calculations. Systematic errors for both experiments are shown as a band at the top of the figure.

TABLE I. The ratio $\mu_p G_{E_p}/G_{M_p}$ with statistical uncertainty (1σ) Δ_{stat} and systematic uncertainty Δ_{syst} . $\langle Q^2 \rangle$ is the value of Q^2 weighted averaged over the acceptance, and ΔQ^2 is the Q^2 acceptance (1σ).

$\langle Q^2 \rangle \pm \Delta Q^2$ (GeV ²)	$\mu_p G_{E_p}/G_{M_p}$	Δ_{stat}	Δ_{syst}
3.50 ± 0.23	0.571	0.072	0.007
3.97 ± 0.26	0.482	0.052	0.008
4.75 ± 0.30	0.382	0.053	0.011
5.54 ± 0.34	0.273	0.087	0.028

dipole fringe field model and to the bending angle in the dispersive plane. Systematic errors associated with proton momentum, electron beam energy, and electron scattering angle give smaller contributions. No radiative corrections have been applied to the ratio, as no full calculation of polarization observables for ep scattering exists. Afanasev *et al.* [28] have calculated the single photon emission corrections to the two polarization observables in hadronic variables. The two corrections are of the same sign, negative, and are each of the order of 1%; thus they largely cancel when one takes the ratio. Other contributions due to two photon-exchange, virtual Compton scattering and interference terms are expected to be at the percent level [29].

A straight line fit has been applied to the ratio $\mu_p G_{E_p}/G_{M_p}$ in the range $0.5 < Q^2 < 5.6$ GeV²:

$$\mu_p \frac{G_{E_p}}{G_{M_p}} = 1 - 0.13(Q^2 - 0.04). \quad (4)$$

Using this Q^2 dependence as a constraint on G_{E_p} , the Rosenbluth separation data have been reanalyzed. This brings a correction of the order of 1.5% to 3% to the magnetic form factor [30].

Also shown in Fig. 2 are the results of some theoretical calculations which discuss possible interpretations of a decrease of the ratio $\mu_p G_{E_p}/G_{M_p}$. Several authors have studied different effects within the framework of the constituent quark model (CQM); all emphasize the necessity of both kinematic and dynamic relativistic corrections. Frank, Jennings, and Miller [26], in their study of nuclear medium effects on nucleon electromagnetic form factors, used Schlumpf's light-front wave function in an early relativistic CQM [31] to compute the free proton elastic form factors (dashed curve). Based on the data of Ref. [11], Cardarelli and Simula [24] show that a suppression of the ratio can be expected in the CQM, if the relativistic effects generated by the SU(6) symmetry breaking caused by the Melosh rotations of the constituent spins are taken into account. Their prediction is shown using pointlike quark constituents (dotted curve) and constituent quark form factors (solid curve). Wagenbrunn *et al.* [23] (thin long-dashed curve) reach a reasonable agreement with all electroweak nucleon form factors in their point-form spectator approximation (PFSA) prediction of the Goldstone boson exchange CQM [32]. Other types of models try to

describe the dynamic features of the nucleon. Holzwarth [25] (thick long-dashed curve) uses a relativistic chiral soliton model, which gives remarkable agreement with the data. Lomon [22] used the world data, including Ref. [11], to perform a fit within the vector meson dominance (VMD) model, where the ρ meson contribution is determined by dispersion relations (dot-dashed curve). It is worthwhile to note that while some models can reproduce the observed behavior of $\mu_p G_{E_p}/G_{M_p}$, they are all based on effective theories and have parameters that can be adjusted to fit the data. No model so far can accurately describe all form factors of the nucleon, as is necessary to fully understand the strong interaction. The result can also be expressed in terms of the non-spin-flip Dirac form factor F_{1p} , and spin-flip Pauli form factor F_{2p} , given by

$$F_{1p} = \frac{G_{E_p} + \tau G_{M_p}}{1 + \tau}; \quad F_{2p} = \frac{G_{M_p} - G_{E_p}}{\kappa_p(1 + \tau)}, \quad (5)$$

where κ_p is the anomalous magnetic moment of the proton, and $\tau = Q^2/4m^2$. The ratio F_{2p}/F_{1p} directly follows from G_{E_p}/G_{M_p} . In Fig. 3a, the results are compared with the pQCD predictions [9] that the asymptotic behavior of the form factors is $F_{1p} \propto 1/Q^4$ and $F_{2p} \propto 1/Q^6$, so that $Q^2 F_{2p}/F_{1p}$ would reach a constant value at high enough Q^2 . The data clearly indicate that this asymptotic regime has not been reached yet. Based on the results of Ref. [11], Ralston *et al.* [33] postulated a different scaling behavior, where F_{2p}/F_{1p} goes as $1/\sqrt{Q^2}$ instead of $1/Q^2$, arguing that it corresponds to the pQCD expectation if one takes into account contributions to the proton quark wave function from states with nonzero orbital angular momentum. The ratio




FIG. 3. Same legend as Fig. 2, for (a) $Q^2 F_{2p}/F_{1p}$ and (b) $\sqrt{Q^2} F_{2p}/F_{1p}$.

$\sqrt{Q^2} \frac{F_{2p}}{F_{1p}}$ is shown in Fig. 3b; a constant value is clearly reached starting at $Q^2 \sim 2 \text{ GeV}^2$.

In conclusion, we have measured G_{E_p}/G_{M_p} by polarization transfer to $Q^2 = 5.6 \text{ GeV}^2$. The ratio obtained in this experiment continues to decrease, as observed first in Ref. [11]. Extrapolation of the linear trend indicates that the electric form factor would cross zero at $Q^2 \sim 7.7 \text{ GeV}^2$. This result also reveals a flattening of the ratio $\sqrt{Q^2} F_{2p}/F_{1p}$ starting at $Q^2 \sim 2 \text{ GeV}^2$. A measurement of G_{E_p}/G_{M_p} to yet higher Q^2 is planned in the near future [34].

The Collaboration thanks the Hall A technical staff and the Jefferson Lab Accelerator Division for their outstanding support during the experiment. The Southern Universities Research Association (SURA) operates the Thomas Jefferson National Accelerator Facility for the United States Department of Energy under Contract No. DE-AC05-84ER40150. This work was also supported by the U.S. National Science Foundation and Department of Energy, the Italian Istituto Nazionale di Fisica Nucleare (INFN), the French Commissariat à l'Energie Atomique (CEA) and Centre National de la Recherche Scientifique (CNRS-IN2P3), the Natural Sciences and Engineering Research Council of Canada (NSERC), the EEC Grants No. INTAS 99-00125 for the Kharkov Institute of Physics and Technology, and No. CRDF UP2-2271, and the Swedish Natural Science Research Council.

*Currently at Thomas Jefferson National Accelerator Facility, Newport News, VA 23606.

†On leave of absence from Institute for Nuclear Research and Nuclear Energy, Sofia, Bulgaria.

‡Currently at George Washington University, Washington, D.C. 20052.

- [1] M.N. Rosenbluth, Phys. Rev. **79**, 615 (1950).
 [2] J. Litt *et al.*, Phys. Lett. **31B**, 40 (1970).
 [3] Ch. Berger *et al.*, Phys. Lett. **35B**, 87 (1971).
 [4] L.E. Price *et al.*, Phys. Rev. D **4**, 45 (1971).
 [5] W. Bartel *et al.*, Nucl. Phys. **B58**, 429 (1973).
 [6] A.F. Sill *et al.*, Phys. Rev. D **48**, 29 (1993).
 [7] R.C. Walker *et al.*, Phys. Rev. D **49**, 5671 (1994).
 [8] L. Andivahis *et al.*, Phys. Rev. D **50**, 5491 (1994).
 [9] G.P. Lepage and S.J. Brodsky, Phys. Rev. D **22**, 2157 (1980).
 [10] A.I. Akhiezer and M.P. Rekalo, Sov. J. Part. Nucl. **3**, 277 (1974); R. Arnold, C. Carlson, and F. Gross, Phys. Rev. C **23**, 363 (1981).
 [11] M.K. Jones *et al.*, Phys. Rev. Lett. **84**, 1398 (2000); V. Punjabi *et al.* (to be published).
 [12] B. Milbrath *et al.*, Phys. Rev. Lett. **80**, 452 (1998); **82**, 2221(E) (1999).
 [13] S. Dieterich *et al.*, Phys. Lett. B **500**, 47 (2001).
 [14] O. Gayou *et al.*, Phys. Rev. C **64**, 038202 (2001).
 [15] See <http://www.jlab.org/~moller>.
 [16] N. Fallete *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **459**, 412 (2001).
 [17] Stéphanie Escoffier, Thèse de doctorat, CEA Saclay, 2001.
 [18] See <http://www.jlab.org/Hall-A/equipment/HRS.html>; B. D. Anderson *et al.* (to be published).
 [19] M.K. Jones *et al.*, in *Intersections between Particle and Nuclear Physics*, edited by T. W. Donnelly, AIP Conf. Proc. No. 412 (AIP, New York, 1997), p. 342; L. Bimbot *et al.* (to be published).
 [20] K. Makino and M. Berz, COSY INFINITY version 8, Nucl. Instrum. Methods Phys. Res., Sect. A **427**, 338 (1999).
 [21] D. Besset *et al.*, Nucl. Instrum. Methods **166**, 512 (1979).
 [22] E.L. Lomon, Phys. Rev. C **64**, 035204 (2001).
 [23] R.F. Wagenbrunn *et al.*, Phys. Lett. B **511**, 33 (2001); e-print hep-ph/0108271.
 [24] F. Cardarelli and S. Simula, Phys. Rev. C **62**, 065201 (2000); S. Simula, e-print nucl-th/0105024.
 [25] G. Holzwarth, Z. Phys. A **356**, 339 (1996).
 [26] M.R. Frank, B.K. Jennings, and G.A. Miller, Phys. Rev. C **54**, 920 (1996).
 [27] L. Pentchev *et al.*, Jlab Technical Note No. TN-01-052, 2001.
 [28] A. V. Afanasev *et al.*, Phys. Lett. B **514**, 269 (2001).
 [29] M.P. Rekalo and E. Tomasi-Gustafsson, e-print nucl-th/0105002; A. V. Afanasev (private communication).
 [30] E.J. Brash, A. Kozlov, Sh. Li, and G. Huber (to be published).
 [31] F. Schlumpf, e-print hep-ph/9405284.
 [32] L. Ya. Glozman *et al.*, Phys. Rev. D **58**, 094030 (1998).
 [33] J. Ralston *et al.*, in *Intersections between Particle and Nuclear Physics*, AIP Conf. Proc. No. 549 (AIP, Melville, NY, 2000), p. 302; (private communication).
 [34] JLab experiment E01-109, C. F. Perdrisat, V. Punjabi, E. J. Brash, and M. K. Jones, spokespersons.