

Dynamically Stabilized Bright Solitons in a Two-Dimensional Bose-Einstein Condensate

Hiroki Saito and Masahito Ueda

Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan

(Received 3 September 2002; published 30 January 2003)

We demonstrate that a matter-wave bright soliton can be stabilized in 2D free space by causing the strength of interactions to oscillate rapidly between repulsive and attractive by using, e.g., Feshbach resonance.

DOI: 10.1103/PhysRevLett.90.040403

PACS numbers: 03.75.Kk, 05.30.Jp, 05.45.Yv, 34.50.-s

A pendulum with an oscillating pivot can have a stable configuration with its bob situated above the pivot (the inverted pendulum [1]). This well-known but surprising phenomenon is attributed to a net stabilizing force produced by alternating stabilizing and destabilizing forces at a frequency much faster than the natural oscillation frequency of the fixed-pivot pendulum. Similar physics has been employed to focus beams of charged particles in a synchrotron (alternating-gradient focusing [2]) and to trap ions in the Paul trap [3]. In this Letter, we present a novel application of such a stabilizing mechanism to producing a bright soliton in a two-dimensional (2D) Bose-Einstein condensate (BEC). By a “bright” soliton we mean a stable, solitary wave whose density is greater than the background one.

In 1D, when the interatomic interaction is attractive, bright solitons can be stabilized even without a trapping potential [4,5]. In 2D or 3D free space, however, the kinetic pressure and the attractive force cannot balance, and the condensate always collapses or expands. This can be understood by the following simple argument. When the characteristic size of the condensate is R , the kinetic and interaction energies are proportional to R^{-2} and $-R^{-d}$ in d dimensions, and an effective potential for R is the sum of these energies. The effective potential, therefore, has a minimum only for $d = 1$. Here we present a novel method to stabilize a soliton in 2D by causing the interaction to oscillate rapidly using, e.g., Feshbach resonance [6,7].

The system considered here is a BEC confined in a quasi-2D axisymmetric trap [8], where the axial confinement energy $\hbar\omega_z$ is much larger than the radial confinement and interaction energies. We assume that the condensate wave function ψ is always in the ground state of the harmonic potential with respect to the z direction, and that the dynamics are effectively 2D in the x - y plane; we will justify these assumptions later. We let the radial confinement frequency $\omega_{\perp}(t)$ and the s -wave scattering length $a(t)$ vary in time. The system is then described by the Gross-Pitaevskii (GP) equation:

$$i \frac{\partial \psi}{\partial t} = -\frac{1}{2} \nabla^2 \psi + \frac{\omega_{\perp}^2(t)}{2} r^2 \psi + g(t) |\psi|^2 \psi, \quad (1)$$

where $r^2 \equiv x^2 + y^2$ and $g(t) \equiv (8\pi m \omega_z / \hbar)^{1/2} N a(t)$ describes the strength of interactions. In Eq. (1), length, time, frequency, and ψ are measured in units of $d_0 \equiv [\hbar / (m \omega_{\perp 0})]^{1/2}$, $\omega_{\perp 0}^{-1}$, $\omega_{\perp 0}$, and $N^{1/2} / d_0$, where $\omega_{\perp 0} \equiv \omega_{\perp}(0)$.

We make the strength of interaction oscillate rapidly at frequency Ω . The oscillation of the scattering length with $\Omega \approx \omega_{\perp}$ is studied in Ref. [9] in a different context. Experimentally, the speed of the change in the strength of interaction using Feshbach resonance is limited by that of the applied magnetic field $\sim 1 \text{ G}/\mu\text{s}$ [10], which is sufficient for our purpose. In order to avoid nonadiabatic disturbances that destabilize a soliton state, we gradually switch on the interaction and simultaneously turn off the radial confinement potential as

$$g(t) = f(t)(g_0 + g_1 \sin \Omega t), \quad (2)$$

$$\omega_{\perp}^2(t) = 1 - f(t), \quad (3)$$

where $f(t)$ is a ramp function given by

$$f(t) = \begin{cases} t/T & (0 \leq t \leq T) \\ 1 & (t > T). \end{cases} \quad (4)$$

The interaction and the confinement potential can be changed independently using an optical trap and magnetic-field-induced Feshbach resonance.

We numerically solve the GP equation (1) with time-dependent parameters (2)–(4) using the Crank-Nicholson scheme [11]. The initial state is assumed to be the non-interacting ground state in the presence of the radial confinement potential with $\omega_{\perp 0}$. We gradually increase the strength of interaction and switch off the trap according to the linear ramp (4) with $T = 20$. Figure 1 demonstrates the dynamic stabilization of a soliton in 2D, where the parameters used are $g_0 = -2\pi$, $g_1 = 8\pi$, $\Omega = 40$ [Fig. 1(a)], and $\Omega = 30$ [Fig. 1(b)]. Since the value of $g_0 = -2\pi$ exceeds the critical strength of attractive interaction for the collapse $g_{\text{cr}} \approx -5.8$ [12], the condensate would collapse if the oscillating term $g_1 \sin \Omega t$ were absent. Even after the radial confinement potential vanishes ($t > 20$), the soliton shape is maintained as shown in the insets of Fig. 1. The plots of the peak density $|\psi(r=0)|^2$ and the monopole moment $\langle r \rangle = \int r |\psi|^2 d\mathbf{r}$ rapidly oscillate at frequency Ω . We note here that the dynamics of




FIG. 1. Time evolution of the peak density $|\psi(r=0)|^2$ and the monopole moment $\langle r \rangle = \int r |\psi|^2 dr$. The interaction $g(t) = -2\pi + 8\pi \sin\Omega t$ is switched on and simultaneously the radial confinement potential is switched off from $t = 0$ to 20 according to Eq. (4), where $\Omega = 40$ in (a) and $\Omega = 30$ in (b). The plots oscillate rapidly at Ω (which is beyond the resolution limit of the graph), and the vertical width of the plots represents the amplitude of the rapid oscillations. The insets show the density profiles $|\psi(r)|^2$ at $t = 50$ (solid curves) and $t = 0$ (dashed curves).

BEC can be separated into two parts: a rapidly oscillating part with small amplitude and a slow, smoothly varying part. After the rapid oscillations are averaged out, the width and the peak density are fairly constant as seen in Fig. 1(a). However, when the nonadiabaticity of ramp function $f(t)$ is not negligible, the lowest breathing mode (but no other radial modes) is excited as illustrated in Fig. 1(b), where the breathing mode is caused by an effective potential due to the oscillating interaction [see Eqs. (9)–(11)]. The amplitude of the excitation decreases with increasing ramp time T . Such excitations can be suppressed by choosing the appropriate ramp functions for $g(t)$ and $\omega_{\perp}(t)$. We have confirmed that no

symmetry-breaking mode grows in the azimuthal direction. The system is therefore free from such modulational instabilities.

We checked the validity of the quasi-2D approximation by numerically integrating the GP equation in 3D axisymmetric systems with large asymmetry parameters $\omega_z/\omega_{\perp} \sim 50$ and found that soliton stabilization occurs without axial modes being excited. Our prediction of dynamically stabilizing a bright soliton in 2D is, therefore, experimentally feasible with an oblate trap as realized by the MIT group [8].

To understand the behavior in Fig. 1, we employ a variational method with a Gaussian wave function [9,13,14]

$$\psi(r, t) = \frac{1}{\sqrt{\pi}R(t)} \exp\left[-\frac{r^2}{2R^2(t)} + i\frac{\dot{R}(t)}{2R(t)}r^2\right], \quad (5)$$

where $R(t)$ is the variational parameter that characterizes the size of the condensate, and the imaginary term in the exponent describes mass current. Substituting Eq. (5) into the action that derives the GP equation (1) and minimizing the action with respect to $R(t)$, we obtain the equation of motion for $R(t)$ as

$$\ddot{R}(t) = \frac{1}{R^3(t)} + \frac{g_0 + g_1 \sin\Omega t}{2\pi R^3(t)}, \quad (6)$$

where we set $f(t) = 1$. We separate $R(t)$ into the slowly varying part $R_0(t)$ and the rapidly oscillating part $\rho(t)$ as $R(t) = R_0(t) + \rho(t)$ [1]. When $\Omega \gg 1$, $\rho(t)$ becomes of the order of Ω^{-2} , and we keep the terms of the order of up to Ω^{-2} in the following analysis. Substituting this $R(t)$ into Eq. (6), we obtain the equations of motion for the rapidly and slowly varying parts as

$$\ddot{\rho}(t) = \frac{g_1}{2\pi R_0^3(t)} \sin\Omega t, \quad (7)$$

$$\ddot{R}_0(t) = \frac{g_0 + 2\pi}{2\pi R_0^3(t)} - \frac{3g_1}{2\pi R_0^4(t)} \overline{\rho(t) \sin\Omega t}, \quad (8)$$

where the overline indicates the time average of the rapid oscillation. Equation (7) yields $\rho(t) = -g_1/[2\pi\Omega^2 R_0^3(t)] \sin\Omega t$. Substituting this into Eq. (8), we obtain the equation of motion for the slowly varying part as

$$\ddot{R}_0 = -\frac{\partial}{\partial R_0} \left(\frac{g_0 + 2\pi}{4\pi R_0^2} + \frac{g_1^2}{16\pi\Omega^2 R_0^6} \right) \equiv -\frac{\partial U(R_0)}{\partial R_0}. \quad (9)$$

The minimum of the effective potential U is attained at

$$R_{\min}^4 = -\frac{3g_1^2}{4\pi\Omega^2(g_0 + 2\pi)}, \quad (10)$$

and the frequency of small oscillations (breathing mode) around the minimum is given by

$$\omega_{\text{br}}^2 = \frac{8\Omega^2}{3g_1^2} (g_0 + 2\pi)^2. \quad (11)$$

The Gaussian approximation thus indicates that a soliton is stable for $g_0 < -2\pi$.

The physical mechanism of soliton stabilization can be understood from the above discussion. In the inverted pendulum, the interplay between the micromotion of the bob and the force gradient (i.e., stronger oscillating force for larger deviation from the equilibrium position) produces a pseudopotential. Since the pseudopotential is proportional to the square of the amplitude of the oscillating force [1], a potential barrier is formed around the equilibrium position, thereby preventing the pendulum from swinging down. Such a mechanism also stabilizes a BEC in a double-well potential with oscillating interaction [15] and an optical beam propagating in nonlinear medium with an alternating nonlinearity [16]. In the present case, the oscillating “force” for R is given by $g_1 \sin \Omega t / (2\pi R^3)$, which becomes larger for smaller R . This force gradient and the micromotion of R produce a pseudopotential proportional to the square of the force $\propto R^{-6}$, which prevents the system from collapsing by counteracting the $-R^{-2}$ term that describes the attractive interaction.

In order to stabilize the soliton, $|g_0|$ must exceed the critical value of collapse, $|g_{\text{cr}}| \simeq 5.8$, which is smaller than that obtained with the Gaussian approximation, $|g_{\text{cr}}| = 2\pi$, since the Gaussian wave function underestimates the peak density [14]. In fact, the GP equation predicts that a soliton state for $g_0 = -2\pi$ is stable (Fig. 1). Although the Gaussian approximation does not accurately describe the exact soliton stability condition, Eqs. (10) and (11) capture the g_1 and Ω dependences of R_{min} and ω_{br} . Figure 2(a) illustrates the monopole moment $\langle r \rangle$ versus $(g_1/\Omega)^{1/2}$, where the circles are obtained by varying g_1 and the squares by varying Ω . We note that the plots are fairly well proportional to $(g_1/\Omega)^{1/2}$, in agreement with Eq. (10). Figure 2(b) shows the frequencies ω_{br} of the slow oscillations [as shown in Fig. 1(b)] versus Ω/g_1 . The plots are linear in Ω/g_1 , which is consistent with Eq. (11).

The appropriate parameter range for stabilizing solitons with size $R \simeq 1$ is found to be $0.4 \lesssim g_1/\Omega \lesssim 1.2$ and $g_0 \simeq -2\pi$ for Ω much larger than the breathing-mode frequency ω_{br} . For $\Omega \lesssim 10$, the rapid oscillations are disturbed by nonadiabatic slow dynamics, and the soliton becomes unstable. In analogy with the inverted pendulum problem, this corresponds to a situation in which the pendulum bob falls a long way during a single pivot cycle at a low pivot oscillation frequency. Thus, the effective force fails to stabilize the pendulum bob. When g_1/Ω is small or $|g_0|$ is large, the effective force is not sufficient to prevent the atoms from accumulating at the center. As a result, the peak density first grows, the condensate then expands due to the R^{-6} potential, and subsequently most




FIG. 2. (a) $(g_1/\Omega)^{1/2}$ dependence of the monopole moment $\langle r \rangle$ and (b) Ω/g_1 dependence of the breathing-mode frequency ω_{br} , where $g_0 = -2\pi$. The circles are obtained by varying g_1 for $\Omega = 30$, and the squares by varying Ω for $g_1 = 8\pi$. The dashed lines show $1.06(g_1/\Omega)^{1/2}$ in (a) and $0.32\Omega/g_1$ in (b), which are determined to best fit the plots.

of the expanded atoms accumulate at the center due to the $-R^{-2}$ potential. Thus, the condensate repeatedly contracts and expands. In each expansion, atoms that are elastically scattered with high energy cannot return to the soliton region due to the absence of the external confinement potential, and the condensate gradually decays.

In conclusion, we have demonstrated that a matter-wave bright soliton can be stabilized in 2D free space by oscillating the strength of the interaction around an attractive value $g_0 < g_{\text{cr}}$ with an amplitude $g_1 > |g_0|$. The rapid oscillation of interaction produces an effective barrier that prevents the condensate from collapsing and stabilizes a soliton. This novel technique of increasing dimensions of a matter-wave bright soliton might be applied for quantum information processing using BEC solitons on a microchip substrate [17]. It merits further study to examine whether BEC “droplets” in 3D can be created with appropriate parameters and ramp schemes.

This work was supported by a Grant-in-Aid for Scientific Research (Grant No. 11216204) and Special Coordination Funds for Promoting Science and Technology by the Ministry of Education, Science, Sports, and Culture of Japan, by the Toray Science Foundation, and by the Yamada Science Foundation.

Note added.—After submission of our Letter, a paper appeared [18] which reached the same conclusion that a bright soliton can be stabilized in 2D with a similar scheme.

[1] L. D. Landau and E. M. Lifshitz, *Mechanics* (Pergamon, Oxford, 1960).

- [2] R. P. Feynman, R. B. Leighton, and M. L. Sands, *The Feynman Lectures on Physics* (Addison-Wesley, Reading, MA, 1964), Vol. II, Chap. 29.
- [3] H. G. Dehmelt, *Adv. At. Mol. Phys.* **3**, 53 (1967).
- [4] L. Khaykovich, F. Schreck, G. Ferrari, T. Bourdel, J. Cubizolles, L. D. Carr, Y. Castin, and C. Salomon, *Science* **296**, 1290 (2002).
- [5] K. E. Strecker, G. B. Partridge, A. G. Truscott, and R. G. Hulet, *Nature (London)* **417**, 150 (2002).
- [6] S. Inouye, M. R. Andrews, J. Stenger, H.-J. Miesner, D. M. Stamper-Kurn, and W. Ketterle, *Nature (London)* **392**, 151 (1998).
- [7] S. L. Cornish, N. R. Claussen, J. L. Roberts, E. A. Cornell, and C. E. Wieman, *Phys. Rev. Lett.* **85**, 1795 (2000).
- [8] A. Görlitz, J. M. Vogels, A. E. Leanhardt, C. Raman, T. L. Gustavson, J. R. Abo-Shaer, A. P. Chikkatur, S. Gupta, S. Inouye, T. Rosenband, and W. Ketterle, *Phys. Rev. Lett.* **87**, 130402 (2001).
- [9] F. Kh. Abdullaev, J. C. Bronski, and R. M. Galimzyanov, *cond-mat/0205464*.
- [10] N. R. Claussen, E. A. Donley, S. T. Thompson, and C. E. Wieman, *Phys. Rev. Lett.* **89**, 010401 (2002).
- [11] P. A. Ruprecht, M. J. Holland, K. Burnett, and M. Edwards, *Phys. Rev. A* **51**, 4704 (1995).
- [12] S. K. Adhikari, *Phys. Lett. A* **265**, 91 (2000).
- [13] V. M. Pérez-García, H. Michinel, J. I. Cirac, M. Lewenstein, and P. Zoller, *Phys. Rev. Lett.* **77**, 5320 (1996); *Phys. Rev. A* **56**, 1424 (1997).
- [14] H. Saito and M. Ueda, *Phys. Rev. A* **63**, 043601 (2001).
- [15] F. Kh. Abdullaev and R. A. Kraenkel, *Phys. Lett. A* **272**, 395 (2000).
- [16] I. Towers and B. A. Malomed, *J. Opt. Soc. Am. B* **19**, 537 (2002).
- [17] W. Hänsel, P. Hommelhoff, T. W. Hänsch, and J. Reichel, *Nature (London)* **413**, 498 (2001).
- [18] F. Kh. Abdullaev, J. G. Caputo, R. A. Kraenkel, and B. A. Malomed, *Phys. Rev. A* **67**, 013605 (2003).