

**Erratum: Sigma-Nucleus Potential in  $A = 28$**   
**[Phys. Rev. Lett. 89, 072301 (2002)]**

H. Noumi, P. K. Saha, D. Abe, S. Ajimura, K. Aoki, H. C. Bhang, T. Endo, Y. Fujii, T. Fukuda, H. C. Guo, K. Imai, O. Hashimoto, H. Hotchi, E. H. Kim, J. H. Kim, T. Kishimoto, A. Krutenkova, K. Maeda, T. Nagae, M. Nakamura, H. Outa, M. Sekimoto, T. Saito, A. Sakaguchi, Y. Sato, R. Sawafta, Y. Shimizu, T. Takahashi, L. Tang, H. Tamura, K. Tanida, T. Watanabe, H. H. Xia, S. H. Zhou, L. H. Zhu, and X. F. Zhu

(Received 12 December 2002; published 28 January 2003)

DOI: 10.1103/PhysRevLett.90.049902

PACS numbers: 21.80.+a, 13.75.Ev, 25.80.Hp, 25.80.Nv, 99.10.Cd

In Fig. 3 and its inset the vertical scales should be reduced by a factor of 4. This plotting error affects only the figure. All relevant quantities in the text and in the table are correct as published. We regret the oversight.

The corrected version of Fig. 3 is reproduced here. This correction does not affect any results or conclusions of the published paper.




FIG. 3. Inclusive ( $\pi^-$ ,  $K^+$ ) spectrum on Si at  $\theta_K = 6^\circ \pm 2^\circ$ . The curves are the calculated spectra for the repulsive (solid) and shallow (dashed)  $\Sigma$ -nucleus potentials, fitted to the measured spectrum. A value of the scaling factor  $\kappa$  and  $\chi^2$  per degree of freedom are shown for each fitting. See text for details.