

Solar Flare Intermittency and the Earth's Temperature Anomalies

Nicola Scafetta^{1,2} and Bruce J. West^{1,2,3}

¹*Pratt School EE Department, Duke University, P.O. Box 90291, Durham, North Carolina 27708, USA*

²*Physics Department, Duke University, Durham, North Carolina 27708, USA*

³*Mathematics Division, Army Research Office, Research Triangle Park, North Carolina 27709, USA*

(Received 6 March 2003; published 17 June 2003)

We argue that Earth's short-term temperature anomalies and the solar flare intermittency are linked. The analysis is based upon the study of the scaling of both the spreading and the entropy of the diffusion generated by the fluctuations of the temperature time series. The joint use of these two methods evidences the presence of a Lévy component in the temporal persistence of the temperature data sets that corresponds to the one that would be induced by the solar flare intermittency. The mean monthly temperature data sets cover the period from 1856 to 2002.

DOI: 10.1103/PhysRevLett.90.248701

PACS numbers: 94.10.Dy, 05.45.Tp, 47.27.Nz, 95.75.Wx

The historical recognition that the Sun warms the earth has suggested a direct connection between the average global temperature and solar activity. Consequently, any significant changes in solar activity should result in equivalent changes in the earth's global temperature. The literature on the solar influence on the earth's temperature is quite extensive [1–3], indicating the importance of the problem and that there are many issues that require further investigation. Herein we address the relation between the statistics of solar flare activity and the fluctuations in the earth's global temperature.

The dynamics of the Sun's surface is turbulent, as is evidenced by changes in solar flare activity, with 11- or 22-year solar cycles [2,3] and strong erratic fluctuations associated with solar flare intermittency [3–6]. Solar irradiance changes in accordance with the frequency of solar flares because each flare releases more energy than the background irradiance [3]. This time variation in the frequency of solar flares induces a similar pseudoperiodic cycle in the earth's average temperature, as well as produces trends that move the global temperature up or down for tens or even hundreds of years [1–3]. However, it is less evident that short-term (weekly and monthly) changes in the global temperature are tied to solar activity whose short-time fluctuations would have the intermittent dynamics of solar flares. Traditional measures, such as cross-correlation functions, would not show the connection between short-time fluctuations in the global temperature and solar flare activity, because of the strong nonlinear hydrodynamic interactions across the earth's surface. For example, the hydrodynamic interaction of the atmosphere over land and water would suppress any direct correlation between the intermittent Sun's irradiance and the earth regions' short-time response.

This Letter focuses on an alternate approach to establishing the connection between the Sun's irradiance and the earth's temperature fluctuations. A link between the two phenomena is detected through a detailed scaling analysis of the time series for the earth's temperature and the time series for the solar flare frequency.

Considering a solar flare as an event, the time series for the number of solar flares has been interpreted as a waiting time distribution function between events. The solar flare waiting time distribution function is determined to be an inverse power-law probability density function (*iplpdf*) [4–6]. Such dynamical stochastic processes can be described by generalizations of random walks. A *Lévy flight*, for example, is such a process with a diverging second moment. A *Lévy walk*, on the other hand, visits the same spatial sites as does a Lévy flight, but each step takes a finite time and the second moment is finite. The time necessary to complete a step in a Lévy walk is specified by an inverse power-law waiting time distribution function, as first noticed by Shlesinger *et al.* [7]. It has been determined [4] that a Lévy walk can describe the intermittent solar flare signal. Scafetta *et al.* [8,9] established that, in general, the presence of a Lévy-walk process in a given time series can be detected by the joint use of two separate scaling techniques, the diffusion entropy analysis (DEA), and standard deviation analysis (SDA). We apply the same approach to the analysis of temperature data sets and compare them with the Lévy-walk statistics induced by solar flare intermittency.

We study the earth's temperature anomalies both globally and locally, that is, North-South hemispheres and Land-Ocean regions (see Fig. 1). The technical term *temperature anomalies* denotes temperature departures from the 1961–1990 mean temperature value. These data are recognized in the geophysical community as among the most accurate data files for global air temperature, and global sea-surface temperature (SST). The basic data set of global Earth's temperature anomalies (HadCRUT) is a combination of Land air temperature anomalies [10] (CRUTEM1) and sea-surface temperature anomalies [11] on a $5^\circ \times 5^\circ$ grid-box basis. The merging of the two data sets is discussed in Parker *et al.* [11] and more recently in Jones *et al.* [12]. The Land surface time series are calculated using data from the Global Historical Climatology Network (Version 2) and sea-surface temperature anomalies from the United




FIG. 1. Global and local Earth temperature anomalies in degrees Celsius (years: 1856–2002).

Kingdom MOHSST data set and the NCEP Optimum Interpolated SSTs (Version2). The data that we analyze are mean monthly temperature anomalies downloaded from Climatic Research Unit, United Kingdom [13] (Global and North-South temperatures, 1856–2002) and from National Climatic Data Center, U.S.A. [14] (Land-Ocean temperatures, 1880–2002).

The analysis of the above data sets is done with both DEA and SDA techniques. These methods are based on the prescription that the numbers in a time series $\{\xi_i\}$ are the fluctuations of a diffusion trajectory; see Refs. [4,8,9] for details. Therefore, we shift our attention from the time series $\{\xi_i\}$ to the probability density function (pdf) $p(x, t)$ of the corresponding diffusion process. Here x denotes the variable collecting the fluctuations and is referred to as the diffusion variable. The scaling property of $p(x, t)$ takes the form

$$p(x, t) = \frac{1}{t^\delta} F\left(\frac{x}{t^\delta}\right), \quad (1)$$

where δ is the scaling exponent. The DEA [8] is based on the evaluation of the Shannon entropy, $S(t)$, using the pdf (1). If the scaling condition of Eq. (1) holds true, it is easy to prove that the entropy is

$$S(t) = - \int_{-\infty}^{\infty} p(x, t) \ln[p(x, t)] = A + \delta \ln(t), \quad (2)$$

where A is a constant. The SDA [8], instead, is based on the evaluation of the standard deviation $D(t)$ using the same pdf (1) and yields

$$D(t) = \sqrt{\langle x^2; t \rangle - \langle x; t \rangle^2} \propto t^H, \quad (3)$$

where H is the Hurst exponent [8,15].

The Lévy-walk statistics is a particular form of anomalous diffusion obtained by generalizing the central limit theorem [16]. While all theoretical details can be found in Refs. [4,8,9] and in the enclosed references, here we show the properties of a Lévy walk using a model that simulates solar flare intermittency. We generate an artificial sequence $\{\tau_j\}$ distributed according to an *iplpdf*,

$$\psi(\tau) \propto \frac{1}{(T + \tau)^\mu}, \quad (4)$$

where T is a positive constant. In the computer-generated example, we use $T = 90$ and $\mu = 2.5$. We use $2 < \mu < 3$, so as to violate the constraint of a finite variance on the central limit theorem. This assumption is necessary to obtain Lévy-walk statistics. The sequence of waiting times $\{\tau_j\}$ is used to generate a time series $\{\xi_i\}$ given by $\xi_i = 1$ only at the occurrence of the events, that is, for $i = [\sum_{j=1}^m \tau_j]$; $[a]$ is the integer part of a . However, in the interval between two events, that is, for $[\sum_{j=1}^m \tau_j] < i < [\sum_{j=1}^{m+1} \tau_j]$, we assume $\xi_i = 0$. As shown in Refs. [4,8,9], by using results of Refs. [16–18] the time series $\{\xi_i\}$ generates a diffusion process with a scaling exponent $\delta = 1/(\mu - 1)$. Moreover, this particular approach to Lévy statistics yields a diffusion process with finite second moments, as in the Lévy walk [7]. Consequently, it is possible to evaluate the scaling exponent of the standard deviation, H , via Eq. (3). Figure 2(a) shows the first 1650 data points in the computer-generated sequence $\{\xi_i\}$, which correspond to the first 5.5 time units, of a 510 000 long data sequence $\{\xi_i\}$. In Fig. 2(b), we show the sequence of frequencies $\{f_i\}$ of impulses for each time unit where we use 1 time unit = 300 natural units of τ .

The analysis is based on the comparison of the scaling exponents δ and H measured by DEA and SDA, respectively. It has been shown [8,9] that, if a time series is characterized by Gaussian statistics, the two scaling exponents are identical; that is, $\delta = H$. If, instead, the process under study is characterized by a Lévy walk, the two scaling exponents δ and H can be both related to the exponent μ of the waiting time *iplpdf* of the underlying intermittent process generating the Lévy statistics. In the Lévy-walk case, the scaling exponent H is slightly larger than δ and the two exponents are related to μ and one another via the following relation [8,9]:

$$\delta = \frac{1}{3 - 2H} = \frac{1}{\mu - 1}. \quad (5)$$




FIG. 2. (a) Computer-generated intermittent sequences of impulses. (b) The frequency of impulses for 1700 time units.

We refer to (5) as the Lévy-walk diffusion relation. The assessment of the property expressed by Eq. (5), that is, the comparison between the value of the scaling exponents δ and H measured for the earth's temperature data sets and their relation with the scaling exponents δ and μ measured for the hard x-ray solar flares waiting time series [4], is used in this Letter to verify the conjecture that the earth's global temperature inherits the intermittent Lévy-like nature of solar flares, within the temporal range from a few weeks to a few months.

In the computer-generated model, we interpret the fluctuations in the frequency of the pulses shown in Fig. 2(b) as inducing a similar memory pattern in the monthly mean global temperature fluctuations. In fact, a higher solar flare frequency implies a higher monthly solar activity. The DEA and SDA of the computer-generated data are shown in Fig. 3. These data are fit with Eqs. (2) and (3) yielding the two scaling coefficients $\delta = 0.67$ and $H = 0.75$. These coefficients are consistent with $\mu = 2.5$ obtained by using Eq. (5); the known input value for this realization of the model. The large picture in Fig. 3 refers to the analysis of the original 510 000 data-point sequence $\{\xi_{ij}\}$. The inset refers to the analysis of 1700 data-point frequency sequence $\{f_i\}$ shown in Fig. 2(b), a length compatible with the length of the mean monthly temperature data sets here analyzed. The two graphs in Fig. 3 show very good agreement with Lévy-walk theory, that is, with Eq. (5), because both time series produce similar diffusion trajectories. In particular, the small inset of Fig. 3 shows the typical *bifurcation* associated to the two scaling laws that characterize Lévy-walk statistics and that the statistics generated by only $N = 1700$ frequency data points is rich enough to obtain a satisfactory pdf with Lévy-walk scaling properties.

Figure 4 shows the numerical results by applying DEA and SDA to the temperature fluctuation data of the four

regions of the earth. On the ordinate axis, we plot $\ln[D(t)/D(1)]$ and $S(t) - S(1)$; thus, all curves start from 0. The scaling exponents δ and H are obtained by fitting the first ten points. This analysis for the Global temperature (not shown in the figure) yields $\delta = 0.90 \pm 0.02$ and $H = 0.95 \pm 0.02$. The high values of the exponents imply a strong persistence in the temperature fluctuations. Both scaling exponents δ and H are slightly larger for the South hemisphere than for the North hemisphere and for the Ocean than for the Land. A reasonable explanation is that the Ocean is an almost stationary system with a very high effective heat capacity. The Land, instead, is subject to stronger random fluctuations (as Fig. 1 shows) due to a lower effective heat capacity and a higher morphological variability due, for example, to the presence of deserts, forests, mountains, and valleys. Finally, we note that the standard deviation scaling exponents H are larger than the diffusion entropy scaling exponents δ and seem to fulfill the Lévy-walk diffusion relation (5) within the accuracy of our statistical analysis.

The scaling behavior in the temperature anomalies can be related to the intermittency of the solar flare activity through the mechanism suggested by the model depicted in Figs. 2 and 3. In fact, the waiting time distribution of solar flares is characterized by an *iplpdf* of the type of Eq. (4) with $\mu = 2.14 \pm 0.05$ [4]. This value of μ would imply a Lévy walk with $\delta = 0.88 \pm 0.02$ and $H = 0.93 \pm 0.02$ (5) for a shuffled data set that would destroy any temporal correlation among flares and conserve only the Lévy component; see Ref. [4] about the effect of the shuffling on these data. We observe that the difference of the above two scaling exponent values gives $H - \delta = 0.05 \pm 0.02$; that is a value compatible with the difference between H and δ measured for all the temperature data sets. In particular, $H - \delta = 0.05 \pm 0.02$ for the Global temperature and for both South and Ocean regions and $H - \delta = 0.04 \pm 0.02$ for North and Land regions.




FIG. 3. DEA and SDA of the computer-generated Lévy-walk sequences shown in Figs. 2(a) and 2(b). The two straight lines show the two theoretical scaling coefficients $\delta = 0.67$ and $H = 0.75$ for $\mu = 2.5$.




FIG. 4. DEA and SDA of the North-South hemisphere and Land-Ocean region temperature anomalies. The scaling exponents H and δ are reported in the figure; the error is ± 0.02 .




FIG. 5. DEA of the unshuffled waiting time of solar flares sequence during the period 1991–2000. The value of the scaling exponent is $\delta = 0.94 \pm 0.02$.

To understand the actual values that we measure for H and δ , we first notice that the waiting time sequence of solar flares is weakly correlated [4]. Therefore, the effective value of δ for the original unshuffled sequence is slightly larger than that calculated via the waiting time *iplpdf* exponent μ for flares [4], $\delta = 0.94 \pm 0.02$, as Fig. 5 shows. We observe that this value of δ is perfectly compatible with $\delta = 0.93 \pm 0.02$ measured for the Ocean temperature. The Ocean, in fact, is the region of the earth that should best mirror solar activity because it is the largest and most homogeneous region of the earth with a very high effective heat capacity; properties that retain correlated events in the temperature. Also we observe that the occurrence of uncorrelated events would slightly reduce the Lévy-like memory and increase the Gaussianity of the data. This might explain why the scaling exponents H and δ as well as their difference slightly decrease (0.04 against 0.05) between the North and Land regions. Finally, Fig. 5 shows that the scaling properties for the waiting time of solar flares last at least 200 consecutive flares and the waiting time for clusters of 200 solar flares may last for several months. The largest waiting time between two consecutive solar flares, during the period 1991–2000, is almost 3 months [4], and on average there are almost 60 solar flares per month. Therefore, the temporal resolution of these clusters is compatible with the temporal range of scaling of the temperature data sets that last at least 1 to almost 20 months, as Fig. 4 shows.

In conclusion, the affinity of the scaling exponents obtained through our analysis suggests that the earth's temperature anomalies inherit a Lévy-walk memory component from the intermittency of solar flares. We obtain reliable scaling properties within a short-time interval, ranging from a few weeks to a few months.

Moreover, the joint use of SDA and DEA has proven to be very useful in detecting a significant Lévy component in the time series and, in general, it may suggest the presence of a link between a phenomenon under study and an intermittent process characterized by a waiting time *iplpdf*.

N.S. thanks the ARO for support under Grant No. DAAG5598D0002.

-
- [1] W. Soon, S. Baliunas, E. S. Posmentier, and P. Okeke, *New Astronomy* **4**, 563 (2000); M. Sharma, *Earth Planet. Sci. Lett.* **199**, 459 (2002); M. Lockwood, *J. Geophys. Res.* **106**, 16021 (2001); W. Soon, E. Posmentier, and S. Baliunas, *Ann. Geophys. Atm. Hydr. Space Sci.* **18**, 583 (2000).
 - [2] T. Baranyi, A. Ludmany, and H. Coffey, *Geophys. Res. Lett.* **25**, 2269 (1998); D. Shindell, D. Rind, N. Balachandran, J. Lean, and P. Lonergan, *Science* **284**, 305 (1999); M. Lockwood, R. Stamper, and M. N. Wild, *Nature (London)* **399**, 437 (1999).
 - [3] J. D. Fix, *Astronomy, Journey to the Cosmic Frontier* (McGraw-Hill, New York, 2001), 2nd ed.
 - [4] P. Grigolini, D. Leddon, and N. Scafetta, *Phys. Rev. E* **65**, 046203 (2002).
 - [5] G. Boffetta, V. Carbone, P. Giuliani, P. Veltri, and A. Vulpiani, *Phys. Rev. Lett.* **83**, 4662 (1999).
 - [6] P. Giuliani, V. Carbone, P. Veltri, G. Boffetta, and A. Vulpiani, *Physica (Amsterdam)* **280A**, 75 (2000).
 - [7] M. F. Shlesinger, B. J. West, and J. Klafter, *Phys. Rev. Lett.* **58**, 1100 (1987).
 - [8] N. Scafetta and P. Grigolini, *Phys. Rev. E* **66**, 036130 (2002).
 - [9] N. Scafetta, V. Latora, and P. Grigolini, *Phys. Rev. E* **66**, 031906 (2002).
 - [10] P. D. Jones, *J. Clim.* **7**, 1794 (1994).
 - [11] D. E. Parker, P. D. Jones, A. Bevan, and C. K. Folland, *J. Geophys. Res.* **99**, 14373 (1994).
 - [12] P. D. Jones, T. J. Osborn, K. R. Briffa, C. K. Folland, E. B. Horton, L. V. Alexander, D. E. Parker, and N. A. Rayner, *J. Geophys. Res.* **106**, 3371 (2001).
 - [13] Climatic Research Unit, United Kingdom, <http://www.cru.uea.ac.uk>
 - [14] National Climatic Data Center, U.S.A., <http://lwf.ncdc.noaa.gov>
 - [15] B. B. Mandelbrot, *The Fractal Geometry of Nature* (Freeman, New York, 1983).
 - [16] B. V. Gnedenko and A. N. Kolomogorov, *Limit Distributions for Sums of Random Variables* (Addison-Wesley, Reading, MA, 1954).
 - [17] J. Klafter, M. F. Shlesinger, and G. Zumofen, *Phys. Today* **49**, No. 2, 33 (1996); G. Zumofen, J. Klafter, and M. F. Shlesinger, *Lect. Notes Phys.* **519**, 15 (1998).
 - [18] W. Feller, *Trans. Am. Math. Soc.* **67**, 98 (1949).