

Relation between Proton and Neutron Asymptotic Normalization Coefficients for Light Mirror Nuclei and its Relevance to Nuclear Astrophysics

N. K. Timofeyuk,¹ R. C. Johnson,¹ and A. M. Mukhamedzhanov²

¹*Department of Physics, University of Surrey, Guildford, Surrey GU2 7XH, England, United Kingdom*

²*Cyclotron Institute, Texas A&M University, College Station, Texas 77843, USA*

(Received 2 June 2003; published 4 December 2003)

We show how the charge symmetry of strong interactions can be used to relate the proton and neutron asymptotic normalization coefficients (ANCs) of the one-nucleon overlap integrals for light mirror nuclei. This relation extends to the case of real proton decay where the mirror analog is a virtual neutron decay of a loosely bound state. In this case, a link is obtained between the proton width and the squared ANC of the mirror neutron state. The relation between mirror overlaps can be used to study astrophysically relevant proton capture reactions based on information obtained from transfer reactions with stable beams.

DOI: 10.1103/PhysRevLett.91.232501

PACS numbers: 24.50.+g, 27.20.+n, 27.30.+t

The astrophysical S -factor associated with the peripheral proton capture reaction $B(p, \gamma)A$ at stellar energies is well known [1] to be related to the asymptotic normalization coefficient (ANC) of the virtual decay $A \rightarrow B + p$. The same ANCs play a crucial role in other peripheral processes such as transfer reactions whose cross sections are significantly higher and therefore more easily measurable than those of the direct capture processes at astrophysically relevant energies [1]. The study of ANCs of astrophysical interest is a new and rapidly developing direction in modern experimental nuclear physics [2,3]. However, in order to exploit these ideas to determine the ANCs for light proton-rich nuclei of importance to nuclear astrophysics, the corresponding transfer reactions often require the use of weak radioactive beams which generally involves more difficult and less accurate experiments than are possible with stable beams. The higher intensities of stable beams means that they can be used at energies below the Coulomb barrier where the sensitivity to optical potentials, which are the main uncertainty of ANCs determined from transfer reactions, is minimised. We point out here that the ANC of the virtual neutron decay of the nucleus mirror to A , which may be susceptible to study with stable beams, is related in a model independent way by the charge symmetry of nuclear forces to the ANC of the corresponding proton decay of A . We propose to exploit this new insight to predict peripheral reaction cross sections in stars.

An asymptotic normalization coefficient (ANC) is one of the fundamental characteristics of the virtual decay of a nucleus into two clusters and is equivalent to the coupling constants in particle physics [4]. When multiplied by a trivial factor, it equals the on-shell amplitude for the virtual decay into two clusters, and it determines the large distance behavior of the projection of the bound state wave function of the nucleus onto a binary channel.

In earlier work [5,6], the ANCs for the one-nucleon virtual decays of the mirror pairs ${}^8\text{B}$ - ${}^8\text{Li}$ and ${}^{12}\text{N}$ - ${}^{12}\text{B}$

were studied in a microscopic approach. The calculated ANCs themselves depended strongly on the choice of the nucleon-nucleon (NN) force but the ratios of ANCs for mirror pairs were practically independent of the choice of the NN force. This observation is based thus far entirely on the calculations using detailed models of nuclear structure. We now show that it follows naturally as a consequence of the charge symmetry of nuclear forces [7].

The ANC C_{lj} for the one-nucleon virtual decay $A \rightarrow B + N$ is defined via the tail of the overlap integral $I_{lj}(r)$ between the wave functions of nuclei A and $B = A - 1$, where l is the orbital momentum and j is the total relative angular momentum between B and N . Asymptotically, this overlap behaves as

$$\sqrt{A}I_{lj}(r) \approx C_{lj} \frac{W_{-\eta, l+1/2}(2\kappa r)}{r}, \quad r \rightarrow \infty, \quad (1)$$

where $\kappa = (2\mu\epsilon/\hbar^2)^{1/2}$, ϵ is the one-nucleon separation energy, $\eta = Z_B Z_N e^2 \mu / \hbar^2 \kappa$, μ is the reduced mass for the $B + N$ system, and W is the Whittaker function. It follows from [4,5,8] that C_{lj} can be expressed in terms of the many-body wave functions of the nuclei A and B :

$$C_{lj} = -\frac{2\mu\sqrt{A}}{\hbar^2} \times \langle [\varphi_l(i\kappa r) Y_l(\hat{\mathbf{r}}) \otimes \chi_{1/2}]_j \otimes \Psi_{J_B} \rangle_{J_A} \langle \hat{\mathbf{V}} | \Psi_{J_A} \rangle, \quad (2)$$

where

$$\varphi_l(i\kappa r) = e^{-i\sigma_l} F_l(i\kappa r) / \kappa r, \quad (3)$$

F_l is the regular Coulomb wave function at imaginary momentum $i\kappa$, $\sigma_l = \arg \Gamma(l + 1 + i\eta)$, r is the distance between N and the center of mass of B , and

$$\hat{\mathbf{V}} = \sum_{i=1}^{A-1} V_{NN}(|\mathbf{r}_i - \mathbf{r}_A|) + \Delta V_{\text{coul}} = \hat{\mathbf{V}}_N + \Delta V_{\text{coul}}, \quad (4)$$

$$\Delta V_{\text{coul}} = \sum_{i=1}^{A-1} \frac{e_i e_A}{|\mathbf{r}_i - \mathbf{r}_A|} - \frac{Z_B e_A e}{r}. \quad (5)$$

Here e_i (e_A) is the charge of the i th (A th) nucleon, Z_B is the charge of the residual nucleus B , and V_{NN} is the two-body nuclear NN potential. If the separated nucleon is a neutron, φ_l is replaced by the Bessel function $j_l(i\kappa r)$.

ANCs can be obtained from Eq. (2) using wave functions which model the structure of nuclear interior well, for example, from the oscillator shell model [9]. The incorrect behavior of these model wave functions at large distances plays a minor role because of the presence of the short range NN potential on the right-hand side of Eq. (2) [8]. We have performed such calculations for several $0p$ shell nuclei, some of which are of astrophysical importance, using fixed $0\hbar\omega$ wave functions obtained in [10] and various NN potentials from literature [11]. The oscillator radii were chosen to provide correct sizes for the nuclei considered. In these calculations, mirror nuclei have exactly the same wave functions but, of course, the mirror ANCs are different because of different functions $\varphi_l(i\kappa r)$ involved. The $|C_{ij}|^2$ values change by a factor of 2 for different NN potential choice, but the ratio $\mathcal{R} = |C_p/C_n|^2$, where C_p and C_n are the proton and neutron ANCs for mirrors and, hence, may refer to different nuclei, changes by less than 4% for each mirror pair of overlaps (see Table I and [17]).

The observed effect has the following explanation. We first replace ΔV_{coul} by $V_{\text{coul}}(r) - Z_B e_A e/r$, where $V_{\text{coul}}(r)$ is the monopole Coulomb interaction of the A th nucleon with the nucleus B . This ignores higher multipole components of ΔV_{coul} . Equation (2) can then be replaced exactly by a formula in which ΔV_{coul} is removed from the matrix element and $\varphi_l(r)$ is replaced by $\varphi_l^{\text{mod}}(r)$. The latter is defined as the regular solution of the Schrödinger equation with the potential $V_{\text{coul}}(r)$ and which is normal-

ized so that $\varphi_l(r) = \varphi_l^{\text{mod}}(r)$ outside the charge radius of B . Inside the charge radius, the potential $V_{\text{coul}}(r)$ varies little over the nuclear volume and can be replaced by a constant equal to the separation energies difference $\epsilon_n - \epsilon_p$. Hence, in the nuclear interior $r \leq R_N$, which is all that matters on the right-hand side of Eq. (2), we can use

$$\varphi_l^{\text{mod}}(r) = \frac{F_l(i\kappa_p r_N)}{\kappa_p R_N j_l(i\kappa_n R_N)} j_l(i\kappa_n r), \quad r \leq R_N, \quad (6)$$

where $i\kappa_p$ and $i\kappa_n$ are determined by the proton and neutron separation energies ϵ_p and ϵ_n . Using Eq. (6) in the modified Eq. (2) and making the assumption that the difference between the wave functions for mirror pairs in the nuclear interior can be ignored, we find

$$\mathcal{R} \approx \mathcal{R}_0 = \left| \frac{F_l(i\kappa_p R_N)}{\kappa_p R_N j_l(i\kappa_n R_N)} \right|^2. \quad (7)$$

\mathcal{R}_0 depends on the NN force only implicitly through R_N .

The values of \mathcal{R}_0 , presented in Table I, have been calculated for $R_N = 1.3B^{1/3}$. They change by less than 2%, when R_N is varied from 2.5 to 4.5 fm in each case, and are smaller by less than 7% than the \mathcal{R} values obtained from microscopic calculations. Equation (7) correctly predicts the dependence of \mathcal{R} on neutron and proton separation energies. The tendency of \mathcal{R}_0 to underestimate \mathcal{R} can be attributed to the contributions from the r^{-2} and r^{-3} multipoles of ΔV_{coul} . When these multipoles are excluded from the microscopic calculations, the \mathcal{R} values decrease and become equal to \mathcal{R}_0 within the uncertainty in its definition.

In practice, overlap integrals for transfer reactions are frequently modeled as normalized single-particle wave functions times spectroscopic factors S , so that $C_{p(n)} = \sqrt{S_{p(n)}} b_{p(n)}$, where $b_{p(n)}$ is the single-particle proton (neutron) ANC. The derivation above shows that the result [Eq. (7)] is valid for $|b_p/b_n|^2$ if we assume that the

TABLE I. Ratio \mathcal{R} of the proton and neutron squared ANCs for mirror overlap integrals calculated with Gogny, Pires, and De Tourreil (GPT) [12], Volkov V1 [13], Brink-Boeker B1 [14], and four versions of the M3Y effective NN potentials [15,16]. M3Y(R), M3Y(HJ), and M3Y(P) were fitted to the oscillator G -matrix elements of the Reid, Hamada-Johnston, and Paris NN potentials, respectively, and M3Y(E) was fitted to the oscillator G -matrix elements derived from the NN scattering data. Analytical estimates \mathcal{R}_0 are also shown.

Overlap	Mirror overlap	j	GPT	V1	B1	M3Y(E)	M3Y(R)	M3Y(P)	M3Y(HJ)	\mathcal{R}_0
$\langle {}^6\text{Li} {}^7\text{Be} \rangle$	$\langle {}^6\text{Li} {}^7\text{Li} \rangle$	1/2	1.05	1.05	1.05	1.05	1.05	1.05	1.05	1.05
		3/2	1.05	1.05	1.05	1.05	1.05	1.05	1.05	1.05
$\langle {}^7\text{Be} {}^8\text{B} \rangle$	$\langle {}^7\text{Li} {}^8\text{Li} \rangle$	1/2	1.18	1.17	1.18	1.20	1.17	1.17	1.19	1.12
		3/2	1.20	1.20	1.20	1.22	1.19	1.20	1.22	1.12
$\langle {}^{11}\text{C} {}^{12}\text{N} \rangle$	$\langle {}^{11}\text{B} {}^{12}\text{B} \rangle$	1/2	1.45	1.42	1.44	1.45	1.42	1.42	1.43	1.37
		3/2	1.45	1.44	1.46	1.46	1.42	1.43	1.46	1.37
$\langle {}^{12}\text{C} {}^{13}\text{N} \rangle$	$\langle {}^{12}\text{C} {}^{13}\text{C} \rangle$	1/2	1.26	1.24	1.24	1.26	1.23	1.24	1.25	1.19
$\langle {}^{14}\text{N} {}^{15}\text{O} \rangle$	$\langle {}^{14}\text{N} {}^{15}\text{N} \rangle$	1/2	1.53	1.50	1.51	1.53	1.51	1.51	1.52	1.48
		3/2	1.54	1.51	1.51	1.57	1.51	1.52	1.56	1.48
$\langle {}^{15}\text{N} {}^{16}\text{O} \rangle$	$\langle {}^{15}\text{O} {}^{16}\text{O} \rangle$	1/2	1.55	1.54	1.54	1.57	1.54	1.55	1.56	1.52

TABLE II. Squared ratio $(b_{\max}/b_{\min})^2$ of the maximal and minimal values of b , average ratio of squared ANCs $\langle \mathcal{R}_b \rangle$, analytical estimates \mathcal{R}_0 , and experimental ratios \mathcal{R}^{exp} . Where several experimental values of ANCs are available, we take their average. Also shown are proton ϵ_p and neutron ϵ_n separation energies (in MeV), number of nodes n , and orbital momentum l .

Overlap	ϵ_p	Mirror overlap	ϵ_n	nl	$(\frac{b_{\max}}{b_{\min}})^2$	$\langle \mathcal{R}_b \rangle$	\mathcal{R}_0	\mathcal{R}^{exp}	Ref. for C_p^{exp}	Ref. for C_n^{exp}
$\langle {}^7\text{Be} {}^8\text{B} \rangle$	0.137	$\langle {}^7\text{Li} {}^8\text{Li} \rangle$	2.033	$0p$	1.23	1.01 ± 0.01	1.12	1.08 ± 0.15	[19]	[20]
$\langle {}^{11}\text{C} {}^{12}\text{N} \rangle$	0.601	$\langle {}^{11}\text{B} {}^{12}\text{B} \rangle$	3.370	$0p$	1.67	1.30 ± 0.02	1.37	1.28 ± 0.29	[21]	[22]
$\langle {}^{14}\text{N} {}^{15}\text{O}(\frac{3}{2}^+) \rangle$	0.507	$\langle {}^{14}\text{N} {}^{15}\text{N}(\frac{3}{2}^+) \rangle$	3.026	$1s$	1.68	3.62 ± 0.03	4.09			
$\langle {}^{15}\text{N} {}^{16}\text{O} \rangle$	12.128	$\langle {}^{15}\text{O} {}^{16}\text{O} \rangle$	15.664	$0p$	2.55	1.55 ± 0.02	1.52			
$\langle {}^{16}\text{O} {}^{17}\text{F}(\frac{5}{2}^+) \rangle$	0.601	$\langle {}^{16}\text{O} {}^{17}\text{O}(\frac{5}{2}^+) \rangle$	4.144	$0d$	2.15	1.21 ± 0.03	1.21	1.33 ± 0.20	[23–26]	[27]
$\langle {}^{16}\text{O} {}^{17}\text{F}(\frac{1}{2}^+) \rangle$	0.106	$\langle {}^{16}\text{O} {}^{17}\text{O}(\frac{1}{2}^+) \rangle$	3.273	$1s$	1.56	702 ± 4	796			
$\langle {}^{22}\text{Mg} {}^{23}\text{Al} \rangle$	0.123	$\langle {}^{22}\text{Ne} {}^{23}\text{Ne} \rangle$	4.419	$0d$	1.50	2.67×10^4	2.61×10^4			
$\langle {}^{26}\text{Si} {}^{27}\text{P} \rangle$	0.859	$\langle {}^{26}\text{Mg} {}^{27}\text{Mg} \rangle$	6.443	$1s$	1.80	40.3 ± 1.1	43.3			

single-particle wave functions in the interior and the nuclear single-particle potentials are the same for p and n . The ratio $\mathcal{R}_b = (b_p/b_n)^2$ is therefore expected to have only weak dependence on these potentials. We have verified this for a range of potentials chosen to simultaneously reproduce fixed proton and neutron binding energies. The individual ANCs b_n and b_p vary by up to a factor of 2 but the ratio \mathcal{R}_b is stable to within 3% with an average which agrees with Eq. (7) [18]. If we assume that the spectroscopic factors S_p and S_n are equal for mirror pairs, then we have an alternative way of estimating \mathcal{R} . Note, however, that our derivation of Eq. (7) involves fewer assumptions than in this alternative approach and in fact does not appeal to the concept of spectroscopic factor at all. Our approach is therefore much more general and provides a basis for further refinement of the value of the ratio \mathcal{R} predicted by theory. For the mirror pairs ${}^8\text{B}$ – ${}^8\text{Li}$ and ${}^{12}\text{N}$ – ${}^{12}\text{B}$, and ${}^{17}\text{F}$ – ${}^{17}\text{O}$, where the experimental values of the proton C_p^{exp} and neutron C_n^{exp} ANCs are simultaneously available, both $\langle \mathcal{R}_b \rangle$ and \mathcal{R}_0 agree with $\mathcal{R}^{\text{exp}} = |C_p^{\text{exp}}/C_n^{\text{exp}}|^2$ within the error bars (see Table II).

Near the edge of stability, where neutron separation energies become very small, the corresponding mirror proton states manifest themselves as resonances. The width Γ_p of a narrow proton resonance is related to the ANC of the Gamow wave function for this resonance by the equation $\Gamma_p = \mu/\kappa_p |C_p|^2$ [28]. The ANC C_p can be calculated from Eqs. (2) and (3) using the regular Coulomb function $F_l(\kappa_p r)$ of a real argument [4].

Therefore, a link must exist between Γ_p and the ANC of mirror neutron bound states. The ratio $\mathcal{R}_\Gamma = \Gamma_p/|C_n|^2$ is then approximated by an equation similar to Eq. (7):

$$\mathcal{R}_\Gamma \approx \mathcal{R}_0^{\text{res}} = \frac{\kappa_p}{\mu} \left| \frac{F_l(\kappa_p R_N)}{\kappa_p R_N j_l(i\kappa_n R_N)} \right|^2. \quad (8)$$

Alternatively, \mathcal{R}_Γ can be approximated by the single-particle ratio $\mathcal{R}_\Gamma^{\text{s.p.}} = \Gamma_p^{\text{s.p.}}/b_n^2$ if the spectroscopic factors and single-particle potential wells for mirror bound-unbound pairs are assumed equal. We have calculated $\mathcal{R}_\Gamma^{\text{s.p.}}$ for the ${}^8\text{B}(1^+)$, ${}^{12}\text{N}(2^+)$, ${}^{13}\text{N}(\frac{1}{2}^+)$, and ${}^{13}\text{N}(\frac{5}{2}^+)$ resonances using a set of two-body Woods-Saxon potentials which reproduce both the separation energy of the loosely bound neutron and the position of the mirror proton resonance. In the case of $l \neq 0$, for different choice of the two-body potentials the ratios $\mathcal{R}_\Gamma^{\text{s.p.}}$ change by about 3% while $\Gamma_p^{\text{s.p.}}$ changes by up to a factor of 2 (see Table III). This is the same as in the case of bound mirror pairs of overlaps. However, for $l = 0$, where the centrifugal barrier is absent and nonresonant contributions are larger, the change in $\mathcal{R}_\Gamma^{\text{s.p.}}$ is larger and reaches 11%. The average value of $\mathcal{R}_\Gamma^{\text{s.p.}}$ agrees with $\mathcal{R}_0^{\text{res}}$ for the $l = 2$ resonance ${}^{13}\text{N}(\frac{5}{2}^+)$ but is smaller than $\mathcal{R}_0^{\text{res}}$ by 16%, 20%, and 37% for ${}^8\text{B}(1^+)$, ${}^{12}\text{N}(2^+)$, and ${}^{13}\text{N}(\frac{1}{2}^+)$, respectively. The $\mathcal{R}_0^{\text{res}}$ values themselves are quite stable with respect to different choice of R_N except in the case of the $l = 0$ resonance ${}^{13}\text{N}(\frac{1}{2}^+)$, where the uncertainty of $\mathcal{R}_0^{\text{res}}$ is 5%.

TABLE III. The ratio $\Gamma_p^{\text{max}}/\Gamma_p^{\text{min}}$ of the maximal and minimal proton widths, average ratio of $\mathcal{R}_\Gamma^{\text{s.p.}}$, analytical estimates $\mathcal{R}_0^{\text{res}}$, and experimental ratios $\mathcal{R}_\Gamma^{\text{exp}}$. Where several experimental values of ANCs are available, we take their average.

Proton resonance	Bound mirror analog	l	$\Gamma_p^{\text{max}}/\Gamma_p^{\text{min}}$	$\langle \mathcal{R}_\Gamma^{\text{s.p.}} \rangle$	$\mathcal{R}_0^{\text{res}}$	$\mathcal{R}_\Gamma^{\text{exp}}$	Ref. for Γ_p^{exp}	Ref. for C_n^{exp}
${}^8\text{B}(1^+, 0.774)$	${}^8\text{Li}(1^+, 0.980)$	1	1.43	$(1.70 \pm 0.03) \times 10^{-3}$	2.03×10^{-3}	$(2.29 \pm 0.40) \times 10^{-3}$	[29]	[20]
${}^{12}\text{N}(2^+, 0.960)$	${}^{12}\text{B}(2^+, 0.953)$	1	1.61	$(1.22 \pm 0.01) \times 10^{-5}$	1.42×10^{-5}			
${}^{13}\text{N}(\frac{1}{2}^+, 2.36)$	${}^{13}\text{C}(\frac{1}{2}^+, 3.09)$	0	1.55	$(5.98 \pm 0.32) \times 10^{-5}$	8.5×10^{-5}	$(4.57 \pm 0.57) \times 10^{-5}$	[30]	[22,31]
${}^{13}\text{N}(\frac{5}{2}^+, 3.55)$	${}^{13}\text{C}(\frac{5}{2}^+, 3.85)$	2	2.01	$(1.37 \pm 0.03) \times 10^{-2}$	1.42×10^{-2}	$(1.06 \pm 0.21) \times 10^{-2}$	[30]	[22]

The proton widths of ${}^8\text{B}(1^+)$, ${}^{13}\text{N}(\frac{1}{2}^+)$, and ${}^{13}\text{N}(\frac{5}{2}^+)$, and neutron ANCs for their mirror states, are known experimentally. The ratios $\mathcal{R}_\Gamma^{\text{exp}} = \Gamma_p^{\text{exp}}/|C_n^{\text{exp}}|^2$ for these states are shown in Table III. In all these cases, the single-particle approximation $\mathcal{R}_\Gamma \approx \mathcal{R}_\Gamma^{\text{s.p.}}$ is not confirmed. For ${}^8\text{B}(1^+)$, $\mathcal{R}_\Gamma^{\text{exp}}$ is larger than $\mathcal{R}_\Gamma^{\text{s.p.}}$ and agrees with $\mathcal{R}_0^{\text{res}}$ but, for ${}^{13}\text{N}(\frac{1}{2}^+)$ and ${}^{13}\text{N}(\frac{5}{2}^+)$, $\mathcal{R}_\Gamma^{\text{exp}}$ is significantly lower than $\mathcal{R}_\Gamma^{\text{s.p.}}$ and $\mathcal{R}_0^{\text{res}}$. This result suggests that estimates based on the relation $\Gamma_p = S_p \Gamma_p^{\text{s.p.}}$ and the assumption $S_p = S_n$ can be unreliable.

The present work confirms the existence of a link between proton and neutron mirror ANCs both for bound-bound and bound-unbound mirror pairs. Therefore, neutron ANCs obtained with stable beams can be used to predict cross sections of low-energy direct and resonance proton capture reactions. Although more accurate theoretical ratios for \mathcal{R} and \mathcal{R}_Γ are required for these purposes, the estimates $\langle \mathcal{R}_b \rangle$, \mathcal{R}_0 , $\mathcal{R}_\Gamma^{\text{s.p.}}$, and $\mathcal{R}_0^{\text{res}}$ of the present paper can already be used in some cases. In fact, the ratio \mathcal{R} has already been used to predict the direct ${}^{11}\text{C}(p, \gamma){}^{12}\text{N}$ capture cross sections in [6] and the results obtained there are in a good agreement with the predictions based on proton ANCs recently measured in [21]. Also, the astrophysical S factor for the ${}^7\text{Be}(p, \gamma){}^8\text{B}$ reaction has been calculated in [20] based on the $\langle \mathcal{R}_b \rangle$ estimate and experimentally measured neutron ANC in ${}^8\text{Li}$. Another example is the proton width of the ${}^{12}\text{N}(2^+)$ resonance for which only an upper limit of 20 keV is available. Using the neutron ANC for the mirror ${}^{12}\text{B}(2^+)$ state from [22], we can predict that Γ_p is equal to 5.9 ± 1.0 or 6.9 ± 1.2 keV for the $\mathcal{R}_\Gamma \approx \langle \mathcal{R}_\Gamma^{\text{s.p.}} \rangle$ and $\mathcal{R}_\Gamma \approx \mathcal{R}_0^{\text{res}}$ assumptions, respectively. These values are less uncertain than the currently available experimental limit $\Gamma_p < 20$ keV.

Among other cases of astrophysical interest is the astrophysical S factor for the direct capture reaction ${}^{14}\text{N}(p, \gamma){}^{15}\text{O}(\frac{3}{2}^+)$, which is mainly responsible for the energy production in the CNO cycle. The ${}^{15}\text{O}(\frac{3}{2}^+)$ state is separated from the neighboring ${}^{15}\text{O}(\frac{5}{2}^+)$ state by only 70 KeV, which influences the precision of measurements involving this state. The spacing between the mirror ${}^{15}\text{N}(\frac{3}{2}^+)$ and ${}^{15}\text{N}(\frac{5}{2}^+)$ states is larger and therefore the ANC for the $\langle {}^{14}\text{N} | {}^{15}\text{N}(\frac{3}{2}^+) \rangle$ overlap integral can be determined using neutron transfer reactions to higher accuracy than the ${}^{15}\text{O}(\frac{3}{2}^+)$ ANC. Also, direct contributions to the cross sections of the ${}^{22}\text{Mg}(p, \gamma){}^{23}\text{Al}$ and ${}^{26}\text{Si}(p, \gamma){}^{27}\text{P}$ reactions, involving proton-rich radioactive nuclei, could be calculated through the mirror neutron ANCs which can be determined using stable targets ${}^{22}\text{Ne}$ and ${}^{26}\text{Mg}$. These reaction are relevant to the nucleosynthesis in novae and are being intensively investigated.

This work has been supported by the U.K. EPSRC Grant No. GR/M/82141, the U.S. Department of Energy under Grant No. DE-FG03-93ER40773 and the U.S.

National Science Foundation under Grant No. PHY-0140343. N.T. is grateful to B.V. Danilin for some useful discussions.

-
- [1] H. M. Xu *et al.*, Phys. Rev. Lett. **73**, 2027 (1994).
 - [2] R. F. Casten and B. M. Sherill, Prog. Part. Nucl. Phys. **45**, S171 (2000).
 - [3] S. Kubono, Prog. Theor. Phys. Suppl. **146**, 237 (2002).
 - [4] L. D. Blokhintsev, I. Borbely, and E. I. Dolinskii, Sov. J. Part. Nucl. **8**, 485 (1977).
 - [5] N. K. Timofeyuk, Nucl. Phys. **A632**, 38 (1998).
 - [6] N. K. Timofeyuk and S. B. Igamov, Nucl. Phys. **A713**, 217 (2003).
 - [7] We believe that charge symmetry rather than full charge independence is involved because mirror nuclei have the same number of $n-p$ pairs.
 - [8] A. M. Mukhamedzhanov and N. K. Timofeyuk, Sov. J. Nucl. Phys. **51**, 431 (1990).
 - [9] The oscillator shape of the single-particle wave functions makes the correct treatment of the center of mass possible, which is crucial for the nuclei considered here.
 - [10] A. N. Boyarkina, *Structure of p-Shell Nuclei* (Moscow State University, Moscow, 1973).
 - [11] The more consistent approach used in [5], in which the same NN potential is used both in Eq. (2) and in the shell-model Hamiltonians, leads to similar results for the ratio of the mirror ANCs.
 - [12] D. Gogny, P. Pires, and R. De Tourreil, Phys. Lett. **32B**, 591 (1970).
 - [13] A. B. Volkov, Nucl. Phys. **74**, 33 (1965).
 - [14] D. M. Brink and E. Boeker, Nucl. Phys. **A91**, 1 (1967).
 - [15] G. Bertsch *et al.*, Nucl. Phys. **A284**, 399 (1977).
 - [16] N. Anantaraman, H. Toki, and G. F. Bertsch, Nucl. Phys. **A398**, 269 (1983).
 - [17] We have found a mistake in the computer code for proton ANCs used in Refs. [5,6]. The corrected values are given in Table I.
 - [18] The difference between the average $\langle \mathcal{R}_b \rangle$ and \mathcal{R}_0 is largest for weakly bound states. We will return to the reason for this elsewhere.
 - [19] A. Azhari *et al.*, Phys. Rev. C **63**, 055803 (2001).
 - [20] L. Trache *et al.*, Phys. Rev. C **67**, 062801(R) (2003).
 - [21] X. Tang *et al.*, Phys. Rev. C **67**, 015804 (2003).
 - [22] Z. H. Liu *et al.*, Phys. Rev. C **64**, 034312 (2001).
 - [23] H. T. Fortune *et al.*, Phys. Rev. C **12**, 1723 (1975).
 - [24] J. Verotte *et al.*, Nucl. Phys. **A571**, 1 (1994).
 - [25] S. V. Artemov *et al.*, Phys. At. Nucl. **59**, 428 (1996); Phys. At. Nucl. **63**, 1763 (2000).
 - [26] C. A. Gagliardi *et al.*, Phys. Rev. C **59**, 1149 (1999).
 - [27] S. Burzynski *et al.*, Nucl. Phys. **A399**, 230 (1983).
 - [28] A. M. Mukhamedzhanov and R. E. Tribble, Phys. Rev. C **59**, 3418 (1999).
 - [29] F. Ajzenberg-Selove, Nucl. Phys. **A490**, 1 (1988).
 - [30] F. Ajzenberg-Selove, Nucl. Phys. **A523**, 1 (1991).
 - [31] N. Imai *et al.*, Nucl. Phys. **A688**, 281c (2001).