

Erratum: Atom Movement in In_3La Studied via Nuclear Quadrupole Relaxation [Phys. Rev. Lett. 92, 225901 (2004)]

Matthew O. Zacate, Aurélie Favrot, and Gary S. Collins

(Received 17 June 2004; published 22 July 2004)

DOI: 10.1103/PhysRevLett.93.049903

PACS numbers: 66.30.-h, 76.80.+y, 82.80.Ej, 99.10.Cd

We wish to make an important distinction between the reorientation frequency of the electric field gradient (EFG) and tracer jump frequency. The symbol w was mistakenly used to represent both frequencies in the Letter. The reorientation frequency of the EFG from one orientation to a *single* other orientation, written as w in [1,2], is written as w_{EFG} below. For a jump on the In sublattice in In_3La , there are two possible EFG reorientations. The tracer jump frequency w , defined in Eq. (1) of the Letter, therefore is equal to $2w_{\text{EFG}}$. Results in the Letter need to be amended as follows. The data in Fig. 3 and fitted prefactors are for w_{EFG} instead of w . Tracer jump frequencies w are 2 times larger and the jump frequency extrapolated to room temperature is 800 Hz.

[1] A. Baudry and P. Boyer, *Hyperfine Interact.* **35**, 803 (1987).

[2] W. E. Evenson, J. A. Gardner, R. W. Wang, H.-Y. Su, and A. G. McKale, *Hyperfine Interact.* **62**, 283 (1990).