

## Signature of the Electron-Electron Interaction in the Magnetic-Field Dependence of Nonlinear $I$ - $V$ Characteristics in Mesoscopic Systems

B. Spivak<sup>1</sup> and A. Zyuzin<sup>2</sup>

<sup>1</sup>Physics Department, University of Washington, Seattle, WA 98195, USA

<sup>2</sup>A. F. Ioffe Institute, 194021 St. Petersburg, Russia

(Received 19 April 2004; published 23 November 2004)

We show that the nonlinear  $I$ - $V$  characteristics of mesoscopic samples with metallic conductivity should contain parts which are linear in the magnetic-field and quadratic in the electric field. These contributions to the current are entirely due to the electron-electron interaction and consequently they are proportional to the electron-electron interaction constant. We also note that both the amplitude and the sign of the nonlinear part of the current exhibit random oscillations as a function of temperature.

DOI: 10.1103/PhysRevLett.93.226801

PACS numbers: 73.40.-c, 73.43.-f

According to Onsager, the linear conductance  $G(H)$  of a conductor measured by the two-probe method must be an even function of the magnetic-field  $H$  [1]:

$$G(H) = G(-H). \quad (1)$$

Equation (1) is a consequence of general principles: the time reversal symmetry and the positive sign of the entropy production. Therefore, it holds in all conductors. It is possible, however, that the nonlinear  $I$ - $V$  characteristics of conductors contain parts odd in  $H$ . In particular, one can have contributions to the total current through a sample, which are linear in  $H$  and quadratic in the voltage across the sample  $V$ :

$$I_{(nl)} = \alpha V^2 H. \quad (2)$$

Since  $H$  is an axial vector and the current is a polar one, the coefficient  $\alpha$  can be nonzero only in noncentrosymmetric media. In the case of bulk noncentrosymmetric crystals, terms in  $I$ - $V$  characteristics that are linear in  $H$  have been investigated both theoretically, by using Boltzmann kinetic equation, and experimentally (see, for example, Ref. [2]). In the case of chiral carbon nanotubes, a classical theory of this effect was discussed in Ref. [3].

In this Letter we study this effect at small temperatures and in mesoscopic disordered samples, where all possible symmetries are broken. In this situation, all electron transport effects have quantum interference nature. The theory of nonlinear characteristics of mesoscopic metallic samples was developed in the approximation of noninteracting electrons [4,5]. It is important, however, that in this approximation  $\alpha = 0$  and magnetic-field dependence of the  $I$ - $V$  characteristics is an even function of  $H$ . Therefore, the coefficient  $\alpha$  in Eq. (2) should be proportional to the electron-electron interaction constant  $\beta$ , which is defined by the interacting part of the electron Hamiltonian

$$H_{(int)} = \frac{\beta}{\nu} \int d\mathbf{r} \Psi(\mathbf{r}) \Psi^*(\mathbf{r}) \Psi(\mathbf{r}) \Psi^*(\mathbf{r}). \quad (3)$$

Here  $\nu$  is the electron density of states. Thus, in principle, by measuring the current in Eq. (2) one can measure the electron-electron interaction constant  $\beta$ .

Let us consider a two-dimensional sample shown in the insert of Fig. 1 and assume that the magnetic field is perpendicular to the plane and that the characteristic size of the sample  $L \gg l$  is much larger than the electron elastic mean free path  $l$ . At low temperatures the main contributions to both mesoscopic fluctuations of the conductance  $\delta G = G - \langle G \rangle$  and the nonlinear current Eq. (2)




FIG. 1. Solid lines correspond to electron Green's functions; thin dashed lines correspond to the correlation function of the random scattering potential  $\langle u(\mathbf{r})u(\mathbf{r}') \rangle$ ; thick dashed lines correspond to the electron-electron interaction  $\beta/\nu$ . Symbols  $E, J, n$  correspond to electric field, current density, and electron density, respectively. Diagrams (a) and (b) describe the correlation function  $\langle G(H)G(0) \rangle$ . Parallel and antiparallel directions of arrows in the ladder parts of these diagrams correspond to the Cooperon and Diffusons, respectively. The diagram (c) describes  $\langle I_{(nl)}^2 \rangle$  in the case when  $\beta = 0$ . The diagram (d) describes Eqs. (4), (12), and (13). The insert shows a schematic picture of the sample.

are due to electron interference effects. As usual in such situations,  $\alpha$  is a random sample-specific quantity with zero average  $\langle \alpha \rangle = 0$ . To characterize  $\alpha$  one has to calculate the variance  $\langle \alpha^2 \rangle$ . Here the brackets  $\langle \rangle$  denote averaging over realizations of a random white noise scattering potential characterized by a correlation function  $\langle u(\mathbf{r})u(\mathbf{r}') \rangle = \frac{\pi}{im^2} \delta(\mathbf{r} - \mathbf{r}')$ , where  $m$  is the electron mass.

A diagrammatic verification of Eq. (1) in the situation when interference corrections to conductance are significant is not entirely trivial. One has to show that the correlation function of conductances  $\langle \delta G(H) \delta G(0) \rangle$  is an even function of the magnetic field. This involves calculation of diagrams shown in Fig. 1(a) and 1(b) [6]. We will use a standard diagrammatic technique for averaging over random realizations of the scattering potential [7]. The correlation function of “Diffuson” and “Cooperon” propagators are represented by the ladder diagrams in Fig. 1(a) and 1(b). The Diffuson is the ladder part of the diagram in Fig. 1(a) where the direction of arrows of the electron Green function is antiparallel, while the Cooperon is the ladder in Fig. 1(b) with parallel directions of the arrows. They contain parts linear in the magnetic field, which are equal in magnitude and of different signs. Thus these contributions to the total current through the sample cancel. To verify that  $\alpha = 0$  in the approximation of noninteracting electrons, one has to calculate diagrams for  $\langle (I_{nl})^2 \rangle$  shown in Fig. 1(c). In this case, Diffuson and Cooperon contributions linear in  $H$  cancel each other as well. The fact that  $\alpha = 0$  when  $\beta = 0$  is also quite obvious in the framework of the Landauer scheme of the conductance calculation.

In the approximation linear in  $\beta$  the variance

$$\langle \alpha^2 \rangle = \beta^2 \frac{e^2}{\nu^2 \Gamma^4 A^2} \left( \frac{e^2}{h} \right)^2 \left( \frac{L^2}{\Phi_0} \right)^2 \quad (4)$$

is given by diagrams shown in Fig. 1(d). Here  $\Gamma = \hbar D/L^2$  and  $D/L^2$  is the inverse lifetime of an electron in the sample,  $A$  is the area of the sample, and  $D = \nu_F l/2$  is the electron diffusion coefficient. Equation (4) is valid when  $eV \ll \Gamma$  and  $\Phi \ll \Phi_0$ , where  $\Phi = HA$  and  $\Phi_0$  are the magnetic flux through the sample area and the flux quantum, respectively. For simplicity we consider the short range  $e$ - $e$  interaction described by Eq. (3). In this case the Hartree term is two times larger than the exchange term and we consider only the diagram shown in Fig. 1(d).

In the case of the Coulomb interaction between electrons at high electron densities we have

$$\beta = e^2 r_D \nu, \quad (5)$$

where  $r_D$  is the electron screening radius. To get Eq. (5) one has to calculate diagrams shown in Fig. 61 in Ref. [7].

The effect described by Eq. (4) is quite different from conventional effects in bulk crystals which can be described by the Boltzmann kinetic equation [2,3]. The latter effects are determined by relaxation processes in

materials with complicated band structures and for this reason they are proportional to the relaxation rates (or proportional to  $\beta^2$ ), while Eq. (2) is proportional to  $\beta$ .

The qualitative explanation of Eq. (4) is the following. The mesoscopic fluctuations of the current density inside the sample are due to random interference of electron waves traveling along different diffusive paths. Though the total current through the sample should be an even function of  $H$ , the local current densities contain a part proportional to  $H$ . For example, the part of the current density proportional to  $H$  can be characterized as a “Hall current density.” To avoid confusion, we would like to mention that this “Hall component” is connected with the electric field in a highly nonlocal way and has a random direction. By the same token, in random system there is a part of electron density

$$\delta n(\mathbf{r}) \sim VH, \quad (6)$$

which is proportional to  $H$  and  $V$  [8]. Existence of fluctuations of the density of the form Eq. (6) is a consequence of the fact that contributions linear in  $H$  of diagrams Fig. 1(a) and 1(b) to  $\delta n(\mathbf{r})$  do not cancel. For our estimate it is enough to consider only Diffuson contribution.

We note also that density fluctuations Eq. (6) are different from Friedel oscillations in disordered samples, which are an even function of  $H$ . In the Hartree approximation there is an additional scattering potential

$$\delta u_e(\mathbf{r}, V, H) = \frac{\beta}{2\nu} \delta n(\mathbf{r}, V, H) \quad (7)$$

associated with the fluctuations of the electron density. Thus we can write an expression for a nonlinear part of the total current through the sample in the form

$$I_{nl} = G_{(nl)}[V, T, H, \{\delta u_e(\mathbf{r}, V, H)\}]V, \quad (8)$$

where the nonlinear conductance  $G_{(nl)}$ , generally speaking, depends on the realization of  $u_e(\mathbf{r})$ . It can be regarded as linear conductance at given  $\delta u_e(\mathbf{r})$ .

The sensitivity of the sample conductance to a change in the scattering potential  $\delta u_e(\mathbf{r})$  has been considered in [9,10]. Generally speaking, the mesoscopic part of the conductance  $G_{(nl)} - \langle G_{(nl)} \rangle$  depends on all spatial harmonics of  $\delta u_e(\mathbf{r})$ . However, the main contribution to the change of the conductance comes from zero harmonics of the potential

$$\delta \bar{u}_e(V, H) = \frac{\beta}{\nu A} \int \delta n(\mathbf{r}, V, H) d\mathbf{r}. \quad (9)$$

This can be verified by making calculations similar to those in [11]. This is also related to the long range character of the correlation function of the part of the electron densities, which are proportional to  $V$  [8]. A  $\delta u_e(\mathbf{r})$  dependence of  $\langle G_{(nl)} \rangle$  is due to the energy dependence of

the electron mean free time. This dependence is small and can be neglected.

Expanding Eq. (8) in terms of  $\bar{u}_e$  and taking into account that in the main approximation  $\bar{u}_e$  and  $G_{(nl)}$  are uncorrelated, we get

$$\langle I_{(nl)}^2 \rangle = \langle (dG_{(nl)}/d\bar{u}_e)^2 \rangle \langle \bar{u}_e^2 \rangle V^2. \quad (10)$$

According to [10]  $\langle (dG_{(nl)}/d\bar{u}_e)^2 \rangle = (e^2/h)^2/\Gamma^2$ . Calculating the correlation function

$$\langle [\delta\bar{u}_e(H, V)]^2 \rangle = \frac{\beta^2}{\nu^2} \frac{1}{A^2} \frac{\Phi^2}{\Phi_0} \frac{|eV|^2}{\Gamma} \quad (11)$$

we arrive at Eq. (4).

At this point we would like to mention that on a qualitative level the effect considered above can be also described in the framework of the Landauer scheme. To do so one has to combine results of [12,13].

Equation (4) is valid at small temperatures  $\Gamma \gg T$ . At finite temperature, the quantity  $\langle \alpha^2(T) \rangle$  decreases with  $T$ . At  $T \gg \Gamma$ ,

$$\langle \alpha^2(T) \rangle \sim \langle \alpha^2(0) \rangle \frac{\Gamma^2}{T^2}. \quad (12)$$

We stress that the temperature dependence of  $\alpha(T)$  is nonmonotonic:  $\alpha(T)$  exhibits random oscillations in magnitude and sign, superimposed on the average decay. One can see this by calculating the quantity

$$\langle \alpha(T)\alpha(0) \rangle \sim \langle \alpha^2(0) \rangle \frac{\Gamma^2}{T^2}. \quad (13)$$

Note that Eqs. (12) and (13) have the same temperature dependence, which is impossible without oscillations of the sign of  $\alpha(T)$  [14]. In case of monotonic  $\alpha(T)$  dependence, one would expect that  $\langle \alpha(T)\alpha(0) \rangle \sim 1/T$ .

In the case of high magnetic-field  $\Phi > \Phi_0$ , (but still  $eV \ll \Gamma$ ), the part of the current which is asymmetric in  $H$  and quadratic in  $V$  exhibits random oscillations as a function of  $\Phi$ . These oscillations, typical for mesoscopic systems, have a characteristic period  $\Phi_0$  and the amplitude

$$\langle [I(H) - I(-H)]^2 \rangle = \beta^2 \left( \frac{eV}{\Gamma} \right)^2 \frac{1}{(\nu\Gamma A)^2} \left( \frac{e^2 V}{h} \right)^2. \quad (14)$$

Finally, we mention that there are no mechanisms contributing to Eq. (2) other than the mechanism considered above. For example, at finite  $V$  there is a new channel of electron transmission through the sample when an incident electron is transmitted into two electrons and a hole. The probability of such a process has a component which is linear in  $H$ . Its magnitude can be estimated in a way similar to estimating the electron-electron scattering rate of quasiparticles in a uniform Fermi liquid. As a result, this component is proportional to  $V^2$ . Thus the

magnitude of the asymmetric-in- $H$  part of the current associated with such process is proportional to  $V^3 H$ .

The effect discussed above was recently observed experimentally in GaAs quantum dots [15]. During the preparation of the manuscript we became aware of similar unpublished results by Sanchez and Büttiker [16].

This work was supported in part by the National Science Foundation under Contract No. DMR-0228104. We thank M. Büttiker, C. M. Marcus, and D. M. Zumbuhl, for useful discussions.

- 
- [1] L. D. Landau and E. M. Lifshitz, *Statistical Physics* (Butterworth, Washington, DC, 1980).
  - [2] B. I. Sturman and V. M. Fridkin, *The Photovoltaic and Photorefractive Effects in Non-Centrosymmetric Materials* (Gordon and Breach, New York, 1992); G. A. Rikken, J. Fölling, and P. Wyder, *Phys. Rev. Lett.* **87**, 236602 (2002).
  - [3] E. Ivchenko and B. Spivak, *Phys. Rev. B* **66**, 155404 (2002).
  - [4] A. Larkin and D. Khmel'nitskii, *JETP* **91**, 1815 (1986); *Phys. Scr.* **T14**, 4 (1986).
  - [5] V. Falko and D. Khmel'nitskii, *Sov. Phys. JETP* **68**, 186 (1989).
  - [6] B. L. Altshuler and D. E. Khmel'nitskii, *JETP Lett.* **42**, 359 (1985).
  - [7] A. A. Abrikosov, L. P. Gorkov, and I. Dzyaloshinski, *Methods of Quantum Field Theory in Statistical Physics* (Dover, New York, 1975).
  - [8] A. Zyuzin and B. Spivak, *Sov. Phys. JETP* **66**, 560 (1987); A. Yu. Zyuzin and B. Spivak, *Mesoscopic Fluctuations of Current Density in Disordered Conductors*, in *Mesoscopic Phenomena in Solids*, edited by B. Altshuler, P. Lee, and R. Webb (Elsevier Science, Holland, 1991), Vol. 30.
  - [9] P. A. Lee and A. D. Stone, *Phys. Rev. Lett.* **55**, 1622 (1985).
  - [10] B. Altshuler and B. Spivak, *JETP Lett.* **42**, 447 (1986).
  - [11] B. Spivak and A. Zyuzin, *J. Opt. Soc. Am. B* **21**, 177 (2004).
  - [12] M. Büttiker, *J. Phys. Condens. Matter* **5**, 9361 (1993).
  - [13] M. Büttiker, *Europhys. Lett.* **35**, 523 (1996).
  - [14] We note that a similar phenomenon exists in the case of mesoscopic oscillations of linear conductance. Namely, the quantity  $\Delta G = [G(H, T) - G(0, T)]$  exhibits random oscillations both in magnitude and in sign as a function of  $T$ . One can see this by calculating the variances  $\langle [\Delta G(T)]^2 \rangle \sim \langle \Delta G(T)\Delta G(0) \rangle \sim 1/T$ . Both quantities have the same  $T$  dependence, which is impossible if sign  $\Delta G(T)$  does not oscillate as a function of  $T$ . As far as we know, this fact has not been discussed in the literature.
  - [15] D. M. Zumbuhl, C. M. Marcus, M. P. Hanson, and A. C. Gossard (to be published).
  - [16] D. Sanchez and M. Büttiker (to be published).