

Gravitational-Wave Emission from Rotating Gravitational Collapse in Three Dimensions

L. Baiotti,¹ I. Hawke,^{2,4} L. Rezzolla,^{1,3} and E. Schnetter²

¹*SISSA, International School for Advanced Studies and INFN, Via Beirut 2, 34014 Trieste, Italy*

²*Max-Planck-Institut für Gravitationsphysik, Albert-Einstein-Institut, 14476 Golm, Germany*

³*Department of Physics, Louisiana State University, Baton Rouge, Louisiana 70803 USA*

⁴*School of Mathematics, University of Southampton, Southampton SO17 1BJ, United Kingdom*

(Received 12 October 2004; published 8 April 2005)

We present the first three-dimensional (3D) calculations of the gravitational-wave emission in the collapse of uniformly rotating stars to black holes. The initial models are polytropes which are dynamically unstable and near the mass-shedding limit. The waveforms have been extracted using a gauge-invariant approach and reflect the properties of both the initial stellar models and of newly produced black holes, being in good qualitative agreement with those computed in previous 2D simulations. The wave amplitudes, however, are about 1 order of magnitude smaller, giving, for a source at 10 kpc, a signal-to-noise ratio $S/N \sim 0.25$ for LIGO-VIRGO and $S/N \lesssim 4$ for LIGO II.

DOI: 10.1103/PhysRevLett.94.131101

PACS numbers: 04.30.Db, 04.25.Dm, 04.70.Bw, 97.60.Jd

The study of the gravitational collapse of rotating stars to black holes is a cornerstone of any theory of gravity and a long-standing problem in general relativity. Over the years, numerous approaches have been applied and several different techniques developed to tackle this problem which is not just academic. Indeed, important issues in relativistic astrophysics awaiting clarification, such as the mechanism responsible for γ -ray bursts, may be unveiled with a more detailed understanding of the physics of gravitational collapse in rotating and magnetized stars. Furthermore, the study of gravitational collapse will provide the waveforms and the energetics of one of the most important sources of gravitational radiation.

In the absence of symmetries, no analytic solutions are known that describe gravitational collapse to a black hole, and it is through numerical relativity simulations that one can hope to improve our knowledge of this process under realistic conditions. However, the modeling of black hole spacetimes with collapsing matter sources in multidimensions is one of the most formidable efforts of numerical relativity. This is due both to the inherent difficulties and complexities of the system of equations to be solved and to the vast computational resources needed in three-dimensional evolutions.

In addition to the technical difficulties due to the accurate treatment of the hydrodynamics involved in the collapse, the precise calculation of the gravitational radiation emitted in the process is particularly challenging as the energy released in gravitational waves is much smaller than the total rest-mass energy of the system. Indications of the difficulties inherent to the problem of calculating the gravitational-wave emission in rotating gravitational collapse have emerged in the first and only work, which dates back almost 20 years [1]. In 1985, in a landmark work in numerical relativity, Stark and Piran used their axisymmetric general-relativistic code to evolve rotating configurations and to compute the gravitational radiation produced

by their collapse to black holes. The results referred to initial configurations consisting of polytropic stars which underwent collapse after the pressure was reduced by a factor ranging from 60% up to 99% for the rapidly rotating models. The initial data effectively consisted of spherically symmetric solutions with a uniform rotation simply “added” on. Not being stationary solutions of the Einstein equations, these stars could reach dimensionless spins up to $a \equiv J/M^2 = 0.94$, with J and M the angular momentum and mass of the star, respectively.

Overall, their investigation revealed that while the nature of the collapse depended on the parameter a , the form of the waves remained roughly the same over the entire range of the values of a , with the amplitude increasing with a . Particularly important was evidence that, despite the complex matter dynamics during the collapse, the gravitational-wave emission could essentially be related to the oscillations of a perturbed black hole spacetime.

In recent years many studies have extended to three spatial dimensions the investigation of gravitational collapse to black holes [2–4]. Despite the improvements in the evolution of the hydrodynamics and of the spacetime achieved by these simulations, none of them has addressed the problem of the gravitational-wave emission. The reason for this can be found in the small amplitude of the signal, often below the truncation error of the 3D simulations, but most importantly in the fact that all of the above calculations made use of Cartesian grids with uniform spacing. With the computational resources currently available, this choice places the outer boundaries too close to the source to detect gravitational radiation.

Using a recently developed code for the solution of the Einstein equations in nonvacuum spacetimes, the WHISKY code, we have investigated the collapse of rapidly rotating relativistic stars to Kerr black holes [5]. An important aspect of these simulations is the detailed study of the geometrical and dynamical properties of both the *apparent*

and *event* horizons, allowing for the determination of both the mass and angular momentum of the black hole with an accuracy unprecedented for a 3D simulation. In turn, these measures have set upper limits on the energy and angular momentum lost during the collapse in the form of gravitational radiation, the first such estimates coming from 3D calculations.

However, as in previous works (e.g., Refs. [2–4]), the simulations reported in [5] made use of numerical grids with uniform spacing and thus with outer boundaries very close to the initial position of the stellar surface. With these restrictions, the gravitational radiation extracted does not provide interesting information besides the obvious change in the quadrupole moment of the background spacetime. As we show below, the use of progressive mesh-refinement (PMR) techniques has removed these restrictions, enabling us to place the outer boundaries of the computational domain at very large distances from the collapsing star. This conceptually simple but technologically challenging improvement has two important physical consequences. First, it reduces the influence of inaccurate boundary conditions at the outer boundaries of the domain while retaining the required accuracy in the region where the black hole forms. Second, it allows the wave zone to be included in the computational domain and thus to extract the gravitational waves produced in the collapse.

In practice, we have adopted a Berger-Oliger prescription for the refinement of meshes on different levels [6] and used the numerical infrastructure described in [7]. In addition to this, we have also implemented a simplified form of adaptivity in which new refined levels are added at predefined positions during the evolution. More specifically, given an initial stellar model of mass M and equatorial coordinate radius R , our initial grid consists of four levels of refinement, with the innermost one covering the star with a typical resolution of $\Delta x \sim 0.17M$ and with the outermost having a typical resolution of $\Delta x \sim 1.38M$ and extending up to $\sim 82.5M \approx 20.9R$. As the collapse proceeds and the star occupies smaller portions of the computational domain, three more refined levels are added one by one, nested in the four original ones. By the time the simulation is terminated at $\sim 81.5M$, the finest typical spatial resolution is $\Delta x \sim 0.02M$. A detailed discussion of the grid and of its evolution will be given in [8].

The initial data for our simulations is described in [5] and consists of axisymmetric rotating relativistic stars, calculated as equilibrium solutions of the Einstein equations in a compactified domain and in polar coordinates. For a direct comparison with the results in [1] and because no shocks are observed during the collapse, the stars are modeled with a polytropic equation of state (EOS) $p = K\rho^\Gamma$, with $\Gamma = 2$ and with the polytropic constant which is initially $K_{\text{ID}} = 100$. Once secularly unstable solutions are found along sequences of fixed angular momentum or fixed rest mass, we consider initial models that have the same axis ratios but slightly larger central energy densities and are dynamically unstable. Hereafter we restrict the discus-

sion to the collapse of the most rapidly rotating dynamically unstable model, namely, model D4 in [5], which represents a star of mass $M = 1.861M_\odot$, circumferential equatorial radius $R_e = 14.25$ km and rotating close to the mass-shedding limit with $a = 0.54$ (cf. Table I of [5]). The discussion of the emission from stellar models rotating at smaller velocities will be presented in [8].

Although model D4 is dynamically unstable, it is very close to a stationary solution, and we trigger its collapse by reducing the pressure by $\lesssim 2\%$, with a change in the value of the polytropic constant. After the perturbation is introduced, the constraint equations are again solved to enforce the constraint violation to be at the truncation-error level. As a contrast, in Ref. [1], rapid rotation was added to a spherically symmetric solution and the large pressure depletion inevitably produced a dependence of the results on the amount of pressure reduction. Hereafter we concentrate on the gravitational-wave emission, but a discussion of the dynamics of matter and trapped surfaces can be found in [5].

While several different methods are possible for the extraction of the gravitational-radiation content in numerical spacetimes, we have adopted a gauge-invariant approach in which the spacetime is matched with the nonspherical perturbations of a Schwarzschild black hole (see Refs. [9–11] for applications to Cartesian coordinate grids). In practice, a set of “observers” is placed on 2-spheres of fixed coordinate radius r_{ex} , where they extract the gauge-invariant, odd $Q_{\ell m}^{(o)}$ and even-parity $\Psi_{\ell m}^{(e)}$ metric perturbations [12]. Here ℓ, m are the indices of the angular decomposition, and we usually compute modes up to $\ell = 5$ with $m = 0$; modes with $m \neq 0$ are essentially zero because of the high degree of axisymmetry in the collapse. Validations of this approach in 3D vacuum spacetimes can be found in Refs. [10,13,14], while its use with matter sources has first been reported in [15].

Although the position of such observers is arbitrary and the information they record must be the same for waves extracted in the wave zone, we place our observers between $40M$ and $50M$ from the center of the grid so as to maximize the length of the extracted waveform. We note that, while a similar choice was made in Ref. [1], it still provides only an approximate description of what would be observed at spatial infinity.

Using the odd and even-parity perturbations $Q_{\ell m}^\times = \lambda Q_{\ell m}^{(o)}$ and $Q_{\ell m}^+ = \lambda \Psi_{\ell m}^{(e)}$, where $\lambda \equiv \sqrt{2(\ell+2)!/(\ell-2)!}$, we report in Fig. 1 the lowest-order multipoles for $Q_{\ell m}^\times$ with the offset produced by the stellar quadrupole removed [8]. The left panel, in particular, refers to the $\ell = 2$ mode as extracted by four different observers at increasing distances and expressed in retarded time. The right panel instead refers to the $\ell = 4$ mode, with the inset giving a comparison between the two modes and showing that the gravitational-wave signal is essentially quadrupolar, with the $\ell = 2$ mode being about an order of magnitude larger than the $\ell = 4$ mode.




FIG. 1. The left panel shows the $\ell = 2$, even-parity perturbation as extracted by observers at different positions r_{ex} expressed in retarded time. The right panel refers to the $\ell = 4$ mode, with the inset offering a comparison in the amplitudes of the two modes.

The very good overlap of the waveforms measured at different positions is important evidence that the extraction has been performed in the wave zone, since the invariance under a retarded-time scaling is a property of the solutions of a wave equation. The overlap disappears if the outer boundary is too close or when the waves are extracted at smaller radii. A similar overlap is seen also for the $\ell = 4$ mode (not shown).

Another indication that the waveforms in Fig. 1 are an accurate description of the gravitational radiation produced by the collapse comes by analyzing their power spectra. The collapse can be viewed as the rapid transition between the spacetime of the initial equilibrium star and the spacetime of the produced rotating black hole. It is natural to expect that the waveforms produced in this process will reflect the basic properties of both spacetimes and, in particular, the fundamental frequencies of oscillation. We validate this in Fig. 2, where we show the power spectral densities (PSD) of the waveforms of the metric perturbations Q_{20}^+ and Q_{40}^+ reported in Fig. 1 (the units on the y axis are arbitrary). The upper panel of Fig. 2, in particular, shows the PSD of Q_{20}^+ and compares it with the frequencies of the $\ell = 2, m = 0$ quasinormal mode (QNM) of a Kerr black hole with $M = 1.861M_{\odot}$ and $a = 0.6$ [16] (dashed line at 6.7 kHz) as well as with the first w_{II} “interface” mode for a typical compact star with $M = 1.27M_{\odot}$ and $R = 8.86$ km (dotted line at 8.8 kHz) [17]. Similarly, the lower panel of Fig. 2, shows the PSD of Q_{40}^+ , comparing it with the $\ell = 4, m = 0$ QNM of a Schwarzschild black hole [17] (dashed line at 14.0 kHz) and the first w_1 “curvature” mode [17] (dotted line at 12.8 kHz) [18]. It is apparent that both peaks in the PSDs are rather far from the maximum sensitivity area of modern interferometric and bar detectors.

Although the waveforms have very short duration with very broad PSDs, Fig. 2 shows that these have strong and narrow peaks (a similar behavior can be shown to be present also for the odd-parity modes [8]). Indeed, the excellent agreement between the position of these peaks and the fundamental frequencies of the vacuum and non-

vacuum spacetimes is an important confirmation of the robustness of the results obtained.

Using the extracted gauge-invariant quantities, it is also possible to calculate the transverse traceless (TT) gravitational-wave amplitudes in the two polarizations h_+ and h_{\times} as

$$h_+ - ih_{\times} = \frac{1}{2r} \sum_{\ell, m} \left(Q_{\ell m}^+ - i \int_{-\infty}^t Q_{\ell m}^{\times}(t') dt' \right) {}_{-2}Y^{\ell m}, \quad (1)$$

where ${}_{-2}Y^{\ell m}$ is the $s = -2$ spin-weighted spherical harmonic. Because of the small amplitude of higher-order modes, the TT wave amplitudes can be simply expressed as $h_+ \simeq h_+(Q_{20}^+, Q_{40}^+)$ and $h_{\times} \simeq h_{\times}(Q_{30}^{\times}, Q_{50}^{\times})$, where $Q_{30}^{\times} \gg Q_{50}^{\times}$. Their waveforms are shown in Fig. 3 for the detector at $r_{\text{ex}} = 37.1M$ and for two different inclination angles [8]. Note that the amplitudes in the cross polarization are about 1 order of magnitude smaller than those in




FIG. 2. Power spectra of the waveforms reported in Fig. 1. The dashed vertical lines indicate the frequencies of the QNMs of a black hole, while the dotted ones indicate the w_1 and w_{II} modes of a typical star.




FIG. 3. TT gravitational-wave amplitudes extracted by the detector at $r_{\text{ex}} = 37.1M$ and for two different inclination angles.

the plus polarization, with the maximum amplitudes in a ratio $|(r/M)h_x|_{\text{max}}/|(r/M)h_+|_{\text{max}} \approx 0.06$. Here the odd-parity perturbations, which are zero in spacetimes with axial and equatorial symmetries, are just the result of the coupling, induced by the rotation, with the even-parity perturbations.

A precise comparison of the amplitudes in Fig. 3 with the corresponding ones calculated in [1] is made difficult by the differences in the choice of initial data and, in particular, by the impossibility of reaching $a \gtrsim 0.54$ when modeling consistently stationary polytropes in uniform rotation. However, when interpolating the results in [1] for the relevant values of a , we find a very good agreement in the form of the waves, but also that our estimates are about 1 order of magnitude smaller, with $|(r/M)h_+|_{\text{max}} \approx 0.00225$. Furthermore, we observe the amplitude of the gravitational waves to increase with the pressure reduction, suggesting that the origin of the difference is related mainly to this [8].

Following Ref. [19] and considering the optimal sensitivity of VIRGO for the burst signal only, we set an upper limit for the characteristic amplitude produced in the collapse of a rapidly and uniformly rotating polytropic star at 10 kpc to be $h_c = 5.77 \times 10^{-22}(M/M_\odot)$ at a characteristic frequency $f_c = 931$ Hz. In the case of Laser Interferometer Gravitational Wave Observatory (LIGO) I, instead, we obtain $h_c = 5.46 \times 10^{-22}(M/M_\odot)$ at $f_c = 531$ Hz. In both cases, the signal-to-noise ratio is $S/N \sim 0.25$, but this can grow to be $\lesssim 4$ in the case of LIGO II. These ratios could be increased considerably with the detection of the black hole ringing following the initial burst [8]. Computing the emitted power as

$$\frac{dE}{dt} = \frac{1}{32\pi} \sum_{\ell, m} \left(\left| \frac{dQ_{\ell m}^+}{dt} \right|^2 + |Q_{\ell m}^\times|^2 \right), \quad (2)$$

the total energy lost to gravitational radiation is $E =$

$1.45 \times 10^{-6}(M/M_\odot)$. This is about 2 orders of magnitude smaller than the estimate made in [1] for a star with $a = 0.54$, but larger than the energy losses computed recently in the collapse of rotating stellar cores to proto-neutron stars [20].

In conclusion, we have presented the first waveforms from the gravitational collapse of rapidly rotating stars to black holes using 3D grids with Cartesian coordinates. The great potential shown by the PMR techniques employed here opens the way to a number of applications that would be otherwise intractable with uniform Cartesian grids. Work is in progress to consider initial models with realistic EOSs or in differential rotation, for which values $a \gtrsim 1$ can be reached and more intense gravitational radiation is expected.

We thank V. Ferrari, K. Kokkotas, N. Stergioulas, O. Zanotti for discussions, and C. Ott and B. Zink for help with the PMR algorithm. The computations were performed on *Albert100* at the University of Parma and *Peyote* at the Albert Einstein Institute. E. S. was supported by the DFG's SFB TR/7.

-
- [1] R. F. Stark and T. Piran, Phys. Rev. Lett. **55**, 891 (1985); in *Proceedings of the Workshop Dynamical Spacetimes and Numerical Relativity, Philadelphia* (Cambridge University Press, Cambridge, U.K., 1986), pp. 40–73.
 - [2] M. Shibata, T. W. Baumgarte, and S. L. Shapiro, Phys. Rev. D **61**, 044012 (2000).
 - [3] M. Shibata, Phys. Rev. D **67**, 024033 (2003); *Astrophys. J.* **595**, 992 (2003).
 - [4] M. D. Duez, S. L. Shapiro, and H. J. Yo, Phys. Rev. D **69**, 104016 (2004).
 - [5] L. Baiotti *et al.*, Phys. Rev. D **71**, 024035 (2005).
 - [6] M. J. Berger and J. Olinger, J. Comput. Phys. **53**, 484 (1984).
 - [7] E. Schnetter, S. H. Hawley, and I. Hawke, *Classical Quantum Gravity* **21**, 1465 (2004).
 - [8] L. Baiotti *et al.* (to be published).
 - [9] M. E. Rupright, A. M. Abrahams, and L. Rezzolla, Phys. Rev. D **58**, 044005 (1998).
 - [10] K. Camarda and E. Seidel, Phys. Rev. D **59**, 064019 (1999).
 - [11] G. Allen, K. Camarda, and E. Seidel, gr-qc/9806036.
 - [12] V. Moncrief, Ann. Phys. (N.Y.) **88**, 323 (1974).
 - [13] L. Rezzolla *et al.*, Phys. Rev. D **59**, 064001 (1999).
 - [14] J. Baker *et al.*, Phys. Rev. D **62**, 127701 (2000).
 - [15] J. A. Font *et al.*, Phys. Rev. D **65**, 084024 (2002).
 - [16] E. W. Leaver, Proc. R. Soc. London A **402**, 285 (1985).
 - [17] K. D. Kokkotas and B. G. Schmidt, *Living Rev. Relativity* **2**, 2 (1999).
 - [18] The frequencies for the QNMs of the black hole and for the w modes of the star are those most easily obtained from tabulated values in the literature and serve here as indicative references.
 - [19] K. S. Thorne, in *300 Years of Gravitation*, edited by S. W. Hawking and W. Israel (Cambridge University Press, Cambridge, U.K., 1987).
 - [20] E. Müller *et al.*, *Astrophys. J.* **603**, 221 (2004).