

$s\bar{s}$ Component of the Proton and the Strangeness Magnetic Moment

B. S. Zou*

Institute of High Energy Physics, CAS, P.O. Box 918, Beijing 100049, China

D. O. Riska†

Helsinki Institute of Physics and Department of Physical Sciences, POB 64, 00014 University of Helsinki, Finland

(Received 25 February 2005; published 11 August 2005)

A complete analysis is given of the implications of the empirical indications for a positive strangeness magnetic moment μ_s of the proton on the possible configurations of the $uuds\bar{s}$ component of the proton. A positive value for μ_s is obtained in the $s\bar{s}$ configuration where the $uuds$ subsystem is in an orbitally excited state with $[4]_{FS}[22]_F[22]_S$ flavor-spin symmetry, which is likely to have the lowest energy. The configurations in which the \bar{s} is orbitally excited, which include the conventional $K^+\Lambda^0$ configuration, with the exception of that in which the $uuds$ component has spin 2, yield negative values for μ_s . The hidden strangeness analogues of recently proposed quark cluster models for the θ^+ pentaquark give differing signs for μ_s .

DOI: 10.1103/PhysRevLett.95.072001

PACS numbers: 12.39.-x, 13.40.Em, 14.20.Dh

Three recent experiments on parity violation in electron-proton scattering suggest that the strangeness magnetic moment of the proton μ_s is positive [1–3]. In contrast, theoretical calculations have led to negative values for this observable, with but few exceptions [4–9]. Here the implications of the empirical result for the configuration of the constituents of the proton are considered by a calculation of μ_s for all positive parity configurations of the $uuds\bar{s}$ system with one constituent in the first orbitally excited state, which may be contained in the proton. The results are given in the form of simple general expressions, in which μ_s is proportional to the $s\bar{s}$ probability of the configuration.

It is shown that μ_s is positive in the $uuds\bar{s}$ configuration, which is likely to have the lowest energy, where the \bar{s} quark is in the ground state and the $uuds$ system is in the P state. If, in contrast, the strange antiquark is in the P state and the 4 quarks are in their ground state, the strangeness magnetic moment is negative (except for the energetically unfavored case where the total spin of the $uuds$ state equals 2). These configurations correspond to that of a fluctuation of the proton into a kaon and a strange hyperon, which is well known to lead to a negative value for the strangeness magnetic moment [10–13].

Several intriguing configurations [14–16] for the system of 4 light flavor quarks and strange antiquarks have been proposed to explain the structure of the (tentative) θ^+ pentaquark. Below their hidden strangeness analogues are considered from the point of view of the strangeness magnetic moment. While the diquark-diquark configurations suggested in Refs. [14,15] with the \bar{s} quark in its ground state yield positive values, the diquark-triquark configuration suggested in Ref. [16] yields a negative value for the strangeness magnetic moment of the proton.

If the $uuds$ quarks are in their ground state, the spatial state is completely symmetric $[4]_X$, and the Pauli principle requires that their flavor-spin state has the mixed symmetry $[31]_{FS}$ so as to combine with the color state $[211]_C$ to form the required total antisymmetry $[1111]$. The different flavor and spin state symmetry configurations that can combine to the required $[31]_{FS}$ mixed symmetry combination are listed in Table I [17]. The requirement of positive parity requires that for these configurations the strange antiquark has to be in the P state.

If, in contrast, the strange antiquark \bar{s} is in its ground state, the $uuds$ system has to be in the P state in order that the combination have positive parity. In this case the symmetry of the spatial state of the 4-quark system is reduced to $[31]_X$. When combined with the mixed symmetry color state $[211]_C$, this allows the flavor-spin combination either to be completely symmetric $[4]_{FS}$ or to have any one of the mixed symmetries $[31]_{FS}$, $[22]_{FS}$, or $[211]_{FS}$. The combinations of flavor and spin symmetry configurations that lead to these symmetries are also listed in Table I [17].

In the quark model the strangeness magnetic moment is defined as the matrix element of the operator

$$\vec{\mu}_s = e \sum_i \frac{\hat{S}_i}{2m_s} (\vec{l}_i + \vec{\sigma}_i), \quad (1)$$

where \hat{S} is the strangeness counting operator, with eigenvalue +1 for s and -1 for \bar{s} quarks, and m_s is the constituent mass of the strange quark.

Consider first the configurations, in which the antiquark is in the P state. For these, the spatial state of the $uuds$ is completely symmetric $[4]_X$ with total orbital angular momentum 0. The corresponding wave functions for a proton with spin-up may then be written in the general form

$$\psi = A_{s\bar{s}} \sum_{a,b,c} \sum_{m,s,M,J,j} (1, 1/2, m, s|J, j) \quad (S, J, M, j|1/2, 1/2) C_{[211]_a, [31]_a}^{[1111]} C_{[F]_b, [S]_c}^{[31]_a} [211]_c(a)[F](b)[S]_M(c) \bar{Y}_{1m} \bar{\chi}_s \varphi(\{r_i\}). \quad (2)$$

Here $[211]_C$, $[F]$, and $[S]$ are the color, flavor, and spin states denoted by their Young patterns, respectively, and the sums over a , b , and c run over the configurations of the $[31][F][S]$ representations of S_4 for which the corresponding Clebsch-Gordan $C_{b,c}^a$ are nonvanishing [18]. The orbital and spin states of the \bar{s} are denoted by \bar{Y}_{1m} and $\bar{\chi}_s$, respectively. $A_{s\bar{s}}$ is the amplitude of the $s\bar{s}$ component, and $\varphi(\{r_i\})$ is a symmetric function of the coordinates of the $uuds\bar{s}$ system.

The configurations in the left column in Table I may be grouped according to their spin symmetries $[22]_S$, $[31]_S$, and $[4]_S$. The uppermost configuration in the left column in Table I, which has the spin symmetry $[22]_S$, is expected to have the lowest energy if the $uuds$ is in its ground state. This is the case if the hyperfine interaction between the quarks is described both by the color magnetic interaction or by the flavor and spin dependent hyperfine interaction $-C \sum_{i<j} \vec{\lambda}_i \cdot \vec{\lambda}_j \vec{\sigma}_i \cdot \vec{\sigma}_j$, where C is a constant with the value $\sim 20\text{--}30$ MeV, and which leads to the empirical ordering of the baryon resonances [17,19]. For the states with this spin symmetry the total spin of the $uuds$ system is $S = 0$. The corresponding strangeness magnetic moment is negative:

TABLE I. Flavor and spin state symmetry configurations of the $uuds$ quark states in the ground state and the first orbitally excited P state. The states are ordered from above after increasing matrix elements of the Casimir operator $-\sum_{i<j} \vec{\lambda}_i \cdot \vec{\lambda}_j \vec{\sigma}_i \cdot \vec{\sigma}_j$, where λ_i are the $SU(3)_F$ generators. These matrix elements are listed in the brackets [17].

$uuds$ ground state	$uuds$ P state
$[31]_{FS}[211]_F[22]_S$ (−16)	$[4]_{FS}[22]_F[22]_S$ (−28)
$[31]_{FS}[211]_F[31]_S$ (−40/3)	$[4]_{FS}[31]_F[31]_S$ (−64/3)
$[31]_{FS}[22]_F[31]_S$ (−28/3)	$[31]_{FS}[211]_F[22]_S$ (−16)
$[31]_{FS}[31]_F[22]_S$ (−8)	$[31]_{FS}[211]_F[31]_S$ (−40/3)
$[31]_{FS}[31]_F[31]_S$ (−16/3)	$[31]_{FS}[22]_F[31]_S$ (−28/3)
$[31]_{FS}[31]_F[4]_S$ (0)	$[31]_{FS}[31]_F[22]_S$ (−8)
$[31]_{FS}[4]_F[31]_S$ (+8/3)	$[4]_{FS}[4]_F[4]_S$ (−8)
	$[22]_{FS}[211]_F[31]_S$ (−16/3)
	$[31]_{FS}[31]_F[31]_S$ (−16/3)
	$[22]_{FS}[22]_F[22]_S$ (4)
	$[211]_{FS}[211]_F[22]_S$ (0)
	$[31]_{FS}[31]_F[4]_S$ (0)
	$[211]_{FS}[211]_F[31]_S$ (8/3)
	$[22]_{FS}[31]_F[31]_S$ (8/3)
	$[31]_{FS}[4]_F[31]_S$ (8/3)
	$[22]_{FS}[22]_F[4]_S$ (4)
	$[211]_{FS}[22]_F[31]_S$ (20/3)
	$[211]_{FS}[211]_F[4]_S$ (8)
	$[211]_{FS}[31]_F[22]_S$ (8)
	$[22]_{FS}[4]_F[22]_S$ (8)
	$[211]_{FS}[31]_F[31]_S$ (32/3)

$$\mu_s = -\frac{1}{3} \frac{m_p}{m_s} P_{s\bar{s}} \quad (3)$$

(in units of nuclear magnetons $\mu_N = e/2m_p$). Here $P_{s\bar{s}} = A_{s\bar{s}}^2$ is the probability of the $s\bar{s}$ component and m_s is the constituent mass of the strange quark.

The following two configurations in the left column in Table I have the spin symmetry $[31]_S$ and thus $S = 1$. For these configurations and all other configurations in the left column that have the same spin symmetry, μ_s may be expressed in the general form

$$\mu_s = -\frac{m_p}{m_s} \left(\frac{7}{6} - \frac{1}{2} \bar{\sigma} \right) P_{s\bar{s}}. \quad (4)$$

Here $\bar{\sigma}$ is the average of σ_z of the s quark in the configuration for $S = 1$ with z projection $M = 1$. As $|\bar{\sigma}| \leq 1$, all of these configurations yield a negative strangeness magnetic moment in apparent conflict with the positive empirical value. The numerical values of $\bar{\sigma}$ are very similar—i.e., $41/72$ and $1/2$ —in the configurations with flavor-spin symmetry $[211]_F[31]_S$ and $[22]_F[31]_S$.

This result that the strangeness magnetic moment is negative if the \bar{s} quark is in the P state generalizes to all other configurations in the left column of Table I except for the energetically unfavored $[31]_{FS}[31]_F[4]_S$ configuration in which the $uuds$ system has total spin 2. For this configuration μ_s takes the value

$$\mu_s = \frac{7}{6} \frac{m_p}{m_s} P_{s\bar{s}}. \quad (5)$$

These results complete the analysis of the configurations with the \bar{s} quark in the P state.

In the configurations where \bar{s} quark is in its ground state, positive parity demands that the $uuds$ system has to be in the $L = 1$ state with mixed spatial symmetry $[31]_X$. In order to construct the corresponding spatial wave functions one may exploit a representation in terms of the following relative coordinates:

$$\begin{aligned} \vec{\xi}_1 &= \frac{1}{\sqrt{2}}(\vec{r}_1 - \vec{r}_2), & \vec{\xi}_2 &= \frac{1}{\sqrt{6}}(\vec{r}_1 + \vec{r}_2 - 2\vec{r}_3), \\ \vec{\xi}_3 &= \frac{1}{\sqrt{12}}(\vec{r}_1 + \vec{r}_2 + \vec{r}_3 - 3\vec{r}_4), & & \\ \vec{\xi}_4 &= \frac{1}{\sqrt{20}}(\vec{r}_1 + \vec{r}_2 + \vec{r}_3 + \vec{r}_4 - 4\vec{r}_5), & & \end{aligned} \quad (6)$$

which when complemented with the center-of-mass coordinate form a complete set of basis vectors.

For these states the wave functions for the configurations in the right columns in Table I may for a proton with spin-up then be written in the general form

$$\psi = A_{s\bar{s}} \sum_{a,b,c,d,e} \sum_{m,S,M,j} (1, S, m, M|J, j) (J, 1/2, j, s|1/2, 1/2) C_{[211]_a, [31]_a}^{[1111]} C_{[31]_b, [FS]_c}^{[31]_a} C_{[F]_d, [S]_e}^{[FS]_c} [211]_C(a)[31]_{X,m}(b)[F]_C(d)[S]_M(e) \bar{\chi}_s \varphi(\{r_i\}). \quad (7)$$

Here J is the total angular momentum of the $uuds$ system. The three components of the spatial state may be described as normalized combinations of a spatially symmetric function that is multiplied by the unit vectors $\hat{\xi}_1$, $\hat{\xi}_2$, and $\hat{\xi}_3$.

The contributions to μ_s from all configurations in the right column of Table I, which have the spatial symmetry $[31]_X$ and spin symmetry $[22]_S$, are positive:

$$\mu_s = \frac{m_p}{m_s} \left(\frac{1}{3} + \frac{2}{3} \bar{\ell} \right) P_{s\bar{s}}. \quad (8)$$

Here $\bar{\ell}$ is the average value of the z components of the orbital angular momentum of the s quark in the $uuds$ system for $m = 1$ ($M = 0$ since $S = 0$ for these configurations). These configurations thus give positive values for μ_s as $|\bar{\ell}| \leq 1/2$.

For the uppermost and energetically most likely configuration in the right column in Table I, $\bar{\ell} = 1/4$, and therefore μ_s is simply

$$\mu_s = \frac{m_p}{2m_s} P_{s\bar{s}}. \quad (9)$$

The following symmetry configuration in the right column of Table I has both spatial and spin symmetry $[31]$. The contribution to μ_s may for all states with both spatial and spin symmetry $[31]$ be expressed as

$$\begin{aligned} \mu_s &= -\frac{m_p}{m_s} P_{s\bar{s}} \quad (J = 0), \\ \mu_s &= \frac{m_p}{m_s} \left(\frac{1}{3} + \frac{1}{3} \bar{\ell} + \frac{1}{3} \bar{\sigma} \right) P_{s\bar{s}} \quad (J = 1). \end{aligned} \quad (10)$$

Of these configurations, that with the lowest energy has flavor-spin symmetry $[4]_{FS}[31]_F[31]_S$. In this configuration when $J = 1$, $\bar{\ell} = 1/4$ and $\bar{\sigma} = -1/18$.

From (7) one may finally infer that the contribution to μ_s from all the configurations in the right column of Table I, which have the spatial symmetry $[31]_X$ and spin symmetry $[4]_S$, are likewise positive and may be expressed in the form

$$\mu_s = \frac{m_p}{m_s} \left(\frac{5}{6} - \frac{1}{3} \bar{\ell} \right) P_{s\bar{s}}. \quad (11)$$

Here the first term is the sum of the contributions of the s and \bar{s} quarks, which are $1/2$ and $1/3$, respectively. In the lowest of these configurations, which also has flavor symmetry $[4]_F$, $\bar{\ell} = 1/4$. In the following one, in order of increasing energy, the flavor symmetry is $[31]_F$ and $\bar{\ell} = 19/54$. This completes the analysis of the configurations in the right column of Table I.

These results reveal that μ_s is positive in the lowest lying 5-quark configurations, in which the strange antiquark \bar{s} is in the ground state. For the lowest configuration

$\mu_s = 0.5m_p/m_s$ times the probability $P_{s\bar{s}}$ for the preformed $s\bar{s}$ component in the proton. This result then leads to $P_{s\bar{s}} \sim 2m_s/m_p \mu_s^{\text{exp}} \sim \mu_s^{\text{exp}}$. For μ_s^{exp} the result of the SAMPLE experiment [1] is $\mu_s = 0.37 \pm 0.2 \pm 0.26 \pm 0.07$.

In order to explain the structure of the (tentative) $\theta^+(uudd\bar{s})$ pentaquark, several clustered quark models have been proposed [14–16]. These quark cluster configurations have implications also for the pentaquark components with hidden strangeness in the nucleons. The magnetic moments of the pentaquark states in the cluster models have been calculated in Ref. [20] for the exotic flavor antidecuplet and normal flavor octet baryons.

In the Jaffe-Wilczek (JW) model [14], the $(uuds\bar{s})$ component in the proton should have the configuration $([ud] \times [us]\bar{s})$ with two scalar diquarks, $[ud]$ and $[us]$, in relative P wave, and the \bar{s} quark in its ground state. The corresponding strangeness magnetic moment is

$$\mu_s = \frac{m_p}{3m_s} \left(1 + \frac{2m_s}{m_{ud} + m_{us}} \right) P_{s\bar{s}}, \quad (12)$$

where m_{ud} and m_{us} are diquark masses for $[ud]$ and $[us]$, respectively.

In the Shuryak-Zahed (SZ) [15] model, the $(uuds\bar{s})$ is composed of one scalar diquark, one tensor diquark, and the \bar{s} antiquark. It has the same flavor configuration as in the JW model. But the orbital excitation $\ell = 1$ is now between the two quarks inside the tensor diquark instead of between two diquarks as in the JW model. The two quarks in the tensor diquark have total spin $S = 1$, which in combination with their relative P wave leads to total angular momentum $J = 1$ of the tensor diquark. In this configuration μ_s is found to be

$$\mu_s = \frac{m_p}{2m_s} \left(1 + \frac{m_q}{3m_{us}} \right) P_{s\bar{s}}. \quad (13)$$

Finally in the Karliner-Lipkin (KL) model [16], the θ^+ pentaquark is composed of a scalar $[ud]$ diquark and a $\{ud\bar{s}\}$ triquark with one orbital angular momentum $L = 1$ between the diquark and the triquark. The total spin of the triquark is one-half with a symmetric spin wave function for the two quarks.

An analogous configuration for the $(uuds\bar{s})$ component in the proton is $[ud]\{us\bar{s}\}$ with $L = 1$ between scalar diquark $[ud]$ and the spin-1/2 triquark $\{ud\bar{s}\}$. For this configuration μ_s is

$$\mu_s = -\frac{1}{3} \frac{m_p}{m_s} P_{s\bar{s}}, \quad (14)$$

which is negative definite and the same as for the $K^+\Lambda^0$ configuration. Another analogous configuration, $[us]ud\bar{s}$,

gives a positive μ_s , but with a magnitude about 100 times smaller than that of the given $[us]ud\bar{s}$. A combination of these two configurations in the flavor SU(3) symmetry [18] still gives a negative definite value for μ_s .

These results for the quark cluster models reveal that the $uuds\bar{s}$ configurations in JW and SZ models give similar and positive values for μ_s , while the configuration in the KL model gives a negative value. A common feature for the JW and SZ models is that in both of these the \bar{s} quark in its ground state and the $uuds$ system has total angular momentum 1. The configurations in KL and the $K^+\Lambda^0$ meson cloud model lack that feature.

This analysis of the $(uuds\bar{s})$ configurations shows that the lowest configurations with the \bar{s} in the ground state give positive values for μ_s , while the lowest configurations with the \bar{s} in the P state give negative values. The empirical indications [1–3] for a positive μ_s implies that the \bar{s} is in the ground state and that the $uuds$ system is orbitally excited. This is at variance with earlier studies on the intrinsic strangeness in the proton based on the $K^+\Lambda^0$ configuration for the strange component [10–13], which leads to negative values for μ_s . It should be instructive to investigate the consequences of the configurations with the \bar{s} in the ground state on other processes relevant to the intrinsic strangeness, such as the NuTeV anomaly problem [21,22].

Equally instructive would be a comparison of the role of the analogous configurations for the nonstrange pentaquark components in nucleon and N^* to that of conventional meson cloud configurations. Thus the observation of an excess \bar{d} over \bar{u} antiquarks in the proton [23], which has been described in terms of a $n\pi^+$ configuration [24], might alternately be described in terms of an $[udud]\bar{d}$ component in the proton. While the $N^*(1440)$ may contain significant $N\pi$ and $\Delta\pi$ components, its unusual properties might also be described in terms of admixtures of colored quark cluster configurations.

A recent chiral quark model calculation for the $uudd\bar{s}$ pentaquark system [25] suggests that the configuration with both $uudd$ and \bar{s} in their ground state gives lower energy than the $[31]_X[4]_{FS}$ configurations. It should be interesting to extend that calculation to the configuration with $uudd$ in the ground state and the \bar{s} in its P -wave state to see if it gives a higher energy than the lowest $[31]_X[4]_{FS}$ state. If so, it might provide an explanation for why the pentaquark component of the proton is in the lowest $[31]_X[4]_{FS}$ state. The pentaquark model with the most favorable configurations of the flavor-spin interaction with one quark in the P state [26] naturally leads to a positive value for μ_s .

In summary, the empirical indications for a positive strangeness magnetic moment of the proton suggest that the $s\bar{s}$ configuration in the proton is such that the \bar{s} is the ground state and the $uuds$ system is in the P state. This suggests that the $qqqq\bar{q}$ components in baryons may be mainly in colored quark cluster configurations rather than

in “meson cloud” configurations. In the configuration with the lowest energy the positive empirical strangeness magnetic moment gives a direct estimate of the probability of the $s\bar{s}$ configuration.

B. S. Z. acknowledges the hospitality of the Helsinki Institute of Physics during the course of this work. D. O. R. acknowledges the hospitality of the W. K. Kellogg Radiation Laboratory, California Institute of Technology. The research was supported in part by the Academy of Finland Grant No. 54038 and the National Natural Science Foundation of China.

*Electronic address: zoubs@ihep.ac.cn

†Electronic address: riska@pcu.helsinki.fi

- [1] D. T. Spayde *et al.*, Phys. Lett. B **583**, 79 (2004).
- [2] K. Aniol *et al.*, Phys. Rev. C **69**, 065501 (2004).
- [3] F. Maas *et al.*, Phys. Rev. Lett. **94**, 152001 (2005).
- [4] H. Weigel *et al.*, Phys. Lett. B **353**, 20 (1995).
- [5] P. Geiger and N. Isgur, Phys. Rev. D **55**, 299 (1997).
- [6] S.-T. Hong, B.-Y. Park, and D. P. Min, Phys. Lett. B **414**, 229 (1997).
- [7] V. Lyubovitskij *et al.*, Phys. Rev. C **66**, 055204 (2002).
- [8] R. Lewis, W. Wilcox, and R. M. Woloshyn, Phys. Rev. D **67**, 013003 (2003).
- [9] A. Silva, H. C. Kim, and K. Goetze, Eur. Phys. J. A **22**, 481 (2004).
- [10] M. Musolf and M. Burkhardt, Z. Phys. C **61**, 433 (1994).
- [11] H. Forkel, F. S. Navarra, and M. Nielsen, Phys. Rev. C **61**, 055206 (2000).
- [12] L. Hannelius and D. O. Riska, Phys. Rev. C **62**, 045204 (2000).
- [13] X. S. Chen *et al.*, Phys. Rev. C **70**, 015201 (2004).
- [14] R. Jaffe and F. Wilczek, Phys. Rev. Lett. **91**, 232003 (2003).
- [15] E. Shuryak and I. Zahed, Phys. Lett. B **589**, 21 (2004).
- [16] M. Karliner and H. Lipkin, Phys. Lett. B **575**, 249 (2003).
- [17] C. Helminen and D. O. Riska, Nucl. Phys. A **699**, 624 (2002).
- [18] J.-Q. Chen, *Group Representation Theory for Physicists* (World Scientific, Singapore, 1989).
- [19] L. Ya. Glozman and D. O. Riska, Phys. Rep. **268**, 263 (1996).
- [20] Y. R. Liu *et al.*, Phys. Rev. C **69**, 035205 (2004).
- [21] G. P. Zeller *et al.*, Phys. Rev. Lett. **88**, 091802 (2002).
- [22] Y. Ding, R. G. Xu, and B. Q. Ma, Phys. Lett. B **607**, 101 (2005); J. Alwall and G. Ingelman, Phys. Rev. D **70**, 111505(R) (2004); F. G. Cao and A. I. Signal, Phys. Lett. B **559**, 229 (2003).
- [23] G. T. Garvey and J. C. Peng, Prog. Part. Nucl. Phys. **47**, 203 (2001), and references therein.
- [24] J. Speth and A. W. Thomas, Adv. Nucl. Phys. **24**, 83 (1997).
- [25] F. Huang *et al.*, Phys. Lett. B **586**, 69 (2004).
- [26] Fl. Stancu and D. O. Riska, Phys. Lett. B **575**, 242 (2003); Fl. Stancu, Phys. Lett. B **595**, 269 (2004); **598**, 295(E) (2004).