

Resonant Fermi Gases with a Large Effective Range

A. Schwenk^{1,*} and C. J. Pethick^{2,†}

¹*Nuclear Theory Center, Indiana University, Bloomington, Indiana 47408, USA*

²*NORDITA, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark*

(Received 18 June 2005; published 10 October 2005)

We calculate the equation of state of a Fermi gas with resonant interactions when the effective range is appreciable. Using an effective field theory for a large scattering length and large effective range, we show how calculations in this regime become tractable. Our results are model independent, and as an application, we study the neutron matter equation of state at low densities of astrophysical interest $0.002 \text{ fm}^{-3} < \rho < 0.02 \text{ fm}^{-3}$, for which the interparticle separation is comparable to the effective range. We compare our simple results with those of conventional many-body calculations.

DOI: [10.1103/PhysRevLett.95.160401](https://doi.org/10.1103/PhysRevLett.95.160401)

PACS numbers: 03.75.Ss, 05.30.Fk, 21.65.+f

The properties of dilute Fermi gases when all two-body scattering lengths a_s are large compared to the interparticle separation r_s and the range of the interaction R is small compared with r_s are universal, in the sense that properties do not depend on details of the interparticle interaction [1]. This is because the only dimensionful scale is the Fermi momentum k_F , and the corresponding energy scale is the Fermi energy $\varepsilon_F = k_F^2/(2m)$, m being the fermion mass. Consequently, all macroscopic observables are given by appropriate powers of k_F or ε_F multiplied by universal factors. For instance, for two spin states with equal populations the energy per particle E/N of cold gases of, e.g., ⁶Li, ⁴⁰K atoms or neutrons under these conditions is given by

$$\frac{E}{N} = \xi \left(\frac{E}{N} \right)_{\text{free}} = \xi \frac{3k_F^2}{10m}, \quad (1)$$

where the universal factor ξ is a number.

Resonant, dilute Fermi gases were realized for the first time by O'Hara *et al.* in 2002 [2]. The factor ξ has been determined experimentally by a number of methods [2–5] in atomic gases where the interaction may be tuned by controlling the magnetic field. The extraction of the equation of state leads to $\xi = 0.51 \pm 0.04$ [3], $\xi \approx 0.7$ [4], and $\xi = 0.27^{+0.12}_{-0.09}$ [5], for temperatures in units of the Fermi temperature $T/T_F \approx 0.05$ (except for $T/T_F \approx 0.6$ in [4]). To date, the most reliable theoretical results are from $T = 0$, fixed-node Green's function Monte Carlo simulations, $\xi = 0.44 \pm 0.01$ [6] and $\xi = 0.42 \pm 0.01$ [7].

The purpose of this Letter is to consider the neutron gas as an example of a Fermi system with a large scattering length. For neutrons, the scattering length is unnaturally large and the currently accepted value is $a_{nn} = -18.5 \pm 0.3 \text{ fm}$ (for a recent review on the experimental situation, see [8]). This is to be compared with the range of nuclear interactions, which is given by the mass of the lightest exchange particle, the pion, and thus $R \sim 1/m_\pi \approx 1.4 \text{ fm}$. The effective range r_e is expected to be approximately charge independent, and therefore we take the neutron-

neutron effective range to be $r_{nn} = 2.7 \text{ fm}$, the same as the neutron-proton one [9]. Consequently, the neutron effective range is also significant, with $r_e m_\pi \approx 2$.

For neutrons, the validity of the universal equation of state applies to densities with $k_F r_e \ll 1$. This restricts neutron densities to $\rho = k_F^3/(3\pi^2) < 10^{-4} \text{ fm}^{-3}$, which is just below neutron drip density $\rho_{nd} = 2.3 \times 10^{-4} \text{ fm}^{-3} \sim 10^{-3} \rho_0$ at which neutrons become unbound. Here $\rho_0 = 0.16 \text{ fm}^{-3}$ is the saturation density of symmetric nuclear matter. Therefore, it is important to generalize the equation of state to resonant Fermi gases with an appreciable effective range. In the regime $k_F r_e \sim 1$, the energy per particle can be expressed as in Eq. (1) but with a system-specific factor $\xi(k_F r_e)$,

$$\frac{E}{N} = \xi(k_F r_e) \frac{3k_F^2}{10m}. \quad (2)$$

This is the case for neutron matter at subnuclear densities $\rho < \rho_0/10$. For the extension Eq. (2) to be valid, the contributions to the energy from higher partial waves and




FIG. 1 (color online). Hartree-Fock results for the energy per particle of neutron matter calculated from $V_{\text{low } k}$ for different cutoffs.

many-body forces must be negligible. This ensures that neutron densities with $k_F r_e \sim 1$ do not probe further details of nuclear forces. In addition, the effective range expansion truncated at r_e , $k \cot \delta(k) = -1/a_s + r_e k^2/2$, must describe the phase shifts $\delta(k)$ for S -wave scattering up to relative momenta $k \sim k_F$. The extension of the universal energy to $k_F r_e \sim 1$ is also valid for trapped Fermi gases where both the scattering length and the effective range are tuned to large values. In this Letter, we calculate $\xi(k_F r_e)$ for resonant Fermi gases with $k_F r_e \gtrsim 1$ and explain how this regime becomes theoretically tractable.

For neutron matter, the non- S -wave contributions to the energy per particle can be estimated from the Hartree-Fock results shown in Fig. 1. The Hartree-Fock calculations are based on the model-independent two-nucleon interaction $V_{\text{low } k}$ for different values of the momentum cutoff Λ . The evolution of $V_{\text{low } k}$ with the cutoff follows a renormalization group (RG) equation that guarantees that nucleon-nucleon phase shifts are cutoff independent. We have previously found that the evolution of all large-cutoff nuclear interactions to $\Lambda \lesssim 2.1 \text{ fm}^{-1}$ leads to the same low-momentum interaction $V_{\text{low } k}$ [10]. Varying the cutoff gives an approximate measure of higher-order contributions beyond Hartree-Fock and the effects of omitted many-body forces, since both are needed to obtain cutoff-independent results. We find in Fig. 1 that the low-density equation of state is cutoff independent for $k_F \lesssim 1.0 \text{ fm}^{-1}$, with negligible $l \geq 1$ contributions for $k_F \lesssim 0.8 \text{ fm}^{-1}$. In addition, the effective range expansion describes the 1S_0 phase shifts well up to relative momenta $k < 1.0 \text{ fm}^{-1}$, with a deviation of -0.4° , 0.2° , and 2.1° from the empirical phase shifts 57.7° , 48.6° , and 39.5° for $k = 0.4, 0.6$, and 0.8 fm^{-1} , respectively. Consequently, the equation of state of neutron matter is given by Eq. (2) for $k_F r_e \lesssim 2$.

We calculate the equation of state Eq. (2) using an effective field theory (EFT) for large scattering length and large effective range. EFT offers a systematic approach to interactions at low energies, and therefore our results are model independent [11]. Under the conditions of interest, both the scattering length and effective range are low-momentum scales $1/a_s \sim 1/r_e \sim Q$. The corresponding EFT is realized by introducing a difermion field d [12] (for a review, see Ref. [13]) with the lowest-order Lagrangian density given by (in units $\hbar = m = 1$)


$$\mathcal{L} = \psi^\dagger \left(i\partial_0 + \frac{\nabla^2}{2} \right) \psi - d^\dagger \left(i\partial_0 + \frac{\nabla^2}{4} - \Delta \right) d - g(d^\dagger \psi \psi + d \psi^\dagger \psi^\dagger). \quad (3)$$

Here ψ denotes the fermion field, and Δ and g are low-energy constants which describe the propagation of the difermion field and its coupling to two fermions, respectively. We also note that $i\partial_0 + \nabla^2/4$ is the operator for the two-body energy corrected for the center-of-mass motion. The EFT Lagrangian Eq. (3) was introduced for nucleon-

nucleon scattering in Ref. [12] following the work of Weinberg [14]. In the difermion EFT, the T matrix for two-body scattering depends only on the energy E of the interacting particles in the center-of-mass system,

$$T(E) = \frac{g^2}{\Delta - E - g^2 I_0(E + i\epsilon)}, \quad (4)$$

with $I_0(E) = \int_0^\Lambda d^3 p / (2\pi)^3 (E - p^2)^{-1}$. This corresponds to summing diagrams where the difermion propagator (represented as a double line) is dressed by fermion loops,



$$+ \dots \quad (5)$$

$I_0(E)$ denotes the loop integral regulated by a momentum cutoff Λ and the on-shell T matrix is given by $4\pi T^{-1}(E = k^2) = 1/a_s - r_e k^2/2 + ik$, which can equivalently be obtained from an energy-dependent potential,

$$V(E) = \frac{g^2}{\Delta - E}, \quad (6)$$

which follows after integrating out the difermion field. Matching the lowest-order low-energy constants Δ and g to the effective range expansion yields

$$\frac{\Delta}{g^2} = \frac{1}{4\pi a_s} - \frac{\Lambda}{2\pi^2} \quad \text{and} \quad \frac{1}{g^2} = \frac{r_e}{8\pi} - \frac{1}{2\pi^2 \Lambda}. \quad (7)$$

Consequently, for systems with large scattering length and large effective range, the potential $V(E)$ scales at low energies $E \sim Q^2$ as $V(E) \sim (Q + \Lambda + Q^2/\Lambda)^{-1}$. The cutoff generates higher-order terms $\sim k^4/\Lambda^3$ in $T^{-1}(E = k^2)$, which are suppressed for $k/\Lambda \ll 1$. Therefore, we consider the lowest-order EFT with large cutoffs. This is not required for $V_{\text{low } k}$, since the RG generates all higher-order contact interactions necessary to maintain cutoff-independent two-body observables. If one rescales the difermion field by $d \rightarrow gd$, the interaction terms become coupling independent and the kinetic term acquires a factor $1/g^2$. For large cutoffs, $1/g^2 \sim r_e$, and therefore the difermion field is an auxiliary field for positive effective range, even if there is a two-body bound state with $a_s > 0$. Integrating out the d field gives the standard contact interaction EFT. For negative effective range the difermion kinetic term has the normal sign and the Lagrangian Eq. (3) is the atom-molecule model. The great advantage of the present formulation is that it enables one to study systems with either positive or negative r_e .

Next, we study the scaling of diagrammatic contributions to the energy. Particle-particle loops are restricted only by the cutoff, and therefore these scale as $\sim \Lambda$ for large cutoffs. By contrast, particle-hole and hole-hole intermediate states scale with the low-momentum scale $\sim Q$. Particle-hole loops, for instance, scale with the density of states $\sim k_F$. Since the lowest-order interaction scales as

$V(E) \sim (Q + \Lambda + Q^2/\Lambda)^{-1}$, cutoff independence for large cutoffs minimally requires summing particle-particle ladders with Pauli blocking, such that the Λ^{-1} scaling of the potential cancels with each loop $\sim \Lambda$ [15]. Neglecting terms which vanish for large cutoffs, we find for T_{med} , the on-shell in-medium T matrix,

$$T_{\text{med}}(E = k^2; P) = 4\pi \left[\frac{1}{a_s} - \frac{r_e k^2}{2} - \frac{k_F + \frac{P}{2}}{\pi} + \frac{k}{\pi} \log \left(\frac{k_F + \frac{P}{2} + k}{k_F + \frac{P}{2} - k} \right) + \frac{k^2 + \frac{P^2}{4} - k_F^2}{\pi P} \log \left(\frac{(k_F + \frac{P}{2})^2 - k^2}{k_F^2 - \frac{P^2}{4} - k^2} \right) \right]^{-1}, \quad (8)$$

where P denotes the particle-pair momentum and we have neglected the imaginary part, as the latter vanishes for $k < \sqrt{k_F^2 - P^2/4}$ and therefore does not contribute to the energy. The resulting in-medium T matrix can be viewed as an effective interaction and scales as $T_{\text{med}}(E = k^2; P) \sim Q^{-1}$, independent of the cutoff. In addition to the scattering length and effective range terms, the inverse effective interaction acquires a density-dependent contribution $-(k_F + P/2)/\pi$ and two logarithmic terms. The first logarithmic term includes the BCS singularity for $P = 0$ and $k = k_F$. The resulting contribution to the energy is integrable and generally small. For neutron matter, realistic calculations of the superfluid pairing gap including polarization effects give $\Delta/\varepsilon_F \sim 0.1$ for densities with $k_F r_e \sim 1$ [16]. We therefore expect corrections to the energy per particle $\xi_{\text{sf}}(k_F r_e) \approx -5/8(\Delta/\varepsilon_F)^2 \approx -0.01$ due to pairing in this regime. The second logarithmic term in Eq. (8) is regular, and for $P = 0$ it simplifies to $-k_F/\pi$.

For large effective range, interactions are weaker at higher momenta and the effect on the in-medium T matrix can be understood qualitatively by inserting average values for the momenta $k^2 = 3/10k_F^2$ and $P^2 = 6/5k_F^2$ into Eq. (8). With this qualitative estimate, the average effective interaction for resonant Fermi gases is given by

$$\overline{T_{\text{med}}} = -\frac{4\pi}{U_0 k_F + \frac{3r_e k_F^2}{20}} = -\frac{4\pi}{U_0 k_F} \frac{1}{1 + 0.27k_F r_e}, \quad (9)$$

with $U_0 = 0.56$. In the universal regime with $k_F r_e \ll 1$, the average effective interaction is given by $\overline{T_{\text{med}}} = -4\pi/(U_0 k_F)$. The remaining contributions to the energy are of particle-hole or hole-hole nature, and scale as $Q \sim k_F$. In this EFT it is thus necessary to sum all diagrams, as the effects of each additional vertex $\sim k_F^{-1}$ combined with the additional loop $\sim k_F$ are of order one.

For Fermi gases with large effective range, the loop scaling is unchanged, but additional T_{med} vertices are now suppressed by $(1 + 0.27k_F r_e)^{-1}$. This implies that corrections to the Hartree-Fock equation of state with T_{med} as effective interaction come with powers of $(1 + 0.27k_F r_e)^{-1}$ relative to the universal case. In perturbation

theory, particle-particle-irreducible contributions to the energy enter in third order. With $\overline{T_{\text{med}}} = 4\pi\overline{a_{\text{med}}}$, we estimate the leading nonladder corrections to the $\xi(k_F r_e)$ obtained from T_{med} at the Hartree-Fock level as $\Delta\xi = 0.077[U_0(1 + 0.27k_F r_e)]^{-3}$ [17]. This provides only a qualitative estimate, since we evaluated the in-medium T matrix for average particle configurations.

To a good approximation, the equation of state of resonant Fermi gases with $k_F r_e \gtrsim 1$ can therefore be calculated from T_{med} at the Hartree-Fock level,

$$\xi_{\text{HF}}(k_F r_e) = 1 + \frac{5}{2\pi^2 k_F^5} \int_0^{2k_F} P^2 dP \times \int_0^{\sqrt{k_F^2 - (P^2/4)}} k^2 dk T_{\text{med}}(k^2; P) \times \min \left[1, \frac{k_F^2 - k^2 - \frac{P^2}{4}}{kP} \right], \quad (10)$$

with $1/a_s = 0$ in Eq. (8). Based on the arguments given above, we expect corrections to ξ beyond Hartree-Fock of order $\Delta\xi = 0.22 - 0.03$ and $\xi_{\text{sf}} \approx -0.01$ due to pairing for densities with $k_F r_e = 1 - 5$. Our results for $\xi_{\text{HF}}(k_F r_e)$ are shown in Fig. 2 with the theoretical error estimate. It is intriguing that we find $\xi_{\text{HF}}(1) \approx 0.47$, which is close to the universal factor $\xi = 0.44 \pm 0.01$ [6]. This implies a cancellation of higher-order contributions for $k_F r_e \ll 1$.

For $k_F r_e \gg 1$, the effective interaction is of the order $\overline{T_{\text{med}}} \sim -(r_e k_F^2)^{-1}$ and interaction effects disappear for infinite effective range. In the latter regime, $\xi \approx 1 + C/(k_F r_e)$, with a coefficient $C < 0$ of order one. Consequently, $\xi \rightarrow 1$ only for very large effective ranges. We find $\xi_{\text{HF}}(10) = 0.79$ and $\xi_{\text{HF}}(100) = 0.96$. If the effective range is negative, the average interaction in the medium is greater than for $r_e = 0$, and higher-order terms are only reduced for very large $|k_F r_e| \gg 1$.

Finally, we take into account a large but finite scattering length, and present in Fig. 3 results for the equation of state




FIG. 2 (color online). Result for $\xi_{\text{HF}}(k_F r_e)$ (solid line) which describes the equation of state of resonant Fermi gases with large effective range $k_F r_e \gtrsim 1$. The dashed lines provide an error estimate for ξ_{HF} as discussed in the text.




FIG. 3 (color online). Comparison of the energy per particle for neutron matter calculated from Eq. (10) for $a_s = -18.5$ fm and $r_e = 2.7$ fm (solid line) to Fermi hypernetted-chain (triangles) [18] and Brueckner results (circles) [19]. We also show the $V_{\text{low } k}$ Hartree-Fock results of Fig. 1. The dashed lines provide an error estimate for ξ_{HF} .

of neutron matter in the regime $k_F r_e \sim 1$. This is straightforward by including the $1/a_s$ term in Eq. (8). Our results are model independent and constrain neutron matter at subnuclear densities $0.002 \text{ fm}^{-3} < \rho < 0.02 \text{ fm}^{-3}$, which is of relevance to the physics of neutron stars and supernovae. We also compare our neutron matter results to many-body calculations using Fermi hypernetted-chain techniques [18] or Brueckner theory [19]. As can be seen from Fig. 3, all microscopic results lie within our theoretical error estimates.

In summary, we have calculated the equation of state of resonant Fermi gases with large effective range $k_F r_e \geq 1$ in the difermion EFT, with particular attention to neutron matter at subnuclear densities. A large magnitude of the effective range for an atomic gas can also be achieved by working with a narrow Feshbach resonance. In the lowest-order EFT, cutoff independence minimally requires the resummation of particle-particle ladders [13], which leads to an effective interaction that becomes weak for all $k_F r_e \geq 1$. The $k_F r_e$ dependence was used to show how resonant Fermi gases with large effective range, such as low-density neutron matter, are theoretically tractable and to estimate the error of our results. Neutron matter equations of state obtained using conventional many-body approaches are consistent with our model-independent difermion EFT results within these errors. For $k_F r_e \sim 1$, the energy is found to be close to the universal regime.

We thank Georg Bruun, Dick Furnstahl, Chuck Horowitz, and Daniel Phillips for useful discussions. The work of A. S. is supported by the U.S. DOE under Grant No. DE-FG02-87ER40365 and the NSF under Grant No. PHY-0244822.

*Electronic address: schwenk@indiana.edu

†Electronic address: pethick@nordita.dk

- [1] Bertsch posed the properties of Fermi gases in this limit as a many-body-theory challenge problem at MBX (2001). The first studies were reported by G. A. Baker, *Int. J. Mod. Phys. B* **15**, 1314 (2001); H. Heiselberg, *Phys. Rev. A* **63**, 043606 (2001).
- [2] K. M. O'Hara, S. L. Hemmer, M. E. Gehm, S. R. Granade, and J. E. Thomas, *Science* **298**, 2179 (2002).
- [3] M. E. Gehm, S. L. Hemmer, S. R. Granade, K. M. O'Hara, and J. E. Thomas, *Phys. Rev. A* **68**, 011401(R) (2003); J. Kinast, A. Turlapov, J. E. Thomas, Q. Chen, J. Stajic, and K. Levin, *Science* **307**, 1296 (2005).
- [4] T. Bourdel, J. Cubizolles, L. Khaykovich, K. M. F. Magalhaes, S. Kokkelmans, G. V. Shlyapnikov, and C. Salomon, *Phys. Rev. Lett.* **91**, 020402 (2003).
- [5] M. Bartenstein, A. Altmeyer, S. Riedl, S. Jochim, C. Chin, J. H. Denschlag, and R. Grimm, *Phys. Rev. Lett.* **92**, 120401 (2004); *cond-mat/0412712*.
- [6] J. Carlson, S.-Y. Chang, V. R. Pandharipande, and K. E. Schmidt, *Phys. Rev. Lett.* **91**, 050401 (2003); S. Y. Chang, J. Carlson, V. R. Pandharipande, and K. E. Schmidt, *Phys. Rev. A* **70**, 043602 (2004).
- [7] G. E. Astrakharchik, J. Boronat, J. Casulleras, and S. Giorgini, *Phys. Rev. Lett.* **93**, 200404 (2004).
- [8] A. Gardestig and D. R. Phillips, *nucl-th/0501049*.
- [9] The neutron-proton effective range parameters are well constrained from the Nijmegen 1993 partial-wave analysis (PWA93): $a_{np} = -23.768 \pm 0.006$ fm and $r_{np} = 2.68 \pm 0.01$ fm [M. C. M. Rentmeester (private communication)].
- [10] S. K. Bogner, T. T. S. Kuo, and A. Schwenk, *Phys. Rep.* **386**, 1 (2003).
- [11] The general problem of calculating the properties of a Fermi gas with resonant interactions in EFT has been considered by T. Schäfer, C.-W. Kao, and S. R. Cotanch, *nucl-th/0504088*, and references therein.
- [12] D. B. Kaplan, *Nucl. Phys.* **B494**, 471 (1997); P. F. Bedaque and U. van Kolck, *Phys. Lett. B* **428**, 221 (1998); S. R. Beane and M. J. Savage, *Nucl. Phys.* **A694**, 511 (2001).
- [13] D. R. Phillips, *Czech. J. Phys.* **52**, B49 (2002).
- [14] S. Weinberg, *Phys. Rev.* **130**, 776 (1963).
- [15] This resummation is not necessary if one evolves the large-cutoff theory to smaller cutoffs using the RG; see S. K. Bogner, A. Schwenk, R. J. Furnstahl, and A. Nogga, *nucl-th/0504043* [*Nucl. Phys. A* (to be published)].
- [16] A. Schwenk, B. Friman, and G. E. Brown, *Nucl. Phys.* **A713**, 191 (2003); see also Fig. 2 in A. Schwenk, *nucl-th/0411070*.
- [17] The third-order nonladder contributions to the energy have been evaluated by Baker; see Eq. (4.84) in G. A. Baker, *Rev. Mod. Phys.* **43**, 479 (1971). Note that for $r_e = 0$, \bar{T}_{med} leads to $\xi_{\text{HF}} = 0.36$ and $\Delta\xi = 0.45$.
- [18] B. Friedman and V. R. Pandharipande, *Nucl. Phys.* **A361**, 502 (1981).
- [19] G. Bao, L. Engvik, M. Hjorth-Jensen, E. Osnes, and E. Østgaard, *Nucl. Phys.* **A575**, 707 (1994).