

Observation of a Resonance $X(1835)$ in $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

M. Ablikim,¹ J. Z. Bai,¹ Y. Ban,¹¹ J. G. Bian,¹ X. Cai,¹ H. F. Chen,¹⁶ H. S. Chen,¹ H. X. Chen,¹ J. C. Chen,¹ Jin Chen,¹ Y. B. Chen,¹ S. P. Chi,² Y. P. Chu,¹ X. Z. Cui,¹ Y. S. Dai,¹⁸ Z. Y. Deng,¹ L. Y. Dong,^{1,*} Q. F. Dong,¹⁴ S. X. Du,¹ Z. Z. Du,¹ J. Fang,¹ S. S. Fang,² C. D. Fu,¹ C. S. Gao,¹ Y. N. Gao,¹⁴ S. D. Gu,¹ Y. T. Gu,⁴ Y. N. Guo,¹ Y. Q. Guo,¹ Z. J. Guo,¹⁵ F. A. Harris,¹⁵ K. L. He,¹ M. He,¹² Y. K. Heng,¹ H. M. Hu,¹ T. Hu,¹ G. S. Huang,^{1,†} X. P. Huang,¹ X. T. Huang,¹² X. B. Ji,¹ X. S. Jiang,¹ J. B. Jiao,¹² D. P. Jin,¹ S. Jin,¹ Yi Jin,¹ Y. F. Lai,¹ G. Li,² H. B. Li,¹ H. H. Li,¹ J. Li,¹ R. Y. Li,¹ S. M. Li,¹ W. D. Li,¹ W. G. Li,¹ X. L. Li,⁸ X. Q. Li,¹⁰ Y. L. Li,⁴ Y. F. Liang,¹³ H. B. Liao,⁶ C. X. Liu,¹ F. Liu,⁶ Fang Liu,¹⁶ H. H. Liu,¹ H. M. Liu,¹ J. Liu,¹¹ J. B. Liu,¹ J. P. Liu,¹⁷ R. G. Liu,¹ Z. A. Liu,¹ F. Lu,¹ G. R. Lu,⁵ H. J. Lu,¹⁶ J. G. Lu,¹ C. L. Luo,⁹ F. C. Ma,⁸ H. L. Ma,¹ L. L. Ma,¹ Q. M. Ma,¹ X. B. Ma,⁵ Z. P. Mao,¹ X. H. Mo,¹ J. Nie,¹ S. L. Olsen,¹⁵ H. P. Peng,¹⁶ N. D. Qi,¹ H. Qin,⁹ J. F. Qiu,¹ Z. Y. Ren,¹ G. Rong,¹ L. Y. Shan,¹ L. Shang,¹ D. L. Shen,¹ X. Y. Shen,¹ H. Y. Sheng,¹ F. Shi,¹ X. Shi,^{11,‡} H. S. Sun,¹ J. F. Sun,¹ S. S. Sun,¹ Y. Z. Sun,¹ Z. J. Sun,¹ Z. Q. Tan,⁴ X. Tang,¹ Y. R. Tian,¹⁴ G. L. Tong,¹ G. S. Varner,¹⁵ D. Y. Wang,¹ L. Wang,¹ L. S. Wang,¹ M. Wang,¹ P. Wang,¹ P. L. Wang,¹ W. F. Wang,^{1,§} Y. F. Wang,¹ Z. Wang,¹ Z. Y. Wang,¹ Zhe Wang,¹ Zheng Wang,² C. L. Wei,¹ D. H. Wei,¹ N. Wu,¹ X. M. Xia,¹ X. X. Xie,¹ B. Xin,^{8,†} G. F. Xu,¹ Y. Xu,¹⁰ M. L. Yan,¹⁶ F. Yang,¹⁰ H. X. Yang,¹ J. Yang,¹⁶ Y. X. Yang,³ M. H. Ye,² Y. X. Ye,¹⁶ Z. Y. Yi,¹ G. W. Yu,¹ C. Z. Yuan,¹ J. M. Yuan,¹ Y. Yuan,¹ S. L. Zang,¹ Y. Zeng,⁷ Yu Zeng,¹ B. X. Zhang,¹ B. Y. Zhang,¹ C. C. Zhang,¹ D. H. Zhang,¹ H. Y. Zhang,¹ J. W. Zhang,¹ J. Y. Zhang,¹ Q. J. Zhang,¹ X. M. Zhang,¹ X. Y. Zhang,¹² Y. Y. Zhang,¹³ Z. P. Zhang,¹⁶ Z. Q. Zhang,⁵ D. X. Zhao,¹ J. W. Zhao,¹ M. G. Zhao,¹⁰ P. P. Zhao,¹ W. R. Zhao,¹ Z. G. Zhao,^{1,||} H. Q. Zheng,¹¹ J. P. Zheng,¹ Z. P. Zheng,¹ L. Zhou,¹ N. F. Zhou,¹ K. J. Zhu,¹ Q. M. Zhu,¹ Y. C. Zhu,¹ Yingchun Zhu,^{1,¶} Y. S. Zhu,¹ Z. A. Zhu,¹ B. A. Zhuang,¹ X. A. Zhuang,¹ and B. S. Zou¹

(BES Collaboration)

¹*Institute of High Energy Physics, Beijing 100049, People's Republic of China*²*China Center for Advanced Science and Technology (CCAST), Beijing 100080, People's Republic of China*³*Guangxi Normal University, Guilin 541004, People's Republic of China*⁴*Guangxi University, Nanning 530004, People's Republic of China*⁵*Henan Normal University, Xinxiang 453002, People's Republic of China*⁶*Huazhong Normal University, Wuhan 430079, People's Republic of China*⁷*Hunan University, Changsha 410082, People's Republic of China*⁸*Liaoning University, Shenyang 110036, People's Republic of China*⁹*Nanjing Normal University, Nanjing 210097, People's Republic of China*¹⁰*Nankai University, Tianjin 300071, People's Republic of China*¹¹*Peking University, Beijing 100871, People's Republic of China*¹²*Shandong University, Jinan 250100, People's Republic of China*¹³*Sichuan University, Chengdu 610064, People's Republic of China*¹⁴*Tsinghua University, Beijing 100084, People's Republic of China*¹⁵*University of Hawaii, Honolulu, Hawaii 96822, USA*¹⁶*University of Science and Technology of China, Hefei 230026, People's Republic of China*¹⁷*Wuhan University, Wuhan 430072, People's Republic of China*¹⁸*Zhejiang University, Hangzhou 310028, People's Republic of China*

(Received 11 August 2005; published 28 December 2005)

The decay channel $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$ is analyzed using a sample of 5.8×10^7 J/ψ events collected with the BESII detector. A resonance, the $X(1835)$, is observed in the $\pi^+\pi^-\eta'$ invariant-mass spectrum with a statistical significance of 7.7σ . A fit with a Breit-Wigner function yields a mass $M = 1833.7 \pm 6.1(\text{stat}) \pm 2.7(\text{syst})$ MeV/ c^2 , a width $\Gamma = 67.7 \pm 20.3(\text{stat}) \pm 7.7(\text{syst})$ MeV/ c^2 , and a product branching fraction $B(J/\psi \rightarrow \gamma X) \cdot B(X \rightarrow \pi^+\pi^-\eta') = [2.2 \pm 0.4(\text{stat}) \pm 0.4(\text{syst})] \times 10^{-4}$. The mass and width of the $X(1835)$ are not compatible with any known meson resonance. Its properties are consistent with expectations for the state that produces the strong $p\bar{p}$ mass threshold enhancement observed in the $J/\psi \rightarrow \gamma p\bar{p}$ process at BESII.

DOI: 10.1103/PhysRevLett.95.262001

PACS numbers: 12.39.Mk, 12.40.Yx, 13.20.Gd, 13.75.Cs

An anomalous enhancement near the mass threshold in the $p\bar{p}$ invariant-mass spectrum from $J/\psi \rightarrow \gamma p\bar{p}$ de-

cays was reported by the BES II experiment [1]. This enhancement was fitted with a subthreshold S -wave Breit-

Wigner resonance function with a mass $M = 1859_{-10}^{+3+5}$ MeV/ c^2 , a width $\Gamma < 30$ MeV/ c^2 (at the 90% C.L.), and a product branching fraction (BF) $B(J/\psi \rightarrow \gamma X) \cdot B(X \rightarrow p\bar{p}) = [7.0 \pm 0.4(\text{stat})_{-0.8}^{+1.9}(\text{syst})] \times 10^{-5}$. This surprising experimental observation has stimulated a number of theoretical speculations [2–7] and motivated further investigations on baryon-antibaryon mass threshold structures, which led to the subsequent experimental observation of a strong $p\bar{\Lambda}$ mass threshold enhancement in $J/\psi \rightarrow pK^-\bar{\Lambda}$ decay [8]. Among various theoretical interpretations of the $p\bar{p}$ mass threshold enhancement, the most intriguing one is that of a $p\bar{p}$ bound state, sometimes called *baryonium* [2,5,9], which has been the subject of many experimental searches [10]. However, it should be noted that many theoretical predictions on the mass and width of a baryonium state depend on the details of models.

The baryonium interpretation of the $p\bar{p}$ mass enhancement requires a new resonance with a mass around 1.85 GeV/ c^2 , and it would be supported by the observation of the resonance in other decay channels. Possible decay modes for a $p\bar{p}$ bound state, suggested in Refs. [4,5], include $\pi^+\pi^-\eta'$. In this Letter, we report an analysis on the $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$ decay channel and the observation of a resonance in the $\pi^+\pi^-\eta'$ mass spectrum with a mass around 1835 MeV/ c^2 , where the η' meson is detected in two decay modes, $\eta' \rightarrow \pi^+\pi^-\eta$ ($\eta \rightarrow \gamma\gamma$) and $\eta' \rightarrow \gamma\rho$. In the following, this resonance is designated as the $X(1835)$. The results reported here are based on a sample of 5.8×10^7 J/ψ decays detected with the upgraded Beijing Spectrometer (BESII) at the Beijing Electron-Positron Collider (BEPC).

BESII is a large solid-angle magnetic spectrometer that is described in detail in Ref. [11]. Charged particle momenta are determined with a resolution of $\sigma_p/p = 1.78\%\sqrt{1+p^2}$ (GeV/ c^2) in a 40-layer cylindrical main drift chamber (MDC). Particle identification is accomplished by specific ionization (dE/dx) measurements in the MDC and time-of-flight (TOF) measurements in a barrel-like array of 48 scintillation counters. The dE/dx resolution is $\sigma_{dE/dx} = 8.0\%$; the TOF resolution is measured to be $\sigma_{\text{TOF}} = 180$ ps for Bhabha events. Outside of the time-of-flight counters is a 12-radiation-length barrel shower counter (BSC) comprised of gas tubes interleaved with lead sheets. The BSC measures the energies and directions of photons with resolutions of $\sigma_E/E \approx 21\%/\sqrt{E}$ (GeV), $\sigma_\phi = 7.9$ mrad, and $\sigma_z = 2.3$ cm. The iron flux return of the magnet is instrumented with three double layers of counters that are used to identify muons. In this analysis, a GEANT3-based Monte Carlo (MC) package with detailed consideration of the detector performance is used. The consistency between data and MC has been carefully checked in many high-purity physics channels, and the agreement is reasonable [12].

For the $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$ ($\eta' \rightarrow \pi^+\pi^-\eta$, $\eta \rightarrow \gamma\gamma$) channel, candidate events are required to have four charged tracks, each of which is well fitted to a helix that is within the polar angle region $|\cos\theta| < 0.8$ and with a transverse momentum larger than 70 MeV/ c . The total charge of the four tracks is required to be zero. For each track, the TOF and dE/dx information are combined to form particle identification confidence levels for the π , K , and p hypotheses; the particle type with the highest confidence level is assigned to each track. At least three of the charged tracks are required to be identified as pions. Candidate photons are required to have an energy deposit in the BSC greater than 60 MeV and to be isolated from charged tracks by more than 5° ; the number of photons is required to be three. A four-constraint (4C) energy-momentum conservation kinematic fit is performed to the $\pi^+\pi^-\pi^+\pi^-\gamma\gamma\gamma$ hypothesis, and the χ_{4C}^2 is required to be less than 8 and also less than the χ^2 for the kinematically similar $K^+K^-\pi^+\pi^-\gamma\gamma\gamma$ hypothesis. An η signal is evident in the $\gamma\gamma$ invariant-mass distribution of all $\gamma\gamma$ pairings [Fig. 1(a)]. In order to reduce combinatorial backgrounds from $\pi^0 \rightarrow \gamma\gamma$ decays, we require that the invariant masses of all $\gamma\gamma$ pairings are greater than 0.22 GeV/ c^2 . Candidate η mesons are selected by requiring $|M_{\gamma\gamma} - m_\eta| < 0.05$ GeV/ c^2 . The events are then subjected to a five-constraint (5C) fit where the invariant mass of the $\gamma\gamma$ pair associated with the η is constrained to m_η , and $\chi_{5C}^2 < 15$ is required. The 5C fit improves the $M_{\pi^+\pi^-\eta}$ mass resolution from 20 MeV/ c^2 (for the 4C fit) to 7 MeV/ c^2 . Figure 1(b) shows the $\pi^+\pi^-\eta$ invariant-mass distribution after the 5C fit, where a clear η' signal is visible. For η' candidates, we select $\pi^+\pi^-\eta$ combinations with $|M_{\pi^+\pi^-\eta} - m_{\eta'}| < 0.015$ GeV/ c^2 . In a small fraction of events, more than one combination passes the above selection. In these cases, the combination with $M_{\pi^+\pi^-\eta}$ closest to η' mass is used [13]. The $\pi^+\pi^-\eta'$ invariant-mass spectrum for the selected events is shown in Fig. 1(c), where a peak at a mass around 1835 MeV/ c^2 is observed.

For the $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$ ($\eta' \rightarrow \gamma\rho$) channel, events with four charged tracks (with zero net charge) and two photons are selected. At least three of the charged tracks are required to be identified as pions. These events are subjected to a 4C kinematic fit to the $\pi^+\pi^-\pi^+\pi^-\gamma\gamma$ hypothesis, and the χ_{4C}^2 is required to be less than 8 and less than the χ^2 for the $K^+K^-\pi^+\pi^-\gamma\gamma$ hypothesis. At this stage of the analysis, the primary remaining background contributions are due to $J/\psi \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$, $J/\psi \rightarrow \pi^+\pi^-\pi^+\pi^-\eta$, and $J/\psi \rightarrow \omega$ ($\omega \rightarrow \gamma\pi^0$) $\pi^+\pi^-\pi^+\pi^-$; these produce peaks at m_{π^0} , m_η , and m_ω in the $\gamma\gamma$ invariant-mass distribution shown in Fig. 2(a). We suppress these backgrounds by rejecting events with $M_{\gamma\gamma} < 0.22$ GeV/ c^2 , $|M_{\gamma\gamma} - m_\eta| < 0.05$ GeV/ c^2 , or 0.72 GeV/ $c^2 < M_{\gamma\gamma} < 0.82$ GeV/ c^2 . To select ρ and η' signals, all $\pi^+\pi^-$ and $\gamma\pi^+\pi^-$ combinations are consid-




FIG. 1 (color online). Invariant-mass distributions for selected $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$ ($\eta' \rightarrow \pi^+\pi^-\eta$, $\eta \rightarrow \gamma\gamma$) candidate events. (a) The invariant-mass distribution of $\gamma\gamma$ pairs. (b) The $\pi^+\pi^-\eta'$ invariant-mass distribution. (c) The $\pi^+\pi^-\eta'$ invariant-mass distributions; the open histogram is data and the shaded histogram is $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$ phase-space MC events (with arbitrary normalization).

ered. The $\gamma\pi^+\pi^-$ invariant-mass distribution shows an η' signal [Fig. 2(b)]. We require that $|M_{\pi^+\pi^-} - m_\rho| < 0.2 \text{ GeV}/c^2$ and $|M_{\gamma\pi^+\pi^-} - m_{\eta'}| < 0.025 \text{ GeV}/c^2$. If more than one combination passes these criteria, the combination with $M_{\gamma\pi^+\pi^-}$ closest to $m_{\eta'}$ is selected [13]. For this channel, there is also a distinct peak near $1835 \text{ MeV}/c^2$ in the $\pi^+\pi^-\eta'$ invariant-mass spectrum [Fig. 2(c)].

To ensure that the peak near $1835 \text{ MeV}/c^2$ is not due to background, we have made extensive studies of potential background processes using both data and MC. Non- η' processes are studied with η' mass-sideband events. The




FIG. 2 (color online). Invariant-mass distributions for the selected $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$ ($\eta' \rightarrow \gamma\rho$) candidate events. (a) The invariant-mass distribution for $\gamma\gamma$ pairs. (b) The $\gamma\pi^+\pi^-$ invariant-mass distribution. (c) The $\pi^+\pi^-\eta'$ invariant-mass distributions; the open histogram is data and the shaded histogram is from $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$ phase-space MC events (with arbitrary normalization).

main background channel $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$ and other background processes with multiphotons and/or with kaons are reconstructed with the data. In addition, we also checked for possible backgrounds with a MC sample of 60×10^6 J/ψ decays generated by the LUND model [14]. None of these background sources produce a peak around $1835 \text{ MeV}/c^2$ in the $\pi^+\pi^-\eta'$ invariant-mass spectrum.

Figure 3 shows the $\pi^+\pi^-\eta'$ invariant-mass spectrum for the combined $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$ ($\eta' \rightarrow \pi^+\pi^-\eta$) and $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$ ($\eta' \rightarrow \gamma\rho$) samples [i.e., the sum of the histograms in Figs. 1(c) and 2(c)]. This spectrum is fitted




FIG. 3. The $\pi^+\pi^-\eta'$ invariant-mass distribution for selected events from both the $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$ ($\eta' \rightarrow \pi^+\pi^-\eta$, $\eta' \rightarrow \gamma\gamma$) and $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$ ($\eta' \rightarrow \gamma\rho$) analyses. The bottom panel shows the fit (solid curve) to the data (points with error bars); the dashed curve indicates the background function.

with a Breit-Wigner (BW) function convolved with a Gaussian mass resolution function (with $\sigma = 13 \text{ MeV}/c^2$) to represent the $X(1835)$ signal plus a smooth polynomial background function. The mass and width obtained from the fit (shown in the bottom panel in Fig. 3) are $M = 1833.7 \pm 6.1 \text{ MeV}/c^2$ and $\Gamma = 67.7 \pm 20.3 \text{ MeV}/c^2$. The signal yield from the fit is 264 ± 54 events with a confidence level 45.5% ($\chi^2/\text{d.o.f.} = 57.6/57$) and $-2 \ln L = 58.4$. A fit to the mass spectrum without a BW signal function returns $-2 \ln L = 126.5$. The change in $-2 \ln L$ with $\Delta(\text{d.o.f.}) = 3$ corresponds to a statistical significance of 7.7σ for the signal.

Using MC-determined selection efficiencies of 3.72% and 4.85% for the $\eta' \rightarrow \pi^+\pi^-\eta$ and $\eta' \rightarrow \gamma\rho$ modes, respectively, we determine a product BF of

$$B(J/\psi \rightarrow \gamma X(1835)) \cdot B(X(1835) \rightarrow \pi^+\pi^-\eta') \\ = (2.2 \pm 0.4) \times 10^{-4}.$$

The consistency between the two η' decay modes is checked by fitting the distributions in Figs. 1(c) and 2(c) separately with the method described above. The fit to Fig. 1(c) gives $M = 1827.4 \pm 8.1 \text{ MeV}/c^2$ and $\Gamma = 54.2 \pm 34.5 \text{ MeV}/c^2$ with a statistical significance of 5.1σ . From the 68 ± 26 signal events obtained from the fit, the product BF is $B(J/\psi \rightarrow \gamma X(1835)) \cdot B(X(1835) \rightarrow \pi^+\pi^-\eta') = (1.8 \pm 0.7) \times 10^{-4}$. Similar results are ob-

tained if we apply only a 4C kinematic fit in this analysis. For the fit to Fig. 2(c), the mass and width are determined to be $M = 1836.3 \pm 7.9 \text{ MeV}/c^2$ and $\Gamma = 70.3 \pm 23.1 \text{ MeV}/c^2$ with a statistical significance of 6.0σ . For this mode alone, the signal yield of 193 ± 43 signal events corresponds to $B(J/\psi \rightarrow \gamma X(1835)) \cdot B(X(1835) \rightarrow \pi^+\pi^-\eta') = (2.3 \pm 0.5) \times 10^{-4}$. The $X(1835)$ mass, width, and product BF values determined from the two η' decay modes separately are in good agreement with each other.

The systematic uncertainties on the mass and width are determined by varying the functional form used to represent the background, the fitting range of the mass spectrum, the mass calibration, and possible biases due to the fitting procedure. The latter are estimated from differences between the input and output mass and width values from MC studies. The total systematic errors on the mass and width are 2.7 and 7.7 MeV/c^2 , respectively. The systematic error on the branching fraction measurement comes mainly from the uncertainties of MDC simulation (including systematic uncertainties of the tracking efficiency and the kinematic fits), the photon detection efficiency, the particle identification efficiency, the η' decay branching fractions to $\pi^+\pi^-\eta$ and $\gamma\rho$, the background function parametrization, the fitting range of the mass spectrum, the requirements on numbers of photons, the invariant-mass distributions of $\gamma\gamma$ pairs in the two analyses, the $\pi^+\pi^-$ invariant-mass distribution in $\eta' \rightarrow \gamma\pi^+\pi^-$ decays, MC statistics, the total number of J/ψ events [15], and the unknown spin-parity of the $X(1835)$. For the latter, we use the difference between phase space and a $J^{PC} = 0^{-+}$ hypothesis for the $X(1835)$. The total relative systematic error on the product branching fraction is 20.2%.

In summary, the decay channel $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$ is analyzed using two η' decay modes, $\eta' \rightarrow \pi^+\pi^-\eta$ and $\eta' \rightarrow \gamma\rho$. A resonance, the $X(1835)$, is observed with a high statistical significance of 7.7σ in the $\pi^+\pi^-\eta'$ invariant-mass spectrum. From a fit with a Breit-Wigner function, the mass is determined to be $M = 1833.7 \pm 6.1(\text{stat}) \pm 2.7(\text{syst}) \text{ MeV}/c^2$, the width is $\Gamma = 67.7 \pm 20.3(\text{stat}) \pm 7.7(\text{syst}) \text{ MeV}/c^2$, and the product branching fraction is $B(J/\psi \rightarrow \gamma X) \cdot B(X \rightarrow \pi^+\pi^-\eta') = [2.2 \pm 0.4(\text{stat}) \pm 0.4(\text{syst})] \times 10^{-4}$. The mass and width of the $X(1835)$ are not compatible with any known meson resonance [16]. In Ref. [16], the candidate closest in mass to the $X(1835)$ is the (unconfirmed) $2^{-+} \eta_2(1870)$ with $M = 1842 \pm 8 \text{ MeV}/c^2$. The width of this state, $\Gamma = 225 \pm 14 \text{ MeV}/c^2$, is considerably larger than that of the $X(1835)$ (see also [17], where the 2^{-+} component in the $\eta\pi\pi$ mode of J/ψ radiative decay has a mass $1840 \pm 15 \text{ MeV}/c^2$ and a width $170 \pm 40 \text{ MeV}/c^2$).

We examined the possibility that the $X(1835)$ is responsible for the $p\bar{p}$ mass threshold enhancement observed in radiative $J/\psi \rightarrow \gamma p\bar{p}$ decays [1]. It has been pointed out that the S -wave BW function used for the fit in Ref. [1]

should be modified to include the effect of final-state interactions (FSI) on the shape of the $p\bar{p}$ mass spectrum [6,7]. Redoing the S -wave BW fit to the $p\bar{p}$ invariant-mass spectrum of Ref. [1], including the zero isospin, S -wave FSI factor of Ref. [7], yields a mass $M = 1831 \pm 7 \text{ MeV}/c^2$ and a width $\Gamma < 153 \text{ MeV}/c^2$ (at the 90% C.L.) [systematic uncertainties are not included in the error of the mass and the upper limit of the width. In contrast to Ref. [7], the isospin = 1 FSI factor is not used to redo the fit since the isospin = 1 states are strongly suppressed in J/ψ radiative decays]; these values are in good agreement with the mass and width of $X(1835)$ reported here. Moreover, according to Ref. [5], the $\pi\pi\eta'$ decay mode is expected to be strong for a $p\bar{p}$ bound state. Thus, the $X(1835)$ resonance is a prime candidate for the source of the $p\bar{p}$ mass threshold enhancement in the $J/\psi \rightarrow \gamma p\bar{p}$ process. In this case, the J^{PC} and I^G of the $X(1835)$ could only be 0^{-+} and 0^+ , which can be tested in future experiments. Also in this context, the relative $p\bar{p}$ decay strength is quite strong: $B(X \rightarrow p\bar{p})/B(X \rightarrow \pi^+\pi^-\eta') \sim 1/3$ [the product BF determined from the fit that includes FSI effects on the $p\bar{p}$ mass spectrum is within the systematic errors of the result reported in Ref. [1]]. Since decays to $p\bar{p}$ are kinematically allowed only for a small portion of the high-mass tail of the resonance and have very limited phase space, the large $p\bar{p}$ branching fraction implies an unusually strong coupling to $p\bar{p}$, as expected for a $p\bar{p}$ bound state [9,18]. However, other possible interpretations of the $X(1835)$ that have no relation to the $p\bar{p}$ mass threshold enhancement are not excluded.

We thank many Chinese theorists for their helpful discussions and suggestions and Dr. A. Sibirtsev *et al.* for providing the FSI functions. The BES Collaboration acknowledges the staff of BEPC for the excellent performance of the machine. This work is supported in part by the National Natural Science Foundation of China under Contracts No. 10491300, No. 10225524, No. 10225525, No. 10425523, the Chinese Academy of Sciences under Contract No. KJ 95T-03, the 100 Talents Program of CAS under Contracts No. U-11, No. U-24, No. U-25, and the Knowledge Innovation Project of CAS under Contracts No. KJCX2-SW-N10, No. U-602, No. U-34 (IHEP); by the National Natural Science Foundation of China under Contract No. 10175060 (USTC); and by the Department of Energy under Contract No. DE-FG04ER41291 (University of Hawaii).

*Present address: Iowa State University, Ames, IA 50011-3160, USA.

†Present address: Purdue University, West Lafayette, IN 47907, USA.

‡Present address: Cornell University, Ithaca, NY 14853, USA.

§Present address: Laboratoire de l'Accélérateur Linéaire, Orsay, F-91898, France.

||Present address: University of Michigan, Ann Arbor, MI 48109, USA.

¶Present address: DESY, D-22607, Hamburg, Germany.

- [1] J.Z. Bai *et al.* (BES Collaboration), Phys. Rev. Lett. **91**, 022001 (2003).
- [2] A. Datta and P.J. O'Donnell, Phys. Lett. B **567**, 273 (2003); M.L. Yan *et al.*, Phys. Rev. D **72**, 034027 (2005); B. Loiseau and S. Wycech, Phys. Rev. C **72**, 011001 (2005).
- [3] J. Ellis, Y. Frishman, and M. Karliner, Phys. Lett. B **566**, 201 (2003); J. L. Rosner, Phys. Rev. D **68**, 014004 (2003).
- [4] C. S. Gao and S. L. Zhu, Commun. Theor. Phys. **42**, 844 (2004).
- [5] G. J. Ding and M. L. Yan, Phys. Rev. C **72**, 015208 (2005).
- [6] B. S. Zou and H. C. Chiang, Phys. Rev. D **69**, 034004 (2004).
- [7] A. Sibirtsev *et al.*, Phys. Rev. D **71**, 054010 (2005).
- [8] M. Ablikim *et al.* (BES Collaboration), Phys. Rev. Lett. **93**, 112002 (2004).
- [9] I. S. Shapiro, Phys. Rep. **35**, 129 (1978); C. B. Dover and M. Goldhaber, Phys. Rev. D **15**, 1997 (1977).
- [10] For recent reviews of this subject, see E. Klempt *et al.*, Phys. Rep. **368**, 119 (2002); J.-M. Richard, Nucl. Phys. B, Proc. Suppl. **86**, 361 (2000).
- [11] J.Z. Bai *et al.* (BES Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **458**, 627 (2001).
- [12] M. Ablikim *et al.* (BES Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **552**, 344 (2005).
- [13] The selection criteria are optimized by maximizing $N_s/\sqrt{N_{\text{tot}}}$, where N_s is the expected number of $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$ signal events and N_{tot} is the total expected number of signal and background events. In order not to bias the $\pi^+\pi^-\eta'$ invariant-mass spectrum, we do not include any resonance in the optimization.
- [14] J. C. Chen *et al.*, Phys. Rev. D **62**, 034003 (2000).
- [15] S. S. Fang *et al.*, High Energy Phys. Nucl. Phys. **27**, 277 (2003).
- [16] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. B **592**, 1 (2004).
- [17] J. Z. Bai *et al.* (BES Collaboration), Phys. Lett. B **446**, 356 (1999).
- [18] S. L. Zhu and C. S. Gao, hep-ph/0507050.