

Search for the Higgs Boson in $H \rightarrow WW^{(*)}$ Decays in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV

- V. M. Abazov,³⁵ B. Abbott,⁷² M. Abolins,⁶³ B. S. Acharya,²⁹ M. Adams,⁵⁰ T. Adams,⁴⁸ M. Agelou,¹⁸ J.-L. Agram,¹⁹ S. H. Ahn,³¹ M. Ahsan,⁵⁷ G. D. Alexeev,³⁵ G. Alkhazov,³⁹ A. Alton,⁶² G. Alverson,⁶¹ G. A. Alves,² M. Anastasoaie,³⁴ T. Andeen,⁵² S. Anderson,⁴⁴ B. Andrieu,¹⁷ Y. Arnoud,¹⁴ M. Arov,⁵¹ A. Askew,⁴⁸ B. Åsman,⁴⁰ A. C. S. Assis Jesus,³ O. Atramentov,⁵⁵ C. Autermann,²¹ C. Avila,⁸ F. Badaud,¹³ A. Baden,⁵⁹ L. Bagby,⁵¹ B. Baldin,⁴⁹ P. W. Balm,³³ P. Banerjee,²⁹ S. Banerjee,²⁹ E. Barberis,⁶¹ P. Bargassa,⁷⁷ P. Baringer,⁵⁶ C. Barnes,⁴² J. Barreto,² J. F. Bartlett,⁴⁹ U. Bassler,¹⁷ D. Bauer,⁵³ A. Bean,⁵⁶ S. Beauceron,¹⁷ M. Begalli,³ M. Begel,⁶⁸ A. Bellavance,⁶⁵ S. B. Beri,²⁷ G. Bernardi,¹⁷ R. Bernhard,^{49,*} I. Bertram,⁴¹ M. Besançon,¹⁸ R. Beuselinck,⁴² V. A. Bezzubov,³⁸ P. C. Bhat,⁴⁹ V. Bhatnagar,²⁷ M. Binder,²⁵ C. Biscarat,⁴¹ K. M. Black,⁶⁰ I. Blackler,⁴² G. Blazey,⁵¹ F. Blekman,⁴² S. Blessing,⁴⁸ D. Bloch,¹⁹ U. Blumenschein,²³ A. Boehlein,⁴⁹ O. Boeriu,⁵⁴ T. A. Bolton,⁵⁷ F. Borcherding,⁴⁹ G. Borissov,⁴¹ K. Bos,³³ T. Bose,⁶⁷ A. Brandt,⁷⁵ R. Brock,⁶³ G. Brooijmans,⁶⁷ A. Bross,⁴⁹ N. J. Buchanan,⁴⁸ D. Buchholz,⁵² M. Buehler,⁵⁰ V. Buescher,²³ S. Burdin,⁴⁹ S. Burke,⁴⁴ T. H. Burnett,⁷⁹ E. Busato,¹⁷ C. P. Buszello,⁴² J. M. Butler,⁶⁰ J. Cammin,⁶⁸ S. Caron,³³ W. Carvalho,³ B. C. K. Casey,⁷⁴ N. M. Cason,⁵⁴ H. Castilla-Valdez,³² S. Chakrabarti,²⁹ D. Chakraborty,⁵¹ K. M. Chan,⁶⁸ A. Chandra,²⁹ D. Chapin,⁷⁴ F. Charles,¹⁹ E. Cheu,⁴⁴ D. K. Cho,⁶⁰ S. Choi,⁴⁷ B. Choudhary,²⁸ T. Christiansen,²⁵ L. Christofek,⁵⁶ D. Claes,⁶⁵ B. Clément,¹⁹ C. Clément,⁴⁰ Y. Coadou,⁵ M. Cooke,⁷⁷ W. E. Cooper,⁴⁹ D. Coppage,⁵⁶ M. Corcoran,⁷⁷ A. Cothenet,¹⁵ M.-C. Cousinou,¹⁵ B. Cox,⁴³ S. Crépé-Renaudin,¹⁴ D. Cutts,⁷⁴ H. da Motta,² M. Das,⁵⁸ B. Davies,⁴¹ G. Davies,⁴² G. A. Davis,⁵² K. De,⁷⁵ P. de Jong,³³ S. J. de Jong,³⁴ E. De La Cruz-Burelo,⁶² C. De Oliveira Martins,³ S. Dean,⁴³ J. D. Degenhardt,⁶² F. Déliot,¹⁸ M. Demarteau,⁴⁹ R. Demina,⁶⁸ P. Demine,¹⁸ D. Denisov,⁴⁹ S. P. Denisov,³⁸ S. Desai,⁶⁹ H. T. Diehl,⁴⁹ M. Diesburg,⁴⁹ M. Doidge,⁴¹ H. Dong,⁶⁹ S. Doulas,⁶¹ L. V. Dudko,³⁷ L. Duflot,¹⁶ S. R. Dugad,²⁹ A. Duperrin,¹⁵ J. Dyer,⁶³ A. Dyshkant,⁵¹ M. Eads,⁶⁵ D. Edmunds,⁶³ T. Edwards,⁴³ J. Ellison,⁴⁷ J. Elmsheuser,²⁵ V. D. Elvira,⁴⁹ S. Eno,⁵⁹ P. Ermolov,³⁷ J. Estrada,⁴⁹ H. Evans,⁶⁷ A. Evdokimov,³⁶ V. N. Evdokimov,³⁸ J. Fast,⁴⁹ S. N. Fataki,⁶⁰ L. Feligioni,⁶⁰ A. V. Ferapontov,³⁸ T. Ferbel,⁶⁸ F. Fiedler,²⁵ F. Filthaut,³⁴ W. Fisher,⁴⁹ H. E. Fisk,⁴⁹ I. Fleck,²³ M. Fortner,⁵¹ H. Fox,²³ S. Fu,⁴⁹ S. Fuess,⁴⁹ T. Gadfort,⁷⁹ C. F. Galea,³⁴ E. Gallas,⁴⁹ E. Galyaev,⁵⁴ C. Garcia,⁶⁸ A. Garcia-Bellido,⁷⁹ J. Gardner,⁵⁶ V. Gavrilov,³⁶ A. Gay,¹⁹ P. Gay,¹³ D. Gelé,¹⁹ R. Gelhaus,⁴⁷ K. Genser,⁴⁹ C. E. Gerber,⁵⁰ Y. Gershtein,⁴⁸ D. Gillberg,⁵ G. Ginther,⁶⁸ T. Golling,²² N. Gollub,⁴⁰ B. Gómez,⁸ K. Gounder,⁴⁹ A. Goussiou,⁵⁴ P. D. Grannis,⁶⁹ S. Greder,³ H. Greenlee,⁴⁹ Z. D. Greenwood,⁵⁸ E. M. Gregores,⁴ Ph. Gris,¹³ J.-F. Grivaz,¹⁶ S. Grünendahl,⁴⁹ M. W. Grünewald,³⁰ G. Gutierrez,⁴⁹ P. Gutierrez,⁷² A. Haas,⁶⁷ N. J. Hadley,⁵⁹ S. Hagopian,⁴⁸ J. Haley,⁶⁶ I. Hall,⁷² R. E. Hall,⁴⁶ C. Han,⁶² L. Han,⁷ K. Hanagaki,⁴⁹ K. Harder,⁵⁷ A. Harel,²⁶ R. Harrington,⁶¹ J. M. Hauptman,⁵⁵ R. Hauser,⁶³ J. Hays,⁵² T. Hebbeker,²¹ D. Hedin,⁵¹ J. M. Heinmiller,⁵⁰ A. P. Heinson,⁴⁷ U. Heintz,⁶⁰ C. Hensel,⁵⁶ G. Hesketh,⁶¹ M. D. Hildreth,⁵⁴ R. Hirosky,⁷⁸ J. D. Hobbs,⁶⁹ B. Hoeneisen,¹² M. Hohlfeld,²⁴ S. J. Hong,³¹ R. Hooper,⁷⁴ P. Houben,³³ Y. Hu,⁶⁹ J. Huang,⁵³ V. Hynek,⁹ I. Iashvili,⁴⁷ R. Illingworth,⁴⁹ A. S. Ito,⁴⁹ S. Jabeen,⁵⁶ M. Jaffré,¹⁶ S. Jain,⁷² V. Jain,⁷⁰ K. Jakobs,²³ C. Jarvis,⁵⁹ A. Jenkins,⁴² R. Jesik,⁴² K. Johns,⁴⁴ M. Johnson,⁴⁹ A. Jonckheere,⁴⁹ P. Jonsson,⁴² A. Juste,⁴⁹ D. Käfer,²¹ S. Kahn,⁷⁰ E. Kajfasz,¹⁵ A. M. Kalinin,³⁵ J. Kalk,⁶³ D. Karmanov,³⁷ J. Kasper,⁶⁰ I. Katsanos,⁶⁷ D. Kau,⁴⁸ R. Kaur,²⁷ R. Kehoe,⁷⁶ S. Kermiche,¹⁵ S. Kesisoglou,⁷⁴ A. Khanov,⁷³ A. Kharchilava,⁵⁴ Y. M. Kharzeev,³⁵ H. Kim,⁷⁵ T. J. Kim,³¹ B. Klima,⁴⁹ J. M. Kohli,²⁷ J.-P. Konrath,²³ M. Kopal,⁷² V. M. Korablev,³⁸ J. Kotcher,⁷⁰ B. Kothari,⁶⁷ A. Koubarovsky,³⁷ A. V. Kozelov,³⁸ J. Kozminski,⁶³ A. Kryemadhi,⁷⁸ S. Krzywdzinski,⁴⁹ Y. Kulik,⁴⁹ A. Kumar,²⁸ S. Kunori,⁵⁹ A. Kupco,¹¹ T. Kurča,²⁰ J. Kvita,⁹ S. Lager,⁴⁰ N. Lahrichi,¹⁸ G. Landsberg,⁷⁴ J. Lazoflores,⁴⁸ A.-C. Le Bihan,¹⁹ P. Lebrun,²⁰ W. M. Lee,⁴⁸ A. Leflat,³⁷ F. Lehner,^{49,*} C. Leonidopoulos,⁶⁷ V. Lesne,¹³ J. Leveque,⁴⁴ P. Lewis,⁴² J. Li,⁷⁵ Q. Z. Li,⁴⁹ J. G. R. Lima,⁵¹ D. Lincoln,⁴⁹ S. L. Linn,⁴⁸ J. Linnemann,⁶³ V. V. Lipaev,³⁸ R. Lipton,⁴⁹ L. Lobo,⁴² A. Lobodenko,³⁹ M. Lokajicek,¹¹ A. Lounis,¹⁹ P. Love,⁴¹ H. J. Lubatti,⁷⁹ L. Lueking,⁴⁹ M. Lynker,⁵⁴ A. L. Lyon,⁴⁹ A. K. A. Maciel,⁵¹ R. J. Madaras,⁴⁵ P. Mättig,²⁶ C. Magass,²¹ A. Magerkurth,⁶² A.-M. Magnan,¹⁴ N. Makovec,¹⁶ P. K. Mal,²⁹ H. B. Malbouisson,³ S. Malik,⁶⁵ V. L. Malyshev,³⁵ H. S. Mao,⁶ Y. Maravin,⁴⁹ M. Martens,⁴⁹ S. E. K. Mattingly,⁷⁴ R. McCarthy,⁶⁹ R. McCroskey,⁴⁴ D. Meder,²⁴ A. Melnitchouk,⁶⁴ A. Mendes,¹⁵ L. Mendoza,⁸ M. Merkin,³⁷ K. W. Merritt,⁴⁹ A. Meyer,²¹ J. Meyer,²² M. Michaut,¹⁸ H. Miettinen,⁷⁷ J. Mitrevski,⁶⁷ J. Molina,³ N. K. Mondal,²⁹ J. Monk,⁴³ R. W. Moore,⁵ T. Moulik,⁵⁶ G. S. Muanza,²⁰ M. Mulders,⁴⁹ L. Mundim,³ Y. D. Mutaf,⁶⁹ E. Nagy,¹⁵ M. Naimuddin,²⁸ M. Narain,⁶⁰ N. A. Naumann,³⁴ H. A. Neal,⁶² J. P. Negret,⁸ S. Nelson,⁴⁸ P. Neustroev,³⁹ C. Noeding,²³ A. Nomerotski,⁴⁹ S. F. Novaes,⁴ T. Nunnemann,²⁵ E. Nurse,⁴³ V. O'Dell,⁴⁹ D. C. O'Neil,⁵ V. Oguri,³ N. Oliveira,³ N. Oshima,⁴⁹ G. J. Otero y Garzón,⁵⁰ P. Padley,⁷⁷ N. Parashar,⁵⁸ S. K. Park,³¹ J. Parsons,⁶⁷ R. Partridge,⁷⁴ N. Parua,⁶⁹ A. Patwa,⁷⁰ G. Pawloski,⁷⁷ P. M. Perea,⁴⁷ E. Perez,¹⁸ P. Pétroff,¹⁶

- M. Petteni,⁴² R. Piegaia,¹ M.-A. Pleier,⁶⁸ P. L. M. Podesta-Lerma,³² V. M. Podstavkov,⁴⁹ Y. Pogorelov,⁵⁴ M.-E. Pol,²
 A. Pompoš,⁷² B. G. Pope,⁶³ W. L. Prado da Silva,³ H. B. Prosper,⁴⁸ S. Protopopescu,⁷⁰ J. Qian,⁶² A. Quadt,²² B. Quinn,⁶⁴
 K. J. Rani,²⁹ K. Ranjan,²⁸ P. A. Rapidis,⁴⁹ P. N. Ratoff,⁴¹ S. Reucroft,⁶¹ M. Rijssenbeek,⁶⁹ I. Ripp-Baudot,¹⁹
 F. Rizatdinova,⁷³ S. Robinson,⁴² R. F. Rodrigues,³ C. Royon,¹⁸ P. Rubinov,⁴⁹ R. Ruchti,⁵⁴ V. I. Rud,³⁷ G. Sajot,¹⁴
 A. Sánchez-Hernández,³² M. P. Sanders,⁵⁹ A. Santoro,³ G. Savage,⁴⁹ L. Sawyer,⁵⁸ T. Scanlon,⁴² D. Schaile,²⁵
 R. D. Schamberger,⁶⁹ Y. Scheglov,³⁹ H. Schellman,⁵² P. Schieferdecker,²⁵ C. Schmitt,²⁶ C. Schwanenberger,²²
 A. Schwartzman,⁶⁶ R. Schwienhorst,⁶³ S. Sengupta,⁴⁸ H. Severini,⁷² E. Shabalina,⁵⁰ M. Shamim,⁵⁷ V. Shary,¹⁸
 A. A. Shchukin,³⁸ W. D. Shephard,⁵⁴ R. K. Shivpuri,²⁸ D. Shpakov,⁶¹ R. A. Sidwell,⁵⁷ V. Simak,¹⁰ V. Sirotenko,⁴⁹
 P. Skubic,⁷² P. Slattery,⁶⁸ R. P. Smith,⁴⁹ K. Smolek,¹⁰ G. R. Snow,⁶⁵ J. Snow,⁷¹ S. Snyder,⁷⁰ S. Söldner-Rembold,⁴³
 X. Song,⁵¹ L. Sonnenschein,¹⁷ A. Sopczak,⁴¹ M. Sosebee,⁷⁵ K. Soustruznik,⁹ M. Souza,² B. Spurlock,⁷⁵ N. R. Stanton,⁵⁷
 J. Stark,¹⁴ J. Steele,⁵⁸ K. Stevenson,⁵³ V. Stolin,³⁶ A. Stone,⁵⁰ D. A. Stoyanova,³⁸ J. Strandberg,⁴⁰ M. A. Strang,⁷⁵
 M. Strauss,⁷² R. Ströhmer,²⁵ D. Strom,⁵² M. Strovink,⁴⁵ L. Stutte,⁴⁹ S. Sumowidagdo,⁴⁸ A. Szajder,³ M. Talby,¹⁵
 P. Tamburello,⁴⁴ W. Taylor,⁵ P. Telford,⁴³ J. Temple,⁴⁴ M. Titov,²³ M. Tomoto,⁴⁹ T. Toole,⁵⁹ J. Torborg,⁵⁸ S. Towers,⁶⁹
 T. Trefzger,²⁴ S. Trincaz-Duvoid,¹⁷ D. Tsybychev,⁶⁹ B. Tuchming,¹⁸ C. Tully,⁶⁶ A. S. Turcot,⁴³ P. M. Tuts,⁶⁷ L. Uvarov,³⁹
 S. Uvarov,³⁹ S. Uzunyan,⁵¹ B. Vachon,⁵ P. J. van den Berg,³³ R. Van Kooten,⁵³ W. M. van Leeuwen,³³ N. Varelas,⁵⁰
 E. W. Varnes,⁴⁴ A. Vartapetian,⁷⁵ I. A. Vasilyev,³⁸ M. Vaupel,²⁶ P. Verdier,²⁰ L. S. Vertogradov,³⁵ M. Verzocchi,⁴⁹
 F. Villeneuve-Seguier,⁴² J.-R. Vlimant,¹⁷ E. Von Toerne,⁵⁷ M. Vreeswijk,³³ T. Vu Anh,¹⁶ H. D. Wahl,⁴⁸ L. Wang,⁵⁹
 J. Warchol,⁵⁴ G. Watts,⁷⁹ M. Wayne,⁵⁴ M. Weber,⁴⁹ H. Weerts,⁶³ N. Wermes,²² M. Wetstein,⁵⁹ A. White,⁷⁵ V. White,⁴⁹
 D. Wicke,⁴⁹ D. A. Wijngaarden,³⁴ G. W. Wilson,⁵⁶ S. J. Wimpenny,⁴⁷ M. Wobisch,⁴⁹ J. Womersley,⁴⁹ D. R. Wood,⁶¹
 T. R. Wyatt,⁴³ Y. Xie,⁷⁴ Q. Xu,⁶² N. Xuan,⁵⁴ S. Yacoob,⁵² R. Yamada,⁴⁹ M. Yan,⁵⁹ T. Yasuda,⁴⁹ Y. A. Yatsunenko,³⁵
 Y. Yen,²⁶ K. Yip,⁷⁰ H. D. Yoo,⁷⁴ S. W. Youn,⁵² J. Yu,⁷⁵ A. Yurkewicz,⁶⁹ A. Zabi,¹⁶ A. Zatserklyaniy,⁵¹ M. Zdrazil,⁶⁹
 C. Zeitnitz,²⁴ D. Zhang,⁴⁹ T. Zhao,⁷⁹ Z. Zhao,⁶² B. Zhou,⁶² J. Zhu,⁶⁹ M. Zielinski,⁶⁸ D. Ziemska,⁵³ A. Zieminski,⁵³
 R. Zitoun,⁶⁹ V. Zutshi,⁵¹ and E. G. Zverev³⁷

(D0 Collaboration)

¹Universidad de Buenos Aires, Buenos Aires, Argentina²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil⁴Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil⁵University of Alberta, Edmonton, Alberta, Canada,

Simon Fraser University, Burnaby, British Columbia, Canada,

York University, Toronto, Ontario, Canada,

and McGill University, Montreal, Quebec, Canada

⁶Institute of High Energy Physics, Beijing, People's Republic of China⁷University of Science and Technology of China, Hefei, People's Republic of China⁸Universidad de los Andes, Bogotá, Colombia⁹Center for Particle Physics, Charles University, Prague, Czech Republic¹⁰Czech Technical University, Prague, Czech Republic¹¹Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic¹²Universidad San Francisco de Quito, Quito, Ecuador¹³Laboratoire de Physique Corpusculaire, IN2P3-CNRS, Université Blaise Pascal, Clermont-Ferrand, France¹⁴Laboratoire de Physique Subatomique et de Cosmologie, IN2P3-CNRS, Université de Grenoble 1, Grenoble, France¹⁵CPPM, IN2P3-CNRS, Université de la Méditerranée, Marseille, France¹⁶IN2P3-CNRS, Laboratoire de l'Accélérateur Linéaire, Orsay, France¹⁷LPNHE, IN2P3-CNRS, Universités Paris VI and VII, Paris, France¹⁸DAPNIA/Service de Physique des Particules, CEA, Saclay, France¹⁹IReS, IN2P3-CNRS, Université Louis Pasteur, Strasbourg, France,

and Université de Haute Alsace, Mulhouse, France

²⁰Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude Bernard, Villeurbanne, France²¹III. Physikalisch Institut A, RWTH Aachen, Aachen, Germany²²Physikalisches Institut, Universität Bonn, Bonn, Germany²³Physikalisches Institut, Universität Freiburg, Freiburg, Germany²⁴Institut für Physik, Universität Mainz, Mainz, Germany²⁵Ludwig-Maximilians-Universität München, München, Germany²⁶Fachbereich Physik, University of Wuppertal, Wuppertal, Germany²⁷Panjab University, Chandigarh, India

- ²⁸*Delhi University, Delhi, India*
- ²⁹*Tata Institute of Fundamental Research, Mumbai, India*
- ³⁰*University College Dublin, Dublin, Ireland*
- ³¹*Korea Detector Laboratory, Korea University, Seoul, Korea*
- ³²*CINVESTAV, Mexico City, Mexico*
- ³³*FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands*
- ³⁴*Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands*
- ³⁵*Joint Institute for Nuclear Research, Dubna, Russia*
- ³⁶*Institute for Theoretical and Experimental Physics, Moscow, Russia*
- ³⁷*Moscow State University, Moscow, Russia*
- ³⁸*Institute for High Energy Physics, Protvino, Russia*
- ³⁹*Petersburg Nuclear Physics Institute, St. Petersburg, Russia*
- ⁴⁰*Lund University, Lund, Sweden, Royal Institute of Technology and Stockholm University, Stockholm, Sweden,
and Uppsala University, Uppsala, Sweden*
- ⁴¹*Lancaster University, Lancaster, United Kingdom*
- ⁴²*Imperial College, London, United Kingdom*
- ⁴³*University of Manchester, Manchester, United Kingdom*
- ⁴⁴*University of Arizona, Tucson, Arizona 85721, USA*
- ⁴⁵*Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA*
- ⁴⁶*California State University, Fresno, California 93740, USA*
- ⁴⁷*University of California, Riverside, California 92521, USA*
- ⁴⁸*Florida State University, Tallahassee, Florida 32306, USA*
- ⁴⁹*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*
- ⁵⁰*University of Illinois at Chicago, Chicago, Illinois 60607, USA*
- ⁵¹*Northern Illinois University, DeKalb, Illinois 60115, USA*
- ⁵²*Northwestern University, Evanston, Illinois 60208, USA*
- ⁵³*Indiana University, Bloomington, Indiana 47405, USA*
- ⁵⁴*University of Notre Dame, Notre Dame, Indiana 46556, USA*
- ⁵⁵*Iowa State University, Ames, Iowa 50011, USA*
- ⁵⁶*University of Kansas, Lawrence, Kansas 66045, USA*
- ⁵⁷*Kansas State University, Manhattan, Kansas 66506, USA*
- ⁵⁸*Louisiana Tech University, Ruston, Louisiana 71272, USA*
- ⁵⁹*University of Maryland, College Park, Maryland 20742, USA*
- ⁶⁰*Boston University, Boston, Massachusetts 02215, USA*
- ⁶¹*Northeastern University, Boston, Massachusetts 02115, USA*
- ⁶²*University of Michigan, Ann Arbor, Michigan 48109, USA*
- ⁶³*Michigan State University, East Lansing, Michigan 48824, USA*
- ⁶⁴*University of Mississippi, University, Mississippi 38677, USA*
- ⁶⁵*University of Nebraska, Lincoln, Nebraska 68588, USA*
- ⁶⁶*Princeton University, Princeton, New Jersey 08544, USA*
- ⁶⁷*Columbia University, New York, New York 10027, USA*
- ⁶⁸*University of Rochester, Rochester, New York 14627, USA*
- ⁶⁹*State University of New York, Stony Brook, New York 11794, USA*
- ⁷⁰*Brookhaven National Laboratory, Upton, New York 11973, USA*
- ⁷¹*Langston University, Langston, Oklahoma 73050, USA*
- ⁷²*University of Oklahoma, Norman, Oklahoma 73019, USA*
- ⁷³*Oklahoma State University, Stillwater, Oklahoma 74078, USA*
- ⁷⁴*Brown University, Providence, Rhode Island 02912, USA*
- ⁷⁵*University of Texas, Arlington, Texas 76019, USA*
- ⁷⁶*Southern Methodist University, Dallas, Texas 75275, USA*
- ⁷⁷*Rice University, Houston, Texas 77005, USA*
- ⁷⁸*University of Virginia, Charlottesville, Virginia 22901, USA*
- ⁷⁹*University of Washington, Seattle, Washington 98195, USA*

(Received 30 August 2005; published 4 January 2006)

We present a search for the standard model Higgs boson in $H \rightarrow WW^{(*)}$ decays with e^+e^- , $e^\pm\mu^\mp$, and $\mu^+\mu^-$ final states in $p\bar{p}$ collisions at a center-of-mass energy of $\sqrt{s} = 1.96$ TeV. The data, collected from April 2002 to June 2004 with the D0 detector, correspond to an integrated luminosity of $300\text{--}325\text{ pb}^{-1}$, depending on the final state. The number of events observed is consistent with the expectation from backgrounds. Limits from the combination of all three channels on the Higgs boson production cross section times branching ratio $\sigma \times \text{BR}(H \rightarrow WW^{(*)})$ are presented.

In the standard model (SM), the hypothetical Higgs boson is crucial to the understanding of electroweak symmetry breaking (EWSB) and the mass generation of electroweak gauge bosons and fermions. Spontaneous EWSB predicts the existence of this neutral scalar particle with mass M_H , a free parameter in the SM. Direct searches at the CERN e^+e^- collider (LEP) yield a lower limit for the Higgs boson mass of $M_H > 114.4$ GeV [1] at the 95% C.L. Indirect measurements via fits to the electroweak precision data give an upper bound of $M_H < 186$ GeV [2] at the 95% C.L.

In this Letter, we present a search for the Higgs boson in $H \rightarrow WW^{(*)} \rightarrow \ell\nu\ell'\nu'$ ($\ell, \ell' = e, \mu, \tau$) decays with e^+e^- , $e^\pm\mu^\mp$, or $\mu^+\mu^-$ final states. Tau decays are detected in their leptonic decay modes to electrons or muons. This is the first search for the Higgs boson at a hadron collider in this decay channel, which plays an important role in the overall discovery potential of the Higgs boson at the Fermilab Tevatron Collider [3]. We use data collected by the D0 detector between April 2002 and June 2004 in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV of the Fermilab Tevatron Collider. The integrated luminosities are 325 ± 21 , 318 ± 21 , and 299 ± 19 pb $^{-1}$ for the e^+e^- , $e^\pm\mu^\mp$, and $\mu^+\mu^-$ channels, respectively. The differences in the integrated luminosities for various channels are primarily due to different trigger conditions. Next-to-leading order (NLO) calculations [4,5] predict the product of the SM Higgs boson production cross section and the branching ratio $\sigma(p\bar{p} \rightarrow H) \times \text{BR}(H \rightarrow WW^{(*)})$ of 11–250 fb for the Higgs boson masses between 100 and 200 GeV. The dominant contribution to the cross section comes from gluon-gluon fusion. Extensions of the SM including a fourth fermion family [6] predict an enhanced Higgs boson production cross section.

The D0 detector and its data acquisition system are described in detail elsewhere [7]. We briefly describe the main components important to this analysis. The D0 detector has a magnetic central-tracking system, consisting of a silicon microstrip tracker and a central fiber tracker, both located within a 2 T superconducting solenoidal magnet. A liquid-argon and uranium calorimeter has a central section (CC) covering pseudorapidities $|\eta|$ up to ≈ 1.1 [$\eta = -\ln(\tan\theta)$ with polar angle θ], and two end calorimeters extending coverage to $|\eta| \approx 4.2$. A muon system resides beyond the calorimetry, and consists of a layer of tracking detectors and scintillation trigger counters before 1.8 T toroids, followed by two more similar layers after the toroids.

The $H \rightarrow WW^{(*)}$ candidates are selected by single or dilepton triggers using a three-level trigger system [7] and are recorded for off-line analysis. In the off-line analysis, electrons are identified as electromagnetic showers in the calorimeter. These showers are selected by comparing the

longitudinal and transverse shower profiles to those expected of the electrons. The showers must be isolated, deposit most of their energy in the electromagnetic part of the calorimeter, and pass a likelihood criterion that includes a spatial track match and, in the CC region, an E/p requirement, where E is the energy of the calorimeter cluster and p is the momentum of the track. All electrons are required to be in the pseudorapidity range $|\eta| < 3.0$. The transverse momentum measurement of the electrons is based on calorimeter cell and track information.

Muon tracks are reconstructed from hits in the wire chambers and scintillators in the muon system and must match a track in the central tracker. To select isolated muons, the scalar sum of the transverse momentum of all tracks other than that of the muon in a cone of $\mathcal{R} = 0.5$ around the muon track must be less than 4 GeV, where $\mathcal{R} = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2}$ and ϕ is the azimuthal angle. Muon detection is performed over the full coverage of the muon system $|\eta| < 2.0$. Muons from cosmic rays are rejected by requiring a timing criterion on the hits in the scintillator layers as well as applying restrictions on the position of the muon track with respect to the primary vertex.

The decay of the two W bosons into electrons or muons results in three different final states $e^+e^- + X$ (ee channel), $e^\pm\mu^\mp + X$ ($e\mu$ channel), and $\mu^+\mu^- + X$ ($\mu\mu$ channel), each of which consists of two oppositely charged isolated leptons with high transverse momentum and large missing transverse energy, \cancel{E}_T , due to the undetected neutrinos. The selection criteria for each channel were chosen to minimize the cross section upper limit on Higgs boson production expected in the absence of signal. To take into account the signal kinematic characteristics that change with the Higgs boson mass, M_H , some selection cuts are M_H dependent [8]. Six Higgs boson masses from 100 to 200 GeV have been studied.

In all three channels, two leptons originating from the same vertex are required to be of opposite charge, and must have transverse momenta $p_T > 15$ GeV for the leading lepton and $p_T > 10$ GeV for the trailing one (cut 1). Figure 1 shows the good agreement between data and Monte Carlo (MC) simulation in distributions of the azimuthal opening angle $\Delta\phi_{\ell\ell'}$ between the two leptons for the ee (a), the $\mu\mu$ (c), and the $e\mu$ channel (e) after applying the lepton transverse momentum cuts.

In all cases, the background is largely dominated by Z/γ^* production, which is further suppressed by requiring $\cancel{E}_T > 20$ GeV in all three channels (cut 2). Background events are also removed if the \cancel{E}_T has a large contribution from the mismeasurement of jet energy. The fluctuation in the measurement of jet energy in the transverse plane can be approximated by $\Delta E^{\text{jet}} \sin\theta^{\text{jet}}$ where ΔE^{jet} is proportional to $\sqrt{E^{\text{jet}}}$. The opening angle $\Delta\phi(\text{jet}, \cancel{E}_T)$ between




FIG. 1. Distribution of the opening angle $\Delta\phi_{\ell\ell'}$ after applying the initial transverse momentum cuts in the (a) ee , (c) $\mu\mu$, and (e) $e\mu$ channels. (b),(d),(f) The $\Delta\phi_{\ell\ell'}$ distributions after the final selection except for the $\Delta\phi_{\ell\ell'}$ criterion for the ee , $\mu\mu$, and $e\mu$ channels, respectively. The arrows indicate the cut values. The QCD contribution is negligible in (c) and (d).

this projected energy fluctuation and the missing transverse momentum provides a measure of the contribution of the jet to the missing transverse energy. The scaled missing transverse energy defined as

$$\not{E}_T^{Sc} = \frac{\not{E}_T}{\sqrt{\sum_{\text{jets}} [\Delta E^{\text{jet}} \sin\theta^{\text{jet}} \cos\Delta\phi(\text{jet}, \not{E}_T)]^2}} \quad (1)$$

is required to be greater than 15 (cut 3).

The charged lepton system and the neutrinos are emitted mostly back to back, so the invariant mass for the leptons from the Higgs decay is restricted to $M_H/2$. Thus, the invariant mass $m_{\ell\ell'}$ is required to be $m_{\ell\ell'} < M_H/2$ (cut 4). In the ee channel the cut is altered to $m_{ee} < \min(80 \text{ GeV}, M_H/2)$. In the $\mu\mu$ channel a lower cut boundary with $m_{\mu\mu} > 20 \text{ GeV}$ is required to remove events from J/ψ , Υ , and Z/γ^* production. The sum of the p_T of the leptons and \not{E}_T is required to be in the range

$M_H/2 + 20 \text{ GeV} < p_T^{\ell_1} + p_T^{\ell_2} + \not{E}_T < M_H$ for the ee and $e\mu$ channel and $M_H/2 + 10 \text{ GeV} < p_T^{\ell_1} + p_T^{\ell_2} + \not{E}_T < M_H$ for the $\mu\mu$ channel (cut 5). The transverse mass, defined as $m_T^{\ell\ell'} = \sqrt{2p_T^{\ell\ell'} \not{E}_T [1 - \cos\Delta\phi(p_T^{\ell\ell'}, \not{E}_T)]}$, with the dilepton transverse momentum $p_T^{\ell\ell'}$, should be in the range $M_H/2 < m_T^{\ell\ell'} < M_H - 10 \text{ GeV}$ (cut 6). The latter two cuts reject events from $W + \text{jet}/\gamma$ and WW production and further reduce backgrounds from Z/γ^* production. Finally, to suppress the background from $t\bar{t}$ production, the scalar sum of the transverse energies of all jets with $E_T^{\text{jet}} > 20 \text{ GeV}$ and $|\eta| < 2.5$, H_T , is required to be less than 100 GeV (cut 7). Remaining Z boson and multijet events can be rejected with a cut on the opening angle, $\Delta\phi_{\ell\ell'} < 2.0$ (cut 8), since most of the backgrounds exhibit a back-to-back topology. This is not the case for Higgs boson decays because of the spin correlations in the decay [9]. Figure 1 shows the distributions of the azimuthal opening angle $\Delta\phi_{\ell\ell'}$ between the two leptons for the ee (b), the $\mu\mu$ (d), and the $e\mu$ channel (f) before applying the final cut on $\Delta\phi_{\ell\ell'}$.

To maximize the sensitivity, the selection in the $\mu\mu$ channel is slightly changed for Higgs boson masses $M_H = 140$ and 160 GeV . For a better Z/γ^* background suppression cuts 4, 5, and 6 are replaced by the following cuts: the invariant mass $m_{\mu\mu}$ should be in the range $20 < m_{\mu\mu} < 80 \text{ GeV}$ (cut 4). Since the momentum resolution is degraded for high p_T tracks, an additional constrained fit is performed to reject events compatible with Z boson production (cut 5). The sum of the muon transverse momenta and the missing transverse energy should be $p_T^{\mu 1} + p_T^{\mu 2} + \not{E}_T > 90 \text{ GeV}$ (cut 6).

The efficiency for $H \rightarrow WW^{(*)} \rightarrow \ell\nu\ell'\nu'$ signal events to pass the acceptance and selection criteria is determined using the PYTHIA 6.2 [10] event generator followed by a detailed GEANT-based [11] simulation of the D0 detector. All trigger, reconstruction, and identification efficiencies are derived from the data. The kinematic acceptance efficiency is derived from MC simulations. The overall detection efficiencies range from $(0.44 \pm 0.03)\%$ to $(3.9 \pm 0.2)\%$ depending on the decay channel and M_H . Table I summarizes these efficiencies.

TABLE I. Overall detection efficiencies (in %) for $H \rightarrow WW^{(*)} \rightarrow \ell\nu\ell'\nu'$ decays for the three channels after all cuts. Quoted are the overall uncertainties, combining statistical and systematic components in quadrature.

$M_H (\text{GeV})$	ee	$e\mu$	$\mu\mu$
100	0.56 ± 0.05	1.02 ± 0.06	0.44 ± 0.03
120	1.18 ± 0.09	2.0 ± 0.1	1.02 ± 0.06
140	1.55 ± 0.08	2.9 ± 0.2	1.34 ± 0.08
160	2.1 ± 0.1	3.9 ± 0.2	2.0 ± 0.1
180	2.1 ± 0.1	3.9 ± 0.2	1.68 ± 0.09
200	1.57 ± 0.09	3.2 ± 0.1	1.53 ± 0.07

TABLE II. Number of signal and background events expected and number of events observed after all selections are applied. Only statistical uncertainties are given.

M_H (GeV)	100	120	140	160	180	200
$H \rightarrow WW^{(*)}$	0.007 ± 0.001	0.125 ± 0.002	0.398 ± 0.008	0.68 ± 0.01	0.463 ± 0.009	0.210 ± 0.004
Z/γ^*	7.9 ± 1.1	7.5 ± 1.0	3.8 ± 0.6	4.0 ± 0.7	6.6 ± 0.9	9.9 ± 1.1
Diboson	4.4 ± 0.2	8.1 ± 0.2	11.7 ± 0.3	12.3 ± 0.3	11.6 ± 0.3	9.6 ± 0.3
$t\bar{t}$	0.03 ± 0.01	0.11 ± 0.02	0.29 ± 0.02	0.47 ± 0.03	0.66 ± 0.05	0.72 ± 0.05
$W + \text{jet}/\gamma$	16.9 ± 2.2	14.2 ± 2.1	5.8 ± 1.2	2.8 ± 0.9	0.7 ± 0.5	0.7 ± 0.5
Multijet	0.6 ± 0.3	0.3 ± 0.1	0.2 ± 0.1	0.2 ± 0.1	0.3 ± 0.1	0.3 ± 0.1
Background sum	29.9 ± 2.5	30.1 ± 2.3	21.8 ± 1.4	19.7 ± 1.2	19.8 ± 1.1	21.2 ± 1.2
Data	27	21	20	19	19	14

Using the NLO cross sections calculated with HIGLU [5] and HDECAY [4] and the branching ratio BR of 0.1068 ± 0.0012 for $W \rightarrow \ell\nu$ [12], the expected number of events for $H \rightarrow WW^{(*)}$ decays from all three channels is $0.68 \pm 0.03(\text{syst}) \pm 0.04(\text{lum})$ events for a Higgs boson mass $M_H = 160$ GeV. The signal expectation for different Higgs masses M_H are given in Table II.

Background contributions from Z/γ^* , $W + \text{jet}/\gamma$, $t\bar{t}$, WW , WZ , and ZZ events are estimated using the PYTHIA event generator normalized to their NLO cross sections [13]. In addition, $W + \text{jet}/\gamma$ contributions are verified using ALPGEN [14]. All events are processed through the full detector simulation. The background due to multijet production, when a jet is misidentified as an electron, is determined from the data using a sample of like-sign dilepton events with inverted lepton quality cuts (called “QCD” in Fig. 1). A summary of the background contributions together with signal expectations and events observed in the data after the final selection is shown in Table II. There is good agreement between the number of events observed in the data and the various backgrounds in all three channels. The largest difference between the data and the background expectation, at $M_H = 120$ GeV, corresponds to a background probability of 6%. The $e\mu$ channel has both the highest signal efficiency and best signal-to-background ratio.

Various sources of systematic uncertainties that affect the background estimation and signal efficiencies have been studied [13]. In these calculations, parameters are varied within $\pm 1\sigma$ of their nominal values, where σ is determined by the corresponding uncertainties. The trigger efficiency, electron and muon identification efficiencies, jet energy scale, electron and muon momentum resolution, parton distribution function (PDF) uncertainty, and cross section calculation of Z/γ^* , WW , and $t\bar{t}$ events contribute to the systematic uncertainties. The total systematic uncertainties for the background estimate and signal efficiencies for the six Higgs boson masses are given in Table III. The largest contribution to the systematic uncertainty on the background for small Higgs boson masses comes from the jet energy scale due to the large $W + \text{jet}/\gamma$ background, whereas for high Higgs boson masses the W boson pair production cross section gives the largest systematic uncertainty. The uncertainty of the PDF is the largest uncertainty of the signal efficiency. Since the Higgs boson is dominantly produced via gluon-gluon fusion, the increased uncertainty introduced by the gluon structure function for large parton momenta is reflected in the PDF uncertainty. The uncertainty of the luminosity measurement is 6.5%.

Since the remaining candidate events after the selection are consistent with the background expectation, limits on the production cross section times branching ratio $\sigma \times \text{BR}(H \rightarrow WW^{(*)})$ are derived using a modified frequentist method described in Ref. [15]. The uncertainty on the background and the expected signal events were determined from the statistical and systematic uncertainties and luminosity uncertainty. Table IV shows the individual

TABLE III. Systematic uncertainties (in %) of the $H \rightarrow WW^{(*)}$ signal efficiencies and of the number of background events (BG), for the ee , $e\mu$, and $\mu\mu$ channels. Uncertainties exclude the uncertainty of the luminosity measurement.

M_H (GeV)	ee		$e\mu$		$\mu\mu$	
	Signal	BG	Signal	BG	Signal	BG
100	8.3	9.5	6.4	11.4	7.8	7.2
120	8.3	8.6	6.7	13.6	7.3	7.5
140	6.4	6.7	6.9	13.6	7.2	8.3
160	6.6	7.3	6.7	12.0	7.1	8.3
180	6.9	10.3	6.6	13.0	7.3	14.6
200	6.8	10.6	6.1	12.3	6.9	18.1

TABLE IV. Expected and observed upper limits at the 95% C.L. for the Higgs boson production cross section times branching ratio $\sigma \times \text{BR}(H \rightarrow WW^{(*)})$ for six values of M_H .

M_H (GeV)	100	120	140	160	180	200
Expected limits (pb)	20.3	9.5	5.9	4.0	3.9	4.5
Observed limits (pb)	18.5	5.6	4.9	3.7	4.1	3.2




FIG. 2 (color online). Expected and observed upper limits on the cross section times branching ratio $\sigma \times \text{BR}(H \rightarrow WW^{(*)})$ at the 95% C.L. together with expectations from standard model Higgs boson production and an alternative model. The LEP limit on the standard model Higgs boson production is taken from [1] and the 4th generation model prediction is described in [6].

expected and observed upper limits on the cross section times branching ratio $\sigma \times \text{BR}(H \rightarrow WW^{(*)})$ for the combination of the three different decay channels for six different Higgs boson masses. The different values of the upper limits are due to different background expectations and signal efficiencies. The best limits are achieved for large Higgs masses since background expectations decrease while signal efficiencies increase. The reasonable agreement between the observed and expected limits shows the reliability of the observed limit, as well as that the data contain no evidence for a signal.

Figure 2 shows the expected and observed cross section limits for $\sigma \times \text{BR}(H \rightarrow WW^{(*)})$ for the different Higgs boson masses compared with predictions from the SM and from an extension including a fourth fermion family [6]. With the current data set, no region of the SM prediction can be excluded.

To conclude, we have searched for the Higgs boson in $H \rightarrow WW^{(*)} \rightarrow \ell\nu\ell'\nu'$ ($\ell, \ell' = e, \mu, \tau$) decays with e^+e^- , $e^\pm\mu^\mp$, or $\mu^+\mu^-$ final states in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV. The data are consistent with the expectation from backgrounds. Since no excess has been observed, limits on the production cross section times branching ratio $\sigma \times \text{BR}(H \rightarrow WW^{(*)})$ have been derived.

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom, and RFBR (Russia); CAPES, CNPq, FAPERJ, FAPESP, and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); PPARC (United Kingdom); MSMT (Czech Republic); CRC Program, CFI, NSERC, and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); Research Corporation, Alexander von Humboldt Foundation, and the Marie Curie Program.

*Permanent address: University of Zurich, Zurich, Switzerland.

- [1] R. Barate *et al.*, Phys. Lett. B **565**, 61 (2003).
- [2] The LEP Collaborations ALEPH, DELPHI, L3, OPAL, the LEP Electroweak Working Group, the SLD Electroweak and Heavy Flavour Groups, hep-ex/0412015 and hep-ex/0509008, updated for the 2005 summer conferences.
- [3] M. Carena *et al.*, hep-ph/0010338.
- [4] A. Djouadi *et al.*, Comput. Phys. Commun. **108**, 56 (1998).
- [5] M. Spira, Report No. DESY T-95-05 (1995); hep-ph/9510347.
- [6] E. Arik *et al.*, Phys. Rev. D **66**, 033003 (2002); O. Cakir and S. Sultansoy, Phys. Rev. D **65**, 013009 (2002).
- [7] V. Abazov *et al.* (D0 Collaboration), physics/0507191 [Nucl. Instrum. Methods Phys. Res., Sect. A (to be published)].
- [8] T. Han *et al.*, Phys. Rev. D **59**, 093001 (1999).
- [9] M. Dittmar *et al.*, Phys. Rev. D **55**, 167 (1997).
- [10] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001).
- [11] R. Brun and F. Carminati, CERN Program Library Long Writeup W5013, 1993 (unpublished).
- [12] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. B **592**, 1 (2004).
- [13] V. M. Abazov *et al.*, Phys. Rev. Lett. **94**, 151801 (2005).
- [14] M. L. Mangano *et al.*, J. High Energy Phys. 07 (2003) 001.
- [15] T. Junk, Nucl. Instrum. Methods Phys. Res., Sect. A **434**, 435 (1999).