

## Resonant Tunneling in Truly Axial Symmetry $Mn_{12}$ Single-Molecule Magnets: Sharp Crossover between Thermally Assisted and Pure Quantum Tunneling

W. Wernsdorfer,<sup>1</sup> M. Murugesu,<sup>2</sup> and G. Christou<sup>2</sup>

<sup>1</sup>Laboratoire L. Néel, associé à l'UJF, CNRS, BP 166, 38042 Grenoble Cedex 9, France

<sup>2</sup>Department of Chemistry, University of Florida, Gainesville, Florida 32611-7200, USA

(Received 19 August 2005; published 8 February 2006)

Magnetization measurements of a truly axial symmetry  $Mn_{12}$ -*t*BuAc molecular nanomagnet with a spin ground state of  $S = 10$  show resonant tunneling. This compound has the same magnetic anisotropy as  $Mn_{12}$ -Ac but the molecules are better isolated and the crystals have less disorder and a higher symmetry. Hysteresis loop measurements at several temperatures reveal a well-resolved step fine structure which is due to level crossings of excited states. All step positions can be modeled by a simple spin Hamiltonian. The results establish a sharp crossover between thermally assisted and pure quantum tunneling, as had been previously predicted.

DOI: 10.1103/PhysRevLett.96.057208

PACS numbers: 75.50.Xx, 75.60.Jk, 75.75.+a, 76.30.-v

$[Mn_{12}O_{12}(O_2CCH_3)_{16}(H_2O)_4] \cdot 2CH_3CO_2H \cdot 4H_2O$ ,  $Mn_{12}$ -Ac for short, was the first single-molecule magnet (SMM), exhibiting slow magnetization relaxation of its  $S = 10$  spin ground state which is split by axial zero-field splitting [1]. It was also the first system to show thermally assisted tunneling of magnetization [2,3]. During the last several years, many more SMMs have been discovered and they are now among the most promising candidates for observing the limits between classical and quantum physics since they have a well-defined structure, spin ground state, and magnetic anisotropy [4–7]. Nevertheless,  $Mn_{12}$ -Ac is still the most widely studied SMM [8–21]. While a rough understanding of the quantum phenomena in  $Mn_{12}$ -Ac was clear from the early studies, a detailed understanding has not yet emerged. For example, current theoretical models assume the presence of quadratic and quartic transverse crystal-field interactions in the spin Hamiltonian, where the former has been ascribed to solvent disorder [20]. However, these interactions, which contain only even powers of the raising and lowering operators, do not provide an explanation for the observation of odd tunneling steps in the hysteresis loops. It has been proposed that easy-axis tilting might give the missing odd transverse interactions [17]. Although such solvent disorder induced tilts exist, the tilt values are still unclear [22]. Hyperfine, dipolar, and Dzyaloshinsky-Moriya interactions were also proposed to be responsible for odd transverse terms [8,9,12].

Other theoretical and experimental studies concern the crossover between thermally assisted and pure quantum tunneling [10–14,21,23]. The crossover can occur in a narrow temperature interval with the field at which the system crosses the anisotropy barrier shifting abruptly with temperature, or the crossover can occur in a broad interval of temperature with this field changing smoothly [14,21,23]. The first studies, published on  $Mn_{12}$ -Ac, gave different answers: [10,11] were in favor of theory and [12] against it. However, later studies showed that significant

distributions of molecular environments are present in  $Mn_{12}$ -Ac crystals [13,20], which render such a detailed study impossible on this compound.

We present here resonant quantum tunneling measurements of a recently discovered analog of  $Mn_{12}$ -Ac, namely,  $[Mn_{12}O_{12}(O_2CCH_2Bu^t)_{16}(CH_3OH)_4] \cdot CH_3OH$ , called  $Mn_{12}$ -*t*BuAc henceforth. We show that this compound has the same magnetic anisotropy as  $Mn_{12}$ -Ac but the molecules are better isolated and the crystals contain less disorder and a higher symmetry. Hysteresis loop measurements at several temperatures reveal a fine structure of steps which is due to the dominating energy level crossings. All step positions can be modeled by a simple spin Hamiltonian. Our results establish a sharp crossover between thermally assisted and pure quantum tunneling, which has been predicted [14,21,23].

$[Mn_{12}O_{12}(O_2CCH_2Bu^t)_{16}(H_2O)_4]$  was prepared by the carboxylate substitution procedure described elsewhere [24], and crystallizes in triclinic space group  $P\bar{1}$  [25]. However, recrystallization from mixed  $CH_3OH/Et_2O$  solvent gave  $Mn_{12}$ -*t*BuAc in tetragonal space group  $I\bar{4}$ . The ground state spin  $S = 10$  was established by magnetization measurements [26]. The molecular structure of  $Mn_{12}$ -*t*BuAc is very similar to that of  $Mn_{12}$ -Ac except that the acetate (Ac) and  $H_2O$  groups have been replaced by  $Bu^tCH_2CO_2$  (*t*BuAc) and  $CH_3OH$  groups, respectively. The increased bulk of the  $Bu^tCH_2CO_2$  groups leads to a unit cell volume for  $Mn_{12}$ -*t*BuAc ( $7.06 \text{ nm}^3$ ) that is almost double that of  $Mn_{12}$ -Ac ( $3.72 \text{ nm}^3$ ), and thus to greater intermolecular separations and decreased intermolecular interactions relative to  $Mn_{12}$ -Ac. In addition, the interstitial  $CH_3OH$  solvent molecules in  $Mn_{12}$ -*t*BuAc are not disordered and neither are they hydrogen-bonding with the  $Mn_{12}$  molecules. As a result, the site symmetry of the latter in  $Mn_{12}$ -*t*BuAc is truly axial with a small distribution of environments. This is in stark contrast to  $Mn_{12}$ -Ac where each of the acetic acid ( $CH_3CO_2H$ ) molecules in the crystal forms a strong  $OH \cdots H$  hydrogen-bond with a  $Mn_{12}$

molecule but will do so with only one of the two  $\text{Mn}_{12}$  molecules next to it. Since each  $\text{Mn}_{12}$  molecule is surrounded by four  $\text{CH}_3\text{CO}_2\text{H}$  molecules, this disorder in the acetic acid orientation leads to the  $\text{Mn}_{12}$  molecules in  $\text{Mn}_{12}\text{-Ac}$  being hydrogen-bonded with  $k$   $\text{CH}_3\text{CO}_2\text{H}$  molecules ( $k = 0-4$ ), with the  $k = 2$  situation also having two possibilities (the two  $\text{CH}_3\text{CO}_2\text{H}$  attached *cis* (adjacent) or *trans* (opposite) about the  $\text{Mn}_{12}$  molecule). Thus, although the  $\text{Mn}_{12}\text{-Ac}$  crystal possesses a crystallographic average symmetry of  $I\bar{4}$ , it contains a mixture of  $\text{Mn}_{12}$  molecules in six different hydrogen-bonded forms [20], and only two of these forms possess axial  $S_4$  site symmetry, the  $k = 0$  and 4 forms. The other forms have lower (rhombic) symmetry. As a result,  $\text{Mn}_{12}\text{-Ac}$  crystals contain  $\text{Mn}_{12}$  molecules with a wide distribution of environments. Consequently, crystals of  $\text{Mn}_{12}\text{-}t\text{BuAc}$  are concluded to be far superior to those of  $\text{Mn}_{12}\text{-Ac}$  for detailed studies. Finally, the axes of the tetragonal site symmetry of each molecule coincide with those of the unit cell, which is not the case for  $\text{Mn}_{12}\text{-Ac}$ .




FIG. 1 (color online). Hysteresis loops of single crystals of (a)–(b)  $\text{Mn}_{12}\text{-}t\text{BuAc}$  and (c)  $\text{Mn}_{12}\text{-Ac}$  molecular clusters at different temperatures and a constant field sweep rate indicated in the figure. The data in (c) were taken from [12]. The loops display a series of steps, separated by plateaus. As the temperature is lowered, there is a decrease in the transition rate due to reduced thermally assisted tunneling. The hysteresis loops become temperature independent below 0.6 K, demonstrating quantum tunneling at the lowest energy levels.

The magnetization measurements were performed by using a magnetometer consisting of a micro-Hall bar [27] on top of which a single crystal of  $\text{Mn}_{12}\text{-}t\text{BuAc}$  was placed. The field was aligned with the easy axis of magnetization using the transverse field method [22]. The applied field  $H_z$  was corrected because the determination of the resonance fields must take into account the internal magnetic field [28].

Figure 1 shows the temperature dependence of the hysteresis loops of  $\text{Mn}_{12}\text{-}t\text{BuAc}$  and  $\text{Mn}_{12}\text{-Ac}$  SMMs. The loops display a series of steps, separated by plateaus. As the temperature is lowered, the hysteresis increases because there is a decrease in the transition rate of thermal assisted tunneling [2,3]. The hysteresis loops become temperature independent below 0.6 K, demonstrating quantum tunneling at the lowest energy levels [10–12]. It is important to note that the loops remain strongly sweep rate dependent below 0.6 K (Fig. 2). Apart from the major steps, these hysteresis loops reveal fine structure in the thermally activated regime which is also strongly sweep rate dependent (Fig. 2). This fine structure was first observed for  $\text{Mn}_{12}\text{-Ac}$  [10–12], see Fig. 1(c), but it is much clearer for  $\text{Mn}_{12}\text{-}t\text{BuAc}$  [Figs. 1(a), 1(b), and 2]. A convenient way of determining the field positions of the steps is to plot the derivative of the magnetization with respect to the applied field (Fig. 3). The step positions, that is the maxima of the relaxation rate, are given by the peaks on the  $dM/dH$  plot.

The simplest model describing the low-temperature spin dynamics of  $\text{Mn}_{12}\text{-}t\text{BuAc}$  has the following spin Hamiltonian

$$\mathcal{H} = -DS_z^2 - BS_z^4 - g_z \mu_B \mu_0 S_z H_z + \mathcal{H}_{\text{trans}}, \quad (1)$$

where  $S_x$ ,  $S_y$ , and  $S_z$  are the three components of the spin operator,  $D$  and  $B$  are the anisotropy constants, the third term is the Zeeman energy associated with an applied field  $H_z$ , and the last term ( $\mathcal{H}_{\text{trans}}$ ) describes small transverse terms containing  $S_x$  and  $S_y$  spin operators. Although




FIG. 2 (color online). Hysteresis loops of single crystals of  $\text{Mn}_{12}\text{-}t\text{BuAc}$  at several field sweep rates and at 2, 0.85, and 0.1 K. The field sweep rates from top to bottom are 0.2, 0.5, 1, 2, 4, 8, 17, 35, and 70 mT/s.




FIG. 3 (color online). (a)–(b) Field derivative of the hysteresis loops of single crystals of  $\text{Mn}_{12}\text{-}t\text{BuAc}$  at different temperatures. The applied field was swept from  $-6$  to  $6$  T at a constant field sweep rate of  $2$  mT/s. Resonant quantum tunneling of magnetization occurs at the peaks of  $dM/dH$ . The corresponding level crossings are labeled with two indexes  $(n:p)$ .

$\mathcal{H}_{\text{trans}}$  produces tunneling, it can be neglected when determining the field positions of the level crossing because it is much smaller than the axial terms. Without  $\mathcal{H}_{\text{trans}}$ , the Hamiltonian is diagonal and the field dependence of the energy levels can be calculated analytically (Fig. 4). The energy level spectrum with  $(2S + 1) = 21$  values can be labeled by the quantum numbers  $m = -10, -9, \dots, 10$ . At  $\vec{H} = 0$ , the levels  $m = \pm 10$  have the lowest energy. When a field  $H_z$  is applied, the energy levels with  $m < 0$  increase, while those with  $m > 0$  decrease (Fig. 4). Therefore, energy levels of positive and negative quantum numbers cross at certain fields. The field position of the crossing of level  $m = -S + p$  with  $m' = S - n - p$  is given by

$$H_{(n:p)} = \frac{n[D + B((-S + p)^2 + (S - n - p)^2)]}{g_z \mu_B \mu_0}, \quad (2)$$

where  $n = -(m + m')$  is the step index and  $p = S + m$  labels the excited states ( $p = 0$  for the ground state,  $p = 1$  for the first excited state, etc.).




FIG. 4 (color online). Zeeman diagram of the 21 levels of the  $S = 10$  manifold of  $\text{Mn}_{12}$  as a function of the field  $H_z$  applied along the easy axis. At  $H_z = 0$ , from bottom to top, the levels are labeled with quantum numbers  $m = \pm 10, \pm 9, \dots, 0$ . The resonant quantum tunneling steps which lead to a step height larger than  $0.02 M_S$  at  $2$  mT/s occur at the indicated level crossings which are labeled with two indexes  $(n:p)$ .

The step positions  $H_{(n:p)}$ , determined from Fig. 3 and which have a step height larger than  $0.03 M_S$  at  $2$  mT/s, are shown in Fig. 5. The horizontal lines indicate the calculated energy level crossing fields using Eq. (2) with  $D = 0.563$  K,  $B = 1.2$  mK, and  $g_z = 2$  where the latter was measured by EPR [29]. These values are very close to those of  $\text{Mn}_{12}\text{-Ac}$  establishing that the magnetic cores of both molecules are comparable. Because the resonance fields of all avoided level crossings are well resolved,  $\text{Mn}_{12}\text{-}t\text{BuAc}$  allows the study of the crossover between thermally as-




FIG. 5 (color online). Temperature dependence of the peak positions of  $dM/dH$  in Fig. 3 at  $2$  mT/s. The horizontal lines indicate the calculated energy level crossing fields. The largest step for each  $n$  are solid dots whereas the others are open dots or crosses for step heights larger or smaller than  $0.03 M_S$ , respectively.




FIG. 6 (color online). Details of hysteresis loops at step  $n = 8$  measured in the presence of four transverse fields  $H_{tr}$  and at temperatures of 0.1, 0.8, 0.9, 0.95, 1.0, 1.05, 1.1, 1.15, and 1.2 K from bottom to top. The field was ramped from  $-6$  to  $6$  T at a rate of  $17$  mT/s. The dominating steps are indicated.

sisted and pure quantum tunneling. The dominant field steps for each step index  $n$  are shown in Fig. 5 by solid circles. Whereas the crossover is gradual for  $n = 9$  and  $10$ , a clear step is seen for  $n = 8$ . Indeed, the quantum step for (8:1) is for all temperatures smaller than either (8:0) or (8:2). Similar results are found for  $n = 7$  and  $6$ . For  $n \leq 5$ , the crossover goes directly from nonmeasurable steps to finite ones with  $p \geq 3$ . Note that these sharp transitions have not been observed in  $\text{Mn}_{12}\text{-Ac}$  [10–12] because significant distributions of molecular environments render such a detailed study impossible [13,20]. It has been predicted that a sharp crossover can be smoothed out by applying a transverse field [14]. This can indeed be observed in  $\text{Mn}_{12}\text{-tBuAc}$ . Figure 6 shows that about  $0.14$  T sufficiently increases the tunnel rate to smooth out the transition. Similar results were found for  $n < 8$ .

Concerning the question of odd tunneling steps, mentioned in the introduction, the fact that both  $\text{Mn}_{12}\text{-Ac}$  and  $\text{Mn}_{12}\text{-tBuAc}$  show such steps in the hysteresis loops suggests that dipolar coupling [8], easy-axis tilting [17], and distributions of molecular environments [20] are not the major contributions. However, asymmetric exchange due to spin-orbit (SO) interactions might be responsible for such tunnel transitions [12]. Indeed, SO interactions are the major source of tunneling in, for example, Co and Ni based molecular systems.

In conclusion, resonant tunneling measurements on a new high symmetry  $\text{Mn}_{12}\text{-tBuAc}$  molecular nanomagnet show levels of detail not possible with  $\text{Mn}_{12}\text{-Ac}$ , as a result of the much less disorder and higher symmetry in the crystals of the former. This has permitted an unprecedented level of analysis of the data to be accomplished, resulting in information not attainable with  $\text{Mn}_{12}\text{-Ac}$ . The crossover between thermally assisted and pure quantum tunneling

can be easily explored and is found to be abrupt or gradual depending on the magnitude and orientation of the applied field. Simulation of the data allows  $D$  and  $B$  to be directly obtained.

This work was supported by the EC-TMR Network QuEMolNa (MRTN-CT-2003-504880), CNRS, Rhône-Alpes funding, and NSF.

- [1] R. Sessoli *et al.*, J. Am. Chem. Soc. **115**, 1804 (1993); R. Sessoli *et al.*, Nature (London) **365**, 141 (1993).
- [2] J.R. Friedman, M.P. Sarachik, J. Tejada, and R. Ziolo, Phys. Rev. Lett. **76**, 3830 (1996).
- [3] L. Thomas *et al.*, Nature (London) **383**, 145 (1996).
- [4] C. Sangregorio *et al.*, Phys. Rev. Lett. **78**, 4645 (1997).
- [5] S.M.J. Aubin *et al.*, J. Am. Chem. Soc. **120**, 839 (1998).
- [6] A. Caneschi *et al.*, J. Magn. Magn. Mater. **200**, 182 (1999).
- [7] J. Yoo *et al.*, Inorg. Chem. **39**, 3615 (2000).
- [8] F. Luis *et al.*, Phys. Rev. B **57**, 505 (1998).
- [9] N.V. Prokof'ev and P.C.E. Stamp, Phys. Rev. Lett. **80**, 5794 (1998).
- [10] A.D. Kent *et al.*, Europhys. Lett. **49**, 521 (2000).
- [11] L. Bokacheva, A.D. Kent, and M.A. Walters, Phys. Rev. Lett. **85**, 4803 (2000).
- [12] I. Chiorescu *et al.*, Phys. Rev. Lett. **85**, 4807 (2000).
- [13] K.M. Mertes *et al.*, Phys. Rev. Lett. **87**, 227205 (2001).
- [14] D.A. Garanin and E.M. Chudnovsky, Phys. Rev. B **65**, 094423 (2002).
- [15] M. Dressel, B. Gorshunov, K. Rajagopal, S. Vongtragool, and A.A. Mukhin, Phys. Rev. B **67**, 060405(R) (2003).
- [16] M. Bal *et al.*, Phys. Rev. B **70**, 140403 (2004).
- [17] S. Takahashi *et al.*, Phys. Rev. B **70**, 094429 (2004).
- [18] K. Park, T. Baruah, N. Bernstein, and M.R. Pederson, Phys. Rev. B **69**, 144426 (2004).
- [19] F. Luis, V. González, A. Millán, and J.L. García-Palacios, Phys. Rev. Lett. **92**, 107201 (2004).
- [20] A. Cornia, R. Sessoli, L. Sorace, D. Gatteschi, A.L. Barra, and C. Daiguebonne, Phys. Rev. Lett. **89**, 257201 (2002).
- [21] E.M. Chudnovsky and D.A. Garanin, Phys. Rev. Lett. **79**, 4469 (1997); D.A. Garanin, X. Martínez Hidalgo, and E.M. Chudnovsky, Phys. Rev. B **57**, 13 639 (1998).
- [22] W. Wernsdorfer, N.E. Chakov, and G. Christou, Phys. Rev. B **70**, 132413 (2004).
- [23] Gwang-Hee Kim, Phys. Rev. B **59**, 11 847 (1999); H.J.W. Mller-Kirsten *et al.*, Phys. Rev. B **60**, 6662 (1999).
- [24] M. Soler *et al.*, Inorg. Chem. **40**, 4902 (2001).
- [25] M. Soler *et al.*, Chem. Commun. (Cambridge) 2672 (2003).
- [26] M. Murugesu *et al.* (to be published).
- [27] K. Petukhov *et al.*, Phys. Rev. B **72**, 052401 (2005).
- [28] The field  $H_z$  seen by a spin is roughly the sum of the applied field  $H_{\text{appl}}$  and the demagnetization field of the crystal. The latter can be determined by sweeping the applied field back and forth over a resonance transition. We found a nearly linear dependence which yields the following field correction:  $\mu_0 H_z = \mu_0 H_{\text{appl}} + 0.04 \text{ T } M/M_S$ .
- [29] S. Hill *et al.*, J. Appl. Phys. **97**, 10M510 (2005).