

Stability of Quantum Breathers

L. S. Schulman* and D. Tolkunov

Physics Department, Clarkson University, Potsdam, New York 13699-5820, USA

E. Mihokova

Institute of Physics, Academy of Sciences of the Czech Republic, Cukrovarnická 10, 162 53 Prague 6, Czech Republic

(Received 14 February 2005; published 13 February 2006)

Using two methods we show that a quantized discrete breather in a 1D lattice is stable. One method uses path integrals and compares correlations for a (linear) local mode with those of the quantum breather. The other takes a local mode as the zeroth order system relative to which numerical, cutoff-insensitive diagonalization of the Hamiltonian is performed.

DOI: [10.1103/PhysRevLett.96.065501](https://doi.org/10.1103/PhysRevLett.96.065501)

PACS numbers: 31.15.Gy, 05.45.-a, 31.70.Hq, 63.20.Pw

Discrete breathers [1] are nondispersive classical excitations that are known to be significant in a variety of natural systems. Recent reviews [2,3] give examples and references. The quantum theory of (discrete) breathers does not yield easily to numerical simulation and is not as fully developed.

Over the past decade there has developed a body of literature [4] in which it is claimed that unlike classical breathers, the quantum version cannot in principle be stable. This conclusion is reached by considering the effect on the phonon field of the rapidly oscillating particle at the core of the breather. Lifetimes have been calculated, and for alkali halides are about 10 ns. Other work, also treating the breather as an external classical force, [5], challenges this estimate and suggests stability in the limit of large systems.

A fully quantum result [6] found eigenstates of a model Hamiltonian with a dropoff in correlation function that is strongly suggestive of breathers, although translational invariance prevents complete localization. As recognized in [6] (and below), such approaches require cutoffs which may affect the conclusion.

In this Letter we make a two-pronged attack. We do a numerical diagonalization, similar to that of [6], incorporating methods to reduce cutoff effects and to provide direct evidence of localization. In addition, we do a path integral calculation, treating the breather fully quantum mechanically, similar to Feynman's polaron [7]. The approximations are different: in one case a cutoff, in the other semiclassical asymptotics.

The Hamiltonian in [6] is

$$H = \sum_{k=0}^N \left\{ \frac{1}{2} p_k^2 + \frac{1}{2} \omega_s^2 x_k^2 + \frac{1}{2} \omega_0^2 (\Delta x)_k^2 + \frac{1}{4} \lambda x_k^4 \right\}, \quad (1)$$

with $x_0 \equiv x_{N+1}$, $(\Delta x)_k \equiv x_k - x_{k+1}$, and $x_k \in \mathbb{R}$. We also consider interactions of the form $\sum (\Delta x)_k^4$, which resembles our own model [8]. An important simplification is to remove all nonlinear interactions except those of particle 0, the one with large amplitude motion. This strategy has been adopted by other authors and we checked whether

this affects the general properties of the classical breather. It does not [9], a consequence of the remarkable breather property, namely, that a central atom vibrates strongly, while its neighbors hardly move (so that nonlinear forces are negligible for them). Claims that the breather decays are based on this same approximation.

We emphasize the reasoning leading from a translationally invariant breather to what is effectively a nonlinear local mode. Although quantum tunneling requires that the breathers form a band, information on the classical breather corresponding to our excitation—including in the presence of moderate noise—indicates that tunneling is small, allowing us to drop translational invariance. This perspective has been adopted by others as well [4,5]. Having done this, we go beyond previous work by quantizing the breather atom itself. In other approaches one has *de facto* a local mode, interacting with the lattice. We retain the nonlinear interactions (implicitly), while also quantizing the principal breather atom.

In both our approaches we make use of a *linear local mode*. If in (the modified) Eq. (1), one replaces $\lambda x_0^4/4$ by $\omega_1^2 x_0^2/2$ with large ω_1 , the resulting system is well understood both classically and quantum mechanically. You have both classical confinement and quantum stability. Our use of the local mode parallels others' treatment of the quantum breather as a phonon field in the presence of rapid classical oscillations. The impact of the breather *cannot* be considered small. They treat that impact by means of a classical oscillating field. We use the quantum local mode. For the path integral, that use is the comparison with local mode correlation functions. For the diagonalization scheme we perturb around that mode.

Path integral approach.—The path integral allows the elimination of quadratic degrees of freedom at the expense of introducing nonlocal self coupling. Dropping quartic terms for all but x_0 , the Lagrangian is

$$\mathcal{L} = \sum_{n=0}^N \{ \dot{x}_n^2 - \omega_0^2 (\Delta x)_n - \omega_s^2 x_n^2 \} / 2 - \lambda x_0^4 / 4 - \mu x_0 (x_m + x_{N+1-m}). \quad (2)$$

The fictitious coupling, $\mu x_0(x_m + x_{N+1-m})$, allows the study of localization for m far from 0. The derivative (at $\mu = 0$) of the propagator provides a $(0-m)$ correlation. By standard methods [7,10,11] we path integrate degrees of freedom 1 to N , which form a *chain* with known normal modes. These modes see a forcing term from the so-far-unintegrated $x_0(t)$. The propagator, G , is a function of the end points of *all* x_k . One takes the matrix element of G in the chain ground state (à la [7]) and divides by corresponding free chain matrix elements. The result is

$$\mathcal{G}(q^f, T; q^i, 0) = \int \mathcal{D}q e^{i/\hbar(S_0 + S_{\text{eff}})}, \quad (3)$$

where $q \equiv x_0$, the action S_0 arises from the original \mathcal{L} sans chain terms, and S_{eff} comes from the integration over chain degrees of freedom. Specifically

$$S_0 = \int dt \{ \dot{q}^2/2 - (\omega_s^2/2 + \omega_0^2)q^2 - \lambda q^4/4 \}, \quad (4)$$

$$S_{\text{eff}} = \int_0^T \int_0^T dt ds K(|t-s|) q(t) q(s), \quad (5)$$

with

$$K(u) = \sum_{n=1,3,\dots}^N \tau_n^2 \frac{\cos \Omega_n (T/2 - u)}{\Omega_n \sin(\frac{\Omega_n T}{2})}, \quad (6)$$

$$\tau_n^2 \equiv \frac{2}{N+1} \left[\omega_0^2 \sin \frac{\pi n}{N+1} + \mu \sin \frac{\pi n m}{N+1} \right]^2, \quad (7)$$

and $\Omega_n^2 = \omega_s^2 + 4\omega_0^2 \sin^2[n\pi/2(N+1)]$ is the spectrum of the chain.

We study \mathcal{G} in the stationary phase approximation; that is, we evaluate $S_0 + S_{\text{eff}}$ along extremal “classical paths.” In principle, one can get the ground state energy by going to large imaginary times, for which $\mathcal{G}(q, -iT; q) \sim |\phi(q)|^2 \exp(-TE_0)$. In practice, $S \rightarrow \text{const}$ and $\partial^2 S / \partial q^i \partial q^f$ gives an exponentially small quantity (as for the harmonic oscillator). Such precision was not numerically possible; in any case, knowing the energy does not establish localization. Note that $\phi(q)$ is the overlap of the ground state of the $(N+1)$ -atom ring with that of the N -atom chain.

To study localization we compare \mathcal{G} with the propagator of a corresponding *local mode*, i.e., a system in which $\lambda q^4/4$ is replaced by $\omega_1^2 q^2/2$ for appropriate ω_1 ($\sim \sqrt{\lambda}$). To derive $\mathcal{G}_{\text{local}}$ we redo the process described above. K is the same. For this system we know that as ω_1 increases, the vibrations are localized near 0. How can that be seen in $\mathcal{G}_{\text{local}}$ or S ? The method is to go to imaginary time and vary μ (the coupling to atom $\#m \approx N/3$).

Before presenting results we comment on numerics. The extremal solves a two-time boundary-value problem with a nonlocal interaction. In the linear case we discretize time, so that d^2/dt^2 becomes a difference matrix, nearly diagonal, but the K matrix is spread out. Call the resulting

operator $B = [d^2/dt^2] + \dots$. Take q to be a column vector, $q_j \leftrightarrow q(t_j)$. For $q(0) = a$, $q(T) = b$, and $\epsilon = t_{k+1} - t_k$, the extremal is $q = B^{-1}q^{(0)}$, where $q_j^{(0)} = -(a, 0, \dots, 0, b)/\epsilon^2$ (cf. [12]). Note that $S = q\dot{q}/2|_0^T$. To deal with computer limits on ϵ , we calculated S for the smallest practical ϵ and for *larger* values. S as a function of ϵ was then extrapolated to $\epsilon = 0$. This technique was validated for the demanding energy calculation, involving the exponentially small $\partial^2 S / \partial a \partial b$. Here the answer is known from the classical chain and ring frequencies. (E_0 also requires a $T \rightarrow \infty$ extrapolation.) The function ϕ , a Gaussian whose spread




FIG. 1. $dS/d\mu$ (essentially a correlation function) has markedly different behavior for large and small end point (“ a ”) values for q . Both for a linear local mode (as a function of ω_1) and for the breather (function of λ), a large value of a demands relatively large correlations when neither the local mode nor the breather is present ($\omega_1 \sim \lambda \sim 0$), but that correlation is wiped out for a large parameter, for which the breather or local mode is effectively decoupled from the rest of the ring.

can be calculated by standard methods, provided another check. Numerically that spread can be extracted by calculating S with varying end points and fitting the slope of \sqrt{S} . This too confirmed the method.

With q^4 the nonlinear, nonlocal two-time boundary-value problem cannot be solved by matrix inversion. Our method was a variation of [7], defining a subsidiary variable $z(t) \equiv \int_0^T \tilde{K}_1(|t-s|)q(s)ds$, where the subscript on \tilde{K}_1 indicates that $\tilde{K} [=iK(-iu)$ with $T \rightarrow -iT$] is replaced by a single “cosh” with optimized parameters, in fact an excellent approximation. This allowed a local boundary-value numerical technique, with a self-consistency demand on $z(0)$. Because of the approximation, we further attempted small variations to lower the true action, but they resulted in essentially no change.

Figure 1 shows the results of changing μ for the local mode (a) and for the breather (b). Localization is deduced from the variation of $dS/d\mu$ as a function of both a ($=q^f = q^i$) and the coupling, ω_1 or λ . All runs are for $T = 2$, which (given the energy scale) allows many states besides the ground state to survive in a spectral sum for \mathcal{G} . Therefore a large a boundary condition selects for the excited states, in particular, for a localized excited state (breather or local mode) if there is one. When the coupling is small, there is no localized state; forcing q to be large forces all atoms on the ring to depart from their usual positions. Hence the large magnitude of $dS/d\mu$ on the left of both figures—for the largest a values. (Small a has almost no effect even for small coupling.) For large coupling, however, there is a localized state: both for the linear (ω_1) and nonlinear (λ) systems, forcing q to depart from 0 has almost no effect on a distant atom. As further checks, in [9] we show that $dS/d\mu \rightarrow 0$ for $\lambda \rightarrow \infty$, and that for both linear and nonlinear cases $dS/d\mu$ has essentially the same behavior as T grows [13].

Diagonalizing a truncated Hamiltonian.—The Hamiltonian (1) is made finite dimensional by using a phonon basis and imposing a cutoff on the level of phonon excitation. A low cutoff is needed because of the proliferation of dimensions in the (implicit) tensor product of phonon operators. The same problem was faced in [6] and we alleviate it by the following strategy. Instead of perturbing around free modes, we use modes of a fictitious (linear) system with a local mode at site 0. This reduces the amplitudes of other modes in the quantum breather. Equation (1), with nonlinearity confined to x_0 , is written

$$H = \sum_{k=0}^N \left\{ \frac{1}{2} p_k^2 + \frac{1}{2} \omega_s^2 x_k^2 + \frac{1}{2} \omega_0^2 (\Delta x)_k^2 \right\} + \frac{1}{2} \omega_1^2 x_0^2 + V_I, \quad (8)$$

with $V_I \equiv \lambda x_0^4/4 - \omega_1^2 x_0^2/2$ the perturbation. A variation on Eq. (1) uses a nonlinear *coupling*, that is, terms $\lambda(\Delta x)_k^4/4$. Here too, we drop nonlinear terms except those (two) that include x_0 , and add and subtract corresponding

quadratics. As for (1), the classical dynamics is substantially the same.

We outline the calculation: The classical *local* mode problem is solved, giving rise to (truncated) creation and annihilation operators in terms of which the perturbation is expressed. The operator $\hat{x}_0 = \sum (a_\ell + a_\ell^\dagger) u_\ell(0) / \sqrt{2\Omega_\ell}$, where u_ℓ is the ℓ th mode of the system and $\ell = 0$ is the local mode. The key to ω_1 's effectiveness lies in $\{u_\ell(0)\}$. For $\omega_1 = 0$ these are all of about the same magnitude (for $N = 20$ they are ~ 0.3 for those not zero by symmetry). But with $\omega_1 = 3$, $u_0(0) \sim 0.988$, while the (nonzero) others average ~ 0.04 . This severely reduces the amplitudes of phonons other than the local mode in the eigenstates of the full Hamiltonian.

With $\lambda = 8$, $\omega_0 = 1$, and $\omega_s = 1$ (used in [6]), and for $\omega_1 = 2.5$ the true eigenfunction having greatest overlap with the first excited state of the local mode is shown in Table I. Clearly the local mode dominates. The next largest component is the thrice excited local mode, which is merely a shape adjustment. Other modes barely make the 10^{-3} level. Note that the highest excitation level for *other* phonons is 3, indicating that a cutoff of 6 is safe. In fact, even to probability 10^{-8} there is no excitation higher than 3 except for the local mode.

As a check of cutoff sensitivity we repeated this calculation with a cutoff of 8 (but the local mode still at 13). The results are in the last column of Table I. There is little sensitivity to the change—not only in the probabilities but in the composition of the state. Increasing ring size and reducing the cutoff preserves the pattern. For a ring of size 8 and a cutoff of 5 (10 for the local mode), the state was again dominated (probability 0.992) by the local mode, with the next contributor an excited local mode. Going to yet larger rings (size 10, and lower cutoff) preserves the pattern, with the first nonlocal mode phonon

TABLE I. Principal components of the breather state. The first 4 columns refer to the 4 symmetric phonons in a 6-atom ring. (N atoms $\Rightarrow [N/2] + 1$ symmetric modes.) Row 1: frequencies (local mode is highest). Subsequent rows: number-operator values. Fifth column: norm squared of the mode (\log_{10} in parentheses). Cutoffs: local mode 13, others 6. For this state, first order perturbation theory is good to 0.3%. The last column reports the same calculation with cutoff 8.

0.749	0.987	1.19	2.04	Probability (6)	Probability (8)
0	0	0	1	0.9947	0.9952
0	0	0	3	3.315 (−3)	3.321 (−3)
0	1	0	0	1.122 (−3)	0.848 (−3)
0	0	1	0	5.748 (−4)	4.345 (−4)
1	0	0	0	1.545 (−4)	1.167 (−4)
0	1	0	2	1.004 (−4)	0.754 (−4)
1	0	0	2	2.866 (−5)	2.152 (−5)
0	0	1	2	2.303 (−5)	1.728 (−5)
0	0	0	5	3.670 (−6)	3.627 (−6)




FIG. 2. Survival as a function of time; the initial state is the singly excited local mode.

contributing at probability level 10^{-4} . Other breather states yield the same result: accuracy of perturbation theory, insensitivity to cutoff change, and eigenfunction domination by local modes.

For nearest-neighbor nonlinearity the story is the same. Systematic study of a size 8 ring with cutoffs of 6, 8 and 12, showed that the local mode dominated the Hamiltonian's ground state.

The message of this dominance is that the eigenstates of the Hamiltonian are well localized, which is to say the quantized breather states are stable.

Quantum time dependence.—Although we have shown the breather to be dominated by the local mode, the point has been raised that by Fermi's golden rule *any* coupling to a continuum implies instability. This is not true. Some years ago it was found that quantum systems with ample continuum coupling nevertheless could survive indefinitely [14]. This departure from ordinary decay depended on smoothness and threshold properties of the coupling. A related matter concerns normalizable bound states in the continuum [15].

In Fig. 2 we show $|\langle\psi_0|\exp(-iHt/\hbar)|\psi_0\rangle|^2$ for ψ_0 a local mode phonon and H the full Hamiltonian. As in [14], initial decay is followed by stabilization bounded away from zero. In [14] the amplitude was typically a few tens of percent, but here, consistent with Table I, the amplitude stays near one. Despite the many connecting states, after a short time amplitude ceases to leak. The explanation [14] is that the true eigenstate of the Hamiltonian has order unity overlap with the initial state.

Note that stability may not persist in all dimensions; certainly threshold features of the density of states and spectrum are affected by dimension, and the usual intuitions regarding Fermi's golden rule may again hold sway. In [8] we made the point that the symmetry breaking of the Jahn-Teller effect makes this a one-dimensional problem, significantly enhancing the possibility of classical breath-

ers. The same is likely to be true quantum mechanically. It may even be that this plays a role in the temperature-dependent decay of the breather (through an effective increase in dimension), as evidenced by the high-temperature disappearance of anomalous decay in doped alkali halides [16]. This issue should be addressable using the path integral method.

In conclusion, we have shown that although quantum tunneling does in principle convert the classical breather into a Bloch state, when the ability to tunnel is removed, an initially localized pulse of energy can be trapped indefinitely as a quantum excitation.

This work was supported by NSF Grant No. PHY 00 99471 and Czech Grants No. ME587 GA and No. AVCR A1010210.

*Electronic address: schulman@clarkson.edu

- [1] Discrete breathers are a kind of soliton, also known as "intrinsic localized modes."
- [2] D. K. Campbell, S. Flach, and Y. S. Kivshar, *Phys. Today* **57**, No. 1, 43 (2004).
- [3] V. Fleurov, *Chaos* **13**, 676 (2003).
- [4] V. Hizhnyakov, *Phys. Rev. B* **53**, 13 981 (1996); *Europhys. Lett.* **45**, 508 (1999); V. Hizhnyakov and A. Nevedrov, *Phys. Rev. B* **56**, R2908 (1997); V. Hizhnyakov, A. Nevedrov, and A. J. Sievers, *Physica (Amsterdam)* **316B–317B**, 132 (2002).
- [5] S. Flach, V. Fleurov, and A. V. Gorbach, *Phys. Rev. B* **71**, 064302 (2005).
- [6] W. Z. Wang, J. T. Gammel, A. R. Bishop, and M. I. Salkola, *Phys. Rev. Lett.* **76**, 3598 (1996).
- [7] R. P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals* (McGraw-Hill, New York, 1965).
- [8] L. S. Schulman, E. Mihóková, A. Scardicchio, P. Facchi, M. Nikl, K. Polák, and B. Gaveau, *Phys. Rev. Lett.* **88**, 224101 (2002).
- [9] L. S. Schulman, D. Tolkunov, and E. Mihokova, *J. Chem. Phys.* (to be published).
- [10] L. S. Schulman, *Techniques and Applications of Path Integration* (Wiley, New York, 1981) (republished by Dover, New York, 2005). See also the Supplement in the republished version.
- [11] U. Weiss, *Quantum Dissipative Systems* (World Scientific, Singapore, 1999), 2nd ed.
- [12] L. S. Schulman, *J. Math. Phys. (N.Y.)* **15**, 295 (1974).
- [13] The large ω_1 and λ values in the figure do not imply an extended wave function. For the classical local mode, the amplitude (hence the quantum excitation) drops *exponentially* as one moves away from atom 0. See [9].
- [14] B. Gaveau and L. S. Schulman, *J. Phys. A* **28**, 7359 (1995).
- [15] M. Reed and B. Simon, *Methods of Modern Mathematical Physics IV* (Academic Press, San Diego, 1978), Section XIII.13.
- [16] E. Mihóková, L. S. Schulman, M. Nikl, B. Gaveau, K. Polák, K. Nitsch, and D. Zimmerman, *Phys. Rev. B* **66**, 155102 (2002).