

Interaction-Driven Real-Space Condensation

M. R. Evans,^{1,3} T. Hanney,^{1,3} and Satya N. Majumdar^{2,3}

¹*SUPA and School of Physics, University of Edinburgh, Mayfield Road, Edinburgh EH9 3JZ, United Kingdom*

²*Laboratoire de Physique Théorique et Modèles Statistiques, Université Paris-Sud, Bâtiment 100, 91405 Orsay-Cedex, France*

³*Isaac Newton Institute for Mathematical Sciences, 20 Clarkson Road, Cambridge CB3 0EH, United Kingdom*

(Received 3 May 2006; published 7 July 2006)

We study real-space condensation in a broad class of stochastic mass transport models. We show that the steady state of such models has a pair-factorized form which generalizes the standard factorized steady states. The condensation in this class of models is driven by interactions which give rise to a spatially extended condensate that differs fundamentally from the previously studied examples. We present numerical results as well as a theoretical analysis of the condensation transition and show that the criterion for condensation is related to the binding-unbinding transition of solid-on-solid interfaces.

DOI: [10.1103/PhysRevLett.97.010602](https://doi.org/10.1103/PhysRevLett.97.010602)

PACS numbers: 05.70.Fh, 05.40.-a, 05.70.Ln, 64.60.-i

Real-space condensation has been observed in a variety of physical contexts such as cluster aggregation [1], jamming in traffic and granular flow [2,3], and granular clustering [4]. The characteristic feature of these systems is the stochastic transport of some conserved quantity, to be referred to as mass; the condensation transition is manifested when above some critical mass density a single condensate captures a finite fraction of the mass. The condensate corresponds to a dominant cluster or a single large jam in these examples. Perhaps more surprising realizations of condensation are wealth condensation in macroeconomies [5] where the condensate corresponds to a single individual or enterprise owning a finite fraction of the wealth, condensation in growing or rewiring networks where a single hub captures a finite fraction of the links [6], and phase separation dynamics in one-dimensional driven systems where condensation corresponds to the emergence of a macroscopic domain of one phase [7].

Mass transport may be modeled in terms of interacting many-particle systems governed by stochastic dynamical rules. Generically these systems lack detailed balance and thus have nontrivial nonequilibrium steady states. Although our understanding of such steady states is still at an early stage, a class of models has been determined which exhibit a factorized steady state (FSS) [8] which can be written as a product of factors, one factor for each site of the system. This simple form for the steady state has afforded an opportunity to study condensation analytically and has also been used as an approximation to more complicated nonequilibrium steady states. The conditions under which condensation can occur have been determined, leading to conditions on the stochastic mass transport rules for condensation to result [9]. One key feature of the condensate arising in these models is that it forms at a single site.

In the physical systems of the kind described above, generically the stochastic transport rules depend not only on the departure site but also on the surrounding environ-

ment. In general such models do not have FSSs and finding their steady states has remained a challenge.

The purpose of this Letter is twofold. First, we introduce a broad class of mass transport models where the transport rules depend on the environment of the departure site. These models do not have a FSS, yet we can determine their steady states explicitly. The structure of the steady state generalizes the FSS to a pair-factorized steady state (PFSS). Second, we find that the nature of the condensate in a PFSS is strikingly different from that of the FSS: unlike in the FSS, the condensate is spatially extended. This is due to the short-range correlations inherent in the PFSS, but absent in the FSS.

We consider a class of mass transport models on a periodic chain with sites labeled by $i = 1, \dots, L$. At each site resides a non-negative integer number, m_i , of particles each of unit mass. We define particle dynamics such that a particle hops from site i to $i+1$ with a rate $u(m_{i-1}, m_i, m_{i+1})$ (provided $m_i > 0$), so the total mass $\sum_i m_i = M$ is conserved. These dynamics drive a current of particles through the system.

If the hop rate is only a function, $u(m_i)$, of mass at the departure site m_i , the model reduces to the zero-range process [10] which has a FSS. Explicitly, the probability of a configuration $\{m_i\}$ occurring in the steady state is

$$P[\{m_i\}] \propto \prod_{i=1}^L f(m_i) \delta\left(\sum_i m_i - M\right), \quad (1)$$

where $f(m) = 1/\prod_{k=1}^m u(k)$ for $m \geq 1$ and $f(0) = 1$. Thus there is one factor $f(m_i)$ for each site i of the system and the delta function ensures that the total mass is M .

When the hop rates $u(m_{i-1}, m_i, m_{i+1})$ depend on all three arguments, we propose the PFSS as a natural generalization of the FSS which takes the following form: the steady state probability of configuration, $\{m_i\}$, is

$$P[\{m_i\}] = Z_{L,M}^{-1} \prod_{i=1}^L g(m_i, m_{i+1}) \delta\left(\sum_{i=1}^L m_i - M\right). \quad (2)$$

Thus there is one factor $g(m_i, m_{i+1})$ for each pair of neighboring sites. This form has been considered for two-state models (e.g., $m_i = 0, 1$ only) as an improved mean-field approximation [11,12] and also as an exact steady state for particular driven diffusive dynamics [10,13]. The normalization $Z_{L,M}$, which plays a role analogous to the canonical partition function in equilibrium statistical mechanics, is given by

$$Z_{L,M} = \sum_{\{m_i\}} \prod_{i=1}^L g(m_i, m_{i+1}) \delta\left(\sum_{i=1}^L m_i - M\right). \quad (3)$$

Note that in the case $g(m_i, m_{i+1}) = f(m_i)$, for example, the PFSS Eq. (2) reduces to the FSS form (1).

First, we establish that the steady state (2) holds for a broad class of mass transport models. We find that if (though not only if) the hop rates factorize [14],

$$u(m_{i-1}, m_i, m_{i+1}) = \alpha(m_{i-1}, m_i) \beta(m_i, m_{i+1}), \quad (4)$$

then the steady state is of the PFSS form (2) with

$$g(m, n) = \prod_{i=1}^n \alpha(m, i)^{-1} \prod_{j=1}^m \beta(j, 0)^{-1}, \quad (5)$$

for $m, n > 0$ where $g(0, 0) = 1$, provided α and β satisfy the constraint

$$\frac{\alpha(m-1, n)}{\alpha(m, n)} = \frac{\beta(m, n-1)}{\beta(m, n)}. \quad (6)$$

Furthermore, given any form of the weight $g(m, n)$ one can determine the functions α and β through the following recursions:

$$\alpha(l, m) = \frac{g(l, m-1)}{g(l, m)}, \quad \beta(m, n) = \frac{g(m-1, n)}{g(m, n)}. \quad (7)$$

Thus for *every* choice of $g(m, n)$ there exists a stochastic mass transport model which will generate the corresponding PFSS.

We will focus on a particular model which has a PFSS with $g(m, n)$ given by

$$g(m, n) = \exp[-J|m-n| + \frac{1}{2}U_0(\delta_{m,0} + \delta_{n,0})]. \quad (8)$$

One can check from (4) and (7) that the corresponding hop rates are

$$u(l, m, n) = \begin{cases} \exp[-2J + U_0\delta_{m,1}] & \text{for } m \leq l, n, \\ \exp[2J + U_0\delta_{m,1}] & \text{for } m > l, n, \\ \exp[U_0\delta_{m,1}] & \text{otherwise.} \end{cases} \quad (9)$$

Physically, the rate is low if the mass at the departure site is less than the neighboring masses and is high if the mass is larger than the neighboring masses. This tends to flatten the density profile and generates the effective surface tension J in (8) implying short-range correlations between the sites. In addition, isolated particles tend to hop relatively quickly leading to a preference in the steady state weights for

vacant sites. This is reflected by the on-site attractive potential $-U_0\delta_{m_i,0}$ in (8).

The model defined by the hop rates (9) is guaranteed to have a PFSS with $g(m, n)$ in (8). To investigate whether the model allows for a condensation transition as the parameters J, U_0 and the conserved mass density $\rho = M/L$ are varied, we have run Monte Carlo simulations, according to the following prescription. The system is prepared in a random, homogeneous initial condition and evolves under random sequential update. During each time step δt , a site l is selected randomly, and if a particle is present, it is transferred to the neighboring site $l+1$ with probability $u(m_{l-1}, m_l, m_{l+1})\delta t$. L such time steps constitute a single Monte Carlo step.

We find that two phases emerge in the steady state depending on ρ and J (where we set $U_0 = 1$). As illustrated in Fig. 1, at low density the system resides in a fluid phase, in which particles are distributed homogeneously throughout the system. When the density exceeds a critical value $\rho_c(J)$, the system is in a condensed phase wherein a condensate containing the excess mass $(\rho - \rho_c)L$ coexists with a critical background fluid of mass $\rho_c L$. In contrast with the usual condensate that occupies a single site, such as, for example, in a FSS, the condensate here extends over many sites. In fact, the condensate extends over typically $O(L^{1/2})$ sites as shown in Fig. 1(a).

To locate the phase boundary in the ρ - J plane we computed the single-site probabilities $p(m, L)$ that a site contains exactly mass m in the steady state. In the fluid phase $p(m, L)$ decays exponentially for large m , whereas in the condensed phase an additional bump emerges at the large m tail of $p(m, L)$ as illustrated in Fig. 2. The phase boundary in Fig. 3 is determined by the value of J , for fixed ρ , at which a bump in $p(m, L)$ first appears as one increases J . Our theoretical prediction for the phase boundary, presented below, is in excellent agreement with the numerical results.




FIG. 1 (color online). Typical steady state configurations for $L = 1000$ and $J = U_0 = 1$ (for which $\rho_c = 0.51$) in (a) condensed phase, $\rho = 3$, and (b) fluid phase, $\rho = 1/4$. The inset in (a) shows an $L^{1/2}$ dependence (indicated by the solid line) of the condensate width on system size.




FIG. 2 (color online). Single-site distribution $p(m, L)$ for a system of $L = 1000$ sites with $\rho = 3$ and $J = U_0 = 1$. The inset shows the mass at the maximum of the condensate bump as a function of system size L , where the crosses are obtained from Monte Carlo simulation and the solid line shows an $L^{1/2}$ dependence.

The condensate bump in $p(m, L)$ also has an interesting scaling behavior with L . We plot in the inset of Fig. 2 the mass at the maximum of the bump as a function of L and find that it grows as $L^{1/2}$. This implies that a typical site inside the condensate has mass of order $L^{1/2}$. On the other hand, since the condensate carries total excess mass $(\rho - \rho_c)L$, this implies that there are typically of order $L^{1/2}$ sites inside the condensate; i.e., the spatial extent of the condensate is of order $L^{1/2}$.

To find the general conditions under which the steady state (2) may admit condensation, we analyze the grand canonical partition function $Z_L(\mu)$ (the Laplace transform of $Z_{L,M}$ with respect to M) given by

$$Z_L(\mu) = \sum_{\{m_i\}} e^{-\mu \sum_i m_i} \prod_{i=1}^L g(m_i, m_{i+1}), \quad (10)$$

where the chemical potential μ is determined from the condition that

$$\rho = \rho(\mu) \equiv -\frac{1}{L} \frac{\partial \ln Z_L(\mu)}{\partial \mu}, \quad (11)$$

Clearly $\rho(\mu)$ is a decreasing function of μ for $\mu \geq 0$ [15]. If, as $\mu \rightarrow 0$, the function $\rho(\mu) \rightarrow \infty$, then a solution $\mu > 0$ of (11) exists for any $\rho > 0$. This implies from (10) that the single-site mass distribution decays exponentially for large m signifying a fluid phase and there is no condensation. On the other hand, if, as $\mu \rightarrow 0$, the function $\rho(\mu)$ approaches a finite value ρ_c , then a solution of (11) can be found only for $\rho < \rho_c$, implying that the fluid phase exists only for $\rho < \rho_c$. When ρ exceeds the critical particle density ρ_c , there is no solution to (11), implying the onset of condensation wherein the “excess” mass $(\rho - \rho_c)L$ is carried by the condensate.

Thus to determine if there is condensation one needs to analyze (10) and (11) as $\mu \rightarrow 0$. But, for $\mu = 0$, (10) is precisely the grand canonical partition function of a solid-on-solid (SOS) interface model [16,17] where the interface height at site i is equivalent to the mass m_i . Since $m_i \geq 0$, the interface heights are strictly non-negative, implying that the interface grows on a substrate at $m_i = 0$. Thus $\rho(0)$ from (11) corresponds to the average interface height $\langle m_i \rangle$ in this SOS model. If $\rho(0) = \infty$, i.e., there is no condensation transition, the interface is unbound since its mean height is divergent. On the other hand, if there is a condensation transition, in which case $\rho(0) = \rho_c$ is finite, the interface is bound with a finite mean height $\rho(0)$. Therefore the criterion for a condensation transition is that the corresponding interface should be bound. Moreover the critical density ρ_c is given by the mean height of the bound interface.

The phase diagram in Fig. 3 can be exactly calculated using a standard transfer matrix formalism. The partition function (10) may be written as $Z_L(\mu) = \text{Tr}[T^L(\mu)]$ where the elements of the transfer matrix T are $g(m', m)$. In the large L limit, only the eigenvector $|\phi_0\rangle$ of T with the largest eigenvalue λ_0 contributes. The eigenvalue equation reads $\sum_{m=0}^{\infty} g(m', m) \langle m | \phi_\alpha \rangle = \lambda_\alpha \langle m' | \phi_\alpha \rangle$. The eigenvectors are either superpositions of extended states $\langle m | \phi_\alpha \rangle \sim e^{ipm}$ or a bound state $\langle m | \phi_\alpha \rangle \sim t^m$ where $|t| < 1$ [16]. If the spectrum contains a bound state, then the bound state corresponds to the largest eigenvalue. Substituting $\langle m | \phi_0 \rangle = t^m$ into the eigenvalue equation for $m' > 0$ and $m' = 0$ separately yields $t = e^{-J}/(1 - e^{-U_0})$. For the bound state to exist, $|t| < 1$, which implies

$$J > J_c = U_0 - \ln(e^{U_0} - 1). \quad (12)$$

Therefore, for $J < J_c$ the system will not condense at any finite density, implying $\rho_c = \infty$; for $J > J_c$, the system condenses above a finite density given by the mean height in the bound state, $\rho_c = \sum_m m \langle m | \phi_0 \rangle^2 / \sum_m \langle m | \phi_0 \rangle^2$. Using the bound state eigenfunction one finds




FIG. 3 (color online). Phase diagram for $U_0 = 1$. The crosses are data points obtained from Monte Carlo simulation. The solid line gives the theoretical prediction (13). For $J < J_c$ given by (12), condensation does not occur at any density.

$$\rho_c = [e^{2(U-J_c)} - 1]^{-1}. \quad (13)$$

This prediction is in excellent agreement with the numerical data shown in Fig. 3.

Finally, we discuss the condensation transition in a more general PFSS of the form

$$g(m, n) = K(|m - n|) \exp\left\{\frac{U}{2}[U(m) + U(n)]\right\}. \quad (14)$$

Here $K(|m - n|)$ represents the interaction between nearest neighbor masses and $-U(m)$ is an on-site potential. If $K(x)$ decays sufficiently rapidly for large x , as in (8), then condensation is possible if $U(m)$ is positive and localized near $m = 0$. In this case the condensation is interaction driven and the existence of the condensation transition corresponds to having a bound interface. In such cases quite generically the height and width of the condensate are expected to scale as $L^{1/2}$. This follows from a Brownian excursion argument: the localized on-site potential plays no role at sites occupied by the condensate—in the absence of the potential, the problem can be related to a random walk problem where the random walker takes independent steps with length drawn from a distribution $K(x)$ [18]. The shape of the condensate is determined by a single large loop defined by the excursion of the random walker as shown in Fig. 1. The probability that the walker returns to the origin for the first time after N steps scales as $N^{-3/2}$ for sufficiently rapidly decaying $K(x)$. So, the average number of steps until the first return is $\int_1^L N^{-3/2} N dN \sim L^{1/2}$ (the upper cutoff, L , is determined by the maximum number of possible steps). This predicts that the spatial extent of the condensate is $\mathcal{O}(L^{1/2})$. Also, because it is Brownian, the typical height of the excursion, and therefore that of the condensate, scales as $L^{1/2}$. Note that the area under the excursion, equivalent to the mass contained in the condensate, is $\mathcal{O}(L)$, as it should be.

This interaction-driven condensation is rather different from the type exhibited in a FSS. There the function $K(x) = 1$ and a localized on-site attractive potential is no longer capable of driving condensation. Instead, one requires a specific unbounded potential of the form $-U(m) \sim \gamma \ln m$ for large m [10]. Thus in the FSS condensation is potential driven.

To summarize, the steady state (2) extends the class of exactly solvable steady states in nonequilibrium statistical mechanics. The condensed phase which emerges in a PFSS is fundamentally different from that in a FSS. In a PFSS the condensation is interaction driven and the condensate extends spatially over $\mathcal{O}(L^{1/2})$ sites. The explicit form of the single-site mass distribution $p(m, L)$ in the FSS condensed phase has been determined recently [9]. It remains a challenge to compute $p(m, L)$ for the PFSS condensed phase. It would also be of interest to study PFSS in higher dimensions. Finally, we note that the FSS has provided insight

into a number of issues of nonequilibrium statistical physics. Although we have focused here on condensation, the generalization to a PFSS should allow one to address further issues, such as the role of conservation laws [19], disorder [20], boundary-induced phenomena [21], and fluctuation theorems [22].

T. H. thanks the EPSRC for support under Programme Grant No. GR/S10377/01.

-
- [1] S. N. Majumdar, S. Krishnamurthy, and M. Barma, Phys. Rev. Lett. **81**, 3691 (1998).
 - [2] O. J. O'Loan, M. R. Evans, and M. E. Cates, Phys. Rev. E **58**, 1404 (1998).
 - [3] D. Chowdhury, L. Santen, and A. Schadschneider, Phys. Rep. **329**, 199 (2000).
 - [4] D. van der Meer, K. van der Weele, and D. Lohse, Phys. Rev. Lett. **88**, 174302 (2002).
 - [5] Z. Burda *et al.*, Phys. Rev. E **65**, 026102 (2002).
 - [6] S. N. Dorogovstev and J. F. F. Mendes, *Evolution of Networks* (OUP, Oxford, 2003).
 - [7] Y. Kafri, E. Levine, D. Mukamel, G. M. Schütz, and J. Török, Phys. Rev. Lett. **89**, 035702 (2002).
 - [8] M. R. Evans, S. N. Majumdar, and R. K. P. Zia, J. Phys. A **37**, L275 (2004); **39**, 4859 (2006).
 - [9] S. N. Majumdar, M. R. Evans, and R. K. P. Zia, Phys. Rev. Lett. **94**, 180601 (2005); M. R. Evans, S. N. Majumdar, and R. K. P. Zia, J. Stat. Phys. **123**, 357 (2006).
 - [10] M. R. Evans and T. Hanney, J. Phys. A **38**, R195 (2005).
 - [11] D. ben-Avraham and J. Kohler, Phys. Rev. A **45**, 8358 (1992).
 - [12] M. Schreckenberg, A. Schadschneider, K. Nagel, and N. Ito, Phys. Rev. E **51**, 2939 (1995).
 - [13] S. Katz, J. L. Lebowitz, and H. Spohn, J. Stat. Phys. **34**, 497 (1984).
 - [14] M. R. Evans, T. Hanney, and S. N. Majumdar (to be published).
 - [15] Strictly, if $g(m, n)$ contains a pure exponential factor, e.g., $\exp[a(m + n)]$, $Z_L(\mu)$ in (10) would exist for $\mu \geq a$, but this just amounts to a shift in the zero of the chemical potential.
 - [16] S. T. Chui and J. D. Weeks, Phys. Rev. B **23**, 2438 (1981).
 - [17] J. M. J. van Leeuwen and H. J. Hilhorst, Physica (Amsterdam) **107A**, 319 (1981); T. W. Burkhardt, J. Phys. A **14**, L63 (1981).
 - [18] G. Schehr and S. N. Majumdar, Phys. Rev. E **73**, 056103 (2006).
 - [19] S. Grosskinsky and H. Spohn, Bull. Braz. Math. Soc. **34**, 489 (2003); M. R. Evans and T. Hanney, J. Phys. A **36**, L441 (2003).
 - [20] J. Krug, Braz. J. Phys. **30**, 97 (2000); K. Jain and M. Barma, Phys. Rev. Lett. **91**, 135701 (2003).
 - [21] E. Levine, D. Mukamel, and G. M. Schütz, J. Stat. Phys. **120**, 759 (2005).
 - [22] R. J. Harris, A. Rákos, and G. M. Schütz, J. Stat. Mech. (2005) P08003.