

Measurement of the Charge Asymmetry in Semileptonic B_s^0 Decays

V. M. Abazov,³⁵ B. Abbott,⁷⁵ M. Abolins,⁶⁵ B. S. Acharya,²⁸ M. Adams,⁵¹ T. Adams,⁴⁹ E. Aguilo,⁵ S. H. Ahn,³⁰ M. Ahsan,⁵⁹ G. D. Alexeev,³⁵ G. Alkhazov,³⁹ A. Alton,⁶⁴ G. Alverson,⁶³ G. A. Alves,² M. Anastasoiaie,³⁴ L. S. Ancu,³⁴ T. Andeen,⁵³ S. Anderson,⁴⁵ B. Andrieu,¹⁶ M. S. Anzels,⁵³ Y. Arnaud,¹³ M. Arov,⁵² A. Askew,⁴⁹ B. Åsman,⁴⁰ A. C. S. Assis Jesus,³ O. Atramentov,⁴⁹ C. Autermann,²⁰ C. Avila,⁷ C. Ay,²³ F. Badaud,¹² A. Baden,⁶¹ L. Bagby,⁵² B. Baldin,⁵⁰ D. V. Bandurin,⁵⁹ P. Banerjee,²⁸ S. Banerjee,²⁸ E. Barberis,⁶³ P. Bargassa,⁸⁰ P. Baringer,⁵⁸ C. Barnes,⁴³ J. Barreto,² J. F. Bartlett,⁵⁰ U. Bassler,¹⁶ D. Bauer,⁴³ S. Beale,⁵ A. Bean,⁵⁸ M. Begalli,³ M. Begel,⁷¹ C. Belanger-Champagne,⁴⁰ L. Bellantoni,⁵⁰ A. Bellavance,⁶⁷ J. A. Benitez,⁶⁵ S. B. Beri,²⁶ G. Bernardi,¹⁶ R. Bernhard,²² L. Berntzon,¹⁴ I. Bertram,⁴² M. Besançon,¹⁷ R. Beuselinck,⁴³ V. A. Bezzubov,³⁸ P. C. Bhat,⁵⁰ V. Bhatnagar,²⁶ M. Binder,²⁴ C. Biscarat,¹⁹ I. Blackler,⁴³ G. Blazey,⁵² F. Blekman,⁴³ S. Blessing,⁴⁹ D. Bloch,¹⁸ K. Bloom,⁶⁷ A. Boehnlein,⁵⁰ D. Boline,⁶² T. A. Bolton,⁵⁹ G. Borissov,⁴² K. Bos,³³ T. Bose,⁷⁷ A. Brandt,⁷⁸ R. Brock,⁶⁵ G. Brooijmans,⁷⁰ A. Bross,⁵⁰ D. Brown,⁷⁸ N. J. Buchanan,⁴⁹ D. Buchholz,⁵³ M. Buehler,⁸¹ V. Buescher,²² S. Burdin,⁵⁰ S. Burke,⁴⁵ T. H. Burnett,⁸² E. Busato,¹⁶ C. P. Buszello,⁴³ J. M. Butler,⁶² P. Calfayan,²⁴ S. Calvet,¹⁴ J. Cammin,⁷¹ S. Caron,³³ W. Carvalho,³ B. C. K. Casey,⁷⁷ N. M. Cason,⁵⁵ H. Castilla-Valdez,³² S. Chakrabarti,¹⁷ D. Chakraborty,⁵² K. M. Chan,⁷¹ A. Chandra,⁴⁸ F. Charles,¹⁸ E. Cheu,⁴⁵ F. Chevallier,¹³ D. K. Cho,⁶² S. Choi,³¹ B. Choudhary,²⁷ L. Christofek,⁷⁷ D. Claes,⁶⁷ B. Clément,¹⁸ C. Clément,⁴⁰ Y. Coadou,⁵ M. Cooke,⁸⁰ W. E. Cooper,⁵⁰ M. Corcoran,⁸⁰ F. Couderc,¹⁷ M.-C. Cousinou,¹⁴ B. Cox,⁴⁴ S. Crépe-Renaudin,¹³ D. Cutts,⁷⁷ M. Ćwiok,²⁹ H. da Motta,² A. Das,⁶² M. Das,⁶⁰ B. Davies,⁴² G. Davies,⁴³ K. De,⁷⁸ P. de Jong,³³ S. J. de Jong,³⁴ E. De La Cruz-Burelo,⁶⁴ C. De Oliveira Martins,³ J. D. Degenhardt,⁶⁴ F. Déliot,¹⁷ M. Demarteau,⁵⁰ R. Demina,⁷¹ D. Denisov,⁵⁰ S. P. Denisov,³⁸ S. Desai,⁵⁰ H. T. Diehl,⁵⁰ M. Diesburg,⁵⁰ M. Doidge,⁴² A. Dominguez,⁶⁷ H. Dong,⁷² L. V. Dudko,³⁷ L. Duflot,¹⁵ S. R. Dugad,²⁸ D. Duggan,⁴⁹ A. Duperrin,¹⁴ J. Dyer,⁶⁵ A. Dyshkant,⁵² M. Eads,⁶⁷ D. Edmunds,⁶⁵ J. Ellison,⁴⁸ V. D. Elvira,⁵⁰ Y. Enari,⁷⁷ S. Eno,⁶¹ P. Ermolov,³⁷ H. Evans,⁵⁴ A. Evdokimov,³⁶ V. N. Evdokimov,³⁸ L. Feligioni,⁶² A. V. Ferapontov,⁵⁹ T. Ferbel,⁷¹ F. Fiedler,²⁴ F. Filthaut,³⁴ W. Fisher,⁵⁰ H. E. Fisk,⁵⁰ M. Ford,⁴⁴ M. Fortner,⁵² H. Fox,²² S. Fu,⁵⁰ S. Fuess,⁵⁰ T. Gadfort,⁸² C. F. Galea,³⁴ E. Gallas,⁵⁰ E. Galyaev,⁵⁵ C. Garcia,⁷¹ A. Garcia-Bellido,⁸² V. Gavrilov,³⁶ A. Gay,¹⁸ P. Gay,¹² W. Geist,¹⁸ D. Gelé,¹⁸ R. Gelhaus,⁴⁸ C. E. Gerber,⁵¹ Y. Gershtein,⁴⁹ D. Gillberg,⁵ G. Ginther,⁷¹ N. Gollub,⁴⁰ B. Gómez,⁷ A. Goussiou,⁵⁵ P. D. Grannis,⁷² H. Greenlee,⁵⁰ Z. D. Greenwood,⁶⁰ E. M. Gregores,⁴ G. Grenier,¹⁹ Ph. Gris,¹² J.-F. Grivaz,¹⁵ A. Grohsjean,²⁴ S. Grünendahl,⁵⁰ M. W. Grünewald,²⁹ F. Guo,⁷² J. Guo,⁷² G. Gutierrez,⁵⁰ P. Gutierrez,⁷⁵ A. Haas,⁷⁰ N. J. Hadley,⁶¹ P. Haefner,²⁴ S. Hagopian,⁴⁹ J. Haley,⁶⁸ I. Hall,⁷⁵ R. E. Hall,⁴⁷ L. Han,⁶ K. Hanagaki,⁵⁰ P. Hansson,⁴⁰ K. Harder,⁴⁴ A. Harel,⁷¹ R. Harrington,⁶³ J. M. Hauptman,⁵⁷ R. Hauser,⁶⁵ J. Hays,⁴³ T. Hebbeker,²⁰ D. Hedin,⁵² J. G. Hegeman,³³ J. M. Heinmiller,⁵¹ A. P. Heinson,⁴⁸ U. Heintz,⁶² C. Hensel,⁵⁸ K. Herner,⁷² G. Hesketh,⁶³ M. D. Hildreth,⁵⁵ R. Hirosky,⁸¹ J. D. Hobbs,⁷² B. Hoeneisen,¹¹ H. Hoeth,²⁵ M. Hohlfield,¹⁵ K. Holubyev,⁴² S. J. Hong,³⁰ R. Hooper,⁷⁷ P. Houben,³³ Y. Hu,⁷² Z. Hubacek,⁹ V. Hynek,⁸ I. Iashvili,⁶⁹ R. Illingworth,⁵⁰ A. S. Ito,⁵⁰ S. Jabeen,⁶² M. Jaffré,¹⁵ S. Jain,⁷⁵ K. Jakobs,²² C. Jarvis,⁶¹ A. Jenkins,⁴³ R. Jesik,⁴³ K. Johns,⁴⁵ C. Johnson,⁷⁰ M. Johnson,⁵⁰ A. Jonckheere,⁵⁰ P. Jonsson,⁴³ A. Juste,⁵⁰ D. Käfer,²⁰ S. Kahn,⁷³ E. Kajfasz,¹⁴ A. M. Kalinin,³⁵ J. M. Kalk,⁶⁰ J. R. Kalk,⁶⁵ S. Kappler,²⁰ D. Karmanov,³⁷ J. Kasper,⁶² P. Kasper,⁵⁰ I. Katsanos,⁷⁰ D. Kau,⁴⁹ R. Kaur,²⁶ R. Kehoe,⁷⁹ S. Kermiche,¹⁴ N. Khalatyan,⁶² A. Khanov,⁷⁶ A. Kharchilava,⁶⁹ Y. M. Kharzheev,³⁵ D. Khatidze,⁷⁰ H. Kim,³¹ T. J. Kim,³⁰ M. H. Kirby,³⁴ B. Klima,⁵⁰ J. M. Kohli,²⁶ J.-P. Konrath,²² M. Kopal,⁷⁵ V. M. Korablev,³⁸ J. Kotcher,⁷³ B. Kothari,⁷⁰ A. Koubarovsky,³⁷ A. V. Kozelov,³⁸ D. Krop,⁵⁴ A. Kryemadhi,⁸¹ T. Kuhl,²³ A. Kumar,⁶⁹ S. Kunori,⁶¹ A. Kupco,¹⁰ T. Kurča,¹⁹ J. Kvita,⁸ D. Lam,⁵⁵ S. Lammers,⁷⁰ G. Landsberg,⁷⁷ J. Lazoflores,⁴⁹ A.-C. Le Bihan,¹⁸ P. Lebrun,¹⁹ W. M. Lee,⁵⁰ A. Leflat,³⁷ F. Lehner,⁴¹ V. Lesne,¹² J. Leveque,⁴⁵ P. Lewis,⁴³ J. Li,⁷⁸ L. Li,⁴⁸ Q. Z. Li,⁵⁰ S. M. Lietti,⁴ J. G. R. Lima,⁵² D. Lincoln,⁵⁰ J. Linnemann,⁶⁵ V. V. Lipaev,³⁸ R. Lipton,⁵⁰ Z. Liu,⁵ L. Lobo,⁴³ A. Lobodenko,³⁹ M. Lokajicek,¹⁰ A. Lounis,¹⁸ P. Love,⁴² H. J. Lubatti,⁸² M. Lynker,⁵⁵ A. L. Lyon,⁵⁰ A. K. A. Maciel,² R. J. Madaras,⁴⁶ P. Mättig,²⁵ C. Magass,²⁰ A. Magerkurth,⁶⁴ N. Makovec,¹⁵ P. K. Mal,⁵⁵ H. B. Malbouisson,³ S. Malik,⁶⁷ V. L. Malyshev,³⁵ H. S. Mao,⁵⁰ Y. Maravin,⁵⁹ R. McCarthy,⁷² A. Melnitchouk,⁶⁶ A. Mendes,¹⁴ L. Mendoza,⁷ P. G. Mercadante,⁴ M. Merkin,³⁷ K. W. Merritt,⁵⁰ A. Meyer,²⁰ J. Meyer,²¹ M. Michaut,¹⁷ H. Miettinen,⁸⁰ T. Millet,¹⁹ J. Mitrevski,⁷⁰ J. Molina,³ R. K. Mommsen,⁴⁴ N. K. Mondal,²⁸ J. Monk,⁴⁴ R. W. Moore,⁵ T. Moulík,⁵⁸ G. S. Muanza,¹⁹ M. Mulders,⁵⁰ M. Mulhearn,⁷⁰ O. Mundal,²² L. Mundim,³ E. Nagy,¹⁴ M. Naimuddin,²⁷ M. Narain,⁶² N. A. Naumann,³⁴ H. A. Neal,⁶⁴ J. P. Negret,⁷ P. Neustroev,³⁹ C. Noeding,²² A. Nomerotski,⁵⁰ S. F. Novaes,⁴ T. Nunnemann,²⁴ V. O'Dell,⁵⁰ D. C. O'Neil,⁵ G. Obrant,³⁹ C. Ochando,¹⁵ V. Oguri,³ N. Oliveira,³ D. Onoprienko,⁵⁹ N. Oshima,⁵⁰ J. Osta,⁵⁵ R. Otec,⁹ G. J. Otero y Garzón,⁵¹ M. Owen,⁴⁴ P. Padley,⁸⁰

M. Pangilinan,⁶² N. Parashar,⁵⁶ S.-J. Park,⁷¹ S. K. Park,³⁰ J. Parsons,⁷⁰ R. Partridge,⁷⁷ N. Parua,⁷² A. Patwa,⁷³ G. Pawloski,⁸⁰ P. M. Perea,⁴⁸ K. Peters,⁴⁴ Y. Peters,²⁵ P. Pétrouff,¹⁵ M. Petteni,⁴³ R. Piegaia,¹ J. Piper,⁶⁵ M.-A. Pleier,²¹ P. L. M. Podesta-Lerma,³² V. M. Podstavkov,⁵⁰ Y. Pogorelov,⁵⁵ M.-E. Pol,² A. Pompoš,⁷⁵ B. G. Pope,⁶⁵ A. V. Popov,³⁸ C. Potter,⁵ W. L. Prado da Silva,³ H. B. Prosper,⁴⁹ S. Protopopescu,⁷³ J. Qian,⁶⁴ A. Quadt,²¹ B. Quinn,⁶⁶ M. S. Rangel,² K. J. Rani,²⁸ K. Ranjan,²⁷ P. N. Ratoff,⁴² P. Renkel,⁷⁹ S. Reucroft,⁶³ M. Rijssenbeek,⁷² I. Ripp-Baudot,¹⁸ F. Rizatdinova,⁷⁶ S. Robinson,⁴³ R. F. Rodrigues,³ C. Royon,¹⁷ P. Rubinov,⁵⁰ R. Ruchti,⁵⁵ G. Sajot,¹³ A. Sánchez-Hernández,³² M. P. Sanders,¹⁶ A. Santoro,³ G. Savage,⁵⁰ L. Sawyer,⁶⁰ T. Scanlon,⁴³ D. Schaile,²⁴ R. D. Schamberger,⁷² Y. Scheglov,³⁹ H. Schellman,⁵³ P. Schieferdecker,²⁴ C. Schmitt,²⁵ C. Schwanenberger,⁴⁴ A. Schwartzman,⁶⁸ R. Schwienhorst,⁶⁵ J. Sekaric,⁴⁹ S. Sengupta,⁴⁹ H. Severini,⁷⁵ E. Shabalina,⁵¹ M. Shamim,⁵⁹ V. Shary,¹⁷ A. A. Shchukin,³⁸ R. K. Shivpuri,²⁷ D. Shpakov,⁵⁰ V. Siccardi,¹⁸ R. A. Sidwell,⁵⁹ V. Simak,⁹ V. Sirotenko,⁵⁰ P. Skubic,⁷⁵ P. Slattery,⁷¹ R. P. Smith,⁵⁰ G. R. Snow,⁶⁷ J. Snow,⁷⁴ S. Snyder,⁷³ S. Söldner-Rembold,⁴⁴ X. Song,⁵² L. Sonnenschein,¹⁶ A. Sopczak,⁴² M. Sosebee,⁷⁸ K. Soustruznik,⁸ M. Souza,² B. Spurlock,⁷⁸ J. Stark,¹³ J. Steele,⁶⁰ V. Stolin,³⁶ A. Stone,⁵¹ D. A. Stoyanova,³⁸ J. Strandberg,⁶⁴ S. Strandberg,⁴⁰ M. A. Strang,⁶⁹ M. Strauss,⁷⁵ R. Ströhmer,²⁴ D. Strom,⁵³ M. Strovink,⁴⁶ L. Stutte,⁵⁰ S. Sumowidagdo,⁴⁹ P. Svoisky,⁵⁵ A. Sznajder,³ M. Talby,¹⁴ P. Tamburello,⁴⁵ W. Taylor,⁵ P. Telford,⁴⁴ J. Temple,⁴⁵ B. Tiller,²⁴ M. Titov,²² V. V. Tokmenin,³⁵ M. Tomoto,⁵⁰ T. Toole,⁶¹ I. Torchiani,²² T. Trefzger,²³ S. Trincas-Duvoid,¹⁶ D. Tsybychev,⁷² B. Tuchming,¹⁷ C. Tully,⁶⁸ P. M. Tuts,⁷⁰ R. Unalan,⁶⁵ L. Uvarov,³⁹ S. Uvarov,³⁹ S. Uzunyan,⁵² B. Vachon,⁵ P. J. van den Berg,³³ B. van Eijk,³⁵ R. Van Kooten,⁵⁴ W. M. van Leeuwen,³³ N. Varelas,⁵¹ E. W. Varnes,⁴⁵ A. Vartapetian,⁷⁸ I. A. Vasilyev,³⁸ M. Vaupel,²⁵ P. Verdier,¹⁹ L. S. Vertogradov,³⁵ M. Verzocchi,⁵⁰ F. Villeneuve-Seguiet,⁴³ P. Vint,⁴³ J.-R. Vlimant,¹⁶ E. Von Toerne,⁵⁹ M. Voutilainen,⁶⁷ M. Vreeswijk,³³ H. D. Wahl,⁴⁹ L. Wang,⁶¹ M. H. L. S. Wang,⁵⁰ J. Warchol,⁵⁵ G. Watts,⁸² M. Wayne,⁵⁵ G. Weber,²³ M. Weber,⁵⁰ H. Weerts,⁶⁵ N. Vermes,²¹ M. Wetstein,⁶¹ A. White,⁷⁸ D. Wicke,²⁵ G. W. Wilson,⁵⁸ S. J. Wimpenny,⁴⁸ M. Wobisch,⁵⁰ J. Womersley,⁵⁰ D. R. Wood,⁶³ T. R. Wyatt,⁴⁴ Y. Xie,⁷⁷ S. Yacoob,⁵³ R. Yamada,⁵⁰ M. Yan,⁶¹ T. Yasuda,⁵⁰ Y. A. Yatsunenko,³⁵ K. Yip,⁷³ H. D. Yoo,⁷⁷ S. W. Youn,⁵³ C. Yu,¹³ J. Yu,⁷⁸ A. Yurkewicz,⁷² A. Zatserklyaniy,⁵² C. Zeitnitz,²⁵ D. Zhang,⁵⁰ T. Zhao,⁸² B. Zhou,⁶⁴ J. Zhu,⁷² M. Zielinski,⁷¹ D. Zieminska,⁵⁴ A. Zieminski,⁵⁴ V. Zutshi,⁵² and E. G. Zverev³⁷

(D0 Collaboration)

¹Universidad de Buenos Aires, Buenos Aires, Argentina²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil⁴Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil⁵University of Alberta, Edmonton, Alberta, Canada,

Simon Fraser University, Burnaby, British Columbia, Canada,

York University, Toronto, Ontario, Canada,

and McGill University, Montreal, Quebec, Canada

⁶University of Science and Technology of China, Hefei, People's Republic of China⁷Universidad de los Andes, Bogotá, Colombia⁸Center for Particle Physics, Charles University, Prague, Czech Republic⁹Czech Technical University, Prague, Czech Republic¹⁰Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic¹¹Universidad San Francisco de Quito, Quito, Ecuador¹²Laboratoire de Physique Corpusculaire, IN2P3-CNRS, Université Blaise Pascal, Clermont-Ferrand, France¹³Laboratoire de Physique Subatomique et de Cosmologie, IN2P3-CNRS, Université de Grenoble 1, Grenoble, France¹⁴CPPM, IN2P3-CNRS, Université de la Méditerranée, Marseille, France¹⁵Laboratoire de l'Accélérateur Linéaire, IN2P3-CNRS et Université Paris-Sud, Orsay, France¹⁶LPNHE, IN2P3-CNRS, Universités Paris VI and VII, Paris, France¹⁷DAPNIA/Service de Physique des Particules, CEA, Saclay, France¹⁸IPHC, IN2P3-CNRS, Université Louis Pasteur, Strasbourg, France,

and Université de Haute Alsace, Mulhouse, France

¹⁹Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude Bernard, Villeurbanne, France²⁰III. Physikalisches Institut A, RWTH Aachen, Aachen, Germany²¹Physikalisches Institut, Universität Bonn, Bonn, Germany²²Physikalisches Institut, Universität Freiburg, Freiburg, Germany²³Institut für Physik, Universität Mainz, Mainz, Germany²⁴Ludwig-Maximilians-Universität München, München, Germany²⁵Fachbereich Physik, University of Wuppertal, Wuppertal, Germany

- ²⁶Panjab University, Chandigarh, India
²⁷Delhi University, Delhi, India
²⁸Tata Institute of Fundamental Research, Mumbai, India
²⁹University College Dublin, Dublin, Ireland
³⁰Korea Detector Laboratory, Korea University, Seoul, Korea
³¹SungKyunKwan University, Suwon, Korea
³²CINVESTAV, Mexico City, Mexico
³³FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands
³⁴Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands
³⁵Joint Institute for Nuclear Research, Dubna, Russia
³⁶Institute for Theoretical and Experimental Physics, Moscow, Russia
³⁷Moscow State University, Moscow, Russia
³⁸Institute for High Energy Physics, Protvino, Russia
³⁹Petersburg Nuclear Physics Institute, St. Petersburg, Russia
⁴⁰Lund University, Lund, Sweden,
Royal Institute of Technology and Stockholm University, Stockholm, Sweden,
and Uppsala University, Uppsala, Sweden
⁴¹Physik Institut der Universität Zürich, Zürich, Switzerland
⁴²Lancaster University, Lancaster, United Kingdom
⁴³Imperial College, London, United Kingdom
⁴⁴University of Manchester, Manchester, United Kingdom
⁴⁵University of Arizona, Tucson, Arizona 85721, USA
⁴⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA
⁴⁷California State University, Fresno, California 93740, USA
⁴⁸University of California, Riverside, California 92521, USA
⁴⁹Florida State University, Tallahassee, Florida 32306, USA
⁵⁰Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA
⁵¹University of Illinois at Chicago, Chicago, Illinois 60607, USA
⁵²Northern Illinois University, DeKalb, Illinois 60115, USA
⁵³Northwestern University, Evanston, Illinois 60208, USA
⁵⁴Indiana University, Bloomington, Indiana 47405, USA
⁵⁵University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵⁶Purdue University Calumet, Hammond, Indiana 46323, USA
⁵⁷Iowa State University, Ames, Iowa 50011, USA
⁵⁸University of Kansas, Lawrence, Kansas 66045, USA
⁵⁹Kansas State University, Manhattan, Kansas 66506, USA
⁶⁰Louisiana Tech University, Ruston, Louisiana 71272, USA
⁶¹University of Maryland, College Park, Maryland 20742, USA
⁶²Boston University, Boston, Massachusetts 02215, USA
⁶³Northeastern University, Boston, Massachusetts 02115, USA
⁶⁴University of Michigan, Ann Arbor, Michigan 48109, USA
⁶⁵Michigan State University, East Lansing, Michigan 48824, USA
⁶⁶University of Mississippi, University, Mississippi 38677, USA
⁶⁷University of Nebraska, Lincoln, Nebraska 68588, USA
⁶⁸Princeton University, Princeton, New Jersey 08544, USA
⁶⁹State University of New York, Buffalo, New York 14260, USA
⁷⁰Columbia University, New York, New York 10027, USA
⁷¹University of Rochester, Rochester, New York 14627, USA
⁷²State University of New York, Stony Brook, New York 11794, USA
⁷³Brookhaven National Laboratory, Upton, New York 11973, USA
⁷⁴Langston University, Langston, Oklahoma 73050, USA
⁷⁵University of Oklahoma, Norman, Oklahoma 73019, USA
⁷⁶Oklahoma State University, Stillwater, Oklahoma 74078, USA
⁷⁷Brown University, Providence, Rhode Island 02912, USA
⁷⁸University of Texas, Arlington, Texas 76019, USA
⁷⁹Southern Methodist University, Dallas, Texas 75275, USA
⁸⁰Rice University, Houston, Texas 77005, USA
⁸¹University of Virginia, Charlottesville, Virginia 22901, USA
⁸²University of Washington, Seattle, Washington 98195, USA

(Received 8 January 2007; published 11 April 2007)

We have performed the first direct measurement of the time-integrated flavor untagged charge asymmetry in semileptonic B_s^0 decays $A_{\text{SL}}^{s,\text{unt}}$ by comparing the decay rate of $B_s^0 \rightarrow \mu^+ D_s^- \nu X$, where $D_s^- \rightarrow \phi \pi^-$ and $\phi \rightarrow K^+ K^-$, with the charge-conjugate \bar{B}_s^0 decay rate. This sample was selected from 1.3 fb^{-1} of data collected by the D0 experiment in run II of the Fermilab Tevatron collider. We obtain $A_{\text{SL}}^{s,\text{unt}} = [1.23 \pm 0.97(\text{stat}) \pm 0.17(\text{syst})] \times 10^{-2}$. Assuming that $\Delta m_s/\bar{\Gamma}_s \gg 1$, this result can be translated into a measurement of the CP -violating phase in B_s^0 mixing: $\Delta\Gamma_s/\Delta m_s \tan\phi_s = [2.45 \pm 1.93(\text{stat}) \pm 0.35(\text{syst})] \times 10^{-2}$.

DOI: 10.1103/PhysRevLett.98.151801

PACS numbers: 13.20.He, 11.30.Er, 12.15.Hh, 14.40.Nd

This Letter presents the first measurement of a time-integrated flavor untagged charge asymmetry $A_{\text{SL}}^{s,\text{unt}}$ in semileptonic B_s^0 decays. This asymmetry is defined as

$$A_{\text{SL}}^{s,\text{unt}} = \frac{N(\mu^+ D_s^-) - N(\mu^- D_s^+)}{N(\mu^+ D_s^-) + N(\mu^- D_s^+)}, \quad (1)$$

where $N(\mu^\pm D_s^\mp)$ is the number of decays $B_s^0 \rightarrow \mu^\pm D_s^\mp \nu X$ integrated over the B_s^0 lifetime. This asymmetry is called untagged because the initial flavor of the B_s^0 meson is not determined. $A_{\text{SL}}^{s,\text{unt}}$ is related to CP violation in B_s^0 mixing [1] and can be expressed through the parameters of the B_s^0 mass matrix as [2]

$$A_{\text{SL}}^{s,\text{unt}} = \frac{1}{2} \frac{x_s^2 + y_s^2}{1 + x_s^2} \frac{\Delta\Gamma_s}{\Delta m_s} \tan\phi_s, \quad (2)$$

where $\Delta\Gamma_s$ (Δm_s) is the width (mass) difference between the mass eigenstates in the B_s^0 system, $x_s = \Delta m_s/\bar{\Gamma}_s$, $y_s = \Delta\Gamma_s/(2\bar{\Gamma}_s)$, where $\bar{\Gamma}_s$ is the average width in the B_s^0 system, and ϕ_s is a CP -violating phase. The standard model (SM) predicts a very small value for this asymmetry $2 \times A_{\text{SL}}^{s,\text{unt}} = a_{\text{SL}}^s = (0.21 \pm 0.06) \times 10^{-4}$ [3], while the contribution of new physics can increase its magnitude up to the 1% level [4–6].

This measurement was performed using a large sample of semileptonic B_s^0 decays collected by the D0 experiment at the Fermilab Tevatron collider in $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ and follows closely the procedure used in the estimate of the dimuon asymmetry described in Ref. [7]. The data correspond to an integrated luminosity of approximately 1.3 fb^{-1} . The D0 detector is described in detail elsewhere [8]. The detector components most important to this analysis are the central tracking and muon systems. The central tracking system consists of a silicon microstrip tracker (SMT) and a central fiber tracker (CFT), both located within a 2 T superconducting solenoidal magnet, with designs optimized for tracking and vertexing for pseudorapidities of $|\eta| < 3$ and $|\eta| < 2.5$, respectively [9]. The outer muon system, with coverage for $|\eta| < 2$, consists of a layer of tracking detectors and scintillation trigger counters in front of 1.8 T iron toroids, followed by two similar layers after the toroids [10]. The polarities of the solenoid and toroids are reversed regularly during data taking, so that the four solenoid-toroid polarity combinations are exposed to approximately the same inte-

grated luminosity. The direct and reverse magnetic fields in the magnet were measured to be equal to within 0.1%. The reversal of magnet polarities is essential to reduce the detector-related systematics in asymmetry measurements and is fully exploited in this analysis.

The asymmetry $A_{\text{SL}}^{s,\text{unt}}$ was measured using the decay $B_s^0 \rightarrow \mu D_s \nu X$, with $D_s \rightarrow \phi \pi$, $\phi \rightarrow K^+ K^-$. The selection of this final state is described in detail in Ref. [11]. No explicit trigger requirement was made, although most of the sample was collected with single-muon triggers. Muons were required to have transverse momentum $p_T(\mu) > 2 \text{ GeV}/c$ and momentum $p(\mu) > 3 \text{ GeV}/c$, to have hits in both the CFT and the SMT, and to have measurements in at least two layers of the muon system. All reconstructed charged particles in the event were clustered into jets [12], and the D_s candidate was reconstructed from three tracks found in the same jet as the reconstructed muon. Oppositely charged particles with $p_T > 0.7 \text{ GeV}/c$ were assigned the kaon mass and were required to have an invariant mass $1.004 < M(K^+ K^-) < 1.034 \text{ GeV}/c^2$, consistent with that of a ϕ meson. The third track was required to have $p_T > 0.5 \text{ GeV}/c$, a charge opposite to that of the muon charge, and was assigned the pion mass. The three tracks were required to have hits in the CFT and the SMT and to form a common D_s vertex using the algorithm described in detail in Ref. [13]. To reduce combinatorial background, the D_s vertex was required to have a positive displacement in the transverse plane, relative to the $p\bar{p}$ collision point (or primary vertex), with at least 4σ significance. The cosine of the angle between the D_s momentum and the direction from the primary vertex to the D_s vertex was required to be greater than 0.9. The trajectories of the muon and D_s candidates were required to originate from a common B_s^0 vertex, and the (μD_s) system was required to have an invariant mass between 2.6 and 5.4 GeV/c^2 .

To further improve the B_s^0 signal selection, a likelihood ratio method [14] was utilized. Using background sidebands (B) and sideband-subtracted signal (S) distributions in the data, probability distributions were found for a number of discriminating variables. These variables were the angle between the D_s and K momenta in the $K^+ K^-$ center-of-mass frame, the isolation of the (μD_s) system, the χ^2 of the D_s vertex, the invariant masses $M(\mu D_s)$ and $M(K^+ K^-)$, and $p_T(K^+ K^-)$. The isolation was defined as the ratio of the sum of the momentum of the tracks used to reconstruct the signal divided by the total momentum of

the tracks contained within a cone with $\sqrt{\Delta\eta^2 + \Delta\phi^2} < 0.5$ centered on the direction of the μD_s system. The final requirement on the combined selection likelihood ratio variable was chosen to maximize the predicted ratio $S/\sqrt{S+B}$.

For this analysis, we required the B_s^0 vertex to have a positive displacement from the primary vertex to suppress the combinatoric background from the process $c\bar{c}(b\bar{b}) \rightarrow \mu D_s \nu X$, with the D_s originating from a b or c quark and the muon arising from another quark. The invariant mass distribution $M(\phi\pi)$ for the selected events is shown in Fig. 1. The low and high peaks correspond, respectively, to (μD) , mostly due to B^0 , and (μD_s) , mostly due to B_s^0 . The curve represents a fit to the $M(\phi\pi)$ spectrum. A single Gaussian was sufficient to describe the $D \rightarrow \phi\pi$ decay and a double Gaussian to describe the $D_s \rightarrow \phi\pi$ decay, and the background was modeled by an exponential. The total number of events passing all cuts in the D_s mass peak is $27\,300 \pm 300(\text{stat})$.

To measure $A_{\text{SL}}^{s,\text{unt}}$, both physics and detector effects contributing to the possible imbalance of events with positively and negatively charged muons must be taken into account. One physics source of asymmetry is CP violation in semileptonic B decays. In addition, forward-backward charge asymmetry of events produced in the proton-antiproton collisions can also be present. Detector effects can give rise to an artificial asymmetry if, e.g., the reconstruction efficiencies of positively and negatively charged particles are different. However, a positively charged particle produces the same track as a negatively charged particle in the detector with reversed magnet polarity. Therefore, almost all detector effects can be canceled, provided the fractions of events with opposite magnet polarities are approximately the same. This is the case in this analysis, where the exposures are the same within 1%.

According to the method described in Ref. [7], the event sample was divided into eight subsamples corresponding to all possible combinations of the toroid polarity $\beta = \pm 1$,




FIG. 1. The invariant mass distribution $M(\phi\pi)$ for the selected B_s^0 candidates. The curve shows the result of fit by a function described in the text.

the sign of the pseudorapidity of the $(\mu\phi\pi)$ system $\gamma = \pm 1$, and the sign of the muon charge $q = \pm 1$. The number of (μD_s) events in each subsample was obtained by a fit to the mass distribution $M(\phi\pi)$ using the same function as for the whole sample. For the cross-check, we also extracted the numbers of (μD) and background events from the fit. The widths and positions of the (μD_s) and (μD) peaks, the relative fractions of the two Gaussians describing the (μD_s) peak, as well as the background slope were fixed to the values obtained from the fit to the total $M(\phi\pi)$ distribution. The numbers of (μD_s) and (μD) events, $n_q^{\beta\gamma}(D_s)$ and $n_q^{\beta\gamma}(D)$, along with the number of the background events in the fitting range 1.75–2.30 GeV/c^2 , $n_q^{\beta\gamma}(\text{bkg})$, for each subsample is given in Table I.

The fitted numbers of (μD_s) [(μD) , background] events were used to disentangle the physics asymmetries and the detector effects. The $n_q^{\beta\gamma}$ can be expressed through the physics and the detector asymmetries as follows [7]:

$$n_q^{\beta\gamma} = \frac{1}{4} N \epsilon^\beta (1 + qA)(1 + q\gamma A_{\text{fb}})(1 + \gamma A_{\text{det}}) \times (1 + q\beta\gamma A_{\text{ro}})(1 + q\beta A_{q\beta})(1 + \beta\gamma A_{\beta\gamma}). \quad (3)$$

Here N is the total number of (μD_s) [(μD) , background] events; ϵ^β is the fraction of integrated luminosity with toroid polarity β ($\epsilon^+ + \epsilon^- = 1$); A is the integrated charge asymmetry to be measured; A_{fb} is the forward-backward asymmetry; A_{det} is the detector asymmetry for particles emitted in the forward and backward direction; A_{ro} is the range-out asymmetry that accounts for the change in acceptance of muons which bend towards the beam line and those which bend away from the beam line; $A_{q\beta}$ is the detector asymmetry which accounts for the change in the muon reconstruction efficiency when the toroid polarity is reversed; $A_{\beta\gamma}$ accounts for any detector-related forward-backward asymmetries that remain after the toroid polarity flip.

Since the system (3) contains eight equations, all six asymmetries together with N and ϵ^+ can be extracted for each of the three types of the events. Results are presented in Table II separately for (μD_s) and (μD) events and the

TABLE I. The numbers of events $n_q^{\beta\gamma}(D_s)$ [$n_q^{\beta\gamma}(D)$] in the D_s [D] mass peak and in the background $n_q^{\beta\gamma}(\text{bkg})$ for eight subsamples.

Subsample: $\beta\gamma q$	$n_q^{\beta\gamma}(D_s)$ (events)	$n_q^{\beta\gamma}(D)$ (events)	$n_q^{\beta\gamma}(\text{bkg})$ (events)
+++	3216 ± 76	907 ± 55	9797 ± 124
+-+	3586 ± 79	965 ± 56	$10\,387 \pm 127$
++-	3391 ± 78	1037 ± 57	$10\,390 \pm 127$
+--	3225 ± 76	963 ± 55	9832 ± 124
-++	3616 ± 80	1003 ± 57	$10\,508 \pm 128$
--+	3370 ± 77	801 ± 54	9987 ± 125
-+-	3353 ± 77	831 ± 55	$10\,215 \pm 125$
---	3532 ± 79	1116 ± 59	$10\,701 \pm 129$

TABLE II. The physics and detector asymmetries for (μD_s) , (μD) , and background events. Uncertainties are statistical.

	(μD_s)	(μD)	Background
N	$27\,289 \pm 220$	7623 ± 162	$81\,817 \pm 357$
ϵ^+	0.492 ± 0.004	0.510 ± 0.011	0.494 ± 0.002
A	0.0102 ± 0.0081	-0.0345 ± 0.0211	-0.0056 ± 0.0045
A_{fb}	-0.0046 ± 0.0081	0.0480 ± 0.0210	-0.0020 ± 0.0043
A_{det}	-0.0051 ± 0.0081	-0.0072 ± 0.0212	0.0001 ± 0.0044
A_{ro}	-0.0352 ± 0.0081	-0.0819 ± 0.0209	-0.0263 ± 0.0044
$A_{\beta\gamma}$	-0.0097 ± 0.0081	0.0104 ± 0.0213	-0.0010 ± 0.0044
$A_{q\beta}$	0.0030 ± 0.0081	0.0014 ± 0.0212	0.0046 ± 0.0044

background. The physics asymmetries A and A_{fb} for background events are consistent with zero. This is an important test for this method, since the precision of the asymmetry measurement for the background events is much higher than that of the signal due to the larger statistics. The largest detector asymmetry for all three types of the events is the range-out asymmetry.

It can be seen from (3) that if $\epsilon^+ = \epsilon^- = 1/2$, and the asymmetries A , A_{fb} , A_{det} , A_{ro} , $A_{q\beta}$, and $A_{\beta\gamma}$ are small, each of them can be obtained independently by the appropriate division of the entire sample of events into two parts. For example, the asymmetry A can be obtained by dividing the sample according to the charge of muon. For such a division, and neglecting the second order terms, we obtain

$$A = \frac{n_+ - n_-}{n_+ + n_-}, \quad n_q = \sum_{\beta, \gamma = -1}^{+1} n_q^{\beta\gamma} \simeq \frac{1}{2} N(1 + qA). \quad (4)$$

This observation explains, in particular, the similar values of statistical uncertainties for all asymmetries in Table II.

The resulting charge asymmetry of (μD_s) events is $A = 0.0102 \pm 0.0081(\text{stat})$. It is related to $A_{\text{SL}}^{s, \text{unt}}$ via $A = f_s A_{\text{SL}}^{s, \text{unt}} + f_d A_{\text{SL}}^{d, \text{unt}}$, where $f_s (f_d)$ is the fraction of $B_s^0 (B_d^0) \rightarrow \mu D_s \nu X$ decays in the (μD_s) sample. $A_{\text{SL}}^{d, \text{unt}}$ may arise only from $B_d^0 \rightarrow DD_s$ decay, the fraction of which in the (μD_s) sample was found to be small, at the level of $(4 \pm 1)\%$. Additionally, the value of $A_{\text{SL}}^{d, \text{unt}}$ is strongly constrained experimentally [15,16] to be close to zero. Therefore, the time-integrated $A_{\text{SL}}^{d, \text{unt}}$ component can be neglected. The fraction of B_s^0 decays f_s was determined as follows. The decays $B_s^0 \rightarrow \mu D_s \nu X$ and $B_s^0 \rightarrow \tau D_s \nu X \rightarrow \mu D_s \nu X$ were considered as a signal. The decays $B_s^0 \rightarrow D_s D_s X$, with $D_s \rightarrow \mu \nu X$, are not flavor-specific and, hence, were considered as a background. The decays $B_d^0 \rightarrow DD_s X$ were also included in the background. In addition, the process $c\bar{c}(b\bar{b}) \rightarrow \mu D_s \nu X$ was taken into account. This background produces a pseudovertex, which peaks around the primary interaction point. It is reduced by approximately 50% by requiring a positive displacement of the (μD_s) vertex.

All processes were simulated using the EVTGEN [17] generator interfaced to PYTHIA [18] and followed by full modeling of the detector response using GEANT [19] and event reconstruction. The branching fractions of B_d^0 decays were taken from Ref. [1], while the contribution of the process $c\bar{c}(b\bar{b}) \rightarrow \mu D_s \nu X$ was measured directly in our data to be $(5.9 \pm 1.7)\%$. With these assumptions, $(83.2 \pm 3.3)\%$ of the selected sample of (μD_s) events is composed of semileptonic B_s^0 decays. The uncertainty on this value comes from the uncertainties on the branching ratios of the contributing B decays and the uncertainty on the fraction of the $c\bar{c}(b\bar{b}) \rightarrow \mu D_s \nu X$ process in the sample. Taking into account the sample composition, the measured integrated charge asymmetry of semileptonic B_s^0 decay is found to be $A_{\text{SL}}^{s, \text{unt}} = [1.23 \pm 0.97(\text{stat})] \times 10^{-2}$.

The following sources of systematic uncertainty were considered. The final state includes a $K^+ K^-$ pair. Therefore, the charge asymmetry of K meson reconstruction, which arises due to the different interaction cross sections of K^+ and K^- in the detector material, does not contribute to the measured $A_{\text{SL}}^{s, \text{unt}}$. The charge asymmetry of pion reconstruction, however, can contribute. The πd interaction cross sections for positive and negative pions differ by $(1.3 \pm 0.3)\%$ in the range 1–2 GeV/c [20]. Taking into account the amount of material which a pion crosses in the detector, the induced asymmetry due to pion reconstruction was estimated to be 2×10^{-4} . This value was included in the systematic uncertainty.

The uncertainty in the fraction of B_s^0 signal in the (μD_s) sample produces a systematic uncertainty of 1×10^{-3} . This uncertainty also includes a possible residual variation of the signal fraction between subsamples.

The uncertainty due to the fitting procedure was estimated by varying the masses and widths of the peaks, and the slope of the background, by 1 standard deviation. The fitting procedure was also repeated with a single Gaussian describing the D_s peak and with a different fitting range. The resulting change of $A_{\text{SL}}^{s, \text{unt}}$ did not exceed 0.14×10^{-2} , which was used as an estimate of the systematic uncertainty from this source.

The B_s^0 reconstruction efficiency varies with the decay length due to the applied requirements. We verified that

this variation does not bias the result for $A_{\text{SL}}^{s,\text{unt}}$ and the relation (2). In addition, any possible contribution of the B_d^0 charge asymmetry to the measured value was estimated to be negligible.

Adding all contributions into the systematic uncertainty in quadratures, we obtain the resulting value of the time-integrated untagged charge asymmetry:

$$A_{\text{SL}}^{s,\text{unt}} = [1.23 \pm 0.97(\text{stat}) \pm 0.17(\text{syst})] \times 10^{-2}. \quad (5)$$

This is the first direct measurement of $A_{\text{SL}}^{s,\text{unt}}$. It can be seen that the statistical uncertainty dominates and will be improved in the future with the increase of statistics and addition of new decay modes. Using Eq. (2) and assuming that $\Delta m_s/\bar{\Gamma}_s \gg 1$, we obtain

$$\frac{\Delta\Gamma_s}{\Delta m_s} \tan\phi_s = [2.45 \pm 1.93(\text{stat}) \pm 0.35(\text{syst})] \times 10^{-2}. \quad (6)$$

This result is consistent with the SM prediction [3]. Its combination with the measurements of $\Delta\Gamma_s$ [21], Δm_s [11,22], and dimuon charge asymmetry [7] provides a constraint on the CP -violating phase ϕ_s . Such a combination, which is beyond the scope of this Letter, is described and presented in Ref. [23].

We thank the staffs at Fermilab and collaborating institutions and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom, and RFBR (Russia); CAPES, CNPq, FAPERJ, FAPESP, and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); PPARC (United Kingdom); MSMT (Czech Republic); CRC Program, CFI, NSERC, and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); Research Corporation; Alexander von Humboldt Foundation; and the Marie Curie Program.

[1] W.-M. Yao *et al.* (Particle Data Group), J. Phys. G **33**, 1 (2006).

- [2] U. Nierste, hep-ph/0406300.
- [3] M. Beneke, G. Buchalla, A. Lenz, and U. Nierste, Phys. Lett. B **576**, 173 (2003); A. Lenz and U. Nierste, hep-ph/0612167.
- [4] Y. Grossman, Y. Nir, and R. Rattazzi, in *Heavy Flavours II*, edited by A. Buras and M. Lindner (World Scientific, Singapore, 1998).
- [5] F. Gabbiani, E. Gabrieli, A. Masiero, and L. Silvestrini, Nucl. Phys. **B477**, 321 (1996).
- [6] W. S. Hou, M. Nagashima, and A. Soddu, hep-ph/0610385.
- [7] V. Abazov *et al.* (D0 Collaboration), Phys. Rev. D **74**, 092001 (2006).
- [8] V. Abazov *et al.* (D0 Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **565**, 463 (2006).
- [9] Pseudorapidity η is defined as $\eta = -\ln[\tan(\theta/2)]$, where θ is the polar angle of the particle with respect to the proton beam direction.
- [10] V. Abazov *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **552**, 372 (2005).
- [11] V. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **97**, 021802 (2006).
- [12] S. Catani *et al.*, Phys. Lett. B **269**, 432 (1991), ‘‘Durham’’ jets with the p_T cutoff parameter set at 15 GeV/ c .
- [13] J. Abdallah *et al.* (DELPHI Collaboration), Eur. Phys. J. C **32**, 185 (2004).
- [14] G. Borisov, Nucl. Instrum. Methods Phys. Res., Sect. A **417**, 384 (1998).
- [15] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. Lett. **96**, 251802 (2006).
- [16] E. Nakano *et al.* (Belle Collaboration), Phys. Rev. D **73**, 112002 (2006).
- [17] D. J. Lange, Nucl. Instrum. Methods Phys. Res., Sect. A **462**, 152 (2001).
- [18] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001).
- [19] CERN Program Library Long Writeup W5013 (1993); documentation available at <http://www.wasd.web.cern.ch/wwwasd/geant/>.
- [20] A. Carter *et al.*, Phys. Rev. **168**, 1457 (1968).
- [21] V. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **98**, 121801 (2007); $|\Delta\Gamma_s| \cos\phi_s|$ was measured in D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **94**, 101803 (2005); V. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **95**, 171801 (2005).
- [22] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. Lett. **97**, 062003 (2006).
- [23] V. M. Abazov *et al.* (D0 Collaboration), hep-ex/0702030.