

Measurements of CP -Violating Asymmetries in $B^0 \rightarrow a_1^+(1260)\pi^\mp$ Decays

- B. Aubert,¹ M. Bona,¹ D. Boutigny,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ X. Prudent,¹ V. Tisserand,¹ A. Zghiche,¹ E. Grauges,² A. Palano,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴ Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ M. Battaglia,⁶ D. N. Brown,⁶ J. Button-Shafer,⁶ R. N. Cahn,⁶ Y. Groysman,⁶ R. G. Jacobsen,⁶ J. A. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomensky,⁶ G. Kukartsev,⁶ D. Lopes Pegna,⁶ G. Lynch,⁶ L. M. Mir,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,^{6,*} K. Tackmann,⁶ W. A. Wenzel,⁶ P. del Amo Sanchez,⁷ M. Barrett,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷ A. T. Watson,⁷ T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸ T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹ J. P. Burke,⁹ W. N. Cottingham,⁹ D. Walker,⁹ D. J. Asgeirsson,¹⁰ T. Cuhadar-Donszelmann,¹⁰ B. G. Fulsom,¹⁰ C. Hearty,¹⁰ N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹ M. Saleem,¹¹ D. J. Sherwood,¹¹ L. Teodorescu,¹¹ V. E. Blinov,¹² A. D. Bukin,¹² V. P. Druzhinin,¹² V. B. Golubev,¹² A. P. Onuchin,¹² S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² K. Yu Todyshev,¹² M. Bondioli,¹³ M. Bruinsma,¹³ M. Chao,¹³ S. Curry,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³ P. Lund,¹³ M. Mandelkern,¹³ E. C. Martin,¹³ D. P. Stoker,¹³ S. Abachi,¹⁴ C. Buchanan,¹⁴ S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ F. Liu,¹⁵ O. Long,¹⁵ B. C. Shen,¹⁵ L. Zhang,¹⁵ E. J. Hill,¹⁶ H. P. Paar,¹⁶ S. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷ A. Cunha,¹⁷ B. Dahmes,¹⁷ T. M. Hong,¹⁷ D. Kovalskyi,¹⁷ J. D. Richman,¹⁷ T. W. Beck,¹⁸ A. M. Eisner,¹⁸ C. J. Flacco,¹⁸ C. A. Heusch,¹⁸ J. Kroeseberg,¹⁸ W. S. Lockman,¹⁸ T. Schalk,¹⁸ B. A. Schumm,¹⁸ A. Seiden,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸ L. O. Winstrom,¹⁸ E. Chen,¹⁹ C. H. Cheng,¹⁹ A. Dvoretskii,¹⁹ F. Fang,¹⁹ D. G. Hitlin,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ G. Mancinelli,²⁰ B. T. Meadows,²⁰ K. Mishra,²⁰ M. D. Sokoloff,²⁰ F. Blanc,²¹ P. C. Bloom,²¹ S. Chen,²¹ W. T. Ford,²¹ J. F. Hirschauer,²¹ A. Kreisel,²¹ M. Nagel,²¹ U. Nauenberg,²¹ A. Olivas,²¹ J. G. Smith,²¹ K. A. Ulmer,²¹ S. R. Wagner,²¹ J. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² A. Soffer,²² W. H. Toki,²² R. J. Wilson,²² F. Winkelmeier,²² Q. Zeng,²² D. D. Altenburg,²³ E. Feltresi,²³ A. Hauke,²³ H. Jasper,²³ J. Merkel,²³ A. Petzold,²³ B. Spaan,²³ K. Wacker,²³ T. Brandt,²⁴ V. Klose,²⁴ H. M. Lacker,²⁴ W. F. Mader,²⁴ R. Nogowski,²⁴ J. Schubert,²⁴ K. R. Schubert,²⁴ R. Schwierz,²⁴ J. E. Sundermann,²⁴ A. Volk,²⁴ D. Bernard,²⁵ G. R. Bonneau,²⁵ E. Latour,²⁵ Ch. Thiebaut,²⁵ M. Verderi,²⁵ P. J. Clark,²⁶ W. Gradl,²⁶ F. Muheim,²⁶ S. Playfer,²⁶ A. I. Robertson,²⁶ Y. Xie,²⁶ M. Andreotti,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Luppi,²⁷ M. Negrini,²⁷ A. Petrella,²⁷ L. Piemontese,²⁷ E. Prencipe,²⁷ F. Anulli,²⁸ R. Baldini-Ferroli,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ G. Finocchiaro,²⁸ S. Pacetti,²⁸ P. Patteri,²⁸ I. M. Peruzzi,²⁸ M. Piccolo,²⁸ M. Rama,²⁸ A. Zallo,²⁹ A. Buzzo,²⁹ R. Contri,²⁹ M. Lo Vetere,²⁹ M. M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰ J. Wu,³⁰ R. S. Dubitzky,³¹ J. Marks,³¹ S. Schenk,³¹ U. Uwer,³¹ D. J. Bard,³² P. D. Dauncey,³² R. L. Flack,³² J. A. Nash,³² M. B. Nikolich,³² W. Panduro Vazquez,³² P. K. Behera,³³ X. Chai,³³ M. J. Charles,³³ U. Mallik,³³ N. T. Meyer,³³ V. Ziegler,³³ J. Cochran,³⁴ H. B. Crawley,³⁴ L. Dong,³⁴ V. Eyges,³⁴ W. T. Meyer,³⁴ S. Prell,³⁴ E. I. Rosenberg,³⁴ A. E. Rubin,³⁴ A. V. Gritsan,³⁵ A. G. Denig,³⁶ M. Fritsch,³⁶ G. Schott,³⁶ N. Arnaud,³⁷ M. Davier,³⁷ G. Grosdidier,³⁷ A. Höcker,³⁷ V. Lepeltier,³⁷ F. Le Diberder,³⁷ A. M. Lutz,³⁷ S. Pruvot,³⁷ S. Rodier,³⁷ P. Roudeau,³⁷ M. H. Schune,³⁷ J. Serrano,³⁷ V. Sordini,³⁷ A. Stocchi,³⁷ W. F. Wang,³⁷ G. Wormser,³⁷ D. J. Lange,³⁸ D. M. Wright,³⁸ C. A. Chavez,³⁹ I. J. Forster,³⁹ J. R. Fry,³⁹ E. Gabathuler,³⁹ R. Gamet,³⁹ D. E. Hutchcroft,³⁹ D. J. Payne,³⁹ K. C. Schofield,³⁹ C. Touramanis,³⁹ A. J. Bevan,⁴⁰ K. A. George,⁴⁰ F. Di Lodovico,⁴⁰ W. Menges,⁴⁰ R. Sacco,⁴⁰ G. Cowan,⁴¹ H. U. Flaecher,⁴¹ D. A. Hopkins,⁴¹ P. S. Jackson,⁴¹ T. R. McMahon,⁴¹ F. Salvatore,⁴¹ A. C. Wren,⁴¹ D. N. Brown,⁴² C. L. Davis,⁴² J. Allison,⁴³ N. R. Barlow,⁴³ R. J. Barlow,⁴³ Y. M. Chia,⁴³ C. L. Edgar,⁴³ G. D. Lafferty,⁴³ T. J. West,⁴³ J. I. Yi,⁴³ C. Chen,⁴⁴ W. D. Hulsbergen,⁴⁴ A. Jawahery,⁴⁴ C. K. Lae,⁴⁴ D. A. Roberts,⁴⁴ G. Simi,⁴⁴ G. Blaylock,⁴⁵ C. Dallapiccola,⁴⁵ S. S. Hertzbach,⁴⁵ X. Li,⁴⁵ T. B. Moore,⁴⁵ E. Salvati,⁴⁵ S. Saremi,⁴⁵ R. Cowan,⁴⁶ G. Sciolla,⁴⁶ S. J. Sekula,⁴⁶ M. Spitznagel,⁴⁶ F. Taylor,⁴⁶ R. K. Yamamoto,⁴⁶ H. Kim,⁴⁷ S. E. Mclachlin,⁴⁷ P. M. Patel,⁴⁷ S. H. Robertson,⁴⁷ A. Lazzaro,⁴⁸ V. Lombardo,⁴⁸ F. Palombo,⁴⁸ J. M. Bauer,⁴⁹ L. Cremaldi,⁴⁹ V. Eschenburg,⁴⁹ R. Godang,⁴⁹ R. Kroeger,⁴⁹ D. A. Sanders,⁴⁹ D. J. Summers,⁴⁹ H. W. Zhao,⁴⁹ S. Brunet,⁵⁰ D. Côté,⁵⁰ M. Simard,⁵⁰ P. Taras,⁵⁰ F. B. Viaud,⁵⁰ H. Nicholson,⁵¹ N. Cavallo,⁵²[‡] G. De Nardo,⁵² F. Fabozzi,⁵²[‡] C. Gatto,⁵² L. Lista,⁵² D. Monorchio,⁵² P. Paolucci,⁵² D. Piccolo,⁵² C. Sciacca,⁵² M. A. Baak,⁵³ G. Raven,⁵³ H. L. Snoek,⁵³ C. P. Jessop,⁵⁴ J. M. LoSecco,⁵⁴ G. Benelli,⁵⁵ L. A. Corwin,⁵⁵ K. K. Gan,⁵⁵ K. Honscheid,⁵⁵ D. Hufnagel,⁵⁵ H. Kagan,⁵⁵ R. Kass,⁵⁵ J. P. Morris,⁵⁵ A. M. Rahimi,⁵⁵ J. J. Regensburger,⁵⁵ R. Ter-Antonyan,⁵⁵ Q. K. Wong,⁵⁵ N. L. Blount,⁵⁶ J. Brau,⁵⁶ R. Frey,⁵⁶ O. Igolkina,⁵⁶ J. A. Kolb,⁵⁶ M. Lu,⁵⁶ C. T. Potter,⁵⁶ R. Rahmat,⁵⁶ N. B. Sinev,⁵⁶ D. Strom,⁵⁶ J. Strube,⁵⁶ E. Torrence,⁵⁶ A. Gaz,⁵⁷ M. Margoni,⁵⁷ M. Morandin,⁵⁷ A. Pompili,⁵⁷ M. Posocco,⁵⁷ M. Rotondo,⁵⁷ F. Simonetto,⁵⁷ R. Stroili,⁵⁷ C. Voci,⁵⁷ E. Ben-Haim,⁵⁸

H. Briand,⁵⁸ J. Chauveau,⁵⁸ P. David,⁵⁸ L. Del Buono,⁵⁸ Ch. de la Vaissière,⁵⁸ O. Hamon,⁵⁸ B. L. Hartfiel,⁵⁸ Ph. Leruste,⁵⁸ J. Malclès,⁵⁸ J. Ocariz,⁵⁸ L. Gladney,⁵⁹ M. Biasini,⁶⁰ R. Covarelli,⁶⁰ C. Angelini,⁶¹ G. Batignani,⁶¹ S. Bettarini,⁶¹ G. Calderini,⁶¹ M. Carpinelli,⁶¹ R. Cenci,⁶¹ F. Forti,⁶¹ M. A. Giorgi,⁶¹ A. Lusiani,⁶¹ G. Marchiori,⁶¹ M. A. Mazur,⁶¹ M. Morganti,⁶¹ N. Neri,⁶¹ E. Paoloni,⁶¹ G. Rizzo,⁶¹ J. J. Walsh,⁶¹ M. Haire,⁶² J. Biesiada,⁶³ P. Elmer,⁶³ Y. P. Lau,⁶³ C. Lu,⁶³ J. Olsen,⁶³ A. J. S. Smith,⁶³ A. V. Telnov,⁶³ F. Bellini,⁶⁴ G. Cavoto,⁶⁴ A. D'Orazio,⁶⁴ D. del Re,⁶⁴ E. Di Marco,⁶⁴ R. Faccini,⁶⁴ F. Ferrarotto,⁶⁴ F. Ferroni,⁶⁴ M. Gaspero,⁶⁴ P. D. Jackson,⁶⁴ L. Li Gioi,⁶⁴ M. A. Mazzoni,⁶⁴ S. Morganti,⁶⁴ G. Piredda,⁶⁴ F. Polci,⁶⁴ C. Voena,⁶⁴ M. Ebert,⁶⁵ H. Schröder,⁶⁵ R. Waldi,⁶⁵ T. Adye,⁶⁶ G. Castelli,⁶⁶ B. Franek,⁶⁶ E. O. Olaiya,⁶⁶ S. Ricciardi,⁶⁶ W. Roethel,⁶⁶ F. F. Wilson,⁶⁶ R. Aleksan,⁶⁷ S. Emery,⁶⁷ M. Escalier,⁶⁷ A. Gaidot,⁶⁷ S. F. Ganzhur,⁶⁷ G. Hamel de Monchenault,⁶⁷ W. Kozanecki,⁶⁷ M. Legendre,⁶⁷ G. Vasseur,⁶⁷ Ch. Yèche,⁶⁷ M. Zito,⁶⁷ X. R. Chen,⁶⁸ H. Liu,⁶⁸ W. Park,⁶⁸ M. V. Purohit,⁶⁸ J. R. Wilson,⁶⁸ M. T. Allen,⁶⁹ D. Aston,⁶⁹ R. Bartoldus,⁶⁹ P. Bechtle,⁶⁹ N. Berger,⁶⁹ R. Claus,⁶⁹ J. P. Coleman,⁶⁹ M. R. Convery,⁶⁹ J. C. Dingfelder,⁶⁹ J. Dorfan,⁶⁹ G. P. Dubois-Felsmann,⁶⁹ D. Dujmic,⁶⁹ W. Dunwoodie,⁶⁹ R. C. Field,⁶⁹ T. Glanzman,⁶⁹ S. J. Gowdy,⁶⁹ M. T. Graham,⁶⁹ P. Grenier,⁶⁹ V. Halyo,⁶⁹ C. Hast,⁶⁹ T. Hryna,⁶⁹ W. R. Innes,⁶⁹ M. H. Kelsey,⁶⁹ P. Kim,⁶⁹ D. W. G. S. Leith,⁶⁹ S. Li,⁶⁹ S. Luitz,⁶⁹ V. Luth,⁶⁹ H. L. Lynch,⁶⁹ D. B. MacFarlane,⁶⁹ H. Marsiske,⁶⁹ R. Messner,⁶⁹ D. R. Muller,⁶⁹ C. P. O'Grady,⁶⁹ V. E. Ozcan,⁶⁹ A. Perazzo,⁶⁹ M. Perl,⁶⁹ T. Pulliam,⁶⁹ B. N. Ratcliff,⁶⁹ A. Roodman,⁶⁹ A. A. Salnikov,⁶⁹ R. H. Schindler,⁶⁹ J. Schwiening,⁶⁹ A. Snyder,⁶⁹ J. Stelzer,⁶⁹ D. Su,⁶⁹ M. K. Sullivan,⁶⁹ K. Suzuki,⁶⁹ S. K. Swain,⁶⁹ J. M. Thompson,⁶⁹ J. Va'vra,⁶⁹ N. van Bakel,⁶⁹ A. P. Wagner,⁶⁹ M. Weaver,⁶⁹ W. J. Wisniewski,⁶⁹ M. Wittgen,⁶⁹ D. H. Wright,⁶⁹ H. W. Wulsin,⁶⁹ A. K. Yarritu,⁶⁹ K. Yi,⁶⁹ C. C. Young,⁶⁹ P. R. Burchat,⁷⁰ A. J. Edwards,⁷⁰ S. A. Majewski,⁷⁰ B. A. Petersen,⁷⁰ L. Wilden,⁷⁰ S. Ahmed,⁷¹ M. S. Alam,⁷¹ R. Bula,⁷¹ J. A. Ernst,⁷¹ V. Jain,⁷¹ B. Pan,⁷¹ M. A. Saeed,⁷¹ F. R. Wappler,⁷¹ S. B. Zain,⁷¹ W. Bugg,⁷² M. Krishnamurthy,⁷² S. M. Spanier,⁷² R. Eckmann,⁷³ J. L. Ritchie,⁷³ C. J. Schilling,⁷³ R. F. Schwitters,⁷³ J. M. Izen,⁷⁴ X. C. Lou,⁷⁴ S. Ye,⁷⁴ F. Bianchi,⁷⁵ F. Gallo,⁷⁵ D. Gamba,⁷⁵ M. Pelliccioni,⁷⁵ M. Bomben,⁷⁶ L. Bosisio,⁷⁶ C. Cartaro,⁷⁶ F. Cossutti,⁷⁶ G. Della Ricca,⁷⁶ L. Lanceri,⁷⁶ L. Vitale,⁷⁶ V. Azzolini,⁷⁷ N. Lopez-March,⁷⁷ F. Martinez-Vidal,⁷⁷ A. Oyanguren,⁷⁷ J. Albert,⁷⁸ Sw. Banerjee,⁷⁸ B. Bhuyan,⁷⁸ K. Hamano,⁷⁸ R. Kowalewski,⁷⁸ I. M. Nugent,⁷⁸ J. M. Roney,⁷⁸ R. J. Sobie,⁷⁸ J. J. Back,⁷⁹ P. F. Harrison,⁷⁹ T. E. Latham,⁷⁹ G. B. Mohanty,⁷⁹ M. Pappagallo,⁷⁹,⁸ H. R. Band,⁸⁰ X. Chen,⁸⁰ S. Dasu,⁸⁰ K. T. Flood,⁸⁰ J. J. Hollar,⁸⁰ P. E. Kutter,⁸⁰ B. Mellado,⁸⁰ Y. Pan,⁸⁰ M. Pierini,⁸⁰ R. Prepost,⁸⁰ S. L. Wu,⁸⁰ Z. Yu,⁸⁰ and H. Neal⁸¹

(BABAR Collaboration)

¹Laboratoire de Physique des Particules, IN2P3/CNRS et Université de Savoie, F-74941 Annecy-Le-Vieux, France²Facultat de Fisica, Departament ECM, Universitat de Barcelona, E-08028 Barcelona, Spain³Dipartimento di Fisica and INFN, Università di Bari, I-70126 Bari, Italy⁴Institute of High Energy Physics, Beijing 100039, China⁵Institute of Physics, University of Bergen, N-5007 Bergen, Norway⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom⁸Institut für Experimentalphysik I, Ruhr Universität Bochum, D-44780 Bochum, Germany⁹University of Bristol, Bristol BS8 1TL, United Kingdom¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia¹³University of California at Irvine, Irvine, California 92697, USA¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA¹⁵University of California at Riverside, Riverside, California 92521, USA¹⁶University of California at San Diego, La Jolla, California 92093, USA¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA¹⁸University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA¹⁹California Institute of Technology, Pasadena, California 91125, USA²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA²¹University of Colorado, Boulder, Colorado 80309, USA²²Colorado State University, Fort Collins, Colorado 80523, USA²³Institut für Physik, Universität Dortmund, D-44221 Dortmund, Germany²⁴Institut für Kern- und Teilchenphysik, Technische Universität Dresden, D-01062 Dresden, Germany²⁵Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France²⁶University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom²⁷Dipartimento di Fisica and INFN, Università di Ferrara, I-44100 Ferrara, Italy

²⁸*Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy*²⁹*Dipartimento di Fisica and INFN, Università di Genova, I-16146 Genova, Italy*³⁰*Harvard University, Cambridge, Massachusetts 02138, USA*³¹*Physikalisches Institut, Universität Heidelberg, Philosophenweg 12, D-69120 Heidelberg, Germany*³²*Imperial College London, London, SW7 2AZ, United Kingdom*³³*University of Iowa, Iowa City, Iowa 52242, USA*³⁴*Iowa State University, Ames, Iowa 50011-3160, USA*³⁵*Johns Hopkins University, Baltimore, Maryland 21218, USA*³⁶*Institut für Experimentelle Kernphysik, Universität Karlsruhe, D-76021 Karlsruhe, Germany*³⁷*Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, B. P. 34, F-91898 ORSAY Cedex, France*³⁸*Lawrence Livermore National Laboratory, Livermore, California 94550, USA*³⁹*University of Liverpool, Liverpool L69 7ZE, United Kingdom*⁴⁰*Queen Mary, University of London, E1 4NS, United Kingdom*⁴¹*University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom*⁴²*University of Louisville, Louisville, Kentucky 40292, USA*⁴³*University of Manchester, Manchester M13 9PL, United Kingdom*⁴⁴*University of Maryland, College Park, Maryland 20742, USA*⁴⁵*University of Massachusetts, Amherst, Massachusetts 01003, USA*⁴⁶*Laboratory for Nuclear Science, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*⁴⁷*McGill University, Montréal, Québec, Canada H3A 2T8*⁴⁸*Dipartimento di Fisica and INFN, Università di Milano, I-20133 Milano, Italy*⁴⁹*University of Mississippi, University, Mississippi 38677, USA*⁵⁰*Physique des Particules, Université de Montréal, Montréal, Québec, Canada H3C 3J7*⁵¹*Mount Holyoke College, South Hadley, Massachusetts 01075, USA*⁵²*Dipartimento di Scienze Fisiche and INFN, Università di Napoli Federico II, I-80126, Napoli, Italy*⁵³*NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands*⁵⁴*University of Notre Dame, Notre Dame, Indiana 46556, USA*⁵⁵*The Ohio State University, Columbus, Ohio 43210, USA*⁵⁶*University of Oregon, Eugene, Oregon 97403, USA*⁵⁷*Dipartimento di Fisica and INFN, Università di Padova, I-35131 Padova, Italy*⁵⁸*Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie-Paris6, Université Denis Diderot-Paris 7, F-75252 Paris, France*⁵⁹*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*⁶⁰*Dipartimento di Fisica and INFN, Università di Perugia, I-06100 Perugia, Italy*⁶¹*Dipartimento di Fisica, Università di Pisa, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy*⁶²*Prairie View A&M University, Prairie View, Texas 77446, USA*⁶³*Princeton University, Princeton, New Jersey 08544, USA*⁶⁴*Dipartimento di Fisica and INFN, Università di Roma La Sapienza, I-00185 Roma, Italy*⁶⁵*Universität Rostock, D-18051 Rostock, Germany*⁶⁶*Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom*⁶⁷*DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France*⁶⁸*University of South Carolina, Columbia, South Carolina 29208, USA*⁶⁹*Stanford Linear Accelerator Center, Stanford, California 94309, USA*⁷⁰*Stanford University, Stanford, California 94305-4060, USA*⁷¹*State University of New York, Albany, New York 12222, USA*⁷²*University of Tennessee, Knoxville, Tennessee 37996, USA*⁷³*University of Texas at Austin, Austin, Texas 78712, USA*⁷⁴*University of Texas at Dallas, Richardson, Texas 75083, USA*⁷⁵*Dipartimento di Fisica Sperimentale and INFN, Università di Torino, I-10125 Torino, Italy*⁷⁶*Dipartimento di Fisica and INFN, Università di Trieste, I-34127 Trieste, Italy*⁷⁷*IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain*⁷⁸*University of Victoria, Victoria, British Columbia, Canada V8W 3P6*⁷⁹*Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom*⁸⁰*University of Wisconsin, Madison, Wisconsin 53706, USA*⁸¹*Yale University, New Haven, Connecticut 06511, USA*

(Received 21 December 2006; published 2 May 2007)

We present measurements of CP -violating asymmetries in the decay $B^0 \rightarrow a_1^\pm(1260)\pi^\mp$ with $a_1^\pm(1260) \rightarrow \pi^\mp\pi^\pm\pi^\pm$. The data sample corresponds to $384 \times 10^6 B\bar{B}$ pairs collected with the *BABAR* detector at the PEP-II asymmetric B factory at SLAC. We measure the CP -violating asymmetry $\mathcal{A}_{CP}^{a_1\pi} = -0.07 \pm 0.07 \pm 0.02$, the mixing-induced CP violation parameter $S_{a_1\pi} = 0.37 \pm 0.21 \pm 0.07$, the direct

CP violation parameter $C_{a_1\pi} = -0.10 \pm 0.15 \pm 0.09$, and the parameters $\Delta C_{a_1\pi} = 0.26 \pm 0.15 \pm 0.07$ and $\Delta S_{a_1\pi} = -0.14 \pm 0.21 \pm 0.06$. From these measured quantities we determine the angle $\alpha_{\text{eff}} = 78.6^\circ \pm 7.3^\circ$.

DOI: 10.1103/PhysRevLett.98.181803

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh

The angle $\alpha \equiv \arg[-V_{td}V_{tb}^*/V_{ud}V_{ub}^*]$ of the unitarity triangle of the Cabibbo-Kobayashi-Maskawa (CKM) quark-mixing matrix [1] has recently been measured by the *BABAR* and *Belle* Collaborations from time-dependent CP asymmetries in the B^0 decays to $\pi^+\pi^-$ [2], $\rho^\pm\pi^\mp$ [3], and $\rho^+\rho^-$ [4]. The decay B^0 to $a_1\pi$ [5] proceeds dominantly through the $\bar{b} \rightarrow \bar{u}u\bar{d}$ process in the same way as the previously studied modes [6,7]. However, due to the presence of additional loop contributions, these measurements determine an effective value α_{eff} , rather than α itself. This obstacle can be overcome using isospin symmetry [8], with bounds to $\Delta\alpha = \alpha - \alpha_{\text{eff}}$ determined using either an isospin analysis [9] or broken SU(3) flavor symmetry [10]. Because it has the smallest contribution from loop diagrams, the $B^0 \rightarrow \rho^+\rho^-$ decay currently allows the most precise single determination of α [11]. The *BABAR* collaboration recently reported the observation of $B^0 \rightarrow a_1^\pm\pi^\mp$ [12]. The state $a_1^\pm\pi^\mp$, like $\rho^\pm\pi^\mp$, is not a CP eigenstate and four flavor-charge configurations must be considered [$B^0(\bar{B}^0) \rightarrow a_1^\pm\pi^\mp$]. Theoretical bounds on $\Delta\alpha$ in these decay modes based on SU(3) flavor symmetry have been derived in Ref. [7].

In this Letter we report measurements of the CP parameters in the decay $B^0 \rightarrow a_1^\pm\pi^\mp$ with $a_1^\pm \rightarrow \pi^\mp\pi^\pm\pi^\pm$. The analysis is done in the quasi-two-body approximation. Details on the reconstruction and handling of the a_1 meson can be found in Ref. [12]. The data were collected with the *BABAR* detector [13] at the PEP-II asymmetric e^+e^- collider. An integrated luminosity of 349 fb^{-1} , corresponding to $(384 \pm 4) \times 10^6 \bar{B}\bar{B}$ pairs, was recorded near the Y(4S) resonance (“on resonance”) at a center-of-mass (c.m.) energy $\sqrt{s} = 10.58 \text{ GeV}$. An additional 37 fb^{-1} were taken about 40 MeV below this energy (“off resonance”) for the study of continuum background in which a charm or lighter quark pair is produced.

From a candidate $B\bar{B}$ pair we reconstruct a B^0 decaying into the final state $f = a_1\pi$ ($B_{a_1\pi}^0$). We also reconstruct the vertex of the other B meson (B_{tag}^0) and identify its flavor. The difference $\Delta t \equiv t_{a_1\pi} - t_{\text{tag}}$ of the proper decay times of the reconstructed and tag B mesons, respectively, is obtained from the measured distance between the $B_{a_1\pi}^0$ and B_{tag}^0 decay vertices and from the boost ($\beta\gamma = 0.56$) of the e^+e^- system. The Δt distributions are given [7] by

$$\begin{aligned} F_{Q_{\text{tag}}}^{a_1^\pm\pi^\mp}(\Delta t) &= (1 \pm \mathcal{A}_{CP}^{a_1\pi}) \frac{e^{-|\Delta t|/\tau}}{4\tau} \{1 - Q_{\text{tag}}\Delta w \\ &\quad + Q_{\text{tag}}(1 - 2w)[(S_{a_1\pi} \pm \Delta S_{a_1\pi}) \sin(\Delta m_d \Delta t) \\ &\quad - (C_{a_1\pi} \pm \Delta C_{a_1\pi}) \cos(\Delta m_d \Delta t)]\}, \end{aligned} \quad (1)$$

where $Q_{\text{tag}} = 1(-1)$ when the tagging meson B_{tag}^0 is a $B^0(\bar{B}^0)$, τ is the mean B^0 lifetime, Δm_d is the mass difference between the two B^0 mass eigenstates, and the mistag parameters w and Δw are the average and difference, respectively, of the probabilities that a true B^0 is incorrectly tagged as a \bar{B}^0 or vice versa. The time- and flavor-integrated charge asymmetry $\mathcal{A}_{CP}^{a_1\pi}$ measures direct CP violation. The quantities $S_{a_1\pi}$ and $C_{a_1\pi}$ parametrize the mixing-induced CP violation related to the angle α , and flavor-dependent direct CP violation, respectively. The parameter $\Delta C_{a_1\pi}$ describes the asymmetry between the rates $\Gamma(B^0 \rightarrow a_1^+\pi^-) + \Gamma(\bar{B}^0 \rightarrow a_1^-\pi^+)$ and $\Gamma(B^0 \rightarrow a_1^-\pi^+) + \Gamma(\bar{B}^0 \rightarrow a_1^+\pi^-)$, while $\Delta S_{a_1\pi}$ is related to the strong phase difference between the amplitudes contributing to $B^0 \rightarrow a_1\pi$ decays. The parameters $\Delta C_{a_1\pi}$ and $\Delta S_{a_1\pi}$ are insensitive to CP violation. The flavor-tagging algorithm uses six mutually exclusive categories. Its analyzing power is measured to be $(30.4 \pm 0.3)\%$ [14].

Charged particles are detected and their momenta measured by the combination of a silicon vertex tracker (SVT), consisting of five layers of double-sided silicon detectors, and a 40-layer central drift chamber, both operating in the 1.5 T magnetic field of a superconducting solenoid. Charged-particle identification (PID) is provided by the average energy loss (dE/dx) in the tracking devices and by an internally reflecting ring-imaging Cherenkov detector (DIRC) covering the central region. Separation between pions and kaons is achieved at the level of 4 standard deviations (σ) for momenta below 3 GeV, decreasing to 2.5σ at 4 GeV.

Full Monte Carlo (MC) simulations [15] of the signal decay modes, continuum, and $B\bar{B}$ backgrounds are used to establish the event selection criteria. The MC signal events are simulated as B^0 decays to $a_1\pi$ with $a_1 \rightarrow \rho\pi$. For the a_1 meson parameters we take the mass $m_0 = 1230 \text{ MeV}$ and the width $\Gamma_0 = 400 \text{ MeV}$ [16,17].

We reconstruct the decay $a_1 \rightarrow 3\pi$ with the following requirement on the invariant mass: $0.87 < m_{a_1} < 1.8 \text{ GeV}$. The intermediate dipion state is reconstructed with an invariant mass between 0.51 and 1.1 GeV. We impose several PID requirements to ensure the identity of the signal pions. For the decay pion coming from the B meson we require the measured Cherenkov angle to be within -2σ and $+5\sigma$ from the expected value for a pion. This requirement removes 98.6% of the background from a_1K . A B candidate is characterized kinematically by the energy-substituted mass $m_{\text{ES}} = \sqrt{(s/2 + \mathbf{p}_0 \cdot \mathbf{p}_B)^2/E_0^2 - \mathbf{p}_B^2}$ and energy difference $\Delta E = E_B^* - \frac{1}{2}\sqrt{s}$, where the subscripts 0 and B refer to the initial Y(4S) and to the B candidate in

the laboratory frame, respectively, and the asterisk denotes the c.m. frame. The resolutions in m_{ES} and in ΔE are about 3.0 and 20 MeV, respectively. We require $|\Delta E| \leq 0.1$ GeV and $5.25 \leq m_{\text{ES}} \leq 5.29$ GeV. To reduce the number of false B -meson candidates we require that the probability of the B vertex fit be greater than 0.01.

To reject continuum background, we use the angle θ_T between the thrust axis of the B candidate and that of the rest of the tracks and neutral clusters in the event, calculated in the center-of-mass frame. The distribution of $\cos\theta_T$ is sharply peaked near ± 1 for $q\bar{q}$ candidates, which have a jetlike topology, and is nearly uniform for the isotropic B -meson decays. We require $|\cos\theta_T| < 0.65$. The absolute value of the cosine of the angle between the direction of the π meson from $a_1 \rightarrow \rho\pi$ with respect to the flight direction of the B in the a_1 meson rest frame is required to be less than 0.85 to suppress combinatorial background. The distribution of this variable is uniform for signal and peaks near unity for this background. We discriminate further against $q\bar{q}$ background with a Fisher discriminant \mathcal{F} that combines several variables that characterize the production dynamics and energy flow in the event [18]. The remaining continuum background is modeled from off-resonance data.

We use MC simulations of $B^0\bar{B}^0$ and B^+B^- decays to look for $B\bar{B}$ backgrounds, which can come from B decays with or without charmed particles in the final state. Neutral and charged D mesons may contribute to background through particle misidentification or misreconstruction. We remove any combinations of the decay products, including possible additional π^0 , with invariant mass consistent with nominal mass values for $D^\pm \rightarrow K^\mp\pi^\pm\pi^\pm$ or $K_S^0\pi^\pm$ and $D^0 \rightarrow K^\mp\pi^\pm$ or $K^\mp\pi^\pm\pi^0$. The decay mode $B^0 \rightarrow a_2^\pm(1320)\pi^\mp$ has the same final-state particles as the signal. We suppress this decay with the angular variable \mathcal{H} , defined as the cosine of the angle between the normal to the plane of the 3π resonance and the flight direction of the primary pion from B meson evaluated in the 3π resonance rest frame. Since the a_1 and $a_2(1320)$ mesons have spins of 1 and 2, respectively, the distributions of the variable \mathcal{H} for these two resonances differ. We require $|\mathcal{H}| < 0.62$.

We obtain the CP parameters and signal yield from an unbinned extended maximum likelihood (ML) fit with the input observables ΔE , m_{ES} , \mathcal{F} , m_{a_1} , \mathcal{H} , and Δt . We have six fit components in the likelihood: signal, charm and charmless $B\bar{B}$ background, $B^0 \rightarrow a_2^\pm(1320)\pi^\mp$, continuum $q\bar{q}$ background, and nonresonant $\rho\pi\pi$. The charmless component also includes candidates that were incorrectly reconstructed from particles in events that contain a true signal candidate. Based on measurements of branching fractions for similar charmless decays, we assume $\mathcal{B}(B^0 \rightarrow \rho^0\pi^+\pi^-) = (2 \pm 2) \times 10^{-6}$ [19], which corresponds to 19 expected events in the ML fit sample. This yield is fixed in the fit and a systematic uncertainty is assigned to the final results.

The total probability density function (PDF) for the component j and tagging category c in the event i , $\mathcal{P}_{j,c}^i$, is written as a product of the PDFs of the discriminating variables used in the fit. The factored form of the PDF is a good approximation since linear correlations among observables are below 10%. The systematic uncertainty from residual correlations is taken into account in the fit bias. We write the extended likelihood function for all events as

$$\mathcal{L} = \prod_c \exp(-n_c) \prod_i^{N_c} \left[\sum_j n_j f_{j,c} \mathcal{P}_{j,c}^i \right], \quad (2)$$

where n_j is the yield of events of component j , $f_{j,c}$ is the fraction of events of component j for each category c , $n_c = \sum_j f_{j,c} n_j$ is the number of events found by the fit for category c , and N_c is the number of events of category c in the sample. We fix $f_{j,c}$ to $f_{B_{\text{flav}},c}$, the values measured with a large sample of fully reconstructed B^0 decays into flavor eigenstates (B_{flav} sample) [20], for the signal, $\rho\pi\pi$, and $B^0 \rightarrow a_2^\pm(1320)\pi^\mp$ fit components. We fix $f_{j,c}$ to values obtained with MC events for the charmless and charm fit components and allow it to vary for the $q\bar{q}$ component.

The PDF $\mathcal{P}_{\text{sig}}(\Delta t, \sigma_{\Delta t}; c)$, for each category c , is the convolution of $F(\Delta t; c)$ [Eq. (1)] with the signal resolution function (sum of three Gaussians) determined from the B_{flav} sample. The Δt resolution functions for all the other fit components are also modeled with the sum of three Gaussians. For charmless, $B^0 \rightarrow a_2^\pm(1320)\pi^\mp$, and $\rho\pi\pi$ components in the nominal fit to the data we assume $S = 0$, $C = 0$, $\Delta S = 0$, and $\Delta C = 0$, and we vary these parameters when evaluating systematic uncertainties on final results. We use an effective B lifetime for the charmless component as obtained from a fit to MC signal events. The continuum (charm) Δt distributions are parametrized as sums of three Gaussians with parameters determined from a fit to off-resonance (MC) events.

The PDF of the invariant mass of the a_1 meson in signal events is parametrized as a relativistic Breit-Wigner line shape with a mass-dependent width that takes into account the effect of the mass-dependent ρ width [21]. The PDF of the invariant mass of the $a_2(1320)$ meson is parametrized as a relativistic Breit-Wigner function. The m_{ES} and ΔE distributions for signal are parametrized as a sum of two Gaussian distributions. The ΔE distribution for continuum background is parametrized by a linear function, and the combinatorial background in m_{ES} is described by a phase-space-motivated empirical function [22]. We model the Fisher distribution \mathcal{F} using a Gaussian function with different widths above and below the mean. The \mathcal{A} distributions are modeled using polynomials.

The PDF parameters are determined from MC simulated events with the exception of the continuum background, where we use off-resonance data, and of the signal resolution function, where we use the B_{flav} sample. Large data control samples of B decays to charmed final states of

similar topology are used to verify the simulated resolutions in m_{ES} and ΔE . Where the control samples reveal differences between data and MC samples in mass and energy resolution, we shift or scale the resolution used in the likelihood fits.

We test and calibrate the fitting procedure by applying it to ensembles of simulated $q\bar{q}$ experiments drawn from the PDF, into which we have embedded the expected number of signal, charmless, $B^0 \rightarrow a_2^\pm(1320)\pi^\mp$, the charm, and the $\rho\pi\pi$ events randomly extracted from the fully simulated MC samples. The measured quantities $S_{a_1\pi}$, $C_{a_1\pi}$, $\Delta S_{a_1\pi}$, $\Delta C_{a_1\pi}$, and $\mathcal{A}_{CP}^{a_1\pi}$ have been corrected for the fit biases, and a systematic uncertainty equal to half of the bias found in MC simulations is assigned on the final results.

In the fit there are 35 free parameters, including $S_{a_1\pi}$, $C_{a_1\pi}$, $\Delta S_{a_1\pi}$, $\Delta C_{a_1\pi}$, the charge asymmetries for signal and continuum background, five yields, the signal a_1 width, eleven parameters determining the shape of the combinatorial background, and 12 tagging efficiencies for the continuum. The main contributions to the systematic error on the signal parameters are summarized in Table I. We have studied systematic uncertainties arising from several sources: variation of the signal PDF shape parameters within their errors; modeling of the signal Δt distribution; tagging efficiency and mistag rates determined from the B_{flav} sample [20]; uncertainties in Δm_d and τ [16]; uncertainty in fit bias; uncertainty due to CP violation present in the $B\bar{B}$ background, the $a_2^\pm(1320)\pi^\mp$ CP violation; uncertainty due to the interference between $B^0 \rightarrow a_1^\pm\pi^\mp$ and other 4π final states have been estimated with MC simulations; doubly Cabibbo-suppressed (DCS) $b \rightarrow \bar{u}cd$ amplitude for some tagside B decays [23]; SVT alignment; and the particle identification algorithm. Systematic uncertainties due to neglected minor a_1 substructures are negligible. We allow for a CP asymmetry up to 20% in B decays to charmless final states, and up to 50% in B decays to $a_2(1320)\pi$.

TABLE I. Summary of the systematic uncertainties (in units of 10^{-2}).

	$S_{a_1\pi}$	$C_{a_1\pi}$	$\Delta S_{a_1\pi}$	$\Delta C_{a_1\pi}$	$\mathcal{A}_{CP}^{a_1\pi}$
PDF parametrization	4.8	5.3	3.3	5.3	1.5
Signal Δt model	0.2	0.2	0.3	0.1	0.0
Tagging and mistag	0.3	0.2	0.2	0.4	0.1
Δm_d and τ	0.0	0.2	0.3	0.1	0.0
Fit bias	0.8	0.2	0.8	1.0	0.3
$B\bar{B}$ CP violation	4.1	4.3	4.2	4.0	0.5
$a_2^\pm(1320)\pi^\mp$ + interf.	2.8	4.5	3.2	0.6	0.2
DCS decays	0.8	2.2	0.0	2.2	0.1
SVT alignment	1.0	0.6	1.0	0.6	0.0
Particle ID	0.1	0.1	0.1	0.1	0.2
Total	7.0	8.5	6.4	7.1	1.6

From the fit to a sample of 29 300 events, we obtain a signal yield of 608 ± 52 (461 ± 46 have their flavor identified) used to measure the following parameters: $S_{a_1\pi} = 0.37 \pm 0.21 \pm 0.07$, $\Delta S_{a_1\pi} = -0.14 \pm 0.21 \pm 0.06$, $C_{a_1\pi} = -0.10 \pm 0.15 \pm 0.09$, $\Delta C_{a_1\pi} = 0.26 \pm 0.15 \pm 0.07$, $\mathcal{A}_{CP}^{a_1\pi} = -0.07 \pm 0.07 \pm 0.02$. Linear correlations between these fit parameters are small.

The angle α_{eff} can be defined [7] as

$$\alpha_{\text{eff}} = \frac{1}{4} \left[\arcsin \left(\frac{S_{a_1\pi} + \Delta S_{a_1\pi}}{\sqrt{1 - (C_{a_1\pi} + \Delta C_{a_1\pi})^2}} \right) \right. \\ \left. + \arcsin \left(\frac{S_{a_1\pi} + \Delta S_{a_1\pi}}{\sqrt{1 - (C_{a_1\pi} - \Delta C_{a_1\pi})^2}} \right) \right]. \quad (3)$$

Using the measured parameters in this formula, the angle α_{eff} can be extracted up to a 16-fold ambiguity, which can be reduced to a fourfold ambiguity with conservative assumptions based on factorization [7,10]. One of the four solutions, $\alpha_{\text{eff}} = 78.6^\circ \pm 7.3^\circ$, is compatible with the result of standard model [1] based fits. We also determine the two direct CP asymmetries [24] $\mathcal{A}_{a_1\pi}^{+-} = 0.15 \pm 0.16$ and $\mathcal{A}_{a_1\pi}^{-+} = 0.07 \pm 0.25$ and their linear correlation 0.63. Using the published branching fraction [12], we obtain also the following values for the flavor-charge branching fractions [24] (in units of 10^{-6}): $\mathcal{B}(B^0 \rightarrow a_1^+\pi^-) = 17.9 \pm 4.8$, $\mathcal{B}(B^0 \rightarrow a_1^-\pi^+) = 11.4 \pm 4.7$, $\mathcal{B}(\bar{B}^0 \rightarrow a_1^+\pi^-) = 13.0 \pm 4.3$, and $\mathcal{B}(\bar{B}^0 \rightarrow a_1^-\pi^+) = 24.2 \pm 5.8$. The errors are obtained adding in quadrature statistical and systematic errors.

Figure 1 shows distributions of m_{ES} and ΔE , enhanced in signal content by requirements on the signal-to-continuum likelihood ratios using all discriminating variables other than the one plotted. Figure 2 gives the Δt projections and asymmetry for flavor tagged events.

In summary, we have measured the CP -violating asymmetries in $B^0 \rightarrow a_1^\pm(1260)\pi^\mp$ decays and determined the angle α_{eff} . We do not find evidence for direct or mixing-induced CP violation in these decays. Once measurements




FIG. 1 (color online). Projections of (a) ΔE , (b) m_{ES} . Points represent on-resonance data, dotted lines the sum of all backgrounds, and solid lines the full fit function. These plots are made with a cut on the signal likelihood.




FIG. 2 (color online). Projections onto Δt of the data (points) for (a) B^0 and (b) \bar{B}^0 tags, showing the fit function (solid line), and the background function (dotted line), and (c) the asymmetry between B^0 and \bar{B}^0 tags.

of branching fractions for SU(3)-related decays become available, quantitative bounds on $\Delta\alpha$ obtained with the method of Ref. [7] will provide significant constraints on the angle α through the measurement of α_{eff} in $B^0 \rightarrow a_1^\pm(1260)\pi^\mp$ decays.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (U.S.A.), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from CONACYT (Mexico), Marie Curie EIF (European Union), the A.P. Sloan Foundation, the Research Corporation, and the Alexander von Humboldt Foundation.

*Deceased.

[†]Also at: Dipartimento di Fisica, Università di Perugia, Perugia, Italy.

[‡]Also at: Università della Basilicata, Potenza, Italy.

^{\$}Also at: Physics Department, IPPP, Durham University, Durham DH1 3LE, United Kingdom.

- [1] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [2] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **89**, 281802 (2002); Phys. Rev. Lett. **94**, 181802 (2005); Phys. Rev. Lett. **95**, 151803 (2005); Y. Chao *et al.* (*Belle* Collaboration), Phys. Rev. D **69**, 111102 (2004); Phys. Rev. Lett. **94**, 181803 (2005); K. Abe *et al.*, Phys. Rev. Lett. **95**, 101801 (2005); H. Ishino *et al.*, hep-ex/0608035.
- [3] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **91**, 201802 (2003); C. C. Wang *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **94**, 121801 (2005).
- [4] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **93**, 231801 (2004); A. Somov *et al.* (*Belle* Collaboration), Phys. Rev. Lett. **96**, 171801 (2006).
- [5] For the $a_1(1260)$ meson we use the short notation a_1 .
- [6] R. Aleksan *et al.*, Nucl. Phys. **B361**, 141 (1991).
- [7] M. Gronau and J. Zupan, Phys. Rev. D **73**, 057502 (2006).
- [8] M. Gronau and D. London, Phys. Rev. Lett. **65**, 3381 (1990); H. J. Lipkin *et al.*, Phys. Rev. D **44**, 1454 (1991); M. Gronau, Phys. Lett. B **265**, 389 (1991).
- [9] Y. Grossman and H. R. Quinn Phys. Rev. D **58**, 017504 (1998); J. Charles, Phys. Rev. D **59**, 054007 (1999); M. Gronau *et al.*, Phys. Lett. B **514**, 315 (2001).
- [10] M. Gronau and J. Zupan, Phys. Rev. D **70**, 074031 (2004).
- [11] A. Bevan, Mod. Phys. Lett. A **21**, 305 (2006).
- [12] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **97**, 051802 (2006).
- [13] B. Aubert *et al.* (*BABAR* Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [14] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **94**, 161803 (2005).
- [15] S. Agostinelli *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **506**, 250 (2003).
- [16] Y.-M. Yao *et al.* (Particle Data Group), J. Phys. G **33**, 1 (2006).
- [17] David J. Lange, Nucl. Instrum. Methods Phys. Res., Sect. A **462**, 152 (2001).
- [18] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **92**, 061801 (2004).
- [19] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **98**, 111801 (2007).
- [20] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. D **66**, 032003 (2002).
- [21] T. A. Armstrong *et al.* (WA76 Collaboration), Z. Phys. C **48**, 213 (1990).
- [22] See H. Albrecht *et al.* (*ARGUS* Collaboration), Phys. Lett. B **241**, 278 (1990).
- [23] O. Long *et al.*, Phys. Rev. D **68**, 034010 (2003).
- [24] J. Charles *et al.* (CKMfitter Group), Eur. Phys. J. C **41**, 1 (2005).