

Saltation Transport on Mars

Eric J. R. Parteli¹ and Hans J. Herrmann^{2,3}

¹*Institut für Computerphysik, ICP, Universität Stuttgart, Pfaffenwaldring 27, 70569 Stuttgart, Germany*

²*Institut für Baustoffe, ETH Hönggerberg, HIF E 12, CH-8093, Zürich, Switzerland*

³*Departamento de Física, Universidade Federal do Ceará-60455-760, Fortaleza, CE, Brazil*

(Received 2 October 2006; published 11 May 2007)

We present the first calculation of saltation transport and dune formation on Mars and compare it to real dunes. We find that the rate at which grains are entrained into saltation on Mars is 1 order of magnitude higher than on Earth. With this fundamental novel ingredient, we reproduce the size and different shapes of Mars dunes, and give an estimate for the wind velocity on Mars.

DOI: [10.1103/PhysRevLett.98.198001](https://doi.org/10.1103/PhysRevLett.98.198001)

PACS numbers: 45.70.-n, 92.40.Gc, 92.60.Gn

The surprising discovery of sand dunes on Mars by Mariner 9 in 1971 has challenged geologists and planetary physicists over the years with the following enigma: Could Mars dunes have been formed by today's Martian thin atmosphere? Bagnold [1] first noticed that the sand of Aeolian dunes is transported through *saltation*, which consists of grains traveling in a sequence of ballistic trajectories and producing a *splash* of new ejected grains when colliding back onto the soil. But saltation occurs only if the wind friction speed u_* —which, together with the air density ρ_{fluid} , is used to define the air shear stress $\tau = \rho_{\text{fluid}} u_*^2$ —is larger than a minimal threshold u_{*t} . Because the atmospheric density of Mars is almost 100 times lower than on Earth, only winds of strength 1 order of magnitude higher than that on our planet are capable of mobilizing the basaltic grains of mean diameter $d = 500 \mu\text{m}$ [2] which constitute Mars dunes [3,4]. Such winds occasionally can occur on Mars [5–7]. Here we find that the strong erosion due to these winds allows to understand dune formation on the red planet under present conditions.

Once saltation starts, the number of saltating grains first increases exponentially due to the multiplicative process inherent in the splash events. But since the wind loses momentum to accelerate the grains, the flux *saturates* after a transient distance [8–12]. At saturation, the air shear stress τ_a decreases to the threshold $\tau_t = \rho_{\text{fluid}} u_{*t}^2$, while the flux of grains increases to its maximum value. This transient phenomenon introduces the only relevant length scale in the physics of dunes: Any sand patch which is shorter than this saturation distance will be eroded and disappear. On the other hand, large enough sand hills develop a slip face wherever the slope exceeds the angle of repose, $\theta_r = 34^\circ$ [1].

Successful modeling of the formation of dunes was recently achieved [13,14]. The dune model consists of a system of continuum equations in two space dimensions which reproduce the shape of dunes, the wind profile and the sand flux and provide excellent quantitative agreement with measurements [15]. The sand bed can exchange

grains with the moving saltation layer. The local change in the flux, $\vec{\nabla} \cdot \vec{q}$, is proportional to the erosion rate, $\Gamma(x, y)$, which is the difference between the vertical flux of ejected grains and the vertical flux ϕ of grains impacting onto the bed: $\Gamma = \phi(n - 1)$, where n is the average number of splashed grains. At saturation ($\tau_a \approx \tau_t$), n can be written as [13]:

$$n = 1 + \tilde{\gamma} \left(\frac{\tau_a}{\tau_t} - 1 \right), \quad (1)$$

where $\tilde{\gamma}$, the *entrainment rate* of grains into saltation, determines how fast the system reaches saturation [13]. On the other hand, $\phi \equiv |\vec{q}|/\ell$, where ℓ is the average saltation length. The balance $\Gamma(x, y) = \vec{\nabla} \cdot \vec{q}$ yields a differential equation for \vec{q} :

$$\vec{\nabla} \cdot \vec{q} = \frac{1}{\ell_s} |\vec{q}| \left(1 - \frac{|\vec{q}|}{q_s} \right); \quad \ell_s = \frac{\ell}{\tilde{\gamma}} \left[\frac{\tau_t}{\tau - \tau_t} \right], \quad (2)$$

where $q_s = |\vec{q}|(\tau - \tau_t)/(\tau - \tau_a)$ is the saturated flux [13], and $\ell = v_z^{\text{eje}}(2v_g/g)$, where v_z^{eje} is the initial vertical velocity of the grains, v_g their mean velocity and g is gravity. Further, $v_z^{\text{eje}} = \alpha \Delta v_{\text{hor}}$, where Δv_{hor} is the horizontal velocity gain of the particle after one saltation trajectory, and α is an effective restitution coefficient for the grain-bed interaction [13]. In this manner, $\ell = (1/r)[2v_g^2 \alpha/g]$, with $r \equiv v_g/\Delta v_{\text{hor}}$, and the saturation length, ℓ_s , may be then written as

$$\ell_s = \frac{1}{\gamma} \left[\frac{2v_g^2 \alpha/g}{(u_*/u_{*t})^2 - 1} \right], \quad (3)$$

where $\gamma \equiv r\tilde{\gamma}$.

The dune model can be sketched as follows: (i) the wind shear stress over the topography is calculated using the algorithm of Weng *et al.* (1991) [16]. The wind velocity u over a flat ground increases logarithmically with the height z : $u(z) = u_* \kappa^{-1} \ln z/z_0$, where $\kappa = 0.4$ is the von Kármán constant and z_0 is the aerodynamic roughness. A dune introduces a perturbation in the shear stress whose Fourier-transformed components are

$$\hat{\tau}_x = \frac{2h(k_x, k_y)k_x^2}{|k|U^2(l)} \left[1 + \frac{2\ln(\mathcal{L}|k_x|) + 4\epsilon + 1 + i\text{sgn}(k_x)\pi}{\ln(l/z_0)} \right] \quad (4)$$

and

$$\hat{\tau}_y = \frac{2h(k_x, k_y)k_x k_y}{|k|U^2(l)}, \quad (5)$$

where the coordinate axes x and y are parallel, respectively, perpendicular to the wind direction, k_x and k_y are wave numbers, $|k| = \sqrt{k_x^2 + k_y^2}$ and $\epsilon = 0.577216$ (Euler's constant). \mathcal{L} is the horizontal distance between the position of maximum height, H_{\max} , and the position of the windward side where the height is $H_{\max}/2$ [16]. $U(l) = u(l)/u(h_m)$ is the undisturbed wind velocity at height $l = 2\kappa^2 \mathcal{L} / \ln l / z_0$ normalized by the velocity at the reference height $h_m = \mathcal{L} / \sqrt{\log \mathcal{L} / z_0}$, which separates the middle and upper flow layers [16]. The shear stress in the direction i ($i = x, y$) is then given by $\vec{\tau}_i = \hat{i}[\tau_0(1 + \hat{\tau}_i)]$, where τ_0 is the undisturbed shear stress; (ii) next, the sand flux \vec{q} (x, y) is calculated using Eqs. (2) and (3); (iii) the change in surface height $h(x, y)$ is computed from mass conservation: $\partial h / \partial t = -\vec{\nabla} \cdot \vec{q} / \rho_{\text{sand}}$, where $\rho_{\text{sand}} = 0.62\rho_{\text{grain}}$ is the bulk density of the bed [13]; and (iv) if sand deposition leads to slopes that locally exceed the angle of repose, the unstable surface relaxes through avalanches in the direction of the steepest descent, and the separation streamlines are introduced at the dune lee [14]. Each streamline is fitted by a third order polynomial connecting the brink with the ground at the reattachment point [14], and defining the "separation bubble", in which the wind and the flux are set to zero. The model is evaluated by performing steps (i) through (iv) computationally in a cyclic manner.

In this Letter, our aim is to reproduce the shape of Mars dunes using the martian present atmospheric conditions. Most of the relevant model parameters are known for Mars: The pressure, P , and temperature, T , which have been measured in several places on Mars [17], determine ρ_{fluid} . Also known are the average grain diameter d and density, $\rho_{\text{grain}} = 3200 \text{ kg/m}^3$, and gravity, $g = 3.71 \text{ m/s}^2$, while u_{*t} is calculated as in Ref. [4].

One very common type of dune on Mars are *barchans* [18]. They have one slip face and two horns, and propagate on bedrock under conditions of unidirectional wind [1]. Indeed, to develop a slip face, they must reach a minimum size. In the Arkhangelsky Crater on Mars (Fig. 1), the minimal dune width is $W_{\min} \approx 200 \text{ m}$ and the corresponding length $L_{\min} \approx 400 \text{ m}$. On the other hand, W_{\min} is around 13 times the saturation length, ℓ_s [19]. In this manner, W_{\min} yields, through Eq. (3), the wind strength u_*/u_{*t} on Mars.

However, there is *one* unknown quantity for Mars which we need in order to solve the sand transport equations: γ , which appears in Eq. (3). Sauermann *et al.* (2001) [13] obtained $\gamma = 0.2$ for saltation on Earth [13] by comparing




FIG. 1 (color online). Barchans in the Arkhangelsky crater, 41.0° S , 25.0° W on Mars: Mars orbiter camera images on the left (image courtesy of NASA/JPL/MSSS) and calculated dunes on the right. Main plot: L vs W of the Arkhangelsky dunes (circles) and north polar barchans at 77.6° N , 103.6° W (stars). Calculations of Arkhangelsky (north polar) dunes are represented by the continuous (dotted) line, obtained with $u_*/u_{*t} = 1.45$ (1.80). The dashed line in the upper inset corresponds to terrestrial dunes obtained with $u_*/u_{*t} = 1.45$. In the lower inset, we see L/L_{\min} vs H/H_{\min} from calculations of the Arkhangelsky (triangles) and terrestrial dunes (dashed line).

with direct measurements of saturation transients [10,11]. Such measurements are not available for Mars.

We perform our calculations using open boundaries and a constant upwind u_* in x direction at the inlet starting with a Gaussian hill, which evolves until displaying the linear relations between height H , width W and length L of barchans [13,19].

For a constant γ , there is *one* value of u_* which reproduces the minimal dune width $W_{\min} = 200 \text{ m}$ in the Arkhangelsky Crater, where $P = 5.5 \text{ mb}$ and $T = 210 \text{ K}$ [17], which gives $u_{*t} \approx 2.12 \text{ m/s}$. We obtained a surprising result: If we take the same γ as on Earth, the value of u_* which gives $W_{\min} = 200 \text{ m}$ is around 6.0 m/s . This value is far too large to be realistic. The largest peaks of martian u_* range between 2.2 and 4.0 m/s [6,7]. To obtain the correct W_{\min} using u_* within this maximum range, we must take γ at least 1 order of magnitude larger than on Earth. We must find how much the martian $\tilde{\gamma} = dn/d(\tau_a/\tau_t)$ [Eq. (1)] differs from the one on Earth.

Anderson and Haff (1988) [9] showed that the number of splashed grains is proportional to the velocity v_{imp} of the impacting grains. Let us rescale v_{imp} with $v_{\text{eje}} = \sqrt{gd}$, which is the velocity necessary to escape from the sand bed [12]. Further, v_{imp} scales with the mean grain velocity v_g [13]. In this manner, we obtain $\tilde{\gamma} \propto v_g / \sqrt{gd}$.

Moreover, v_g scales with u_{*t} [13] and has only a very weak dependence on u_* which we neglect, and thus $\tilde{\gamma} \propto u_{*t} / \sqrt{gd}$. Since we know that $\gamma = 0.2$ on Earth, where $g = 9.81 \text{ m/s}^2$, $d = 250 \mu\text{m}$ and $u_{*t} = 0.218 \text{ m/s}$ [13], we

TABLE I. Main quantities controlling saltation on Mars and on Earth. u_{*t} is around 80% of the direct entrainment threshold $u_{*ft} = A\sqrt{\rho_{\text{grain}}gd/\rho_{\text{fluid}}}$, where $A = 0.129[(1 + 6.0 \times 10^{-7}/\rho_{\text{grain}}gd^{2.5})^{0.5}/(1.928\text{Re}_{*ft}^{0.092} - 1)^{0.5}]$ [4], $\text{Re}_{*ft} \equiv u_{*ft}d\rho_{\text{fluid}}/\eta$, and η is the viscosity of CO_2 gas.

Barchan field	ρ_{fluid} (kg/m ³)	u_{*t} (m/s)	v_g (m/s)	γ
Arkhangelsky	0.014	2.12	17.8	2.24
77.6° N, 103.6° W	0.022	1.62	12.3	1.71
Earth	1.225	0.22	1.5	0.20

obtain

$$\gamma = 0.045 \frac{u_{*t}}{\sqrt{gd}}. \quad (6)$$

Substituting Eq. (6) into Eq. (3), we obtain a closed set of sand transport equations from which the value of the wind friction speed u_* on Mars can be determined.

Equation (6) gives $\gamma \approx 2.24$ in the Arkhangelsky Crater, and $W_{\text{min}} = 200$ m is reproduced for $u_*/u_{*t} = 1.45$ or $u_* = 3.07$ m/s, which gives $\ell_s \approx 15.5$ m. Figure 1 shows that $u_* = 3.07$ m/s reproduces not only the minimal dune but also the dependence of the dune shape on its size. Furthermore, we tested the scaling relation (6) with a second martian barchan field which is near the north pole, and where $W_{\text{min}} \approx 80$ m. At the location of the field, $P = 8.0$ mb and $T = 190$ K [17], and thus $u_{*t} \approx 1.62$ m/s. From W_{min} , we obtain $\ell_s \approx 6.0$ m, which gives $u_*/u_{*t} \approx 1.8$ or $u_* = 2.92$ m/s. The plot in Fig. 1 shows that the behavior L against W of the barchans in both studied fields is well captured by the model.

It is interesting that the u_* obtained for the north polar field is very similar to that in the Arkhangelsky Crater, although u_{*t} is lower in the north polar field due to the higher ρ_{fluid} (Table I).

Summarizing, we discovered that the rate γ at which grains enter saltation on Mars is about 1 order of magnitude higher than on earth. Taking this finding into account, we found that the wind velocity which reproduces the size and shape of martian barchans is $u_* \approx 3.0 \pm 0.1$ m/s, which is well within the range of estimated wind speeds on Mars [6,7]. We found that Mars grains travel with a velocity v_g 10 times higher than that of terrestrial grains, as observed in wind tunnel experiments for Mars [20]. Such high-speed grains produce much larger splash events than on Earth [21], and lead to higher γ values (Table I).

Our results show that the larger splash events have a crucial implication for the formation of dunes on Mars. While on one hand the lower martian g and ρ_{fluid} result in longer grain trajectories [20], the higher rate at which grains enter saltation on Mars shortens the saturation transient of the flux, which determines the minimal dune size. The distance of flux saturation, λ_s , is around 6 times the characteristic length ℓ_s [13], which effectively increases with ℓ . Here we found that λ_s on Mars is indeed reduced by




FIG. 2 (color online). Dune velocity v as function of dune length L . We see that Mars dunes (filled symbols) move typically 10 times faster than Earth dunes (empty symbols) of same L , obtained with similar values of u_*/u_{*t} as on Mars.

the larger martian splash. This explains the previously reported [22] failure of the scaling $W_{\text{min}} \propto \ell$ for Mars.

The values of u_*/u_{*t} obtained for Mars are within the range of the ones measured in terrestrial barchan fields [15]. Indeed, we see in Fig. 2 that, for the same value of u_*/u_{*t} , Mars barchans would move 10 times faster than those on Earth.

Many exotic and up to now unexplained dune forms have also been observed on Mars. Dunes as those in




FIG. 3. MOC images on top (courtesy of NASA/JPL/MSSS) and calculations obtained with bimodal wind regimes on bottom. A sketch showing the definition of the angle θ_w of the wind direction (arrows) is shown in (a'). The wind changes its direction with frequency $1/T_w$. We chose $T_w = 2.9, 5.8,$ and 0.7 d to obtain dunes in (a'), (b'), and (c'), respectively. The dune in (a') has been obtained with $|\theta_w| = 50^\circ$ and the dune in (b') with $|\theta_w| = 70^\circ$. In (c'), a linear dune obtained with $|\theta_w| = 60^\circ$ decays into barchans after the angle $|\theta_w|$ is reduced to 40° . The dune in (b') elongates to the right further and further with time, which is not observed for dune in (a'). The dune in (c') decays further until only a string of rounded barchans is visible.

TABLE II. For each dune field in Fig. 3, the fluid density ρ_{fluid} and the threshold u_{*t} is calculated from the local pressure and temperature which are taken from Ref. [17]. In spite of the broad range of u_{*t} , all dune forms in Fig. 3 have been obtained with one single value of $u_* = 3.0$ m/s.

Field	Location	ρ_{fluid} (kg/m ³)	u_{*t} (m/s)
Fig. 3(a)	48.6° S, 25.5° W	0.017	1.89
Fig. 3(b)	49.6° S, 352.9° W	0.014	2.06
Fig. 3(c)	76.4° N, 272.9° W	0.03	1.35

Fig. 3 cannot be obtained from calculations with a unidirectional wind. We found that the dune shapes in Fig. 3 may be obtained with a *bimodal* wind regime. In our calculations, the wind *alternates* its direction periodically with frequency $1/T_w$ forming an angle $2|\theta_w|$ as sketched in Fig. 3(a'). In both directions the strength is the same, namely $u_* = 3.0$ m/s. In this manner, the value of u_*/u_{*t} is particular to each field, since u_{*t} depends on the field location (Table II). To simulate the change of wind direction, we rotate the field by an angle $2\theta_w$, keeping the wind direction constant. The separation bubble, thus, adapts to the wind direction after rotation of the field. We use open boundaries as in the calculations of barchan dunes. The initial condition is a Gaussian hill as before, whose volume is taken according to the volume of the dune.

The angle $2\theta_w$ between the wind directions determines which of the different forms in Fig. 3 is obtained. We found that a barchan moving in the resulting wind direction is always obtained if $|\theta_w| < 50^\circ$. If $|\theta_w| \approx 50^\circ$, the dune shape in Fig. 3(a') is achieved. And for $|\theta_w|$ between 50° and 70° , elongated dune forms as those in Fig. 3(b') are obtained, which elongate further and further in time. As $|\theta_w| \rightarrow 90^\circ$, a barchanoidal form of alternating slip face position appears.

Moreover, we found that the structure observed in the dune field of Fig. 3(c) can be obtained by a change in the local wind regime. The dune shape in Fig. 3(c') has been obtained in the following manner: (i) first, an elongated dune form as the one in Fig. 3(b') is formed with an angle $|\theta_w| = 60^\circ$; (ii) next, the angle $|\theta_w|$ has been suddenly reduced to 40° . Thereafter, the linear dune becomes unstable and decays into a string of rounded barchans seen in Fig. 3(c).

Each one of the dune shapes in Fig. 3 can only be achieved if the time T_w is of the order of 1–5 days. If the period is too large, of the order of 1 month, then the dune evolves into a barchanoidal form.

In conclusion, we found from calculations of martian barchans using the present atmospheric conditions of Mars, that the splash on Mars must be about 10 times larger than on Earth. Furthermore, we obtained a general equation for

the entrainment rate of grains into saltation which could be used in future calculations of other dune forms on Mars, Venus, or Titan. We also found that winds on Mars do not exceed 3.0 ± 0.1 m/s, which can explain the shape of elongated dunes formed by a bimodal wind. It would be interesting to make a full microscopic simulation for the saltation mechanism of Mars similar to the one that was recently performed by Almeida *et al.* (2006) [23] to confirm our findings microscopically.

We acknowledge Kenneth Edgett, Keld Rasmussen, Bruno Andreotti, and Orencio Durán for discussions. This research was supported in part by The Max-Planck Prize and the Volkswagenstiftung. E. J. R. Parteli acknowledges support from CAPES-Brasília/Brazil.

-
- [1] R. A. Bagnold, *The Physics of Blown Sand and Desert Dunes* (Methuen, London, 1941).
 - [2] K. S. Edgett and P. R. Christensen, *J. Geophys. Res.* **96**, 22765 (1991).
 - [3] R. Greeley *et al.*, *Geophys. Res. Lett.* **7**, 121 (1980).
 - [4] J. D. Iversen and B. R. White, *Sedimentology* **29**, 111 (1982).
 - [5] J. L. Sutton, C. B. Leovy, and J. E. Tillman, *J. Atmos. Sci.* **35**, 2346 (1978).
 - [6] H. J. Moore, *J. Geophys. Res.* **90**, 163 (1985).
 - [7] R. Sullivan *et al.*, *Nature (London)* **436**, 58 (2005).
 - [8] P. R. Owen, *J. Fluid Mech.* **20**, 225 (1964).
 - [9] R. S. Anderson and P. K. Haff, *Science* **241**, 820 (1988).
 - [10] I. K. McEwan and B. B. Willetts, *Acta Mech. (Suppl.)* **1**, 53 (1991).
 - [11] G. R. Butterfield, in *Turbulence: Perspectives on Flow and Sediment Transport*, edited by N. J. Clifford, J. R. French, and J. Hardisty (John Wiley, New York, 1993), Chap. 13, p. 305.
 - [12] B. Andreotti, *J. Fluid Mech.* **510**, 47 (2004).
 - [13] G. Saueremann, K. Kroy, and H. J. Herrmann, *Phys. Rev. E* **64**, 031305 (2001).
 - [14] K. Kroy, G. Saueremann, and H. J. Herrmann, *Phys. Rev. E* **66**, 031302 (2002).
 - [15] G. Saueremann *et al.*, *Geomorphology* **54**, 245 (2003).
 - [16] W. S. Weng *et al.*, *Acta Mech. (Suppl.)* **2**, 1 (1991).
 - [17] Mars Global Surveyor Radio Science Team. <http://nova.stanford.edu/projects/mgs/>.
 - [18] M. C. Bourke, M. Balme, and J. Zimbelman, *Proc. Lunar Planet. Sci. Conf.* **35** 1453 (2004).
 - [19] E. J. R. Parteli, O. Durán, and H. J. Herrmann, *Phys. Rev. E* **75**, 011301 (2007).
 - [20] B. White, *J. Geophys. Res.* **84**, 4643 (1979).
 - [21] J. Marshall, J. Borucki, and C. Bratton, *Proc. Lunar Planet. Sci. Conf.* **29**, 1131 (1998).
 - [22] K. Kroy, S. Fischer, and B. Obermayer, *J. Phys. Condens. Matter* **17**, S1229 (2005).
 - [23] M. P. Almeida, J. S. Andrade, Jr., and H. J. Herrmann, *Phys. Rev. Lett.* **96**, 018001 (2006).