

Measurement of CP -Violating Asymmetries in $B^0 \rightarrow D^{(*)\pm} D^\mp$

- B. Aubert,¹ M. Bona,¹ D. Boutigny,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ X. Prudent,¹ V. Tisserand,¹ A. Zghiche,¹ J. Garra Tico,² E. Grauges,² L. Lopez,³ A. Palano,³ G. Eigen,⁴ B. Stugu,⁴ L. Sun,⁴ G. S. Abrams,⁵ M. Battaglia,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵ R. N. Cahn,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵ J. A. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ G. Kukartsev,⁵ D. Lopes Pegna,⁵ G. Lynch,⁵ L. M. Mir,⁵ T. J. Orimoto,⁵ M. T. Ronan,^{5,*} K. Tackmann,⁵ W. A. Wenzel,⁵ P. del Amo Sanchez,⁶ C. M. Hawkes,⁶ A. T. Watson,⁶ T. Held,⁷ H. Koch,⁷ B. Lewandowski,⁷ M. Pelizaeus,⁷ T. Schroeder,⁷ M. Steinke,⁷ D. Walker,⁸ D. J. Asgeirsson,⁹ T. Cuhadar-Donszelmann,⁹ B. G. Fulsom,⁹ C. Hearty,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ A. Khan,¹⁰ M. Saleem,¹⁰ L. Teodorescu,¹⁰ V. E. Blinov,¹¹ A. D. Bukin,¹¹ V. P. Druzhinin,¹¹ V. B. Golubev,¹¹ A. P. Onuchin,¹¹ S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ K. Yu. Todyshev,¹¹ M. Bondioli,¹² S. Curry,¹² I. Eschrich,¹² D. Kirkby,¹² A. J. Lankford,¹² P. Lund,¹² M. Mandelkern,¹² E. C. Martin,¹² D. P. Stoker,¹² S. Abachi,¹³ C. Buchanan,¹³ S. D. Foulkes,¹⁴ J. W. Gary,¹⁴ F. Liu,¹⁴ O. Long,¹⁴ B. C. Shen,¹⁴ L. Zhang,¹⁴ H. P. Paar,¹⁵ S. Rahatlou,¹⁵ V. Sharma,¹⁵ J. W. Berryhill,¹⁶ C. Campagnari,¹⁶ A. Cunha,¹⁶ B. Dahmes,¹⁶ T. M. Hong,¹⁶ D. Kovalskyi,¹⁶ J. D. Richman,¹⁶ T. W. Beck,¹⁷ A. M. Eisner,¹⁷ C. J. Flacco,¹⁷ C. A. Heusch,¹⁷ J. Kroseberg,¹⁷ W. S. Lockman,¹⁷ T. Schalk,¹⁷ B. A. Schumm,¹⁷ A. Seiden,¹⁷ D. C. Williams,¹⁷ M. G. Wilson,¹⁷ L. O. Winstrom,¹⁷ E. Chen,¹⁸ C. H. Cheng,¹⁸ F. Fang,¹⁸ D. G. Hitlin,¹⁸ I. Narsky,¹⁸ T. Piatenko,¹⁸ F. C. Porter,¹⁸ R. Andreassen,¹⁹ G. Mancinelli,¹⁹ B. T. Meadows,¹⁹ K. Mishra,¹⁹ M. D. Sokoloff,¹⁹ F. Blanc,²⁰ P. C. Bloom,²⁰ S. Chen,²⁰ W. T. Ford,²⁰ J. F. Hirschauer,²⁰ A. Kreisel,²⁰ M. Nagel,²⁰ U. Nauenberg,²⁰ A. Olivas,²⁰ J. G. Smith,²⁰ K. A. Ulmer,²⁰ S. R. Wagner,²⁰ J. Zhang,²⁰ A. M. Gabareen,²¹ A. Soffer,²¹ W. H. Toki,²¹ R. J. Wilson,²¹ F. Winkelmeier,²¹ Q. Zeng,²¹ D. D. Altenburg,²² E. Feltresi,²² A. Hauke,²² H. Jasper,²² J. Merkel,²² A. Petzold,²² B. Spaan,²² K. Wacker,²² T. Brandt,²³ V. Klose,²³ M. J. Kobel,²³ H. M. Lacker,²³ W. F. Mader,²³ R. Nogowski,²³ J. Schubert,²³ K. R. Schubert,²³ R. Schwierz,²³ J. E. Sundermann,²³ A. Volk,²³ D. Bernard,²⁴ G. R. Bonneau,²⁴ E. Latour,²⁴ V. Lombardo,²⁴ Ch. Thiebaut,²⁴ M. Verderi,²⁴ P. J. Clark,²⁵ W. Gradl,²⁵ F. Muheim,²⁵ S. Playfer,²⁵ A. I. Robertson,²⁵ Y. Xie,²⁵ M. Andreotti,²⁶ D. Bettoni,²⁶ C. Bozzi,²⁶ R. Calabrese,²⁶ A. Cecchi,²⁶ G. Cibinetto,²⁶ P. Franchini,²⁶ E. Luppi,²⁶ M. Negrini,²⁶ A. Petrella,²⁶ L. Piemontese,²⁶ E. Prencipe,²⁶ V. Santoro,²⁶ F. Anulli,²⁷ R. Baldini-Ferroli,²⁷ A. Calcaterra,²⁷ R. de Sangro,²⁷ G. Finocchiaro,²⁷ S. Pacetti,²⁷ P. Patteri,²⁷ I. M. Peruzzi,^{27,†} M. Piccolo,²⁷ M. Rama,²⁷ A. Zallo,²⁷ A. Buzzo,²⁸ R. Contri,²⁸ M. Lo Vetere,²⁸ M. M. Macri,²⁸ M. R. Monge,²⁸ S. Passaggio,²⁸ C. Patrignani,²⁸ E. Robutti,²⁸ A. Santroni,²⁸ S. Tosi,²⁸ K. S. Chaisanguanthum,²⁹ M. Morii,²⁹ J. Wu,²⁹ R. S. Dubitzky,³⁰ J. Marks,³⁰ S. Schenk,³⁰ U. Uwer,³⁰ D. J. Bard,³¹ P. D. Dauncey,³¹ R. L. Flack,³¹ J. A. Nash,³¹ M. B. Nikolich,³¹ W. Panduro Vazquez,³¹ M. Tibbetts,³¹ P. K. Behera,³² X. Chai,³² M. J. Charles,³² U. Mallik,³² N. T. Meyer,³² V. Ziegler,³² J. Cochran,³³ H. B. Crawley,³³ L. Dong,³³ V. Eges,³³ W. T. Meyer,³³ S. Prell,³³ E. I. Rosenberg,³³ A. E. Rubin,³³ A. V. Gritsan,³⁴ Z. J. Guo,³⁴ C. K. Lae,³⁴ A. G. Denig,³⁵ M. Fritsch,³⁵ G. Schott,³⁵ N. Arnaud,³⁶ J. Béquilleux,³⁶ M. Davier,³⁶ G. Grosdidier,³⁶ A. Höcker,³⁶ V. Lepeltier,³⁶ F. Le Diberder,³⁶ A. M. Lutz,³⁶ S. Pruvot,³⁶ S. Rodier,³⁶ P. Roudeau,³⁶ M. H. Schune,³⁶ J. Serrano,³⁶ V. Sordini,³⁶ A. Stocchi,³⁶ W. F. Wang,³⁶ G. Wormser,³⁶ D. J. Lange,³⁷ D. M. Wright,³⁷ I. Bingham,³⁸ C. A. Chavez,³⁸ I. J. Forster,³⁸ J. R. Fry,³⁸ E. Gabathuler,³⁸ R. Gamet,³⁸ D. E. Hutchcroft,³⁸ D. J. Payne,³⁸ K. C. Schofield,³⁸ C. Touramanis,³⁸ A. J. Bevan,³⁹ K. A. George,³⁹ F. Di Lodovico,³⁹ W. Menges,³⁹ R. Sacco,³⁹ G. Cowan,⁴⁰ H. U. Flaecher,⁴⁰ D. A. Hopkins,⁴⁰ S. Paramesvaran,⁴⁰ F. Salvatore,⁴⁰ A. C. Wren,⁴⁰ D. N. Brown,⁴¹ C. L. Davis,⁴¹ J. Allison,⁴² N. R. Barlow,⁴² R. J. Barlow,⁴² Y. M. Chia,⁴² C. L. Edgar,⁴² G. D. Lafferty,⁴² T. J. West,⁴² J. I. Yi,⁴² J. Anderson,⁴³ C. Chen,⁴³ A. Jawahery,⁴³ D. A. Roberts,⁴³ G. Simi,⁴³ J. M. Tuggle,⁴³ G. Blaylock,⁴⁴ C. Dallapiccola,⁴⁴ S. S. Hertzbach,⁴⁴ X. Li,⁴⁴ T. B. Moore,⁴⁴ E. Salvati,⁴⁴ S. Saremi,⁴⁴ R. Cowan,⁴⁵ D. Dujmic,⁴⁵ P. H. Fisher,⁴⁵ K. Koeneke,⁴⁵ G. Sciolla,⁴⁵ S. J. Sekula,⁴⁵ M. Spitznagel,⁴⁵ F. Taylor,⁴⁵ R. K. Yamamoto,⁴⁵ M. Zhao,⁴⁵ Y. Zheng,⁴⁵ S. E. McLachlin,⁴⁶ P. M. Patel,⁴⁶ S. H. Robertson,⁴⁶ A. Lazzaro,⁴⁷ F. Palombo,⁴⁷ J. M. Bauer,⁴⁸ L. Cremaldi,⁴⁸ V. Eschenburg,⁴⁸ R. Godang,⁴⁸ R. Kroeger,⁴⁸ D. A. Sanders,⁴⁸ D. J. Summers,⁴⁸ H. W. Zhao,⁴⁸ S. Brunet,⁴⁹ D. Côté,⁴⁹ M. Simard,⁴⁹ P. Taras,⁴⁹ F. B. Viaud,⁴⁹ H. Nicholson,⁵⁰ G. De Nardo,⁵¹ F. Fabozzi,^{51,†} L. Lista,⁵¹ D. Monorchio,⁵¹ C. Sciacca,⁵¹ M. A. Baak,⁵² G. Raven,⁵² H. L. Snoek,⁵² C. P. Jessop,⁵³ J. M. LoSecco,⁵³ G. Benelli,⁵⁴ L. A. Corwin,⁵⁴ K. Honscheid,⁵⁴ H. Kagan,⁵⁴ R. Kass,⁵⁴ J. P. Morris,⁵⁴ A. M. Rahimi,⁵⁴ J. J. Regensburger,⁵⁴ Q. K. Wong,⁵⁴ N. L. Blount,⁵⁵ J. Brau,⁵⁵ R. Frey,⁵⁵ O. Igonkina,⁵⁵ J. A. Kolb,⁵⁵ M. Lu,⁵⁵ R. Rahmat,⁵⁵ N. B. Sinev,⁵⁵ D. Strom,⁵⁵ J. Strube,⁵⁵ E. Torrence,⁵⁵ N. Gagliardi,⁵⁶ A. Gaz,⁵⁶ M. Margoni,⁵⁶ M. Morandin,⁵⁶ A. Pompili,⁵⁶ M. Posocco,⁵⁶ M. Rotondo,⁵⁶ F. Simonetto,⁵⁶ R. Stroili,⁵⁶ C. Voci,⁵⁶ E. Ben-Haim,⁵⁷ H. Briand,⁵⁷ G. Calderini,⁵⁷ J. Chauveau,⁵⁷ P. David,⁵⁷ L. Del Buono,⁵⁷ Ch. de la Vaissière,⁵⁷ O. Hamon,⁵⁷ Ph. Leruste,⁵⁷ J. Malclès,⁵⁷ J. Ocariz,⁵⁷ A. Perez,⁵⁷ L. Gladney,⁵⁸ M. Biasini,⁵⁹ R. Covarelli,⁵⁹ E. Manoni,⁵⁹ C. Angelini,⁶⁰

G. Batignani,⁶⁰ S. Bettarini,⁶⁰ M. Carpinelli,⁶⁰ R. Cenci,⁶⁰ A. Cervelli,⁶⁰ F. Forti,⁶⁰ M. A. Giorgi,⁶⁰ A. Lusiani,⁶⁰ G. Marchiori,⁶⁰ M. A. Mazur,⁶⁰ M. Morganti,⁶⁰ N. Neri,⁶⁰ E. Paoloni,⁶⁰ G. Rizzo,⁶⁰ J. J. Walsh,⁶⁰ M. Haire,⁶¹ J. Biesiada,⁶² P. Elmer,⁶² Y. P. Lau,⁶² C. Lu,⁶² J. Olsen,⁶² A. J. S. Smith,⁶² A. V. Telnov,⁶² E. Baracchini,⁶³ F. Bellini,⁶³ G. Cavoto,⁶³ A. D'Orazio,⁶³ D. del Re,⁶³ E. Di Marco,⁶³ R. Faccini,⁶³ F. Ferrarotto,⁶³ F. Ferroni,⁶³ M. Gaspero,⁶³ P. D. Jackson,⁶³ L. Li Gioi,⁶³ M. A. Mazzoni,⁶³ S. Morganti,⁶³ G. Piredda,⁶³ F. Polci,⁶³ F. Renga,⁶³ C. Voena,⁶³ M. Ebert,⁶⁴ T. Hartmann,⁶⁴ H. Schröder,⁶⁴ R. Waldi,⁶⁴ T. Adye,⁶⁵ G. Castelli,⁶⁵ B. Franek,⁶⁵ E. O. Olaiya,⁶⁵ S. Ricciardi,⁶⁵ W. Roethel,⁶⁵ F. F. Wilson,⁶⁵ R. Aleksan,⁶⁶ S. Emery,⁶⁶ M. Escalier,⁶⁶ A. Gaidot,⁶⁶ S. F. Ganzhur,⁶⁶ G. Hamel de Monchenault,⁶⁶ W. Kozanecki,⁶⁶ G. Vasseur,⁶⁶ Ch. Yèche,⁶⁶ M. Zito,⁶⁶ X. R. Chen,⁶⁷ H. Liu,⁶⁷ W. Park,⁶⁷ M. V. Purohit,⁶⁷ J. R. Wilson,⁶⁷ M. T. Allen,⁶⁸ D. Aston,⁶⁸ R. Bartoldus,⁶⁸ P. Bechtle,⁶⁸ N. Berger,⁶⁸ R. Claus,⁶⁸ J. P. Coleman,⁶⁸ M. R. Convery,⁶⁸ J. C. Dingfelder,⁶⁸ J. Dorfan,⁶⁸ G. P. Dubois-Felsmann,⁶⁸ W. Dunwoodie,⁶⁸ R. C. Field,⁶⁸ T. Glanzman,⁶⁸ S. J. Gowdy,⁶⁸ M. T. Graham,⁶⁸ P. Grenier,⁶⁸ C. Hast,⁶⁸ T. Hryna'ova,⁶⁸ W. R. Innes,⁶⁸ J. Kaminski,⁶⁸ M. H. Kelsey,⁶⁸ H. Kim,⁶⁸ P. Kim,⁶⁸ M. L. Kocian,⁶⁸ D. W. G. S. Leith,⁶⁸ S. Li,⁶⁸ S. Luitz,⁶⁸ V. Luth,⁶⁸ H. L. Lynch,⁶⁸ D. B. MacFarlane,⁶⁸ H. Marsiske,⁶⁸ R. Messner,⁶⁸ D. R. Muller,⁶⁸ C. P. O'Grady,⁶⁸ I. Ofte,⁶⁸ A. Perazzo,⁶⁸ M. Perl,⁶⁸ T. Pulliam,⁶⁸ B. N. Ratcliff,⁶⁸ A. Roodman,⁶⁸ A. A. Salnikov,⁶⁸ R. H. Schindler,⁶⁸ J. Schwiening,⁶⁸ A. Snyder,⁶⁸ J. Stelzer,⁶⁸ D. Su,⁶⁸ M. K. Sullivan,⁶⁸ K. Suzuki,⁶⁸ S. K. Swain,⁶⁸ J. M. Thompson,⁶⁸ J. Va'vra,⁶⁸ N. van Bakel,⁶⁸ A. P. Wagner,⁶⁸ M. Weaver,⁶⁸ W. J. Wisniewski,⁶⁸ M. Wittgen,⁶⁸ D. H. Wright,⁶⁸ A. K. Yarritu,⁶⁸ K. Yi,⁶⁸ C. C. Young,⁶⁸ P. R. Burchat,⁶⁹ A. J. Edwards,⁶⁹ S. A. Majewski,⁶⁹ B. A. Petersen,⁶⁹ L. Wilden,⁶⁹ S. Ahmed,⁷⁰ M. S. Alam,⁷⁰ R. Bula,⁷⁰ J. A. Ernst,⁷⁰ V. Jain,⁷⁰ B. Pan,⁷⁰ M. A. Saeed,⁷⁰ F. R. Wappler,⁷⁰ S. B. Zain,⁷⁰ W. Bugg,⁷¹ M. Krishnamurthy,⁷¹ S. M. Spanier,⁷¹ R. Eckmann,⁷² J. L. Ritchie,⁷² A. M. Ruland,⁷² C. J. Schilling,⁷² R. F. Schwitters,⁷² J. M. Izzen,⁷³ X. C. Lou,⁷³ S. Ye,⁷³ F. Bianchi,⁷⁴ F. Gallo,⁷⁴ D. Gamba,⁷⁴ M. Pelliccioni,⁷⁴ M. Bomben,⁷⁵ L. Bosisio,⁷⁵ C. Cartaro,⁷⁵ F. Cossutti,⁷⁵ G. Della Ricca,⁷⁵ L. Lanceri,⁷⁵ L. Vitale,⁷⁵ V. Azzolini,⁷⁶ N. Lopez-March,⁷⁶ F. Martinez-Vidal,^{76,§} D. A. Milanes,⁷⁶ A. Oyanguren,⁷⁶ J. Albert,⁷⁷ Sw. Banerjee,⁷⁷ B. Bhuyan,⁷⁷ K. Hamano,⁷⁷ R. Kowalewski,⁷⁷ I. M. Nugent,⁷⁷ J. M. Roney,⁷⁷ R. J. Sobie,⁷⁷ J. J. Back,⁷⁸ P. F. Harrison,⁷⁸ J. Illic,⁷⁸ T. E. Latham,⁷⁸ G. B. Mohanty,⁷⁸ M. Pappagallo,^{78,||} H. R. Band,⁷⁹ X. Chen,⁷⁹ S. Dasu,⁷⁹ K. T. Flood,⁷⁹ J. J. Hollar,⁷⁹ P. E. Kutter,⁷⁹ Y. Pan,⁷⁹ M. Pierini,⁷⁹ R. Prepost,⁷⁹ S. L. Wu,⁷⁹ and H. Neal⁸⁰

(BABAR Collaboration)

¹Laboratoire de Physique des Particules, IN2P3/CNRS et Université de Savoie, F-74941 Annecy-Le-Vieux, France

²Facultat de Fisica, Departament ECM, Universitat de Barcelona, E-08028 Barcelona, Spain

³Dipartimento di Fisica and INFN, Università di Bari, I-70126 Bari, Italy

⁴Institute of Physics, University of Bergen, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Institut für Experimentalphysik 1, Ruhr Universität Bochum, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, California 92697, USA

¹³University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁴University of California at Riverside, Riverside, California 92521, USA

¹⁵University of California at San Diego, La Jolla, California 92093, USA

¹⁶University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁷Institute for Particle Physics, University of California at Santa Cruz, Santa Cruz, California 95064, USA

¹⁸California Institute of Technology, Pasadena, California 91125, USA

¹⁹University of Cincinnati, Cincinnati, Ohio 45221, USA

²⁰University of Colorado, Boulder, Colorado 80309, USA

²¹Colorado State University, Fort Collins, Colorado 80523, USA

²²Institut für Physik, Universität Dortmund, D-44221 Dortmund, Germany

²³Institut für Kern- und Teilchenphysik, Technische Universität Dresden, D-01062 Dresden, Germany

²⁴Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France

²⁵University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

²⁶Dipartimento di Fisica and INFN, Università di Ferrara, I-44100 Ferrara, Italy

²⁷Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy

²⁸Dipartimento di Fisica and INFN, Università di Genova, I-16146 Genova, Italy

²⁹Harvard University, Cambridge, Massachusetts 02138, USA

³⁰*Physikalisches Institut, Universität Heidelberg, Philosophenweg 12, D-69120 Heidelberg, Germany*³¹*Imperial College London, London, SW7 2AZ, United Kingdom*³²*University of Iowa, Iowa City, Iowa 52242, USA*³³*Iowa State University, Ames, Iowa 50011-3160, USA*³⁴*Johns Hopkins University, Baltimore, Maryland 21218, USA*³⁵*Institut für Experimentelle Kernphysik, Universität Karlsruhe, D-76021 Karlsruhe, Germany*³⁶*Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, B.P. 34, F-91898 ORSAY Cedex, France*³⁷*Lawrence Livermore National Laboratory, Livermore, California 94550, USA*³⁸*University of Liverpool, Liverpool L69 7ZE, United Kingdom*³⁹*Queen Mary, University of London, E1 4NS, United Kingdom*⁴⁰*University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom*⁴¹*University of Louisville, Louisville, Kentucky 40292, USA*⁴²*University of Manchester, Manchester M13 9PL, United Kingdom*⁴³*University of Maryland, College Park, Maryland 20742, USA*⁴⁴*University of Massachusetts, Amherst, Massachusetts 01003, USA*⁴⁵*Laboratory for Nuclear Science, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*⁴⁶*McGill University, Montréal, Québec, Canada H3A 2T8*⁴⁷*Dipartimento di Fisica and INFN, Università di Milano, I-20133 Milano, Italy*⁴⁸*University of Mississippi, University, Mississippi 38677, USA*⁴⁹*Physique des Particules, Université de Montréal, Montréal, Québec, Canada H3C 3J7*⁵⁰*Mount Holyoke College, South Hadley, Massachusetts 01075, USA*⁵¹*Dipartimento di Scienze Fisiche and INFN, Università di Napoli Federico II, I-80126, Napoli, Italy*⁵²*NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands*⁵³*University of Notre Dame, Notre Dame, Indiana 46556, USA*⁵⁴*The Ohio State University, Columbus, Ohio 43210, USA*⁵⁵*University of Oregon, Eugene, Oregon 97403, USA*⁵⁶*Dipartimento di Fisica and INFN, Università di Padova, I-35131 Padova, Italy*⁵⁷*Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie-Paris 6, Université Denis Diderot-Paris 7, F-75252 Paris, France*⁵⁸*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*⁵⁹*Dipartimento di Fisica and INFN, Università di Perugia, I-06100 Perugia, Italy*⁶⁰*Dipartimento di Fisica, Scuola Normale Superiore and INFN, Università di Pisa, I-56127 Pisa, Italy*⁶¹*Prairie View A&M University, Prairie View, Texas 77446, USA*⁶²*Princeton University, Princeton, New Jersey 08544, USA*⁶³*Dipartimento di Fisica and INFN, Università di Roma La Sapienza, I-00185 Roma, Italy*⁶⁴*Universität Rostock, D-18051 Rostock, Germany*⁶⁵*Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom*⁶⁶*DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France*⁶⁷*University of South Carolina, Columbia, South Carolina 29208, USA*⁶⁸*Stanford Linear Accelerator Center, Stanford, California 94309, USA*⁶⁹*Stanford University, Stanford, California 94305-4060, USA*⁷⁰*State University of New York, Albany, New York 12222, USA*⁷¹*University of Tennessee, Knoxville, Tennessee 37996, USA*⁷²*University of Texas at Austin, Austin, Texas 78712, USA*⁷³*University of Texas at Dallas, Richardson, Texas 75083, USA*⁷⁴*Dipartimento di Fisica Sperimentale and INFN, Università di Torino, I-10125 Torino, Italy*⁷⁵*Dipartimento di Fisica and INFN, Università di Trieste, I-34127 Trieste, Italy*⁷⁶*IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain*⁷⁷*University of Victoria, Victoria, British Columbia, Canada V8W 3P6*⁷⁸*Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom*⁷⁹*University of Wisconsin, Madison, Wisconsin 53706, USA*⁸⁰*Yale University, New Haven, Connecticut 06511, USA*

(Received 8 May 2007; published 17 August 2007)

We present updated measurements of CP -violating asymmetries in the decays $B^0 \rightarrow D^{*\pm} D^\mp$ and $B^0 \rightarrow D^+ D^-$ using $(383 \pm 4) \times 10^6 B\bar{B}$ pairs collected by the *BABAR* detector at the SLAC PEP-II *B* factory. We determine the time-integrated CP asymmetry $\mathcal{A}_{D^{*\pm} D^\mp} = 0.12 \pm 0.06 \pm 0.02$, and the time-dependent asymmetry parameters to be $C_{D^{*+} D^-} = 0.18 \pm 0.15 \pm 0.04$, $S_{D^{*+} D^-} = -0.79 \pm 0.21 \pm 0.06$, $C_{D^{*-} D^+} = 0.23 \pm 0.15 \pm 0.04$, $S_{D^{*-} D^+} = -0.44 \pm 0.22 \pm 0.06$, $C_{D^+ D^-} = 0.11 \pm 0.22 \pm 0.07$, and $S_{D^+ D^-} = -0.54 \pm 0.34 \pm 0.06$, where the first uncertainty is statistical and the second is systematic.

In the standard model (SM), CP violation arises from a complex phase in the Cabibbo-Kobayashi-Maskawa quark-mixing matrix V [1]. Measurements of CP asymmetries in $B^0 \rightarrow (c\bar{c})K^{(*)0}$ decays [2] by the *BABAR* [3] and *Belle* [4] Collaborations have firmly established this effect and precisely determined the parameter $\sin 2\beta$, where β is $\arg[-V_{cd}V_{cb}^*/V_{td}V_{tb}^*]$. Another way to measure $\sin 2\beta$ is to use decays whose amplitudes are dominated by a tree-level, color-allowed $b \rightarrow c\bar{c}d$ transition, such as $B^0 \rightarrow D^{(*)\pm}D^\mp$. Within the framework of the SM, the time-dependent CP asymmetries of $B^0 \rightarrow D^{(*)\pm}D^\mp$ are directly related to $\sin 2\beta$ when corrections due to penguin diagram contributions are neglected. The penguin-induced corrections have been estimated in models based on the factorization approximation and heavy quark symmetry and are predicted to be a few percent [5,6]. However, contributions from non-SM processes may lead to a large shift [7]. A significant deviation in the $\sin 2\beta$ measurement from that of the $B^0 \rightarrow (c\bar{c})K^{(*)0}$ decays would be evidence involving new physics beyond the SM.

Studies of the CP violation in $b \rightarrow c\bar{c}d$ transitions have been carried out by both the *BABAR* and *Belle* Collaborations. Most recently, the *Belle* Collaboration reported evidence of large direct CP violation in $B^0 \rightarrow D^+D^-$ where $C_{D^+D^-} = -0.91 \pm 0.23 \pm 0.06$ [8], in contradiction to the SM expectation. However, such a large direct CP violation has not been observed in previous measurements with $B^0 \rightarrow D^{(*)\pm}D^{(*)\mp}$ decays, involving the same quark-level weak decay [9–12].

In this Letter, we present an updated measurement of CP -violating asymmetries in the decays $B^0 \rightarrow D^{*\pm}D^-$, $B^0 \rightarrow D^{*-}D^+$, and $B^0 \rightarrow D^+D^-$. The data used in this analysis comprise $(383 \pm 4) \times 10^6$ $Y(4S) \rightarrow B\bar{B}$ decays collected by the *BABAR* detector at the PEP-II storage rings. The *BABAR* detector is described in detail elsewhere [13]. Monte Carlo (MC) simulation based on GEANT4 [14] is used to validate the analysis procedure and to study the relevant backgrounds.

The decay rate $f_+(f_-)$ for a neutral B meson decay to a common final state accompanied by a $B^0(\bar{B}^0)$ tag is given by

$$f_\pm(\Delta t) = e^{-|\Delta t|/\tau_{B^0}} / 4\tau_{B^0} \{ (1 \mp \Delta w) \pm (1 - 2w) \times [S \sin(\Delta m_d \Delta t) - C \cos(\Delta m_d \Delta t)] \}, \quad (1)$$

where $\Delta t \equiv t_{\text{rec}} - t_{\text{tag}}$ is the difference between the proper decay time of the reconstructed B meson (B_{rec}) and that of the tagging B meson (B_{tag}), τ_{B^0} is the B^0 lifetime, and Δm_d is the difference between the heavy and light mass eigenstates determined from the $B^0 - \bar{B}^0$ oscillation frequency [15]. The average mistag probability w describes the effect of incorrect tags, and Δw is the difference between the

mistag probabilities for B^0 and \bar{B}^0 . Since $D^{*\pm}D^-$ and $D^{*-}D^+$ are not CP eigenstates, we can define a time-integrated asymmetry $\mathcal{A}_{D^{*\pm}D^\mp}$ between the rate of $B^0 \rightarrow D^{*\pm}D^-$ and $B^0 \rightarrow D^{*-}D^+$, calculated as

$$\mathcal{A}_{D^{*\pm}D^\mp} = \frac{N_{D^{*\pm}D^-} - N_{D^{*-}D^+}}{N_{D^{*\pm}D^-} + N_{D^{*-}D^+}}, \quad (2)$$

where N is the signal event yield.

For $B^0 \rightarrow D^{*\pm}D^\mp$, the general relations are $S_{D^{*\pm}D^\mp} = -\sqrt{1 - C_{D^{*\pm}D^\mp}^2} \sin(2\beta_{\text{eff}} \pm \delta)$, where δ is the strong phase difference between $B^0 \rightarrow D^{*\pm}D^-$ and $B^0 \rightarrow D^{*-}D^+$ [16]. Under the assumption of negligible penguin contribution, $\beta_{\text{eff}} = \beta$, $\mathcal{A}_{D^{*\pm}D^\mp} = 0$, and $C_{D^{*\pm}D^-} = -C_{D^{*-}D^+}$. For $B^0 \rightarrow D^+D^-$ and in the case of negligible penguin contribution, $C_{D^+D^-}$ measures direct CP violation and is zero, while $S_{D^+D^-}$ is $-\sin 2\beta$.

The selections of $B^0 \rightarrow D^{*\pm}D^\mp$ and $B^0 \rightarrow D^+D^-$ candidates are similar to those of our previous analysis [10]. We reconstruct $D^{*\pm}$ in its decay to $D^0\pi^\pm$. We reconstruct candidates for D^0 and D^+ mesons in the modes $D^0 \rightarrow K^-\pi^+$, $K^-\pi^+\pi^0$, $K^-\pi^+\pi^+\pi^-$, $K_S^0\pi^+\pi^-$, and $D^+ \rightarrow K^-\pi^+\pi^+$, $K_S^0\pi^+$. We reconstruct $B^0 \rightarrow D^+D^-$ candidates only through the decay $D^\pm \rightarrow K^\mp\pi^\pm\pi^\pm$. We require the reconstructed masses of the D^0 and D^+ candidates to be within $20 \text{ MeV}/c^2$ of their respective nominal masses [15], except for the $D^0 \rightarrow K^-\pi^+\pi^0$ candidate, where we use a looser requirement of $40 \text{ MeV}/c^2$. We apply a mass-constrained fit to the selected D^0 and D^+ candidates and combine D^0 candidates with a π^+ track, with momentum below $450 \text{ MeV}/c$ in the $Y(4S)$ frame, to form $D^{*\pm}$ candidates.

We reconstruct the K_S^0 candidates from two oppositely charged tracks with an invariant mass within $20 \text{ MeV}/c^2$ of the nominal K_S^0 mass [15]. The χ^2 probability of the track vertex fit must be greater than 0.1%. We require charged kaon candidates to be identified as such using a likelihood technique based on the Cherenkov angle measured by the Cherenkov detector and the ionization energy loss measured by the charged-particle tracking systems [13]. We form neutral pion candidates from two photons detected in the electromagnetic calorimeter [13], each with energy above 30 MeV. The invariant mass of the pair must be within $30 \text{ MeV}/c^2$ of the nominal π^0 mass [15], and we require their summed energy to be greater than 200 MeV. In addition, we further apply a mass-constrained fit to the π^0 candidates.

To suppress the $e^+e^- \rightarrow q\bar{q}$ ($q = u, d, s$, and c) continuum background, we exploit the contrast between the spherical shape of $B\bar{B}$ events and the more jetlike nature of continuum events. We require the ratio of the second to the zeroth order Fox-Wolfram moments [17] to be less than

0.6. We also use a Fisher discriminant, constructed as an optimized linear combination of 11 event shape variables [18]: the momentum flow in nine concentric cones around the thrust axis of the reconstructed B^0 candidate, the angle between that thrust axis and the beam axis, and the angle between the line of flight of the B^0 candidate and the beam axis. In addition, we employ a combined D flight-length significance variable, derived from the sum of flight lengths of the two D candidates [19], to reduce background.

For each $B^0 \rightarrow D^{(*)\pm} D^\mp$ candidate, we construct a likelihood function $\mathcal{L}_{\text{mass}}$ from the masses and mass uncertainties of the D and D^* candidates [19]. The D mass resolution is modeled by a Gaussian whose variance is determined on a candidate-by-candidate basis from its mass uncertainty before the mass-constrained fit. The $D^* - D$ mass difference resolution is modeled by the sum of two Gaussian distributions whose parameters are determined from simulated events. The values of $\mathcal{L}_{\text{mass}}$ and $\Delta E \equiv E_B^* - E_{\text{beam}}$, the difference between the B^0 candidate energy E_B^* and the beam energy E_{beam} in the $\Upsilon(4S)$ frame, are used to reduce the combinatoric background. From the simulated events, we optimize the maximum allowed values of $-\ln \mathcal{L}_{\text{mass}}$ and $|\Delta E|$ for each individual final state to obtain the highest expected signal significance.

We extract the signal yield from the events satisfying the selection criteria using the energy-substituted mass, $m_{\text{ES}} \equiv \sqrt{E_{\text{beam}}^2 - p_B^{*2}}$, where p_B^* is the B^0 candidate momentum in the $\Upsilon(4S)$ frame. We select the B^0 candidates that have $m_{\text{ES}} \geq 5.23 \text{ GeV}/c^2$. On average, we have 1.5 and 1.1 B^0 candidates per event for $B^0 \rightarrow D^{*\pm} D^\mp$ and $B^0 \rightarrow D^+ D^-$, respectively. If more than one candidate is reconstructed in an event, we select the candidate with the smallest value of $-\ln \mathcal{L}_{\text{mass}}$. Studies using MC samples show that this procedure results in the selection of the correct B^0 candidate more than 95% of the time.

We perform an unbinned maximum likelihood fit to the m_{ES} and Δt distributions to extract the CP asymmetries. We fit the events from $B^0 \rightarrow D^{*+} D^-$ and $B^0 \rightarrow D^{*-} D^+$

decays simultaneously. The probability density function (PDF) of the m_{ES} distribution consists of a Gaussian for the signal and a threshold function [20] for the combinatorial background. We expect some background events to peak in the m_{ES} signal region due to cross feed from other decay modes. We estimate the fraction of events in the signal Gaussian due to this peaking background to be $(8.8 \pm 4.4)\%$ for $B^0 \rightarrow D^{*\pm} D^\mp$ and $(4.8 \pm 7.4)\%$ for $B^0 \rightarrow D^+ D^-$ using detailed MC simulations of inclusive B decays.

The technique used to fit the Δt distribution is analogous to that used in previous *BABAR* measurements described in Refs. [21,22]. We use information from the other B meson in the event to tag the flavor of the fully reconstructed $B^0 \rightarrow D^{(*)\pm} D^\mp$ candidate [21]. The signal Δt PDF in Eq. (1) is convolved with an empirical Δt resolution function [21]. The Δt is calculated from the measured separation Δz between the decay vertices of B_{rec} and B_{tag} along the collision (z) axis [21]. The B_{tag} decay vertex is determined by fitting charged tracks not belonging to the B_{rec} candidate to a common vertex, employing constraints from the beam-spot location and the B_{rec} momentum [21]. Only events with a Δt uncertainty less than 2.5 ps and a measured $|\Delta t|$ less than 20 ps are accepted for the fit to the Δt distribution. Both the signal mistag probability and the Δt resolution function are determined from a large sample of neutral B decays to flavor eigenstates B_{flav} . The combinatoric background Δt distributions are parametrized with an empirical description that includes zero and nonzero lifetime components [21]. The nonzero lifetime background is allowed to have effective CP asymmetries, and these float in the likelihood fit. By default, we assume that the peaking backgrounds have the same Δt PDF as the signal but zero CP asymmetries.

The fits to the data yield 280 ± 19 signal events for $B^0 \rightarrow D^{*+} D^-$, 219 ± 18 signal events for $B^0 \rightarrow D^{*-} D^+$, and 131 ± 14 signal events for $B^0 \rightarrow D^+ D^-$, where the quoted uncertainties are statistical only. In the region of $m_{\text{ES}} > 5.27 \text{ GeV}/c^2$, the signal purity is approximately 41% for $B^0 \rightarrow D^{*+} D^-$, 34% for $B^0 \rightarrow D^{*-} D^+$, and 46% for $B^0 \rightarrow D^+ D^-$. The fitted CP violating parameters are




FIG. 1. Measured distribution of m_{ES} for (a) $B^0 \rightarrow D^{*+} D^-$, (b) $B^0 \rightarrow D^{*-} D^+$, and (c) $B^0 \rightarrow D^+ D^-$ candidates. The solid line is the projection of the fit result and the dotted line represents the background components.




FIG. 2. The distributions of Δt and fit projections for $B^0 \rightarrow D^{*\pm} D^\mp$ (left), $B^0 \rightarrow D^{*-} D^+$ (middle), and $B^0 \rightarrow D^+ D^-$ (right) candidates in the signal region $m_{ES} > 5.27 \text{ GeV}/c^2$ with a B^0 or \bar{B}^0 tag (a)–(c). The raw time-dependent asymmetries ($N_{B^0} - N_{\bar{B}^0}$) / ($N_{B^0} + N_{\bar{B}^0}$) as functions of Δt are also shown (d)–(e).

$$\begin{aligned} \mathcal{A}_{D^{*\pm} D^\mp} &= 0.12 \pm 0.06 \pm 0.02, \\ C_{D^{*\pm} D^\mp} &= 0.18 \pm 0.15 \pm 0.04, \\ S_{D^{*\pm} D^\mp} &= -0.79 \pm 0.21 \pm 0.06, \\ C_{D^{*-} D^+} &= 0.23 \pm 0.15 \pm 0.04, \\ S_{D^{*-} D^+} &= -0.44 \pm 0.22 \pm 0.06, \\ C_{D^+ D^-} &= 0.11 \pm 0.22 \pm 0.07, \\ S_{D^+ D^-} &= -0.54 \pm 0.34 \pm 0.06, \end{aligned} \quad (3)$$

where the first uncertainty is statistical and the second is systematic.

Projections of the fits onto m_{ES} for the three different samples are shown in Fig. 1. Figure 2 shows the Δt distributions and asymmetries in yields between events with B^0 and \bar{B}^0 tags, overlaid with the projection of the likelihood fit result. As a cross-check, we repeat the fit by allowing the B^0 lifetime to float. The obtained lifetime is in good agreement with its world average [15].

The systematic uncertainty of the time-integrated CP asymmetry $\mathcal{A}_{D^{*\pm} D^\mp}$ is dominated by the potential differences in the reconstruction efficiencies of the positively and negatively charged tracks (0.014). Other sources that contribute to the systematic error include the estimate of

the peaking background fraction (< 0.001), the uncertainty in the m_{ES} resolution for the $B^0 \rightarrow D^{*\pm} D^\mp$ signal events (0.005), and a possible fit bias (0.004).

The systematic uncertainties on C and S are evaluated separately for each of the decay modes. Their sources and estimates are summarized in Table I. The systematic uncertainties arise from the amount of possible background that tends to peak under the signal and its CP asymmetry, the assumed parametrization of the Δt resolution function, the possible differences between the B_{flav} and signal mistag fractions, the knowledge of the event-by-event beam-spot position, the uncertainties from the finite MC sample used, the possible interference between the suppressed $\bar{b} \rightarrow \bar{u} c \bar{d}$ and the favored $b \rightarrow c \bar{u} \bar{d}$ amplitudes in some tag-side decays [23], and the uncertainty in the m_{ES} resolution for the signal events. All of the systematic uncertainties are found to be much smaller than the statistical uncertainties.

Since $D^{*\pm} D^\mp$ and $D^{*-} D^+$ are not CP eigenstates, it is also illustrative to express the measured CP -violating parameters C and S in a slightly different parametrization [24]: $C_{D^* D} = (C_{D^{*+} D^-} + C_{D^{*-} D^+})/2$, $\Delta C_{D^* D} = (C_{D^{*+} D^-} - C_{D^{*-} D^+})/2$, $S_{D^* D} = (S_{D^{*+} D^-} + S_{D^{*-} D^+})/2$, and $\Delta S_{D^* D} = (S_{D^{*+} D^-} - S_{D^{*-} D^+})/2$. The quantities $C_{D^* D}$ and $S_{D^* D}$ parametrize flavor-dependent direct CP violation and mixing-induced CP violation re-

TABLE I. Sources of systematic error on time-dependent CP asymmetry parameters for the decays $B^0 \rightarrow D^{*\pm} D^\mp$ and $B^0 \rightarrow D^+ D^-$.

Source	$C_{D^{*\pm} D^\mp}$	$S_{D^{*\pm} D^\mp}$	$C_{D^{*-} D^+}$	$S_{D^{*-} D^+}$	$C_{D^+ D^-}$	$S_{D^+ D^-}$
Peaking backgrounds	0.026	0.041	0.027	0.031	0.044	0.042
Δt resolution parametrization	0.011	0.021	0.013	0.012	0.015	0.020
Mistag fraction differences	0.014	0.011	0.016	0.012	0.023	0.013
Beam-spot position	0.004	0.006	0.007	0.036	0.005	0.002
Δm_d , τ_B	0.002	0.003	0.003	0.004	0.001	0.004
MC statistics	0.011	0.015	0.011	0.015	0.036	0.023
Tag-side interference and others	0.016	0.025	0.017	0.020	0.020	0.013
Total	0.037	0.056	0.040	0.056	0.066	0.055

lated to the angle β , respectively. The parameters ΔC_{D^*D} and ΔS_{D^*D} are insensitive to CP violation. ΔC_{D^*D} describes the asymmetry between the rates $\Gamma(B^0 \rightarrow D^{*+}D^-) + \Gamma(\bar{B}^0 \rightarrow D^{*-}D^+)$ and $\Gamma(B^0 \rightarrow D^{*-}D^+) + \Gamma(\bar{B}^0 \rightarrow D^{*+}D^-)$, while ΔS_{D^*D} is related to the strong phase difference δ . We find

$$\begin{aligned} C_{D^*D} &= 0.21 \pm 0.11 \pm 0.03, \\ S_{D^*D} &= -0.62 \pm 0.15 \pm 0.04, \\ \Delta C_{D^*D} &= -0.02 \pm 0.11 \pm 0.03, \\ \Delta S_{D^*D} &= -0.17 \pm 0.15 \pm 0.04, \end{aligned} \quad (4)$$

where the first uncertainty is statistical and the second is systematic.

In summary, this Letter reports updated measurements of the CP -violating asymmetries for the decays $B^0 \rightarrow D^{*\pm}D^\mp$ and $B^0 \rightarrow D^+D^-$. These measurements supersede the previous *BABAR* results [10], with a more than 50% reduction in the statistical uncertainties. The time-dependent asymmetries are consistent with the SM predictions within their statistical uncertainties. We do not see evidence of large direct CP violation in the decay $B^0 \rightarrow D^+D^-$ as reported by the *Belle* Collaboration [8].

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (U.S.A.), NSERC (Canada), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), MEC (Spain), and STFC (United Kingdom). Individuals have received support from the Marie Curie EIF (European Union) and the A. P. Sloan Foundation.

*Deceased.

[†]Also at: Dipartimento di Fisica, Università di Perugia, Perugia, Italy.

[‡]Also at: Università della Basilicata, Potenza, Italy.

[§]Also at: Facultat de Fisica, Departament ECM, Universitat de Barcelona, E-08028 Barcelona, Spain.

[¶]Also at: IPPP, Physics Department, Durham University, Durham DH1 3LE, United Kingdom.

- [1] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [2] We imply charge conjugate modes throughout the Letter.
- [3] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **89**, 201802 (2002).
- [4] K. Abe *et al.* (*Belle* Collaboration), Phys. Rev. D **66**, 071102 (2002).
- [5] Z. Z. Xing, Phys. Lett. B **443**, 365 (1998).
- [6] Z. Z. Xing, Phys. Rev. D **61**, 014010 (1999).
- [7] Y. Grossman and M. P. Worah, Phys. Lett. B **395**, 241 (1997).
- [8] S. Fratina *et al.* (*Belle* Collaboration), Phys. Rev. Lett. **98**, 221802 (2007).
- [9] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **95**, 151804 (2005).
- [10] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **95**, 131802 (2005).
- [11] T. Aushev *et al.* (*Belle* Collaboration), Phys. Rev. Lett. **93**, 201802 (2004).
- [12] H. Miyake *et al.* (*Belle* Collaboration), Phys. Lett. B **618**, 34 (2005).
- [13] B. Aubert *et al.* (*BABAR* Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [14] S. Agostinelli *et al.* (*GEANT4* Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **506**, 250 (2003).
- [15] W. M. Yao *et al.* (Particle Data Group), J. Phys. G **33**, 1 (2006).
- [16] R. Aleksan, A. Le Yaouanc, L. Oliver, O. Pene, and J. C. Raynal, Phys. Lett. B **317**, 173 (1993).
- [17] G. C. Fox and S. Wolfram, Phys. Rev. Lett. **41**, 1581 (1978).
- [18] D. M. Asner *et al.* (*CLEO* Collaboration), Phys. Rev. D **53**, 1039 (1996).
- [19] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. D **73**, 112004 (2006).
- [20] H. Albrecht *et al.* (*ARGUS* Collaboration), Z. Phys. C **48**, 543 (1990).
- [21] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. D **66**, 032003 (2002).
- [22] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **94**, 161803 (2005).
- [23] O. Long, M. Baak, R. N. Cahn, and D. Kirkby, Phys. Rev. D **68**, 034010 (2003).
- [24] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **91**, 201802 (2003).