

Edge Solitons of Topological Insulators and Fractionalized Quasiparticles in Two Dimensions

Dung-Hai Lee,^{1,2} Guang-Ming Zhang,³ and Tao Xiang^{4,5}

¹Department of Physics, University of California at Berkeley, Berkeley, California 94720, USA

²Material Science Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

³Department of Physics, Tsinghua University, Beijing 100084, China

⁴Institute of Physics, Chinese Academy of Sciences, P.O. Box 603, Beijing 100080, China

⁵Institute of Theoretical Physics, Chinese Academy of Sciences, P.O. Box 2735, Beijing 100080, China

(Received 31 May 2007; published 9 November 2007)

An important characteristic of topological band insulators is the necessary presence of in-gap edge states on the sample boundary. We utilize this fact to show that when the boundary is reconnected with a twist, there are always zero-energy defect states. This provides a natural connection among novel defects in the two-dimensional $p_x + ip_y$ superconductor, the Kitaev model, the fractional quantum Hall effect, and the one-dimensional domain wall of polyacetylene.

DOI: 10.1103/PhysRevLett.99.196805

PACS numbers: 73.43.-f, 71.10.Li

Excitations carrying fractional quantum numbers (e.g., fractional charge), such as the quasiparticles in the fractional quantum Hall effect [1], have always been a subject of interest. In 1976 Jackiw and Rebbi [2] wrote a seminal paper which laid the foundation of charge fractionalization in one spatial dimension. Four years later the influential paper of Su, Schrieffer, and Heeger [3] proposed the “Jackiw-Rebbi soliton” as the charge carrier in doped polyacetylene. Today, all quantum number fractionalization phenomena, such as the fractionalization of magnon into spinons [4], can be attributed to the Jackiw-Rebbi mechanism.

In the last 15 years, starting with the fractional quantum Hall effect, condensed matter physicists stumbled upon several instances where quantum number fractionalization occurs in two spatial dimensions. These include the quasiparticles of the “Pfaffian” quantum Hall state [5], the vortices of a spin-polarized $p_x + ip_y$ superconductor [6–10], and the topological excitations in a spin model proposed by Kitaev [11]. However, what is lacking is a general framework, like the Jackiw-Rebbi theory in one dimension, specifying the condition under which fractionalized excitations will appear. In this Letter, we provide such a mechanism and reveal its connection to the Jackiw-Rebbi theory. In particular, we show that, in two dimensions, fractional charge will naturally appear around defects of “topological” insulators [12–14].

Recently it has been shown that the existence of fractional charge in the quantum Hall effect is connected to the existence of fractionally charged domain walls in certain one-dimensional systems [15–18]. In the following we generalize such a connection and show that when the boundary of a topological insulator, an insulator which necessarily possesses an in-gap edge state, is reconnected with a twist, there are always zero-energy defect states possessing fractional quantum numbers.

Let us begin by considering the Kitaev model [11]. This exactly soluble model describes a honeycomb lattice of quantum one-half spins interacting via three type of

nearest-neighbor interactions [Fig. 1(a)]. The Hamiltonian is given by

$$H = \frac{1}{2} \sum_{n \in w} \sum_{\mu=1,2,3} J_{\mu} \sigma_{n+e_{\mu}}^{\mu} \sigma_n^{\mu}, \quad (1)$$

where w (b) denotes the white (black) sublattice [see Fig. 1(a)], $n + e_{\mu}$ is the nearest neighbors of n along the μ bond, and σ^{μ} are the Pauli matrices. By performing Jordan-Wigner transformation it was shown in Ref. [19] that this model is equivalent to a free Majorana fermion model:

$$H_M = -i \sum_{n \in w} \left\{ \sum_{\mu=1,2} J_{\mu} \gamma_{n+e_{\mu}} \gamma_n + J_3 D_n \gamma_{n+e_3} \gamma_n \right\}, \quad (2)$$

where $D_n = \pm 1$ is a classical Ising variable and γ_n 's are Majorana fermion operators [19]. Since the honeycomb lattice consists of two sublattices, Eq. (2) can be recast into




FIG. 1 (color online). (a) A graphical representation of the Kitaev model. There are two sublattices (white and black), and three types of bonds (labeled by 1, 2, and 3). (b) The graphical representation of $H + H_t$. In the Majorana representation, the three-spin interactions in Eq. (5) become the next neighbor hopping along the zigzag chain in the x direction. The arrow represents the direction in which the second neighbor hopping matrix elements are $-iJ_4$.

$$H_M = \sum_{n \in w, m \in w} \Psi_n^\dagger H_{nm} \Psi_m, \quad (3)$$

where $\Psi_n^\dagger = (\gamma_n, \gamma_{n+e_3})$. In Eq. (3) the 2×2 coupling matrix H_{nm} has the values $H_{nm} = J_3 D_n \sigma_2$ for $n = m$ and $H_{nm} = \mp (iJ_\mu/2) \sigma_\pm$ for $n \neq m$. The upper sign applies if the bond linking m to n is a black-to-white bond, and the lower sign applies if it is a white-to-black bond.

In the ground state, the classical Ising variable takes value $D_n = 1$ modulo a global flip per row [19]. For $D_n = 1$, H_M is translation invariant and can be diagonalized by Fourier transformation. The Bloch matrix of H_M is given by $H_M(\mathbf{k}) = h_1(\mathbf{k})\sigma_1 + h_2(\mathbf{k})\sigma_2$ with

$$\begin{aligned} h_1(\mathbf{k}) &= -J_2 \sin\alpha(\mathbf{k}) + J_1 \sin\beta(\mathbf{k}), \\ h_2(\mathbf{k}) &= J_3 + J_2 \cos\alpha(\mathbf{k}) + J_1 \cos\beta(\mathbf{k}), \end{aligned} \quad (4)$$

and $\alpha(\mathbf{k}) = (\sqrt{3}k_x - 3k_y)/2$, $\beta(\mathbf{k}) = (\sqrt{3}k_x + 3k_y)/2$. It is easy to show that for $J_1 + J_2 \geq J_3 \geq |J_1 - J_2|$ the energy spectrum is gapless. The possible connection to the quantum spin liquids has been discussed in Ref. [20]. In Ref. [11] a magnetic field is introduced to open an excitation gap in this parameter region, and the non-Abelian quasiparticles become low energy excitations of this gapped phase.

Unfortunately the magnetic field spoils the integrability of the model. Here we propose a different way of opening a gap while maintaining the integrability. This is achieved by adding the following three-spin interaction to Eq. (1):

$$H_t = \frac{J_4}{2} \sum_{(ijk) \in \Delta} \sigma_i^2 \sigma_j^3 \sigma_k^1 + \frac{J_4}{2} \sum_{(ijk) \in \nabla} \sigma_i^1 \sigma_j^3 \sigma_k^2. \quad (5)$$

Here (ijk) denote three adjacent sites (with i being the leftmost one) along the zigzag chain running along the x direction. Depending on whether (ijk) form an up-pointing or a down-pointing triangle, we use the first or second term of Eq. (5). In terms of the Majorana fermion operators, this amounts to adding a second nearest-neighbor hopping between sites along the zigzag chain:

$$H_t = -iJ_4 \left(\sum_{i,k \in w} \gamma_i \gamma_k - \sum_{i,k \in b} \gamma_i \gamma_k \right). \quad (6)$$

Then the Bloch matrix becomes $H_M(\mathbf{k}) = h_1(\mathbf{k})\sigma_1 + h_2(\mathbf{k})\sigma_2 + h_3(\mathbf{k})\sigma_3$, where $h_3(\mathbf{k}) = 2J_4 \sin(\sqrt{3}k_x)$. The vector function $\mathbf{h}(\mathbf{k})$ is a continuous mapping from the first Brillouin zone to the space spanned by $\mathbf{h} = (h_1, h_2, h_3)$. The image is a closed two-dimensional manifold (henceforth referred as the h surface). Since the eigenvalues of the Bloch matrix are $\pm |\mathbf{h}(\mathbf{k})|$, it follows that if the h surface contains the origin, the spectrum is gapless; otherwise the spectrum has a gap.

For an h surface not containing the origin, there is an integer topological index

$$\mathcal{P} = \frac{1}{8\pi} \int d^2k \epsilon^{\mu\nu} \hat{\mathbf{h}} \cdot (\partial_{k_\mu} \hat{\mathbf{h}} \times \partial_{k_\nu} \hat{\mathbf{h}}) \quad (7)$$

which counts the number of times the unit vector $\hat{\mathbf{h}}$ wraps around the origin. As shown in Ref. [21], \mathcal{P} is proportional to the well-known ‘‘Thouless-Kohmoto-Nightingale-den Nijs’’ index [22] in the case of two bands. Spectra characterized by different \mathcal{P} are topologically distinct. They cannot be deformed into each other without gap closing. In the parameter regime where the non-Abelian quasiparticle exists $\mathcal{P} = 1$.

Topological excitations of the Kitaev model are created by reversing the sign of D_n 's in H_M along half a row. This is shown by the dashed bonds in Fig. 2(c). In Fig. 2(d) we have shown the result of numerical diagonalization for a system of 1,600 sites with a toric boundary condition. Because of the boundary condition two defects are introduced; they are separated by 100 sites in the x direction. They introduce two midgap states with a tunnel splitting (which is already invisible here) which decreases exponentially with the separation.

Now we switch gears to discuss the vortices in a spin-polarized $p_x + ip_y$ superconductor. Let us consider




FIG. 2 (color online). The h surface for the Kitaev model (a) and the $p_x + ip_y$ superconductor (e). In panels (b) and (f) the h surfaces are dissected to expose the origin (bold dot). (c) Topological defects in the Kitaev model. To the right of the gray-shaded plaquette the sign of D_n 's are reversed. As a result the corresponding vertical hopping matrix elements change sign. They are shown by the dashed bonds. (d) The eigenspectrum associated with two far-separated topological defects in the Kitaev model with $J_1 = J_2 = 1$, $J_3 = 0.2$, and $J_4 = 0.5$. (g) A vortex centered at the gray-shaded plaquette in a $p_x + ip_y$ superconductor. The pairing order parameter is shown for four bonds. (h) The energy spectrum of two far-separated vortex-antivortex pairs with $t = 1$, $\Delta_0 = 0.5$, and $\mu = 0.3$.

this problem on a square lattice. Similar to the Kitaev model, the Hamiltonian can also be written in the form of Eq. (3), except $\Psi_n^\dagger = (c_n^\dagger, c_n)$ and c_n is a fermion operator. The Bloch matrix is characterized by $\mathbf{h}(\mathbf{k}) = [-\Delta_0 \sin k_y, \Delta_0 \sin k_x, -t(\cos k_x + \cos k_y) - \mu]$. Here Δ_0 is the pairing amplitude, t is the hopping integral, and μ is the chemical potential. The h surface is shown in Figs. 2(e) and 2(f). Straightforward calculation shows that $\mathcal{P} = 1$. After a singular gauge transformation, a vortex can be created by reversing the sign of the hopping matrix elements along a cut as shown in Fig. 2(g). Explicit calculation shows that there is also a zero mode associated with each vortex [Fig. 2(h)].

For each free Majorana or Bogoliubov fermion model discussed above, there is a free fermion model with an identical excitation spectrum. To obtain this fermion model we simply replace Ψ_n^\dagger in Eq. (3) by the fermion operator $\Psi_n^\dagger = (c_{1n}^\dagger, c_{2n}^\dagger)$, where 1 and 2 are “flavor” indices (they might represent the two sites in the unit cell of a lattice). This fermion model acts as a representative of all models which share the same H_{nm} (hence the same eigenspectrum). However, while the representative fermion model is global U(1) invariant, the Majorana and Bogoliubov fermion models have only Z_2 symmetry. The fact that in the Z_2 models the particle number is conserved only modulo two is the root of non-Abelian statistics. In the rest of the discussion, we refer to the system described by a gapped free fermion model with nonzero \mathcal{P} as a “topological band insulator.” Thus the representative fermion models for the Kitaev model and the $p_x + ip_y$ superconductor are topological band insulators. Knowing the properties of edge states and defects of the representative fermion model, one can readily deduce the corresponding properties of the Majorana fermion (Kitaev) or the Bogoliubov fermion ($p_x + ip_y$) models with the same H_{nm} . For example, while in the fermion model the edge states are free fermions, and the defect zero modes carry half fermion quantum numbers, those in the Majorana or Bogoliubov fermion model are true Majorana fermions.

In the following we provide a unifying mechanism for the appearance of defect zero mode when the representative fermion model describes a topological band insulator.

As an example, let us consider the fermion representative of the Kitaev model [Fig. 3(a)]. Figure 3(b) shows the gapped spectrum as a function of momentum along the longitudinal circle. Because of its topological nature, if we remove a row of bonds [Fig. 3(c)], in-gap edge states appear [23] as shown in Fig. 3(d). The left and right moving chiral edge fermions, represented by the arrowed-bolded lines in Fig. 3(c), are described by a massless Dirac Hamiltonian in 1D. If we reconnect the two edges, but with weaker bonds, a smaller gap reappears in the edge spectrum [see Figs. 3(e) and 3(f)]. The edge fermions are now described by a massive Dirac Hamiltonian




FIG. 3 (color online). (a) The Kitaev model and (b) its eigenspectrum as a function of wave vector k along the longitudinal direction around $k = \pi$. (c) The system cut open along the longitudinal direction and (d) the corresponding energy spectrum. The arrowed-bolded lines in (c) indicate the edge states are chiral fermions. (e) Hopping between the edge (thick gray vertical bonds) is reintroduced, but with a magnitude smaller than the bulk value. (f) The gapped energy spectrum corresponds to (e). (g) An edge soliton is introduced by reversing the sign for half of the vertical bonds between the edges. The sign reversal is indicated by the solid and dotted vertical bonds. (h) The energy spectrum corresponds to (f). The spectra in (b), (d), and (h) are obtained from H_F with $J_1 = J_2 = 1$, $J_3 = 0.2$, and $J_4 = 0.5$. The spectrum in (f) and (h) was obtained with a restored edge coupling $J_3 = \pm 0.02$.

$$H_E = \int dx (-iv\psi^\dagger \sigma_z \partial_x \psi + m\psi^\dagger \sigma_x \psi), \quad (8)$$

where v is the edge velocity, $\psi^\dagger = (\psi_R^\dagger, \psi_L^\dagger)$ with $\psi_{R/L}^\dagger$ being the right/left fermion creation operators. When the restored bonds have a sign reversal along half a row [Fig. 3(g)], the mass term in Eq. (8) becomes x dependent and changes sign as x goes through the location of the topological defect. This should result in one localized zero mode per defect according to Refs. [2,3]. Figure 3(h) shows that this is indeed true. The presence of such a zero mode, plus the fact that the spectra of Majorana fermion models are $E \leftrightarrow -E$ symmetric, immediately im-




FIG. 4 (color online). The evolution of the h surface as the model in Fig. 1(b) is gradually deformed into Haldane’s model. In constructing the figure we used $J_1 = J_2 = J_3 = 1$. The full strength of the second neighbor hopping is 0.5.

plies that the zero modes are Majorana fermion states in the original Kitaev model. Furthermore, using the argument of Ref. [24] it can be shown that the braiding of such Majorana fermion defects leads to the non-Abelian statistics. Thus, via the mechanism of Refs. [2,3] a *two-dimensional* defect with a fractionalized quantum number has emerged. Its presence is determined by the topological nature of the host bulk band insulator just as the edge states are.

Actually, a similar phenomenon was also found in the continuum theory of Dirac fermions interacting with the topological defects of a Higgs field [25]. However, in all the examples we considered here, the location of the *edge* Dirac point in the momentum space is far away from those of the *bulk* Dirac points. Consequently the theory discussed in Ref. [25] is not applicable here.

As a digression, we now show that the fermion representative of the Kitaev model is topologically equivalent to Haldane’s lattice model for the integer quantum Hall effect [12]. Figure 4 shows the evolution of the h surface by gradually switching off the second neighbor hopping in Haldane’s model which are not contained in Eq. (6). The leftmost column are the h surfaces for the Haldane model (top) and Kitaev (bottom), respectively. In the rest of the figure the surfaces are dissected to reveal the origin (the bold dot). Clearly as we follow the evolution the origin never migrates across the h surface. Thus \mathcal{P} for the two models is the same.

Finally what about the Laughlin quasiparticles? Although fractional quantum Hall liquids are not band insulators they are clearly *topological insulators*. Indeed, as shown by Wen [26], when a quantum Hall liquid on a torus is cut open [Fig. 3(a)], there is “chiral Luttinger liquid” in-gap edge modes. At $1/m$ filling, the edge modes are described by the following free boson Hamiltonian:

$$H_B = \int dx \left\{ \frac{2\pi}{m} \Pi(x)^2 + \frac{m}{8\pi} [\partial_x \phi(x)]^2 \right\}, \quad (9)$$

where Π and ϕ are conjugate boson fields satisfying $[\Pi(x), \phi(y)] = i\delta(x-y)$. To reconnect the edges, a potential $V = -g \int dx \cos(m\phi)$ needs to be added [26]. When g is sufficiently big, a gap opens in the edge spectrum and the ground states become m -fold degenerate. They are characterized by $\langle \phi(x) \rangle = 2\pi l/m$, ($l = 0, \dots, m-1$). In this case an edge soliton is where $\phi(x)$ interpolates between two different ground states, say, $\langle \phi(x) \rangle = 0$ and $2\pi/m$. As shown by Goldstone and Wilczek [27], such a soliton carries a charge $\Delta Q = \Delta\phi/2\pi = 1/m$, precisely the same as that of a Laughlin quasiparticle [1]. Hence the Laughlin quasiparticles are also edge solitons of a topological insulator.

D. H. L. was supported by DOE Contract No. DE-AC02-05CH11231. G. M. Z. and T. X. acknowledge the support from the NSF–China and the national program for basic research.

-
- [1] R. B. Laughlin, Phys. Rev. Lett. **50**, 1395 (1983).
 - [2] R. Jackiw and C. Rebbi, Phys. Rev. D **13**, 3398 (1976).
 - [3] W. P. Su, J. R. Schrieffer, and A. J. Heeger, Phys. Rev. B **22**, 2099 (1980).
 - [4] F. D. M. Haldane, Phys. Rev. Lett. **66**, 1529 (1991).
 - [5] G. Moore and N. Read, Nucl. Phys. **B360**, 362 (1991).
 - [6] N. B. Kopnin and M. M. Salomaa, Phys. Rev. B **44**, 9667 (1991).
 - [7] G. E. Volovik, JETP Lett. **70**, 609 (1999).
 - [8] N. Read and D. Green, Phys. Rev. B **61**, 10267 (2000).
 - [9] M. Stone and R. Roy, Phys. Rev. B **69**, 184511 (2004).
 - [10] S. Tewari, S. Das Sarma, and D.-H. Lee, Phys. Rev. Lett. **99**, 037001 (2007).
 - [11] A. Kitaev, Ann. Phys. (N.Y.) **321**, 2 (2006).
 - [12] F. D. M. Haldane, Phys. Rev. Lett. **61**, 2015 (1988).
 - [13] C. L. Kane and E. J. Mele, Phys. Rev. Lett. **95**, 226801 (2005).
 - [14] S. Murakami, N. Nagaosa, and S. C. Zhang, Science **301**, 1348 (2003).
 - [15] A. Seidel *et al.*, Phys. Rev. Lett. **95**, 266405 (2005).
 - [16] A. Seidel and D.-H. Lee, Phys. Rev. Lett. **97**, 056804 (2006).
 - [17] E. J. Bergholtz and A. Karlhede, Phys. Rev. Lett. **94**, 026802 (2005).
 - [18] E. J. Bergholtz *et al.*, Phys. Rev. B **74**, 081308 (2006).
 - [19] X.-Y. Feng, G.-M. Zhang, and T. Xiang, Phys. Rev. Lett. **98**, 087204 (2007).
 - [20] G. Baskaran, S. Mandal, and R. Shankar, Phys. Rev. Lett. **98**, 247201 (2007).
 - [21] W.-Y. Hsiang and D.-H. Lee, Phys. Rev. A **64**, 052101 (2001).
 - [22] D. J. Thouless *et al.*, Phys. Rev. Lett. **49**, 405 (1982).
 - [23] Y. Hatsugai, Phys. Rev. Lett. **71**, 3697 (1993).
 - [24] D. A. Ivanov, Phys. Rev. Lett. **86**, 268 (2001).
 - [25] C. G. Callan, Jr. and J. A. Harvey, Nucl. Phys. **B250**, 427 (1985).
 - [26] X. G. Wen, Phys. Rev. B **41**, 12838 (1990).
 - [27] J. Goldstone and F. Wilczek, Phys. Rev. Lett. **47**, 986 (1981).