

**Erratum: Interface-sensitive nuclear magnetic resonance at a semiconductor heterojunction using hyperpolarization [Phys. Rev. Materials 1, 074601 (2017)]**

Atsushi Goto, Kenjiro Hashi, Shinobu Ohki, and Tadashi Shimizu



(Received 7 February 2019; published 21 February 2019)

DOI: [10.1103/PhysRevMaterials.3.029901](https://doi.org/10.1103/PhysRevMaterials.3.029901)

There is an inconsistency between the definition of “quadrupolar splitting:  $\nu_Q$ ” and its usage in the text, corrected as follows. These corrections do not affect the overall content or the conclusions of the paper.

(1) In the fourth line from the bottom of the right-hand column on p. 1, the symbol “ $\nu_Q$ ” should be defined in the frequency unit as “ $\nu_Q = e^2 q Q / 2h$ ” rather than in the angular frequency unit “ $\nu_Q = e^2 q Q / 2\hbar$ .”

(2) The descriptions “ $\nu_Q/2$ ” that appear in the following parts of the paper should be replaced by  $\nu_Q$ .

Page 3: The horizontal axis label in Fig. 3(e).

Page 4: The fourth line of the left-hand column and the ninth and 11th lines of the right-hand column.

Page 5: The first and fifth lines of the left-hand column. The equation on the first line should also read “ $\Delta E = \nu_Q(a/\delta L_D)$ ,” removing the erroneous subscript “ $E$ ” in  $\Delta E_E$ .