

## Euclidean operator growth and quantum chaos

Alexander Avdoshkin<sup>1</sup> and Anatoly Dymarsky<sup>2,3</sup>

<sup>1</sup>*Department of Physics, University of California, Berkeley, California 94720, USA*

<sup>2</sup>*Department of Physics and Astronomy, University of Kentucky, Lexington, Kentucky 40506, USA*

<sup>3</sup>*Skolkovo Institute of Science and Technology, Moscow, Russia, 143026*



(Received 3 June 2020; accepted 9 September 2020; published 13 November 2020)

We consider growth of local operators under Euclidean time evolution in lattice systems with local interactions. We derive rigorous bounds on the operator norm growth and then proceed to establish an analog of the Lieb-Robinson bound for the spatial growth. In contrast to the Minkowski case when ballistic spreading of operators is universal, in the Euclidean case spatial growth is system-dependent and indicates if the system is integrable or chaotic. In the integrable case, the Euclidean spatial growth is at most polynomial. In the chaotic case, it is the fastest possible: exponential in 1D, while in higher dimensions and on Bethe lattices local operators can reach spatial infinity in finite Euclidean time. We use bounds on the Euclidean growth to establish constraints on individual matrix elements and operator power spectrum. We show that one-dimensional systems are special with the power spectrum always being superexponentially suppressed at large frequencies. Finally, we relate the bound on the Euclidean growth to the bound on the growth of Lanczos coefficients. To that end, we develop a path integral formalism for the weighted Dyck paths and evaluate it using saddle point approximation. Using a conjectural connection between the growth of the Lanczos coefficients and the Lyapunov exponent controlling the growth of the out-of-time-ordered correlators (OTOCs), we propose an improved bound on chaos valid at all temperatures.

DOI: [10.1103/PhysRevResearch.2.043234](https://doi.org/10.1103/PhysRevResearch.2.043234)

### I. INTRODUCTION AND RESULTS

Operator spreading, or growth, in local systems is a question of primary interest, which encodes transport properties, emergence of chaos and other aspects of many-body quantum dynamics [1–8]. A classic result of Lieb and Robinson [9] (see also Refs. [10,11] for recent progress) establishes that under time evolution the fastest possible spatial spreading of local operators is ballistic. There is no norm growth in this case since the time evolution is unitary. Ballistic spreading of operators, and signals, has been established for many models [3,12–18] and seems to be a universal feature of local systems in any dimensions. At the same time, evolution of local operators in Euclidean time

$$A(-i\beta) = e^{\beta H} A e^{-\beta H}, \quad (1)$$

which we study in this paper, is much more nuanced. Since the Euclidean evolution is not unitary, the norm of  $A(-i\beta)$  quickly grows with  $\beta$ . Moreover, as we explain below, the operator growth is not universal and reflects if the system in question is integrable or chaotic.

We start in Sec. II by deriving a bound on  $|A(t)|$  valid uniformly for  $|t| = \beta$  by expanding (1) in Taylor series

and bounding corresponding nested commutators. For local  $H$ , there is a combinatorial problem of counting contributing nested commutators, which we solve exactly for short range systems defined on Bethe lattices, which includes local systems in 1D. In higher dimensions, we conjecture an asymptotically tight bound. Hence, we expect our bounds on operator norm to be optimal in the class of Hamiltonians we consider—lattice Hamiltonians with local interactions. We find that maximal rate of growth is very different in 1D, where it is at most double-exponential, and in higher dimensions or Bethe lattices, where the norm can become infinite in finite Euclidean time. We extend the analysis to include spatial growth in Sec. III, where we find that in 1D, operators spread at most exponentially, while in higher dimensions, including Bethe lattices, they can reach spatial infinity in finite Euclidean time. When the 1D system is finite, the minimal time necessary for an operator to reach the boundary is logarithmic, which may explain logarithmic convergence of the numerical Euclidean time algorithm proposed in Ref. [19]. We further speculate in Sec. V that the timescale originating from the Euclidean Lieb-Robinson bound might be related to the Thouless energy of the corresponding quantum many-body system [20].

In Sec. IV, the results on norm growth are used to constrain individual matrix elements. We find that matrix elements in energy eigenbasis  $\langle E_i | A | E_j \rangle$  must decay at least exponentially with  $\omega = |E_i - E_j|$ , while in 1D the decay must be faster than exponential, as provided by (48) and (51). We also establish a number of bounds on the auto-correlation function at finite temperature  $C_T(t)$ , and its Fourier transform—the power

---

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

spectrum  $\Phi_T(\omega)$ ,

$$C_T(t) \equiv \text{Tr}(\rho A(t)A) = \int_{-\infty}^{\infty} \Phi_T(\omega) e^{i\omega t} d\omega,$$

$$\rho \propto e^{-H/T}, \text{Tr}(\rho) = 1. \quad (2)$$

The bounds have integral form, see (60), (61) and (63). At the physical level of rigor, they suggest that  $\Phi_T(\omega)$  decreases exponentially with  $\omega$  in  $D \geq 2$ , while in 1D the decay at large frequencies is superexponential. This emphasizes that one-dimensional systems are indeed very special, and many numerical results established for one dimensional systems may not necessarily apply to higher-dimensional systems.

The bound on  $|A(t)|$  established in Sec. II depends only on the absolute value  $|t|$ . Obviously, it is overly conservative for real  $t$  when the time evolution is unitary. We argue, however, in Sec. V, that it does correctly capture the Euclidean growth  $t = -i\beta$  of chaotic systems. We also consider system size dependence of  $|A(t)|$  and find it to be consistent with the eigenstate thermalization hypothesis (ETH). For the integrable systems, we find the growth of  $|A(t)|$  to be much slower than maximal possible, and in particular spatial growth of  $A(-i\beta)$  in this case is not exponential but polynomial.

The bound on  $|A(-i\beta)|$  can be translated into a bound on the growth of Lanczos coefficients  $b_n$ , appearing as a part of the recursion method to numerically compute  $C_T(t)$ . This is provided we assume that asymptotically  $b_n$  is a smooth function of  $n$ . To perform this calculation, we introduce a formalism of summing over weighted Dyck paths in Sec. VI and evaluate the corresponding path integral via saddle point approximation.

The obtained bound on Lanczos coefficients growth (79) is valid at all temperatures. Translating it into a bound on Lyapunov exponent of the out-of-time-ordered correlator (OTOC), we find a new bound on chaos

$$\lambda_{\text{OTOC}} \leq \frac{2\pi T}{1 + 2T\bar{\beta}(T)}, \quad (3)$$

where  $\bar{\beta}$  is such that  $C_T(t)$  is analytic inside the strip  $|\Im(t)| \leq \bar{\beta}(T)$ . For local systems, we find  $\bar{\beta}(T) \geq 2\beta^*$  with  $\beta^*$  given by (33) for all  $T$ . We illustrate this bound for SYK model in Sec. VI, see Fig. 2. We conclude with a discussion in Sec. VII.

## II. BOUND ON OPERATOR NORM GROWTH IN EUCLIDEAN TIME

Our goal in this section is to bound the infinity norm of a local operator evolved in Euclidean time

$$A(-i\beta) = e^{\beta H} A e^{-\beta H}, \quad |A(-i\beta)| \leq |A|f(\beta). \quad (4)$$

Here,  $f(\beta)$  is a bound which depends on the inverse temperature  $\beta$ , the strength of local coupling  $J$  and geometrical properties of the underline lattice model. We argue that our bound (19) (for 1D systems) and (32) (for higher dimensions) is optimal for the class of models characterized by the same strength of the local coupling constant  $J$  and lattice geome-




FIG. 1. One-dimensional lattice with short-range interactions Hamiltonian  $H = \sum_{I=1}^5 h_I$ . Local operator  $A$  sits at a third site counting from the left, between second and third bonds. Bonds highlighted in gray form a lattice animal  $I = 2, 3, 4$ .

try encoded in the Klärner's constant  $\lambda$  and animal histories constant  $\varepsilon$  which we introduce later in this work.

For simplicity, we first consider nearest-neighbor interaction Hamiltonian in 1D

$$H = \sum_{I=1}^L h_I, \quad (5)$$

where each  $h_I$  acts on sites  $I$  and  $I + 1$  and for all  $|h_I| \leq J$  for some  $J$ <sup>1</sup>. Any nearest-neighbor interaction spin chain would be an example. The operator  $A$  will be an one-site operator. An example with  $L + 1 = 6$  sites is shown in Fig. 1.

Euclidean time-evolved  $A(-i\beta)$  can be expanded in Taylor series

$$A(-i\beta) = A + \beta[H, A] + \frac{\beta^2}{2}[H, [H, A]] + \dots \quad (6)$$

Using decomposition (5) operator  $A(-i\beta)$  can be represented as a sum of nested commutators of the form

$$A(-i\beta) = A + \sum_{k=1}^{\infty} \sum_{\{I_1, \dots, I_k\}} [h_{I_k}, [\dots, [h_{I_1}, A]]] \frac{\beta^k}{k!}. \quad (7)$$

Here the sum is over all sets of indexes  $\{I_1, \dots, I_k\}$  which satisfy the following ‘‘adjacency’’ condition: first index  $I_1$  must be adjacent to the site of  $A$ ,  $I_2$  must be adjacent to the endpoints of  $I_1$  (which include the site of  $A$ ),  $I_3$  is adjacent to the endpoints of the union of  $I_1, I_2$ , etc. In other words, any subset of bonds  $I_1, I_2, \dots, I_\ell$  for  $\ell \leq k$  defines a connected cluster. Otherwise, the commutator in (7) vanishes.

A connected cluster of bonds of any particular shape is called a bond lattice animal. In 1D, all lattice animals consisting of  $j$  bonds are easy to classify: they are strings of consecutive bonds from some  $I$  to  $I + j - 1$ . In higher dimensions, the number of different bond lattice animals consisting of  $j$  bonds grows quickly with  $j$ .

Each set  $\{I_1, \dots, I_k\}$  in (7) defines a lattice animal, but the same animal may correspond to different sets. This is because indexes can repeat and appear in different orders, subject to the constraints outlined above. If we think of the set  $\{I_1, \dots, I_k\}$  as a ‘‘word’’ written in terms of ‘‘letters’’  $I_\ell$ , then corresponding lattice animal defines the alphabet.

There is a more nuanced characteristics of index sets from (7), the order in which new indexes appear. Namely, we take

<sup>1</sup>Time-evolved  $A(t)$  will not change if any of the local Hamiltonians  $h_I$  is shifted by a constant. Therefore we define  $h_I$  such that the absolute value of its largest and smallest eigenvalues are the same

a set  $\{I_1, \dots, I_k\}$  and while going from left to right remove indexes which have already appeared. In this way, we obtain a new (shorter) set which also satisfies the adjacency condition. A particular order is called “history.” For example, two sets  $\{2, 3, 2, 4, 3\}$  and  $\{3, 3, 4, 2, 4\}$  define the same lattice animal consisting of bonds  $I = 2, 3, 4$  but different histories,  $\{2, 3, 4\}$  and  $\{3, 4, 2\}$ , correspondingly, see Fig. 1.

Going back to the sum (7), to bound the infinity norm of  $A(-i\beta)$ , we can bound each nested commutator by  $(2J)^k|A|$ . Then

$$|A(-i\beta)| \leq |A|f(\beta), \tag{8}$$

$$f(\beta) = \left( 1 + \sum_{k=1}^{\infty} \sum_{\{I_1, \dots, I_k\}} \frac{(2J|\beta|)^k}{k!} \right), \tag{9}$$

and the nontrivial task is to calculate the number of sets  $\{I_1, \dots, I_k\}$  for any given  $k$ , which satisfy the adjacency condition. Evaluating sum (9) can be split into two major steps. First step is to calculate the total number  $\phi(j)$  of animal histories associated with all possible lattice animals consisting of  $j$  bonds. Second step is to calculate the sum over sets  $\{I_1, \dots, I_k\}$  associated with any given history  $\{J_1, \dots, J_j\}$ .

This last problem can be solved exactly in full generality. Let’s assume we are given a history—a set  $\{J\} = \{J_1, \dots, J_j\}$  which satisfies the adjacency condition. We want to know the number of different sets  $\{I\} = \{I_1, \dots, I_k\}$  for  $k \geq j$  satisfying the adjacency condition such that  $\{J\}$  is the history of  $\{I\}$ . We denote this number by  $S(k, j)$ . An important observation here is that any given set  $\{I\}$  defines a partition of  $\{1, 2, \dots, k\}$  into  $j$  groups labeled by elements from  $\{J\}$  by assigning each number  $1 \leq i \leq k$  to a group specified by  $I_i$ . And vice versa, each partition of  $\{1, 2, \dots, k\}$  into  $j$  groups defines a proper set  $\{I\}$  satisfying the adjacency condition. To see that we need to assign each group a unique label from  $\{J\}$ . We do it iteratively. The element 1 belongs to a group, which will be assigned the label  $J_1$ . Then we consider element 2. If it belongs to the same group labeled by  $J_1$  we move on to element 3, otherwise we assign the group it belongs label  $J_2$ . Then we consider elements 3, 4, and so on. In this way all  $j$  groups will be labeled by the unique elements from  $\{J\}$  such that the adjacency condition is satisfied.

In other words, we have established a one-to-one correspondence between the space of proper sets  $\{I\}$  for the given history  $\{J\}$  with the space of partitions of  $k$  elements into  $j$  groups. The number  $S(k, j)$  of such partitions is the Stirling numbers of the second kind which admits the following representation [21]:

$$S(k, j) = \sum_{s=1}^j \frac{(-1)^{j-s} s^{k-1}}{(j-s)!(s-1)!}. \tag{10}$$

If we introduce the number of proper sets  $\{I_1, \dots, I_k\}$  in (9) consisting of  $k$  bonds by  $\mathcal{N}(k)$ , such that

$$f(\beta) = 1 + \sum_{k=1}^{\infty} \mathcal{N}(k) \frac{(2J|\beta|)^k}{k!}, \tag{11}$$

then  $\mathcal{N}(k)$  and  $\phi(j)$  are related by the Stirling transform,

$$\mathcal{N}(k) = \sum_{j=1}^k S(k, j)\phi(j). \tag{12}$$

The inverse relation is  $\phi(j) = \sum_{k=1}^j s(j, k)\mathcal{N}(k)$ , where  $s(j, k)$  are the Stirling numbers of the first kind. From here in full generality follows [22]:

$$f(\beta) = 1 + \sum_{j=1}^{\infty} \phi(j) \frac{q^j}{j!}, \tag{13}$$

where

$$q := (e^{2|\beta|J} - 1). \tag{14}$$

We will derive this identity below

The expansion in  $q$  (13) has an obvious advantage over (11). Locality is implicit in (11), where the terms at the order  $\beta^k$  come from the lattice animals of all sizes. At the same time (13) makes locality manifest, terms at the order  $q^j$  come only from the lattice animals which have at least  $j$  bonds. This representation therefore can be used to establish Euclidean version of Lieb-Robinson bound, see Sec. III.

To evaluate (13), we still need to know the number of lattice animal histories  $\phi(j)$  for a given  $j$ . In case of 1D systems, those can be calculated exactly, while in higher dimensions we propose an asymptotically tight bound. Hence, we consider these cases separately.

### A. 1D systems

In one dimension, all lattice animals consisting of  $j$  bonds are simply the strings of  $j$  consecutive bonds. There are  $N(j) = j + 1$  such animals which include the site of the operator  $A$ . A convenient way to enumerate them is to count the number of bonds  $j_1$  and  $j_2$ ,  $j_1 + j_2 = j$ , to the left and to the right of  $A$ , respectively. For the given  $j_1, j_2$  there is, obviously, only one animal,  $N(j_1, j_2) = 1$ .

For any given  $j_1, j_2$  we denote by  $h(j_1, j_2)$  the number of histories associated with this animal, i.e., the number of different sets  $\{J\} = \{J_1, \dots, J_j\}$  such that each  $J_i$  belongs to the animal, all  $J_i$  in the set are unique and  $\{J\}$  satisfies the adjacency condition. Each history  $\{J\}$  can be completely parametrized by the order in which the cluster “grew” in left and right directions, for example histories  $\{2, 3, 4\}$  and  $\{3, 4, 2\}$  from Fig. 1 can be parametrized as “left, right, right” and “right, right, left,” correspondingly. In other words, histories with given  $j_1, j_2$  are in one to one correspondence with strings of  $j$  elements, each element being either “left” or “right,” and there are in total  $j_1$  and  $j_2$  elements of each kind. Obviously, the total number of such strings is

$$h(j_1, j_2) = \frac{(j_1 + j_2)!}{j_1! j_2!}. \tag{15}$$

Combining all ingredients together, we find the number of lattice histories for all lattice animals of size  $j$

$$\phi(j_1, j_2) = N(j_1, j_2)h(j_1, j_2) = \frac{(j_1 + j_2)!}{j_1! j_2!}, \tag{16}$$

$$\phi(j) = \sum_{j_1+j_2=j} \phi(j_1, j_2) = 2^j, \tag{17}$$

from (11) and (12), we find in full generality

$$f(\beta) = 1 + \sum_{j \geq 1} \sum_{k=j}^{\infty} \phi(j) S(k, j) \frac{(2|\beta|J)^k}{k!}. \quad (18)$$

By definition  $k \geq j$ . Crucially, expression (10) vanishes for  $1 \leq k < j$ . Therefore the sum over  $k$  can be extended to go from  $k = 1$  and can be easily evaluated,

$$f(\beta) = 1 + \sum_{j \geq 1} \sum_{s=1}^j \frac{(-1)^{j-s} \phi(j)}{(j-s)!(s-1)!} \frac{e^{2\beta J s} - 1}{s}.$$

The sum over  $s$  can be evaluated explicitly, yielding (13).<sup>2</sup> Using the explicit value of  $\phi(j)$  (17), we find

$$f(\beta) = \sum_{j=0}^{\infty} f(j, \beta) = e^{2q}, \quad f(j, \beta) = \frac{(2q)^j}{j!}. \quad (19)$$

Here,  $f(j, \beta)$  is a contribution to the bound coming from the clusters which include at least  $j$  bonds.

This result can be further refined. In (16), we introduced the number of lattice histories associated with the lattice animal which consists of  $j_1$  bonds to the left of  $A$ , and  $j_2$  bonds to the right. Repeating the summation in (18), we readily find

$$f(\beta) = \sum_{j_1, j_2 \geq 0} f(j_1, j_2, \beta), \quad f(j_1, j_2, \beta) = \frac{q^{j_1+j_2}}{j_1! j_2!}. \quad (20)$$

Here  $f(j_1, j_2, \beta)$  is the bound on the norm of the part of  $A(-i\beta)$  supported on the cluster of size  $j_1 + j_2$ . It therefore can be used to obtain the bound in the case of finite 1D lattice, or an infinite 1D lattice with a boundary.

By re-expanding (19) into Taylor series in  $\beta$ ,

$$f(\beta) = \sum_{k=0}^{\infty} \frac{B_k(2)}{k!} (2J|\beta|)^k, \quad (21)$$

where  $B_k$  are Bell polynomials, we find a bound on the norm of individual nested commutators,

$$\underbrace{|[H, [\dots, [H, A]]]}_{k \text{ commutators}}| \leq |A| B_k(2) (2J)^k. \quad (22)$$

<sup>2</sup>As a side note that evaluation of (18) in Sec. II A imply lemma 5 of Ref. [52]. Let us consider a fixed lattice animal consisting of  $j$  bonds, listed in some arbitrary order  $\{J_1, \dots, J_j\}$ . One may want to calculate  $G = \sum_{k \geq j} \sum_{\{I_1, \dots, I_k\}} (2J|\beta|)^k / k!$ , where the sum is over all sets  $\{I_1, \dots, I_k\}$ , where each  $I_i$  belongs to the set  $\{J_1, \dots, J_j\}$ , and each  $J_i$  appears in the set  $\{I_1, \dots, I_k\}$  at least once. This is a simplified version of our main calculation, with the adjacency condition being ignored. It is the sum evaluated in lemma 5 of Ref. [52]. By taking a set  $\{I_1, \dots, I_k\}$  from the sum we can associate to it a set  $\{I_1, I_2, \dots, I_j\}$  by going from the left to the right and removing repeating labels. As a set (i.e., ignoring the order)  $\{I_1, I_2, \dots, I_j\}$  coincides with  $\{J_1, \dots, J_j\}$ . The key point here is the same, the number of sets  $\{I_1, \dots, I_k\}$  associated with the same set  $\{I_1, I_2, \dots, I_j\}$  is equal to  $S(k, j)$ . If we now sum over all sets  $\{I_1, \dots, I_k\}$  associated with a particular  $\{I_1, I_2, \dots, I_j\}$ , this is exactly the sum evaluated in (18) with  $\phi(j) = 1$ . Since there are  $j!$  different permutations of labels in  $\{J_1, \dots, J_j\}$ , and thus  $j!$  sets  $\{I_1, I_2, \dots, I_j\}$  we therefore obtain  $G = q^j$

## B. Bethe lattices

The behavior of  $f(\beta)$  differs drastically in one and higher dimensions. To better understand this difference we consider an “intermediate” scenario of a short range Hamiltonian define on a Bethe lattice of coordination number  $z$  [23]. Namely, we assume that each  $h_I$  from (5) “lives” on a bond and acts on the Hilbert spaces associated with two vertexes adjacent to that bond. For any finite  $k$  in the Taylor series expansion (7), only finite number of bonds are involved and the corresponding lattice animals (clusters) live on the Cayley tree. Thus, similarly to 1D, there are no loops, but the total number of lattice animals consisting of  $j$  bonds grows exponentially,  $N(j) \sim \lambda(z)^j$ ,

$$\ln \lambda(z) = (z-1) \ln(z-1) - (z-2) \ln(z-2). \quad (23)$$

This exponential growth is typical for lattices in higher dimensions  $D > 1$ .

The total number of lattice animal histories  $\phi(j)$  can be calculated exactly in this case (see Appendix A),

$$\phi(j) = (z-2)^j \frac{\Gamma(j+z/(z-2))}{\Gamma(z/(z-2))}, \quad (24)$$

leading to the bound

$$f(\beta) = (1 - (z-2)q)^{-z/(z-2)}. \quad (25)$$

In other words, the total number of histories  $\phi(j)$  grows as a factorial. The same qualitative behavior applies for all higher-dimensional lattices.

As a final remark, we notice that taking  $z \rightarrow 2$  in (25) yields  $f(\beta) = e^{2q}$ , in full agreement with (19).

## C. Higher-dimensional systems

The calculations of previous sections can, in principle, be extended to an arbitrary lattice system, but the number of lattice animal histories is difficult to evaluate exactly. Nevertheless it is known that the number of different lattice animals  $N(j)$  consisting of  $j$  bonds (which include a particular site) grows rapidly in higher dimensions. While the exact formula is not known, the asymptotic growth is known to be exponential, and is controlled by the so-called Klarner’s constant  $\lambda$ ,

$$N(j) \sim \lambda^j. \quad (26)$$

By introducing a sufficiently large but  $j$ -independent constant  $C$  we can uniformly bound the number of lattice animals consisting of  $j$  bonds by<sup>3</sup>

$$N(j) \leq C \lambda^j. \quad (27)$$

The number of histories for any given animal is the number of different sets  $\{J_1, \dots, J_j\}$  where all indexes are distinct, subject to the adjacency condition. Let us denote by  $h(j)$  the average number of histories for all animals consisting of  $j$  bonds. Then, it is trivially bounded by  $h(j) \leq j!$ . It can be

<sup>3</sup>To account for a polynomial pre-exponential factor, coefficient  $\lambda$  in (27) may need to be taken strictly larger than the Klarner’s constant  $\lambda$  in (26).

shown that for sufficiently large  $j$  [24]

$$h(j) \geq \frac{j!}{a^j}, \tag{28}$$

for some  $a > 1$ . We, therefore, conjecture that for higher-dimensional lattices  $h(j)$  is uniformly bounded by

$$h(j) \leq C' \frac{j!}{\varepsilon^j}, \tag{29}$$

for some  $\varepsilon > 1$  and a  $j$ -independent constant  $C' \geq 1$ . This bound is trivially satisfied for  $\varepsilon = 1$ . The nontrivial part here is the expectation that (29) correctly captures the leading (exponential) asymptotic behavior of  $h(j)$  with some  $\varepsilon > 1$ , i.e., (29) is the optimal bound which can not be further improved (excluding polynomial pre-factors). We therefore introduce here the constant  $\varepsilon$  which we call *animal histories* constant and conjecture that it is strictly larger than 1. In the end of this section, we also derive a lower bound on  $\varepsilon/\lambda$ . By combining (27) together with (29)

$$\phi(j) = N(j)h(j) \leq C'(\lambda/\varepsilon)^j j!, \tag{30}$$

we find the bound

$$f(\beta) = \sum_{j=0}^{\infty} f(j, \beta), \quad f(j, \beta) = C'(q/q_0)^j.$$

Here  $f(\beta)$  is defined to be larger than the sum in (9). The coefficient

$$q_0 = \frac{\varepsilon}{\lambda} \tag{31}$$

characterizes lattice geometry. Unlike in 1D, where (19) has an additional factorial suppression factor,  $f(j, \beta)$  in higher dimensions grows exponentially for sufficiently large  $\beta$ . Summing over  $j$  yields

$$f(\beta) = \frac{C'}{1 - q/q_0}. \tag{32}$$

In contrast to 1D, while (19) is finite for all  $\beta$ , (32) is finite only for

$$|\beta| < \beta^* \equiv \ln(1 + q_0)/(2J). \tag{33}$$

While (32) is only a bound on  $f(\beta)$  defined in (9), location of the singularity in both cases is the same because it is only sensitive to the asymptotic behavior of  $N(j)$  and  $h(j)$ .

Expanding (32) in Taylor series

$$f(\beta) = C' \sum_{k=0}^{\infty} \frac{P_k(q_0^{-1})}{k!} (2J|\beta|)^k, \tag{34}$$

where  $P_k$  are the polynomials defined via

$$P_k(x) = \frac{1}{1+x} \left( x(1+x) \frac{\partial}{\partial x} \right)^k (1+x) \tag{35}$$

yields a bound on individual nested commutators

$$|\underbrace{[H, [\dots, [H, A]]]}_{k \text{ commutators}}| \leq |A| C' P_k(q_0^{-1}) (2J)^k. \tag{36}$$

The divergence of bound (32) at  $|\beta| = \beta^*$  is not an artifact of an overly conservative counting, as confirmed by a 2D model introduced in Ref. [24], for which  $|A(-i\beta)|$  is known

to diverge. We will argue in Sec. V that the growth outlined by the bounds (19), (32) reflects actual growth of  $|A(-i\beta)|$  in nonintegrable systems and singularity of (32) at finite  $\beta$  is a sign of chaos. We also note that in case of 1D systems the bound (19) ensures that the operator norm of  $A(t)$  remains bounded for any complex  $t$ . This is consistent with analyticity of correlation functions in 1D [25]. On the contrary, in higher dimensions, physical observables may not be analytic. We discuss the relation between the singularity of  $|A(-i\beta)|$  and nonanalyticity of physical observables due to a phase transition in Sec. V and show that they have different origin.

It is interesting to compare our result for a general lattice in  $D > 2$  with the exact result for Bethe lattices obtained in the previous section. From (24) and (30), we obtain lattices animal histories constant  $\varepsilon$  for Bethe lattices,

$$\varepsilon = \left( \frac{z-1}{z-2} \right)^{z-1}, \quad q_0 = \frac{\varepsilon}{\lambda} = \frac{1}{z-2}. \tag{37}$$

For any  $z \geq 2$ ,  $\varepsilon > 1$  supporting our conjecture that  $\varepsilon$  is always strictly larger than 1. Our universal expression (32) bounds the exact result (25) from above with any  $q_0 < 1/(z-2)$  and sufficiently large  $C'$ .

Bethe lattices provide a lower bound on the combination  $q_0 = \frac{\varepsilon}{\lambda}$  and hence on the critical value  $\beta^*$ . We show in Appendix B that for any lattice of coordination number  $z$ , such that each vertex is attached to at most  $z$  bonds the number of lattice animal histories is bounded by  $\phi(j) \leq (z-2)^j \frac{\Gamma(j+z/(z-2))}{\Gamma(z/(z-2))}$ . We therefore find in full generality

$$q_0 = \frac{\varepsilon}{\lambda} \geq \frac{1}{z-2}. \tag{38}$$

This bound is stronger than any previously known, as we explain below.

To conclude this section, we demonstrate the advantage of counting lattice animal histories as is done in (13) over previously explored approaches. There is a straightforward way to estimate the number of sets  $I_1, \dots, I_k$  in (9) from above by counting the number of ways a new bond can be added to the set at each step. Provided the lattice has coordination number  $z$ , starting from the site of  $A$ , there are  $z$  ways to choose  $I_1$ , at most  $z(2z)$  ways to choose  $I_2$ ,  $z(2z)(3z)$  ways to choose  $I_3$  and so on. As a result we would get an estimate for  $f(\beta)$ ,

$$f(\beta) \leq f_{\text{approx}} = \sum_{k=0}^{\infty} (2J|\beta|)^k z^k = \frac{1}{1 - 2J|\beta|z}. \tag{39}$$

This result was previously obtained in Refs. [26,27]. This gives the following estimate for the location of the pole:

$$|\beta| = \frac{z^{-1}}{2J}. \tag{40}$$

The approximation (40) is naive as it overcounts the number of sets  $\{I_1, \dots, I_k\}$  assuming the underlying cluster is always of size  $k$ . We therefore expect (39) to be weaker than our (32),  $f(\beta) \leq f_{\text{approx}}(\beta)$ , and in particular the location of the singularity (40) to be smaller than  $\beta^*$  defined in (33). This can be written as an inequality

$$\varepsilon/\lambda \geq e^{1/z} - 1, \tag{41}$$

which is indeed satisfied due to (38). The advantage of (38) becomes apparent in the limit  $z \rightarrow 2$  when  $\beta^*$  becomes infinite while (40) remains finite.

A result analogous to (32) has been previously established in Ref. [28], but importantly there  $q_0$  was just inverse of the lattice animal constant, i.e., Klarnar's constant introduced in previous section,  $q_0 = \lambda^{-1}$ . Crucially, we improve this result to account for proper lattice animal histories by introducing  $\varepsilon > 1$  in (31). Without  $\varepsilon$  critical value of  $\beta$  where  $f(\beta)$  diverges is given by  $q_0 = e^{2J\beta} - 1 = \lambda^{-1}$  and, e.g., for a cubic lattice in  $D$  dimensions  $\lambda$  asymptotes to  $2De$  when  $D \rightarrow \infty$  [28,29]. This value is smaller than (40) with  $z = 2D$ , meaning the inequality (41) is not satisfied. To conclude, without taking lattice animal histories into account, even exact value of  $\lambda$  results in a less stringent bound than (40), while our bound is always stronger than that due to (38).

### III. SPATIAL GROWTH IN EUCLIDEAN TIME

While deriving the bound on the norm of local operators evolved in Euclidean time, (19) and (32), we obtained a stronger result—a bound  $f(j, \beta)$  [or  $f(j_1, j_2, \beta)$  in 1D] on spatial growth of  $A(-i\beta)$ . It can be immediately translated into the Euclidean analog of the Lieb-Robinson bound [9] on the norm of the commutator of two spatially separated local operators. If  $B$  is an operator with finite support located distance  $\ell$  away from  $A$  (measured in the Manhattan norm in case of a cubic lattice), then in  $D \geq 2$ ,

$$|[A(i\beta), B]| \leq 2|A||B| \sum_{j=\ell}^{\infty} f(j, \beta) = 2|A||B| \frac{C'(q/q_0)^\ell}{1 - (q/q_0)},$$

where we assumed that  $|\beta| < \beta^*$ . For larger  $|\beta|$  there is no bound as the sum does not converge. This result means that the local operator can spread to the whole system, no matter how large or even infinite that is, in finite Euclidean time  $\beta = \beta^*$ . We will argue in Sec. V that this is the true physical behavior in the chaotic case and therefore the bound can not be improved to get rid of the divergence at  $|\beta| = \beta^*$  in full generality.

In 1D, the situation is very different. Assuming local operator  $B$  is located  $\ell$  bonds away from  $A$ , we find

$$\begin{aligned} |[A(i\beta), B]| &\leq 2|A||B| \sum_{j_1=0, j_2=\ell}^{\infty} f(j_1, j_2, \beta) \\ &= 2|A||B| \frac{e^{2q}}{(\ell-1)!} \int_0^q e^{-t} t^{\ell-1} dt. \end{aligned} \quad (42)$$

(If the system is infinite only in one direction and  $A$  is sitting at the boundary, one factor of  $e^q$  should be removed.) Qualitatively the RHS of (42) behaves as

$$|[A(-i\beta), B]| \leq 2|A||B| \frac{q^\ell}{\ell!} e^q, \quad (43)$$

for  $\ell \gg q + 1$ , and asymptotes to  $2|A||B|e^{2q}$  for  $\ell \ll q + 1$ . This means a local operator spreads exponentially fast, to distances  $\ell \sim e^{2J\beta}$ , in Euclidean time  $\beta$ .

Exponential spreading of operators in 1D seems to be in agreement with the convergence of the Euclidean variational algorithm of Ref. [19] in logarithmic time. The connection

between Euclidean Lieb-Robinson bound and the convergence time is intuitive, but difficult to establish rigorously, in particular, because the latter is sensitive to the choice of initial wave-function. For the integrable models, for which the spreading of operators is at most polynomial (see Sec. V), convergence time might be even shorter because of a well-tuned initial wave function. For the chaotic systems we expect no fine-tuning of the initial state and hence a direct relation between the convergence time and Euclidean Lieb-Robinson bound.

Another possibly intriguing connection is with the studies of Thouless times in chaotic Floquet systems without conserved quantities [20]. There, it was noticed that in 1D Thouless time is logarithmic in system size (see also Ref. [30]), and finite in  $D \geq 2$  (see, however, Ref. [31]). That is exactly the same behavior as in the case of Euclidean operator spreading. One potential interpretation would be that Thouless time can be associated with the slowest Euclidean mode propagating in the system. Under Euclidean time evolution with a time-dependent random Hamiltonian our extension of Lieb-Robinson bound holds. We also surmise that in this case spatial growth of all quantities, including the slowest, is qualitatively and outlined by the bound with some effective  $J, q_0$ . When the system in question has a local conserved quantity, the slowest transport mode is diffusive, leading to  $L^2$  scaling of Thouless time [32]. Thus, to complete this picture it would be necessary to establish that under Euclidean time evolution time necessary for a diffusive mode to travel across the system is the same as in the Minkowski case, i.e.,  $\beta \sim L^2$ , where  $L$  is the system size.

Finally, we notice that the Euclidean analog of the Lieb-Robinson bound in 1D (42) looks similar to the conventional Minkowski bound [11]

$$|[A(t), B]| \leq 2|A||B| \frac{(2Jt)^\ell}{\ell!}, \quad (44)$$

with  $2Jt$  substituted by  $q(\beta)$ .

### IV. CONSTRAINTS ON MATRIX ELEMENTS

#### A. Individual matrix elements

Constraints on the infinity-norm of  $A(i\beta)^\dagger = A(-i\beta)$  provide an upper bound on the magnitude of matrix elements  $A_{ij} = \langle E_i | A | E_j \rangle$  in the energy eigenbasis. Starting from

$$A(-i\beta)_{ij} \equiv \langle E_i | A(-i\beta) | E_j \rangle = A_{ij} e^{\beta(E_i - E_j)}, \quad (45)$$

we find

$$|A_{ij}| \leq e^{-\beta(E_i - E_j)} |A(-i\beta)|. \quad (46)$$

This inequality holds for any  $\beta$  and we therefore can optimize it over  $\beta$ . Using explicit form of the bound (19) in 1D, we find optimal value of  $\beta$  to be (without loss of generality we assumed  $\omega = E_i - E_j \geq 0$ )

$$\beta = \begin{cases} \ln(\frac{\omega}{4J}) / (2J), & \omega \geq 4J, \\ 0, & 4J \geq \omega. \end{cases} \quad (47)$$

This yields

$$|A_{ij}| \leq |A| \kappa(\omega), \quad \omega = |E_i - E_j|, \quad (48)$$

where

$$\kappa(\omega) \equiv \begin{cases} \exp\{2\tilde{\omega}(1 - \ln \tilde{\omega}) - 2\}, & \tilde{\omega} = \omega/(4J) \geq 1, \\ 1, & \tilde{\omega} \leq 1. \end{cases}$$

These results shows that in 1D for large energy difference  $\omega = |E_i - E_j| \gg J$  off-diagonal matrix elements  $A_{ij}$  decay faster than exponential. For  $\omega \leq 4J$ , the bound trivializes to  $|A_{ij}| \leq |A|$ .

In higher dimensions, the bound on  $A_{ij}$  from (46) can not be better than exponential. This is because  $f(\beta)$  is a monotonically increasing function of  $\beta$ , which diverges for some  $|\beta| = \beta^*$ . In particular

$$e^{-\beta\omega} |A(-i\beta)| \geq e^{-\beta^*\omega} |A| \tag{49}$$

for any  $\beta$  and  $\omega \geq 0$ . To find leading exponent, we optimize (46) over  $\beta$  to find

$$\beta = \frac{\ln\left(\frac{\omega(1+q_0^{-1})}{2(\omega+J)}\right)}{2J}, \tag{50}$$

and  $|A_{ij}| \leq |A|\kappa(\omega)$ , where  $\omega = |E_i - E_j|$ ,

$$\kappa(\omega) = C' q_0^{-1} \tilde{\omega} \left(\frac{\tilde{\omega}(1+q_0^{-1})}{1+\tilde{\omega}}\right)^{-1-\tilde{\omega}}, \quad \tilde{\omega} = \omega/(2J).$$

Taking  $\omega \rightarrow \infty$  limit, we find that the asymptotic exponential behavior is given by (49)

$$\kappa(\omega) \lesssim C'' \omega e^{-\beta^*\omega}, \quad \omega \gg J, \tag{51}$$

where  $C''$  is some  $\omega$ -independent constant.

Constraints on individual matrix elements (49) and (49) only depend on energy difference  $\omega$ . In the case when the system satisfies ETH, off-diagonal matrix elements for  $i \neq j$  are known to be exponentially suppressed by the entropy factor,  $|A_{ij}|^2 \sim e^{-S}$ . Therefore, for the chaotic systems, the bound will be trivially satisfied unless  $\omega$  is extensive.

The bound analogous to (51) has previously appeared in Ref. [28], with  $\beta^*$  given by (33) with  $\varepsilon = 1$ .

**B. Constraints on power spectrum**

Bounds on individual matrix elements found above can be extended to the autocorrelation function of a Hermitian local  $A$ ,

$$C(t) \equiv \text{Tr}(\rho A(t)A), \tag{52}$$

and its power spectrum

$$\begin{aligned} \Phi(\omega) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} dt e^{-i\omega t} C(t) \\ &= \sum_{i,j} p_i |A_{ij}|^2 \delta(E_i - E_j - \omega). \end{aligned} \tag{53}$$

Here  $\rho$  is an arbitrary density matrix which commutes with the Hamiltonian,  $\rho = \sum_i p_i |E_i\rangle\langle E_i|$ ,  $\text{Tr} \rho = 1$ .

Although bounds on moments  $M_k$  derived below are universal for all  $\rho$ , in what follows, we will be most interested in the case when  $\rho$  is the Gibbs ensemble  $\rho = e^{-H/T}/Z$ , in which case autocorrelation function and power spectrum will

be denotes by  $C_T$  and  $\Phi_T$ , correspondingly. As a function of complex argument  $C_T$  satisfies

$$C_T(t - i/(2T)) = C_T(-t - i/(2T)), \tag{54}$$

$$C_T(t^*) = (C_T(-t))^*. \tag{55}$$

First we notice that

$$|C(t)| \leq |A(t/2)|^2 \leq |A|^2 f^2(|t|/2), \tag{56}$$

for any complex  $t$ , which guarantees analyticity of  $C(t)$  for 1D systems on the entire complex plane.

Using the bound on individual nested commutators (22) and (36) one can bound the growth of Taylor coefficients of  $C$ ,

$$M_k = \int_{-\infty}^{\infty} \Phi(\omega) \omega^k d\omega = \text{Tr}(\rho \underbrace{[H, [\dots, [H, A]]]}_{k \text{ commutators}} A). \tag{57}$$

To obtain an optimal bound, nested commutators should be split equally between two  $A$ 's using cyclicity of trace

$$|M_{2k+i}| \leq |A|^2 (2J)^{2k+i} R_k R_{k+i}, \quad i = 0, 1. \tag{58}$$

Here  $R_k = B_k(2)$  for infinite 1D system,  $R_k = B_k$  for semi-infinite 1D system with a boundary, and  $R_k = C' P_k(q_0^{-1})$  for  $D \geq 2$ .

Using the asymptotic behavior of Bell polynomials [33]

$$B_n(x) \sim \left(\frac{n(1+o(1))}{e \ln(n/x)}\right)^n, \quad n \gg x, \tag{59}$$

and the Stirling approximation formula, the bound on moments for  $k \gg 1$  can be rewritten as (for the infinite 1D system)

$$|M_k| \leq |A|^2 (2J)^k \left(\frac{k}{2e \ln k}\right)^k \times e^{o(k)}. \tag{60}$$

It is easy to see that the Taylor series of  $C_T(t)$  converges in the whole complex plane, as was pointed out above.

In higher dimensions, to find asymptotic behavior of  $P_k(x)$  for large  $k$ , we use the following representation:

$$P_k(x) = \sum_{j=1}^n j! S(k, j) x^j = \frac{1}{1+x} \sum_{j=1}^{\infty} j^k \left(\frac{x}{1+x}\right)^j.$$

Substituting the sum over  $j$  by an integral and taking saddle point approximation gives

$$|M_k| \leq |A|^2 \left(\frac{q_0}{1+q_0}\right)^2 \left(\frac{k}{2e\beta^*}\right)^k \times e^{o(k)}. \tag{61}$$

Focusing on the case when  $\rho = e^{-H/T}/Z$ , (61) guarantees that Taylor series of  $C_T(t)$  converges absolutely inside the disk  $|t| \leq 2\beta^*$ . By representing  $C_T$  as a sum over individual matrix elements it is easy to see that if the sum for  $C_T(-i\beta)$  is absolutely convergent, then it is absolutely convergent for any  $C_T(t)$ ,  $\Im(t) = -i\beta$ . Therefore  $C_T(t)$  is analytic inside the strip  $2\beta^* > \Im(t) > -2\beta^*$ . Because of reflection symmetry (54) function  $C_T(t)$  must be analytic inside a wider strip

$2\beta^* > \Im(t) > -2\beta^* - 1/T$ <sup>4</sup>. Hence symmetrically ordered autocorrelation function

$$C_T^W(t) \equiv \text{Tr}(\rho^{1/2} A(t) \rho^{1/2} A) = C_T(t - i/(2T)), \quad (62)$$

is analytic inside the strip  $2\beta^* + 1/(2T) > \Im(t) > -2\beta^* - 1/(2T)$ , which is wider than the strip of analyticity of  $C_T(t)$ , and indicates a more rapid exponential decay of the power spectrum  $\Phi_T^W$  of (62) in comparison with  $\Phi_T(\omega)$ .

The logic above is general and does not require any specific details of  $M_k$ . Using reflection symmetry (54), we have shown in full generality that if  $C_T(t)$  develops a singularity at  $t = \pm i2\beta^*$ , then  $C_T^W(t)$  is analytic at least inside the strip  $|\Im(t)| \leq 2\beta^* + 1/(2T)$ .

There is another integral bound on power spectrum, valid for any density matrix  $\rho$  which commutes with  $H$ . By integrating (53), we find the following inequality:

$$\begin{aligned} \int_{\omega}^{\infty} d\omega' \Phi(\omega') &\equiv \sum_{E_i \geq E_j + \omega} p_i |A_{ij}|^2 \\ &= \sum_{E_i \geq E_j + \omega} p_i e^{-2\beta(E_i - E_j)} |A(-i\beta)_{ij}|^2 \\ &\leq \sum_{E_i \geq E_j + \omega} p_i e^{-2\beta\omega} |A(-i\beta)_{ij}|^2 \\ &\leq e^{-2\beta\omega} \sum_{i,j} p_i |A(-i\beta)_{ij}|^2 \\ &= e^{-2\beta\omega} \text{Tr}(\rho A(-i\beta) A(i\beta)) \leq e^{-2\beta\omega} |A(-i\beta)|^2. \end{aligned}$$

Here  $\omega$  is non-negative and in the second equality we used (45) with an arbitrary positive  $\beta$ . Now we can use  $|A(-i\beta)| \leq |A|f(\beta)$  and optimize over  $\beta$ , yielding

$$\int_{\omega}^{\infty} d\omega' \Phi(\omega') \leq |A|^2 \kappa(\omega)^2. \quad (63)$$

Function  $\kappa$  is given by (49) and (51) in  $D = 1$  and  $D \geq 2$ , correspondingly.

When  $\rho$  is maximally mixed state, i.e., temperature  $T$  is infinite, the bound can be strengthened to

$$\int_{|\omega'| \geq \omega} d\omega' \Phi(\omega') \leq |A|^2 \kappa(\omega)^2. \quad (64)$$

We would like to emphasize that all bounds discussed above, i.e., bounds on  $M_k$  and (63), are integral in form. We do not know a rigorous way to directly constrain asymptotic behavior of  $\Phi(\omega)$ . At the same time at physical level of rigor, if we assume that  $\Phi(\omega)$  is a smoothly behaving function at large  $\omega$ , analyticity of  $C(t)$  inside the strip  $|\Im(t)| < 2\beta^*$

<sup>4</sup>If  $1/(2T) \leq 2\beta^*$ , a union of an original strip  $|\Im(t)| < 2\beta^*$  and its reflection around the point  $\beta = -1/(2T)$  is a wider strip  $2\beta^* > \Im(t) > -2\beta^* - 1/T$ . Function  $C_T(t)$  has to be analytic there. If  $1/(2T) > 2\beta^*$  the same union consists of two strips,  $2\beta^* > \Im(t) > -2\beta^*$  and  $2\beta^* - 1/T > \Im(t) > -2\beta^* - 1/T$ . It is easy to show though that  $C_T$  has to be analytic also in between,  $-2\beta^* > \Im(t) > 2\beta^* - 1/T$ . From the definition  $C_T(t) = \text{Tr}(\rho^a A \rho^b A)$ ,  $a = it + 1/T$ ,  $b = it$ , and positivity  $\Re(a), \Re(b) > 0$  it follows that the sum over Hilbert space converges,  $C_T$  is well defined and therefore analytic.

immediately implies that power spectrum in  $D \geq 2$  is exponentially suppressed by

$$|\Phi(\omega)| \lesssim |A|^2 e^{-2\beta^*\omega}, \quad \omega \rightarrow \infty. \quad (65)$$

In 1D, we similarly find superexponential suppression

$$|\Phi(\omega)| \lesssim |A|^2 e^{-\omega(1 + \ln(4J/\omega))/J}, \quad \omega \rightarrow \infty. \quad (66)$$

The bound on moments for large  $k$  (60), (61) and the integral bound (63) for large  $\omega$  follow from here via saddle point approximation.

Superexponential suppression of  $\Phi(\omega)$  emphasizes peculiarity of one-dimensional systems. In particular, it implies that high frequency conductivity [34] and energy absorption [26] for such systems will be superexponentially suppressed. This is a very special behavior, which should be kept in mind in light of the numerical studies, which are often limited to one dimensions, and therefore may not capture correct physical behavior.

An exponential bound on the integral of  $\Phi(\omega)$  was first established in Ref. [26], where the authors also noted superexponential suppression in 1D, albeit without proposing an explicit analytic form.

## V. FINITE SIZE SCALING AND CHAOS

The bounds obtained in Sec. II correctly account for the number of nontrivial nested commutators  $[h_k, [\dots, [h_1, A]]]$  but do not take into account peculiarities of individual local Hamiltonians  $h_l$ . We therefore expect our bound to be strongest possible among the uniform bounds for the entire family of local short-ranged Hamiltonians defined on a particular lattice. We further assumed that each nested commutator is equal to its maximal possible value  $(2J)^k |A|$ . This is certainly too conservative, but for the chaotic systems, i.e. in absence of some additional symmetries, we expect a finite fraction of nested commutators to grow as a power of  $k$ . We therefore expect that for large  $\beta$  our bounds (19) and (32) to correctly describe growth of operator norm in local chaotic systems with some effective values of  $J$ , as it happens in Ref. [24]. In particular, in one dimension, we expect  $|A(-i\beta)|$  to grow double-exponentially, and in higher dimensions we expect  $|A(-i\beta)|$  to diverge at some finite  $\beta^*$ .

We similarly expect the bound on spatial growth outlined in Sec. III to correctly capture the spread of local operators when the system is chaotic. An indirect evidence to support that comes from the numerical results of Ref. [19], i.e., logarithmic convergence time of a numerical Euclidean time algorithm, in agreement with (43).

Below we further outline how  $|A(-i\beta)|$  reflects chaos of the underlying system when the system size is finite. It follows from (13) that for large  $\beta$ , animal histories with the largest number of bonds will dominate,

$$f(\beta) \propto q^j \sim e^{2^j j |\beta|}, \quad (67)$$

where  $j$  is the total number of bonds in the system, i.e.,  $j$  is proportional to the volume. Let us compare this behavior with the growth of the Frobenius norm,

$$C(-i\beta) = \frac{\text{Tr}(A(-i\beta)A)}{\text{Tr}(1)} = \sum_{ij} e^{\beta(E_i - E_j)} \frac{|A_{ij}|^2}{\text{Tr}(1)}. \quad (68)$$



At large  $\beta$ , leading behavior is

$$C(-i\beta) \propto e^{\beta\Delta E}, \tag{69}$$

where  $\Delta E$  is the maximal value of  $\Delta E = E_i - E_j$  such that corresponding matrix element  $A_{ij}$  is not zero. [In other words,  $\Delta E$  is the support of  $\Phi(\omega)$ .] For the chaotic systems satisfying eigenstate thermalization hypothesis, we expect most matrix elements to be nonzero, even for extensive  $\Delta E$ , matching extensive behavior of  $2Jj$  in (67).

Assuming qualitative behavior of (32) is correct for non-integrable systems, going back to thermodynamic limit in  $D \geq 2$ , we expect a singularity of  $|A(-i\beta)|$  and  $C(-2i\beta)$  at some finite  $\beta$ . This singularity has a clear interpretation in terms of  $A$  spreading in the operator space. We first interpret  $(A|B) := \text{Tr}(A^\dagger B)/\text{Tr}(1)$  as a scalar product in the space of all operators and denote corresponding Frobenius norm of  $A$  by  $|A|_F \equiv (|A|)^{1/2}$ . Then if  $A$  were typical, i.e., random in the space of all operators,

$$C(-i\beta) = \text{Tr}(A(-i\beta)A) = |A(-i\beta/2)|_F^2 \frac{Z(\beta)Z(-\beta)}{Z(0)^2},$$

$$Z(\beta) \equiv \text{Tr} e^{-\beta H}.$$

Euclidean time evolution can be split into two parts,  $A(-i(\beta + \beta')) = e^{\beta'H}A(-i\beta)e^{-\beta'H}$  such that

$$C(-i(\beta + \beta')) = (A(-i\beta/2)|e^{i\beta'\text{adj}_H}|A(-i\beta/2)). \tag{70}$$

At time  $\beta = 0$ , we start with a local operator, which is not typical. In principle,  $A(-i\beta/2)$  only explores a particular trajectory in the space of all operators, and therefore can not be fully typical at any  $\beta$ . Yet, if we assume that by the time  $\beta$  the trajectory of  $A$  has explored substantial part of operator space such that  $A(-i\beta/2)$  can be considered typical enough, we obtain

$$C(-i(\beta + \beta')) \approx C(-i\beta) \frac{Z(\beta')Z(-\beta')}{Z(0)^2}. \tag{71}$$

Taking into account that free energy  $\ln(Z)$  is extensive, we immediately see that (71) diverges for any  $\beta' > 0$ . Hence, the singularity of  $C(-i\beta)$  and thus also of  $|A(-i\beta/2)|$  marks the moment when  $A(-i\beta/2)$  becomes typical. This picture is further developed in Ref. [35], where we show that the singularity of  $|A(-i\beta/2)|$  can be associated with delocalization of  $A$  in the Krylov space.

It is interesting to note that since  $C(t)$  is analytic for local one-dimensional systems, for such systems, even nonintegrable,  $A$  never becomes typical and hence these systems can not be regarded as fully chaotic.

We separately remark that the conventional time evolution  $C(t) = (A(t/2)|A(-t/2))$  does not have an interpretation as the Frobenius norm-squared of  $A(t/2)$ , therefore (70) does not apply and even if  $A(t/2)$  becomes sufficiently typical at late  $t$ , the analog of (71) may not hold.

If the system is finite, at large  $\beta$  free energy simply becomes  $\ln Z(\beta)/Z(0) \sim -\beta E_m$ , where  $E_m$  is extensive (minimal or maximal) energy of the system. Hence (71) will be proportional to  $e^{\beta'\Delta E}$ , where  $\Delta E$  is extensive, in full agreement with (69). This gives the following qualitative behavior of  $C(-i\beta)$  when the chaotic system is sufficiently large but finite. For small  $\beta$ ,  $\ln C(-i\beta)$  will behave as  $\propto e^q$  in 1D

and  $\propto \ln(q_0 - q)$  in higher dimensions. This growth will stop at  $\beta \sim \ln(L)$  in 1D or  $\beta \sim \beta^*$  in  $D \geq 2$ , at which point in both cases  $\ln C(-i\beta)$  will be extensive. At later times  $\ln C(-i\beta)$  will grow as  $\beta\Delta E$  with some extensive  $\Delta E$ . In the nonintegrable case, the transition between two regimes, “thermodynamic” when  $C(-i\beta)$  has not yet been affected by the finite system size, and “asymptotic,” is very quick, at most double-logarithmic in 1D.

Behavior of chaotic systems described above should be contrasted with integrable models. In this case, most matrix elements  $A_{ij}$  are zero and for a wide class of systems, including classical spin models and systems with projector Hamiltonians, support of  $\Phi(\omega)$  remains bounded in the thermodynamic limit. (In terms of the Lanczos coefficients, introduced in the next section, this is the case of  $\lambda = 0$ .) For such systems the bounds (19) and (32) will be overly conservative. For sufficiently large systems and large  $\beta$ , we expect (69) with a system size independent  $\Delta E$ . This asymptotic behavior will emerge in finite system-independent Euclidean time. Infinity norm  $|A(-i\beta)|$  will behave similarly. We further can use (69) to estimate the Frobenius norm of nested commutators  $|\underbrace{[H, [\dots, [H, A]]]}_{k \text{ commutators}}|_F \leq |A|\Delta E^k$ . Assuming infinity and

Frobenius norms exhibit qualitatively similar behavior, we can substantially improve the Euclidean analog of the Lieb-Robinson bound

$$|[A(-i\beta), B]| \leq 2|A||B| \sum_{k=\ell}^{\infty} \frac{\beta^k \Delta E^k}{k!} \sim 2|A||B| \frac{\beta^\ell \Delta E^\ell}{\ell!},$$

where last step assumes  $\beta\Delta E \ll \ell$ . This bound has the same structure as the conventional Lieb-Robinson bound in Minkowski space (44). Thus, in the case of noninteracting models or projector Hamiltonians ( $\lambda = 0$  in the language of next section), we find ballistic spreading of operators for any complex  $t$ .

In the case of a general integrable model, the support of  $\Phi(\omega)$  is extensive and the behavior is more intricate. In many explicit examples in the thermodynamic limit  $\Phi(\omega)$  decays as a Gaussian, and  $C(-i\beta) \propto e^{(J\beta)^2}$  with some appropriate local coupling  $J$  [36–40]. (This is the case of  $\lambda = 1$  in terms of the next section. See Appendix C where we derive the Gaussian behavior starting from  $\lambda = 1$ .) Using the same logic as above, this leads to the Euclidean Lieb-Robinson bound of the form

$$|[A(-i\beta), B]| \lesssim 2|A||B| \frac{(\beta J)^{2\ell}}{\ell!}, \tag{72}$$

which indicates a polynomial propagation of the signal  $\ell \propto \beta^2$ . For a finite system of linear size  $L$ , we may expect Gaussian behavior  $C(-i\beta) \propto e^{(J\beta)^2}$  up to the times  $\beta \propto L^{1/2}$ , after which the asymptotic behavior (69) should emerge. Although the model is integrable,  $\Delta E$  is extensive, which implies the transition between “thermodynamic” and “asymptotic” behavior is long and will take up to  $\beta \sim L$ . This indicates the qualitative difference between integrable and nonintegrable (chaotic) models. When the system is finite in both cases the asymptotic behavior is given  $C(-i\beta) \propto e^{\beta\Delta E}$  with an extensive  $\Delta E$  (except for the  $\lambda = 0$  case), but asymptotic behavior will emerge quickly, in finite (for  $D \geq 2$ ) or logarithmic (for  $D = 1$ ) times in the nonintegrable case, while in the inte-

grable case asymptotic behavior will emerge much slower, after polynomial times in  $L$ .

A qualitatively similar picture will also apply if integrability is broken weakly, by a parametrically small coupling. For an operator initially characterized by  $\lambda = 0$ , the correlation function will first exhibit (69) with some subextensive  $\Delta E$ , which will gradually grow to extensive values. It would be interesting to study this transition in detail, to see if the required times may be parametrically longer than  $\beta \sim L$ .

We stress that nonanalyticity of  $C(t)$  at imaginary times is due to  $A(-i\beta)$  becoming typical and is not related to non-analyticity of free energy  $\ln Z(\beta)$  due to a phase transitions at some temperature  $\beta$ . Indeed,  $C(t)$  for the SYK model is known to have a pole at imaginary time [41], while there is no phase transition and free energy is analytic. On the contrary, for the 3d Ising  $\ln Z(\beta)$  is nonanalytic due to a phase transition, but  $C(t)$  is entire, simply because  $A(t)$  explores only a very small part of the corresponding Hilbert space.

In conclusion, we note that the singularity of  $|A(-i\beta)|$  and  $C(-i\beta)$  at finite  $\beta$  in the thermodynamic limit has an IR origin. A straightforward attempt to extend the analysis of this section to field theoretic systems, which can be obtained from lattice systems via an appropriate limit, fails because both  $|A(-i\beta)|$  and  $C(-i\beta)$  are UV-divergent, and this obscures the IR divergence due to chaos. Formulating the criterion of chaos for QFTs using Euclidean operator growth thus remains an open question.

### VI. CONSTRAINTS ON LANCZOS COEFFICIENTS

The bound on power spectrum established in Sec. IV B can be used to constrain the growth of Lanczos coefficients. To remind the reader, Lanczos coefficients  $b_n$  are non-negative real numbers associated with an orthonormal basis in the Krylov space  $A_n$  generated by the action of  $H$  on a given operator  $A_0 = A$ . Starting from a scalar product

$$(A, B) \equiv \text{Tr}(\rho^{1/2} A^\dagger \rho^{1/2} B), \tag{73}$$

and choosing  $A$  normalized such that  $|A|^2 = (A, A) = 1$ , Lanczos coefficients are fixed iteratively from the condition that operators  $A_n$  defined via  $A_{n+1} = ([H, A_n] - b_n A_{n-1})/b_{n+1}$  are orthonormal,  $(A_n, A_m) = \delta_{nm}$ .

An autocorrelation function  $C^W = (A(t), A)$ , defined via scalar product (73), can be parametrized in a number of ways, via its power spectrum  $\Phi^W(\omega)$ , Taylor coefficients (moments)  $M_k$ , or Lanczos coefficients  $b_n$ . Schematically an asymptotic growth of  $b_n$  for large  $n \gg 1$  is related to the behavior of  $M_k$ ,  $k \gg 1$ , high-frequency tail of  $\Phi^W(\omega)$ ,  $\omega \rightarrow \infty$ , or growth of  $C^W(t)$  at the Euclidean time  $t = -i\beta$ . However, the detailed relation is not always trivial. Assuming exponential behavior of power spectrum at large frequencies

$$\Phi^W(\omega) \sim e^{-(\omega/\omega_0)^{2/\lambda}}, \tag{74}$$

it is trivial to obtain the growth of  $M_k$  and  $C^W(\beta)$  by calculating corresponding integrals over  $\omega$  using saddle point approximation. Although much less trivial, but starting from the power spectrum (74), it is also possible to establish an asymptotic behavior of Lanczos coefficients [42]

$$b_n^2 \propto n^\lambda. \tag{75}$$

The converse relations between asymptotic behavior of  $b_n$ ,  $M_k$ ,  $\Phi^W(\omega)$  and  $C^W(-i\beta)$  are much more subtle and may not hold. Thus, we show in Appendix D that smooth asymptotic behavior of  $M_k$  does not imply smooth asymptotic of  $b_n$ .

It was proposed long ago that  $\lambda$  defined in (75) falls into several universality classes, characterizing dynamical systems [38]. In particular, it was observed that  $\lambda = 0$  for noninteracting and  $\lambda = 1$  for interacting integrable models. (It should be noted that since  $\lambda$  characterizes a particular operator, the same system may exhibit several different values of  $\lambda$ .) Recently it was argued in Ref. [7] that  $\lambda = 2$  is a universal behavior in chaotic systems in  $D \geq 2$ .

To thoroughly investigate possible implications of this conjecture, it is desirable to derive the constraints on the behavior of  $C^W$ ,  $\Phi^W$ , and  $M_k$  starting directly from the assumption that  $b_n$  is a smooth function of  $n$  for large  $n$ . In full generality, Lanczos coefficients  $b_n$  are related to the moments  $M_k$  via

$$M_k = \sum_{h_1 \dots h_{k-1}} b_{(h_0+h_1)/2} b_{(h_1+h_2)/2} \dots b_{(h_{k-1}+h_k)/2}. \tag{76}$$

Here the sum is over Dyck paths parameterized by the sets satisfying  $h_0 = h_k = 1/2$ , and  $h_{i+1} = h_i \pm 1$ ,  $h_i > 0$ . Assuming (75) our goal is to deduce an asymptote of  $M_{2k}$  using (76). We develop the approach of summing over weighted Dyck paths in Appendix C. Here we just mention main results. If  $b_n^2$  is asymptotically a smooth function of  $n$ , path integral over Dyck paths can be evaluated via saddle point approximation by identifying a trajectory in the space of indexes, which gives the leading contribution. Thus, if  $b_n$  is smooth,  $M_k$  is also smooth. Furthermore, if  $\lambda = 2$ ,  $b_n^2 \sim \alpha^2 n^2$ , and the leading order behavior is

$$M_{2k} \approx \left(\frac{2\alpha}{\pi}\right)^{2k} (2k)! \tag{77}$$

Thus, starting from the asymptotic behavior  $b_n^2 \propto \alpha^2 n^2$ , we necessarily find that  $C^W$  has a singularity at  $\beta = \pi/(2\alpha)$ , in full agreement with the conjecture of previous section that singularity in Euclidean time is the characteristic property of chaos.

Provided  $C^W(t)$  is analytic inside a strip  $\Im(t) \leq \bar{\beta}_W$  for some  $\bar{\beta}_W$  would immediately imply a bound

$$\alpha \leq \frac{\pi}{2\bar{\beta}_W}. \tag{78}$$

When  $\rho \propto e^{-H/T}$ , provided autocorrelation function  $C_T$  (52) is analytic inside  $|\Im(t)| \leq \bar{\beta}(T)$ , function  $C_T^W$  defined in (62) will be analytic at least inside  $|\Im(t)| \leq \bar{\beta}_W = \bar{\beta}(T) + 1/(2T)$  (see the discussion in Sec. IV B) and therefore

$$\alpha \leq \frac{\pi T}{1 + 2T\bar{\beta}(T)}. \tag{79}$$

We have also established in Sec. II C that  $\bar{\beta}(T) \geq 2\beta^*$  for all  $T$ .

The coefficient  $\alpha$  has been recently conjectured to bound maximal Lyapunov exponent governing exponential growth of the out of time ordered correlation function (OTOC) [7,43],  $\lambda_{\text{OTOC}} \leq 2\alpha$ . This leads to the improved bound on chaos

$$\lambda_{\text{OTOC}} \leq \frac{2\pi T}{1 + 4T\beta^*}, \tag{80}$$




FIG. 2. Lyapunov exponent  $\lambda_{\text{OTOC}}$  for the SYK model as a function of parameter  $v$ , which is related to temperature,  $\pi v T = \cos(\pi v/2)$ . Limit  $v \rightarrow 0$  corresponds to high temperatures,  $v \rightarrow 1$  to small temperatures. Blue line—exact analytic result (81), orange dashed line—improved bound (80) with  $2\beta^* = 1$ , green dotted line—original Maldacena-Shenker-Stanford bound  $2\pi T$ .

which is stronger than the original bound  $\lambda_{\text{OTOC}} \leq 2\pi T$  of Ref. [44]. In the limit of quantum field theory,  $(\beta^*)^{-1}$  will be of the order of UV-cut-off, reducing (80) to the original bound. Yet the new bound is nontrivial for discrete models exhibiting chaos.

To illustrate the improved bound, we plot (80) in Fig. 2 for the SYK model in the large  $q$ -limit<sup>5</sup> against the exact value of  $\lambda_{\text{OTOC}}$ , evaluated in [7,41]. We take  $2\beta^* = 1$  to ensure that the autocorrelation function  $C_T$  is analytic inside  $\Im(t) < 2\beta^* = 1$  for all  $T$ . Temperature  $T$  is parametrized via  $1 \geq v \geq 0$ ,  $\pi v T = \cos(\pi v/2)$  such that the exact Lyapunov exponent is

$$\lambda_{\text{OTOC}} = 2 \cos(\pi v/2). \quad (81)$$

We have emphasized above that for 1D systems with short range interactions  $C_T(t)$  has to be analytic in the entire complex plane. This imposes a bound on the growth of Lanczos coefficients. Assuming  $b_n$  is a smooth function of  $n$  [7] proposed that the asymptotic growth in 1D nonintegral systems will acquire a logarithmic correction

$$b_{n+1} \approx \alpha \frac{n}{\ln(n/n_0)}. \quad (82)$$

Using the integral over weighted Dyck paths in Appendix C, we find this to be consistent with the behavior of  $M_k$  outlined in (60) provided

$$\alpha = \pi J/2. \quad (83)$$

Sum over Dyck paths in the case of  $\lambda = 1$  associated with integrable systems is discussed in Appendix C. Since for the local models  $C^W(t)$  is analytic inside a sufficiently small vicinity of  $t = 0$ , asymptotic behavior with  $\lambda > 2$  in such systems is excluded.

<sup>5</sup>Here,  $q$  is a parameter of SYK model and should not be mixed with  $q(\beta)$  defined in (14).

## VII. CONCLUSIONS

We have derived a number of rigorous bounds on the infinity norm of a local operator evolved in Euclidean time, and extended them to autocorrelation function (2). The novel ingredient of our approach is the counting of *lattice animal histories* and formula (13), using which we solved exactly combinatorial problem of counting nested commutators for Bethe lattices (and establish a correct asymptotic for lattices in  $D \geq 2$ ). Some of the bounds derived in this paper were known before. We improved numerical coefficients, including the location of the singularity  $\beta^*$  in  $D \geq 2$ . Our results are strongest possible among the bounds uniformly valid for all local Hamiltonians characterized by the same  $|h_I| \leq J$  defined on a lattice of a particular geometry.

We have also established Euclidean version of Lieb-Robinson bound on the spatial operator growth. In 1D, operators spread at most exponentially, while in  $D \geq 2$ , operators can reach spatial infinity in finite Euclidean time. When the system is integrable, in all  $D$  operators spread polynomially.

As a main point of this paper, we advocated that Euclidean operator growth reflects chaos in the underlying quantum system. If the system is chaotic, the norm growth and spatial growth are maximal possible and the operator norm diverges at some finite Euclidean time. We interpreted this divergence as a consequence of typicality in Krylov space.

There are several distinct characteristic properties of chaos for many-body quantum systems. One is the eigenstate thermalization hypothesis [45], which is concerned with individual matrix elements. Another popular probe is out of time ordered correlation function, which extends the notion of exponential Lyapunov growth to quantum case. Its use as a characteristic of many-body quantum chaos was pioneered in Refs. [46–48] and brought to the spotlight by applications to quantum gravity [44]. Despite recent efforts [43,49–51] there is no clear understanding of how to relate these two characteristics of chaos to each other. We hope that the Euclidean growth, which on the one hand is related to ETH via the behavior of  $C(-i\beta)$  at large  $\beta$ , see (69), and on the other hand is related to OTOC via the bound (80), may provide such a bridge.

## ACKNOWLEDGMENTS

We would like to thank Dima Abanin, Xiangyu Cao, Nick Hunter-Jones, Vadim Oganessian, and Dan Parker for discussions. We are also grateful to Vladimir Kravtsov, William Berdanier, and Sarang Gopalakrishnan for raising our interest in Bethe lattices and for discussions. AD gratefully acknowledges support from the Simons Center for Geometry and Physics, Stony Brook University at which some of the research for this paper was performed. AD is supported by the National Science Foundation under Grant No. PHY-1720374. AA acknowledges support from NSF Grant No. DMR-1918065 and a Kavli ENSI fellowship at UC Berkeley.

## APPENDIX A: ANIMAL HISTORIES GROWTH ON BETHE LATTICES

We consider Bethe lattice of coordination number  $z$  and would like to calculate the total number of lattice animal




FIG. 3. Example of a lattice animal consisting of  $j = 6$  bonds on the Bethe lattice with the coordination number  $z = 3$ . Vertices attached to only one bond are colored blue, the vertex attached to two bonds is colored red and the vertices attached to three bonds are shown in black. The vector corresponding to this lattice animal is  $a_i = (2, 1, 4, 0)$ . One can check that it satisfies (A2).

histories for all possible lattice animals (clusters) consisting of  $j$  bonds. Each lattice animal can be characterized (nonuniquely) by the vector  $a_i$  for  $i = 0, 1, \dots, z$ , where  $a_i$  is the number of vertices attached to  $z - i$  bonds (of that cluster). We illustrate this definition with an example shown in Fig. 3. Since the considered clusters are connected, either  $a_z = 1$ , in which case  $j = 0$  and  $a_i = 0$  for  $i < z$ , or  $a_z = 0$ .

Consider any given lattice animal history associated with a lattice animal with a given  $a_i$ . We can add one additional bond by attaching it to any vertex, which has less than  $z$  bonds already attached to it. If we attach a bond to a vertex with  $k$  bonds already attached to it, the new lattice animal (and associated lattice animal history), is characterized by a new set

$$a'_i = a_i + e_i^{(k)}, \quad e_i^{(k)} = \delta_{z-1,i} - \delta_{z-k,i} + \delta_{z-k-1,i}. \quad (A1)$$

This equation simply reflects the fact that the new cluster has one more vertex with only 1 bond attached to it, one more vertex with  $(k + 1)$  bonds attached to it, and one less vertex with  $k$  attached bonds.

The total number of bonds  $j$  can be expressed through  $a_i$  as follows:

$$j = \frac{\sum_{i=1}^z i a_i - z}{(z - 2)}. \quad (A2)$$

It can be easily checked that adding one bond via (A1) increases  $j$  by one, and taking  $a_z = 1$ ,  $a_i = 0$  for  $i < z$  gives  $j = 0$ .

We denote the total number of lattice animal histories (associated with all possible animals) consisting of  $j$  bonds by

$\phi(j)$ . (The total number of histories characterized by  $a_i$  can be denoted by  $\phi(a_i)$ . Then  $\phi(j)$  is a sum of  $\phi(a_i)$  over all possible vectors  $a_i$  with non-negative coefficients satisfying (A2).) Given a particular lattice animal history, there are many ways one bond can be added. If we decide to add a bond to vertex which already has  $k < z$  bonds attached to it, we will have  $a_{z-k}$  vertices to choose from and  $(z - k)$  possibilities for each vertex we chose. Hence, in total, each lattice animal history parametrized by  $a_i$  gives rise to

$$\sum_{k=0}^{z-1} (z - k) a_{z-k} = j(z - 2) + z \quad (A3)$$

new animal histories consisting of  $j + 1$  bonds. If we sum over all possible animal histories with  $j$  bounds, we should find the total number of animal histories consisting of  $j + 1$  bonds,

$$\phi(j + 1) = \phi(j)(j(z - 2) + z). \quad (A4)$$

This immediately yields

$$\phi(j) = (z - 2)^j \frac{\Gamma(j + z/(z - 2))}{\Gamma(z/(z - 2))}, \quad (A5)$$

where we additionally required  $\phi(0) = 1$ .

While this is not necessary for the bound on operator norm growth, for completeness we derive the number of lattice animals consisting of  $j$  bonds,  $N(j)$ . We first consider all lattice animals which originate at the same vertex and extend into one particular direction (“branch”) on the Bethe lattice. If the number of such animals is  $n(j)$ , then it must satisfy the recursive relation

$$n(j) = \sum_{j_1 + \dots + j_{z-1} = j-1} n(j_1) \dots n(j_{z-1}). \quad (A6)$$

It reflects the fact that we can “move” the initial point by one bond inside the branch and decompose  $j$  into  $j = \sum_{i=1}^z j_i$ ,  $j_z = 1$ . In (A6), we also use that  $n(1) = 1$ . This gives in full generality

$$n(j) = \frac{\Gamma((z - 1)j + 1)}{\Gamma(j + 1)\Gamma(2 + (z - 2)j)}. \quad (A7)$$

The full number of lattice animals  $N(j)$  is related to  $n(j)$  via

$$N(j) = \sum_{j_1 + \dots + j_{z-1} + j_z = j} n(j_1) \dots n(j_{z-1})n(j_z), \quad (A8)$$

with the total number being

$$N(j) = \frac{z \Gamma((z - 1)j + z)}{\Gamma(j + 1)\Gamma(z + 1 + (z - 2)j)}. \quad (A9)$$

At large  $j$ , this number grows as  $\lambda^j$  with the Klarner’s constant

$$\ln \lambda(z) = (z - 1) \ln(z - 1) - (z - 2) \ln(z - 2). \quad (A10)$$

### APPENDIX B: ANIMAL HISTORIES GROWTH ON ARBITRARY LATTICES IN $D \geq 2$

In this section, we consider an arbitrary lattice of coordination number  $z$ , which means that each vertex is adjacent to at most  $z$  bonds. Similarly to previous section, we will characterize a lattice animal (cluster) by a set of numbers  $a_i$ ,




FIG. 4. Example of a lattice animal consisting of  $j = 8$  bonds on a triangular lattice ( $z = 6$ ). Vertices attached to only one bond are colored blue. A unique vertex attached to two bonds is colored red. Vertices attached to three bonds are shown black. Therefore this lattice animal is characterized by the vector  $a_i = (0, 0, 0, 4, 1, 2, 0)$ . This lattice animal has  $\ell = 2$  loops. One can easily check that both (B1) and (B2) are satisfied.

$i = 0, \dots, z$ , where  $a_i$  is the number of vertices attached to  $z - i$  bonds of that cluster. These notations are illustrated with an example in Fig. 4. Assuming that each lattice vertex is attached to exactly  $z$  bonds, we can immediately find the total number of bonds,

$$j = \frac{\sum_{i=0}^z (z - i)a_i}{2}. \tag{B1}$$

The main difference between general case and the case of Bethe lattices is the possibility of loops. We define the number of loops  $\ell$  of a given lattice animal as the minimal number of bonds which should be removed for the animal to have a tree topology. Then Euler’s characteristic formula gives  $\sum_{i=0}^z a_i - j + \ell = 1$ . From here and (B1), we readily find

$$j = \frac{\sum_{i=1}^z ia_i - z(1 - \ell)}{(z - 2)}, \tag{B2}$$

which is a generalization of (A2).

Let us denote by  $n$  total number of ways one can add a bond to a given animal. This number is the total number of bonds adjacent to the animal but not belonging to it. If each vertex had exactly  $z$  bonds adjacent to it, the sum  $\sum_{k=0}^{z-1} (z - k)a_{z-k}$  counts the number of bonds which can be added to each vertex of the animal. Since some bonds have both ends adjacent to the animal, the sum  $\sum_{k=0}^{z-1} (z - k)a_{z-k}$  includes those bonds twice. Furthermore, since some vertices might actually have less than  $z$  bonds adjacent to them,  $\sum_{k=0}^{z-1} (z - k)a_{z-k}$  provides an upper bound. We therefore have an inequality [compare

with (A3)]

$$n \leq \sum_{k=0}^{z-1} (z - k)a_{z-k} = j(z - 2) + (1 - \ell)z. \tag{B3}$$

Since  $\ell \geq 0$ , we can conclude that in full generality  $n \leq j(z - 2) + z$ . This expression does not depend on any details of the animal, except its size  $j$ . We therefore can bound the growth of animal histories for all animals of size  $j$ ,

$$\phi(j + 1) \leq \phi(j)(j(z - 2) + z), \tag{B4}$$

from where follows the inequality

$$\phi(j) \leq (z - 2)^j \frac{\Gamma(j + z/(z - 2))}{\Gamma(z/(z - 2))}. \tag{B5}$$

### APPENDIX C: INTEGRAL OVER WEIGHTED DYCK PATHS

In the context of recursion method Lanczos coefficients  $b_n$  define tridiagonal Liouvillian matrix

$$\mathcal{L} = \begin{pmatrix} 0 & b_1 & 0 & 0 & \dots \\ b_1 & 0 & b_2 & 0 & \dots \\ 0 & b_2 & 0 & b_3 & \dots \\ 0 & 0 & b_3 & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix} \tag{C1}$$

such that correlation function

$$C^W(-i\beta) \equiv (A(-i\beta), A) = \langle 0 | e^{\mathcal{L}\beta} | 0 \rangle, \tag{C2}$$

where scalar product of operators is defined in (73), and  $\langle 0 | \dots | 0 \rangle$  denotes the upper left corner matrix element. By definition, moments  $M_k$  are Taylor series coefficients of  $C^W$ ,

$$M_k = \langle 0 | \mathcal{L}^k | 0 \rangle. \tag{C3}$$

From here and the tridiagonal form of  $\mathcal{L}$  it follows that

$$M_{2k} = \sum_{h_1, \dots, h_{2k}} \prod_{i=1}^{2k-1} b_{(h_i + h_{i+1})/2}, \tag{C4}$$

while all odd moments vanish (this is also obvious from the symmetry  $C^W(t) = C^W(-t)$ ). The sum above is over the sets  $h_i$  such that  $h_1 = h_{2k} = 1/2$ ,  $h_i > 0$ , and  $h_{i+1} = h_i \pm 1$ .

When  $k$  is large, sum over Dyck paths becomes a path integral, parametrized by a smooth function  $f(t)$ ,  $0 \leq t \leq 1$  [53],

$$h_i = \frac{1}{2} + 2kf(i/(2k)). \tag{C5}$$

Function  $f(t)$  satisfies

$$f(0) = f(1) = 0, \quad |f(t_1) - f(t_2)| \leq |t_1 - t_2|, \quad f(t) \geq 0.$$

Derivative  $f'(t)$  defines an average slope of a “microscopic” Dyck path around index  $i \approx 2kt$ . The path is a sequence of “up” and “down” jumps with the probabilities  $p$  and  $1 - p$ , which vary smoothly, such that  $2p(t) - 1 = f'(t)$ . The number of different “microscopic” paths  $\mathcal{N}[f(t)]$  associated with  $f(t)$  is given by the Shannon

entropy of  $p(t)$ ,

$$\mathcal{N}[f(t)] \approx e^{S_0} \quad S_0 = 2k \int_0^1 dt H(p(t)), \quad (C6)$$

$$H(p) = -p \ln(p) - (1-p) \ln(1-p). \quad (C7)$$

In other words,  $\mathcal{N}[f(t)]$  is the measure in the path integral over  $f(t)$ . To verify this result, we calculate the total number of Dyck paths, which is known to be given by the Catalan number,

$$C_k \approx \int \mathcal{D}f(t) e^{S_0}, \quad (C8)$$

by evaluating corresponding path integral via saddle point approximation. By interpreting  $S_0[f(t)]$  as a classical action, classical EOM is

$$\frac{d}{dt} \arctan(f') = 0. \quad (C9)$$

The only solution satisfying boundary conditions is  $f(t) = 0$ , which gives saddle point value

$$C_k \approx 4^k. \quad (C10)$$

This reproduces correct exponential behavior of Catalan numbers,  $C_k \approx 4^k / (k^{3/2} \pi^{1/2})$ .

Assuming  $b_{n+1} = b(n)$  is a smooth function of index, at least for large  $n$ , sum over weighted Dyck paths (C4) can be represented as an integral

$$M_k \approx \int \mathcal{D}f(t) e^S, \quad (C11)$$

$$S = 2k \int_0^1 dt [H(p(t)) + \ln b(2kf(t))].$$

In case of asymptotic behavior  $b^2(n) = \alpha^2 n^\lambda$ , the EOM is

$$-\frac{f''(t)}{1-f'(t)^2} = \frac{\lambda}{2f(t)}. \quad (C12)$$

For general  $\lambda$ , this equation can be solved in terms of an inverse of the Hypergeometric function. We are most interested in two cases,  $\lambda = 2$  and  $\lambda = 1$ . In the latter case,  $b^2(n) = \alpha^2 n$ , the saddle point trajectory is  $f(t) = t(1-t)$  leading to the asymptotic behavior of moments

$$M_{2k} \approx \left(\frac{2k}{e}\right)^k \alpha^{2k} \approx \frac{(2k)!}{k!} \left(\frac{\alpha^2}{2}\right)^k. \quad (C13)$$

This gives an exponential growth of  $C^W$  at larger  $\beta$ ,

$$C^W(-i\beta) \approx e^{(\alpha\beta)^2/2}. \quad (C14)$$

In the ‘‘chaotic’’ case  $\lambda = 2$  the solution satisfying boundary condition is

$$f(t) = \frac{\sin(\pi t)}{\pi}, \quad (C15)$$

and the saddle point value is

$$M_{2k} \approx \left(\frac{4k\alpha}{e\pi}\right)^{2k} \approx \left(\frac{2\alpha}{\pi}\right)^{2k} (2k)! \quad (C16)$$

Provided  $C^W$  is analytic inside the strip  $\Im(t) < \bar{\beta}$ , the asymptotic growth constant has to be bounded by

$$\alpha \leq \frac{\pi}{2\bar{\beta}}. \quad (C17)$$

Finally we consider the scenario when the growth of Lanczos coefficients acquires logarithmic correction,

$$b(n) = \alpha \frac{n}{\ln(n/n_0)}. \quad (C18)$$

In this case, the action (C11) becomes

$$S = 2k \int_0^1 dt (H((1+f')/2) + \ln(f) - \ln(\ln(2kf/n_0))) + 2k \ln(2k\alpha). \quad (C19)$$

Taking into account only leading term in  $1/\ln(2k/n_0)$  expansion, we obtain effective action

$$S = 2k \int_0^1 dt \left[ H((1+f')/2) + \frac{\lambda}{2} \ln(f) \right] + 2k \ln((2k\alpha)/\ln(2k/n_0)), \quad \lambda = 2(1 - 1/\ln(2k/n_0)). \quad (C20)$$

In other words at leading order, the effect of logarithmic correction is in adjusting the scaling parameter  $\lambda$ . When  $\lambda \approx 2$ , the solution of the EOM (C12) can be found in the power series expansion in  $2 - \lambda$ , with the leading term being simply

$$f = \frac{\sin(\pi t)}{\pi} + O\left(\frac{1}{\ln(2k/n_0)}\right). \quad (C21)$$

At leading order, the  $1/\ln(2k)$  correction to  $f$  does not affect the on-shell value of (C20) evaluated at  $\lambda = 2$  simply because at leading order (C21) is a solution of the EOMs of (C20) with  $\lambda = 2$ . Hence the only correction comes from

$$\delta S = 2k \int_0^1 dt \left(\frac{\lambda}{2} - 1\right) \ln(f), \quad (C22)$$

where  $f$  is given by (C15). Combining all together we find [compare with (C16)]

$$M_{2k} \approx \left(\frac{4k\alpha}{e\pi \ln(2k/n_0)}\right)^{2k} (2\pi)^{2k/\ln(2k/n_0)}. \quad (C23)$$

It is more convenient to work with the logarithm of moments,

$$\frac{\ln M_{2k}}{2k} = \ln\left(\frac{2k\alpha}{e\pi}\right) - \ln \ln(2k/(2\pi n_0)) + o(1/\ln(k)).$$

Comparing this with the asymptotic behavior of moments in 1D (60), we identify  $\alpha = \pi J/2$ , while matching  $n_0$  would exceed the available precision of (60).

#### APPENDIX D: RECONSTRUCTION OF $b_n$ FROM $M_k$

In the previous section, we introduced path integral approach to calculate power spectrum moments  $M_k$  summing over the Dyck paths weighted by products of  $b_n$ . This approach immediately shows that if  $b_n$  is a smooth function of  $n$ , at least for large  $n$ , then  $M_k$  smoothly depends on  $k$  for large  $k$ . Conversely, Lanczos coefficients  $b_n$  can be calculated from




FIG. 5. Lanczos coefficients  $b_n^2$  associated with the moments  $M_{2k} = B_{2k} \equiv B_{2k}(1)$ . Choosing different  $m$  in (D3) leads to a qualitatively similar behavior.

$M_k$  using the following relation:

$$\prod_{i=1}^n b_i^2 = b_1^2 \dots b_n^2 = \frac{\det \mathcal{M}_{n+1}}{\det \mathcal{M}_n}, \quad (\text{D1})$$

where  $\mathcal{M}_n$  is a  $n \times n$  Hankel matrix

$$(\mathcal{M}_n)_{ij} = \begin{cases} M_{i+j-2}, & i+j \bmod 2 = 0, \\ 0, & i+j \bmod 2 = 1. \end{cases} \quad (\text{D2})$$

This expression allows calculating individual  $b_n$  as a ratio of determinants, but it does not guarantee that  $b_n$  will smoothly depend on the index, even if  $M_k$  do. To illustrate that smoothness of  $M_k$  does not imply smoothness of  $b_n$ , we consider a mock autocorrelation function

$$C(-i\beta) = \frac{1}{2}(e^{m(e^\beta-1)} + e^{m(e^{-\beta}-1)}), \quad (\text{D3})$$

inspired by (19). In this case,  $M_{2k} = B_{2k}(m)$  and Lanczos coefficients can be calculated numerically. They exhibit a peculiar behavior: initially  $b_n$  seems to be a smooth function of  $n$ , but starting at some critical  $m$ -dependent value behavior of  $b_n$  for even and odd  $n$  becomes drastically different. For even  $n$ ,  $b_n^2 \propto n^2$ , while for odd  $n$ ,  $b_n^2 \propto n$ . This is shown in Fig. 5. It should be noted that while mock correlation function (D3) exhibits expected behavior along the imaginary axis  $t = -i\beta$ , its behavior along real axis is periodic and hence unphysical. Thus it remains to be seen if for lattice models with local interactions  $b_n$  is always asymptotically smoothly depend of  $n$ , or the behavior can be more complicated.

- 
- [1] D. A. Roberts, D. Stanford, and L. Susskind, *J. High Energy Phys.* **03** (2015) 051.
- [2] D. A. Roberts and D. Stanford, *Phys. Rev. Lett.* **115**, 131603 (2015).
- [3] A. Nahum, S. Vijay, and J. Haah, *Phys. Rev. X* **8**, 021014 (2018).
- [4] V. Khemani, A. Vishwanath, and D. A. Huse, *Phys. Rev. X* **8**, 031057 (2018).
- [5] X.-L. Qi and A. Streicher, *J. High Energy Phys.* **08** (2019) 012.
- [6] G. Bentsen, Y. Gu, and A. Lucas, *Proc. Natl. Acad. Sci.* **116**, 6689 (2019).
- [7] D. E. Parker, X. Cao, A. Avdoshkin, T. Scaffidi, and E. Altman, *Phys. Rev. X* **9**, 041017 (2019).
- [8] J. L. F. Barbón, E. Rabinovici, R. Shir, and R. Sinha, *J. High Energy Phys.* **10** (2019) 264.
- [9] E. H. Lieb and D. W. Robinson, in *Statistical Mechanics* (Springer, Berlin, 1972) pp. 425–431.
- [10] Z. Huang and X.-K. Guo, *Phys. Rev. E* **97**, 062131 (2018).
- [11] C.-F. Chen and A. Lucas, Operator growth bounds from graph theory, [arXiv:1905.03682](https://arxiv.org/abs/1905.03682) [math-ph].
- [12] P. Calabrese and J. L. Cardy, *Phys. Rev. Lett.* **96**, 136801 (2006).
- [13] I. L. Aleiner, L. Faoro, and L. B. Ioffe, *Ann. Phys.* **375**, 378 (2016).
- [14] D. J. Luitz and Y. Bar Lev, *Phys. Rev. B* **96**, 020406(R) (2017).
- [15] A. A. Patel, D. Chowdhury, S. Sachdev, and B. Swingle, *Phys. Rev. X* **7**, 031047 (2017).
- [16] A. Das, S. Chakrabarty, A. Dhar, A. Kundu, D. A. Huse, R. Moessner, S. S. Ray, and S. Bhattacharjee, *Phys. Rev. Lett.* **121**, 024101 (2018).
- [17] T. Rakovszky, F. Pollmann, and C. W. von Keyserlingk, *Phys. Rev. X* **8**, 031058 (2018).
- [18] C. W. von Keyserlingk, T. Rakovszky, F. Pollmann, and S. L. Sondhi, *Phys. Rev. X* **8**, 021013 (2018).
- [19] M. J. S. Beach, R. G. Melko, T. Grover, and T. H. Hsieh, *Phys. Rev. B* **100**, 094434 (2019).
- [20] A. Chan, A. De Luca, and J. T. Chalker, *Phys. Rev. Lett.* **121**, 060601 (2018).
- [21] M. Abramovich and I. Stegun, Washington, DC: US Government Printing Office (1964).
- [22] M. Bernstein and N. J. Sloane, *Linear Algebra its Appl.* **226**, 57 (1995).
- [23] J. Vannimenus, B. Nickel, and V. Hakim, *Phys. Rev. B* **30**, 391 (1984).
- [24] G. Bouch, *J. Math. Phys.* **56**, 123303 (2015).
- [25] H. Araki, *Commun. Math. Phys.* **14**, 120 (1969).
- [26] D. A. Abanin, W. De Roeck, and F. Huveneers, *Phys. Rev. Lett.* **115**, 256803 (2015).
- [27] I. Arad, T. Kuwahara, and Z. Landau, *J. Stat. Mech.: Theory Exp.* **2016**, 033301 (2016).
- [28] T. R. de Oliveira, C. Charalambous, D. Jonathan, M. Lewenstein, and A. Riera, *New J. Phys.* **20**, 033032 (2018).
- [29] Y. M. Miranda, G. Slade, *et al.*, *Electronic Communications Probability* **16**, 129 (2011).
- [30] H. Gharibyan, M. Hanada, S. H. Shenker, and M. Tezuka, *J. High Energy Phys.* **07** (2018) 124 [Erratum: JHEP02, 197 (2019)].
- [31] B. Bertini, P. Kos, and T. Prosen, *Phys. Rev. Lett.* **121**, 264101 (2018).
- [32] A. J. Friedman, A. Chan, A. D. Luca, and J. T. Chalker, Spectral statistics and many-body quantum chaos with conserved charge, *Phys. Rev. Lett.* **123**, 210603 (2019).

- [33] O. Khorunzhiy, On asymptotic behavior of bell polynomials and high moments of vertex degree of random graphs, [arXiv:1904.01339](#) [math.PR].
- [34] S. Mukerjee, V. Oganessian, and D. Huse, *Phys. Rev. B* **73**, 035113 (2006).
- [35] A. Dymarsky and A. Gorsky, *Phys. Rev. B* **102**, 085137 (2020).
- [36] U. Brandt and K. Jacoby, *Z. Phys. B* **25**, 181 (1976).
- [37] J. Perk and H. Capel, *Physica A* **89**, 265 (1977).
- [38] J.-M. Liu and G. Müller, *Phys. Rev. A* **42**, 5854 (1990).
- [39] V. Viswanath and G. Müller, *The Recursion Method: Application to Many-Body Dynamics* (Springer Science & Business Media, Berlin, 2008), Vol. 23 .
- [40] P. Calabrese, F. H. L. Essler, and M. Fagotti, *J. Stat. Mech.: Theory Exp.* (2012) P07016.
- [41] J. Maldacena and D. Stanford, *Phys. Rev. D* **94**, 106002 (2016).
- [42] D. Lubinsky, *Acta Applicandae Mathematica* **33**, 121 (1993).
- [43] C. Murthy and M. Srednicki, *Phys. Rev. Lett.* **123**, 230606 (2019).
- [44] J. Maldacena, S. H. Shenker, and D. Stanford, *J. High Energy Phys.* **08** (2016) 106.
- [45] M. Srednicki, *Phys. Rev. E* **50**, 888 (1994).
- [46] B. V. Fine, T. A. Elsayed, C. M. Kropf, and A. S. de Wijn, *Phys. Rev. E* **89**, 012923 (2014).
- [47] T. A. Elsayed and B. V. Fine, *Phys. Scr.* **T165**, 014011 (2015).
- [48] A. E. Tarkhov and B. V. Fine, *New J. Phys.* **20**, 123021 (2018).
- [49] Y. D. Lensky and X.-L. Qi, *J. High Energy Phys.* **06** (2019) 025.
- [50] L. Foini and J. Kurchan, *Phys. Rev. E* **99**, 042139 (2019).
- [51] A. Chan, A. De Luca, and J. T. Chalker, *Phys. Rev. Lett.* **122**, 220601 (2019).
- [52] M. Kliesch, C. Gogolin, M. J. Kastoryano, A. Riera, and J. Eisert, *Phys. Rev. X* **4**, 031019 (2014).
- [53] A. Okounkov, *Bull. Am. Math. Soc.* **53**, 187 (2016).