

High phase-space density gas of NaCs Feshbach molecules

Aden Z. Lam,¹ Niccolò Bigagli,¹ Claire Warner,¹ Weijun Yuan,¹ Siwei Zhang,¹ Eberhard Tiemann,² Ian Stevenson,¹ and Sebastian Will^{1,*}

¹*Department of Physics, Columbia University, New York, New York 10027, USA*

²*Institut für Quantenoptik, Leibniz Universität Hannover, 30167 Hannover, Germany*



(Received 8 February 2022; accepted 29 March 2022; published 25 April 2022)

We report on the creation of ultracold gases of bosonic Feshbach molecules of NaCs. The molecules are associated from overlapping gases of Na and Cs using a Feshbach resonance at 864.12(5) G. We characterize the Feshbach resonance using bound-state spectroscopy, in conjunction with a coupled-channel calculation. By varying the temperature and atom numbers of the initial atomic mixtures, we demonstrate the association of NaCs gases over a wide dynamic range of molecule numbers and temperatures, reaching 70 nK for our coldest systems and a phase-space density near 0.1. This is an important stepping stone for the creation of degenerate gases of strongly dipolar NaCs molecules in their absolute ground state.

DOI: [10.1103/PhysRevResearch.4.L022019](https://doi.org/10.1103/PhysRevResearch.4.L022019)

The ability to associate weakly bound molecules from ultracold atoms via Feshbach resonances has enabled groundbreaking advances in ultracold quantum science, including the creation of Bose-Einstein condensates (BECs) of Feshbach molecules [1–4], studies of the BEC-BCS crossover in ultracold Fermi gases [5–8], and the realization of polaronic systems [9–11]. Feshbach molecules have also enabled the preparation of ultracold molecules in their absolute ground state [12,13]. Heteronuclear molecules, which have an electric dipole moment in their ground state, are of particular interest. The long-range interactions between dipolar molecules open up new opportunities for the creation of strongly correlated and highly entangled many-body quantum states [14–16], with exciting possibilities for quantum simulation [17–19] and quantum computing [20].

Many applications of dipolar molecules will require quantum degenerate molecular gases. The coldest molecular gases have been created by associating Feshbach molecules from ultracold atomic mixtures that are then transferred to the rovibrational ground state via stimulated Raman adiabatic passage (STIRAP). So far, ultracold gases of heteronuclear Feshbach molecules have been created with KRb [21,22], RbCs [23,24], NaK [25], LiK [26], NaRb [27], and LiNa [28]. Typically, numbers have ranged from 1000 to 20 000 molecules and temperatures from 200 to 1200 nK. Recently, it has been shown that degenerate Fermi gases of KRb [29] and NaK [30] can be directly created through optimized Feshbach association from the initial Bose-Fermi mixture, reaching temperatures of 50 and 100 nK, respectively. The equivalent for heteronu-

clear bosonic molecules, the creation of a molecular BEC, has not been achieved yet. For Bose-Bose mixtures enhanced loss in the vicinity of interspecies Feshbach resonances tends to induce heating [31,32] which so far has limited phase-space densities (PSDs) for bosonic molecular gases to around 0.01 [23,33].

In this Letter, we demonstrate the creation of ultracold ensembles of NaCs Feshbach molecules from overlapping Bose-Bose mixtures of ²³Na and ¹³³Cs and observe phase-space densities near 0.1 (see Fig. 1). We study molecule formation for different temperatures and atom numbers of the initial mixtures, associating molecules from thermal Na and Cs gases, from Na BECs overlapping with thermal Cs gases, and from overlapping BECs. This allows us to control the properties of molecular ensembles over a wide dynamic range, from up to 20 000 molecules at 2 μK to 600 molecules at 70 nK. We find a lifetime of about 6 ms for high-density NaCs ensembles ($\sim 10^{12}$ cm⁻³), providing excellent conditions for the transfer of ultracold NaCs Feshbach molecules into the rovibrational ground state via STIRAP.

NaCs molecules are of special interest due to an exceptionally large dipole moment of $d = 4.6$ D [34] in the rovibrational ground state. For NaCs, the effective range of dipole-dipole interactions, $a_d = md^2/(8\pi\epsilon_0\hbar^2)$, can reach tens of micrometers, an order of magnitude larger than for NaK and two orders of magnitude larger than for KRb (m denotes the molecular mass). In earlier work, NaCs molecules have been created from laser-cooled mixtures of Na and Cs [35–38] and single NaCs molecules have been created in optical tweezer traps [39–41]. The present work constitutes a step towards ultracold bulk gases of NaCs molecules, enabled by our recent demonstration of miscible and overlapping quantum gas mixtures of Na and Cs [42]. Due to its large dipole moment, NaCs is a promising candidate for the realization of novel many-body phases in strongly interacting dipolar quantum matter [15–19], making the pursuit of a BEC of NaCs ground-state molecules an exciting goal.

*sebastian.will@columbia.edu

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.




FIG. 1. Formation of ultracold NaCs Feshbach molecules. (a) Illustration of association of free atoms into weakly bound Feshbach molecules. (b) Absorption images of NaCs molecules after 16 ms of time-of-flight expansion. Images corresponds to the data shown in Fig. 4. Phase-space density increases from left to right. Gravity points in the $-z$ direction. (c) Azimuthally averaged optical density profiles (black points) with Gaussian fits (red line). The molecule temperatures are obtained as discussed in the main text.

Our experiment begins with the preparation of ultracold mixtures of Na and Cs in their hyperfine ground states $|F, m_F\rangle = |1, 1\rangle$ and $|3, 3\rangle$, respectively. Here, F denotes the total angular momentum and m_F its projection on the magnetic field axis. Following evaporative cooling in the magnetic trap, the mixture is transferred into a crossed optical dipole trap operating at 1064 nm and further evaporatively cooled in the presence of a strong magnetic field of $B_{\text{prep}} = 894$ G. In the optical dipole trap, the trap frequencies of Na and Cs are very similar, $\omega^{\text{Na}} \approx 1.08\omega^{\text{Cs}}$, such that the differential gravitational sag is minimal. At the end of evaporation the trap frequencies are $\{\omega_x, \omega_y, \omega_z\} = 2\pi \times \{74(1), 54(1), 332(7)\}$ Hz (measured for Na). At B_{prep} the intra- and interspecies scattering lengths are such that Na and Cs clouds are miscible [42].

First, we identify a Feshbach resonance for molecule formation. Using trap loss spectroscopy, we locate a Feshbach resonance at 864 G with an approximate width of 1 G [43]. This resonance has been theoretically predicted [44] and was recently observed in Ref. [40]. With the coarse location of the resonance known, we associate NaCs molecules via a magnetic field ramp from B_{prep} to 862.8 G [see Fig. 2(a)]. The ramp quickly traverses the region of increasingly attractive Cs-Cs scattering length [see Fig. 2(e)] with a speed of 15 G/ms to suppress three-body losses of Cs, before crossing the Feshbach resonance at a reduced speed of 1 G/ms to ensure adiabatic molecule formation. After association, we apply a magnetic field gradient of 25 G/cm and reduce the depth of the optical dipole trap by a factor of 4. The gradient removes nonassociated Na and Cs atoms from the trap, while levitating NaCs Feshbach molecules [45]. This allows for background-free imaging of molecules [see details in the Supplemental Material (SM) [43]].




FIG. 2. Magnetoassociation of NaCs Feshbach molecules and characterization of the Feshbach resonance. (a) Schematic of the magnetic field ramp for molecule association, dissociation, and subsequent imaging. t_{TOF} denotes the duration of time-of-flight expansion. (b) Molecule dissociation yields the accurate location of the Feshbach resonance. The gray-shaded region indicates $B_{\text{res}} = 864.12(5)$ G. (c) Molecular binding energy as a function of magnetic field. The solid line shows a fit using a coupled-channel calculation. (d) Closed-channel fraction as a function of magnetic field, determined from the coupled-channel calculation. (e) Scattering length of Na-Na (orange) [47], Cs-Cs (red) [48], and Na-Cs (black) as a function of magnetic field.

We determine the resonance position by measuring the magnetic field at which NaCs molecules dissociate using the magnetic field ramp shown in Fig. 2(a) [5,6]. If B_{final} is above the resonance, the molecules are dissociated and detected; if B_{final} is below the resonance, they remain undetected. Figure 2(b) shows the determination of the resonance location at $B_{\text{res}} = 864.12(5)$ G. In addition, we measure the molecular binding energy as a function of magnetic field with the oscillating magnetic field method [46]. We record loss spectra at several magnetic field values (see details in SM [43]) and obtain the data shown in Fig. 2(c). The binding energy shows a linear behavior up to very close to the resonance, which indicates that the molecular state is closed-channel dominated. This is confirmed by a coupled-channel calculation, which allows the calculation of the closed-channel fraction as shown in Fig. 2(d). The measured and calculated parameters are summarized in Table I and show good agreement with Ref. [40]. The Na-Cs scattering length that results from the parametrization of the Feshbach resonance is displayed in Fig. 2(e).

We characterize the properties of the associated molecular clouds in terms of molecule number, temperature, and

TABLE I. Parametrization of the Feshbach resonance using $a(B) = a_{\text{bg}}[1 - \Delta/(B - B_{\text{res}})]$. The error bar in the position reflects the systematic uncertainty in our magnetic field calibration. Theoretical values are obtained from a coupled-channel calculation.

	Experiment	Theory
B_{res} (G)	864.12(5)	864.07(2)
a_{bg} (a_0)	29(4) [42]	31.58(3)
Δ (G)	~ 1	1.30(1)

phase-space density. For the temperature measurement, we develop a protocol that is based on the time-of-flight expansion of dissociated molecules, following the sequence shown in Fig. 2(a). Right after association, the molecules are released from the trap, initiating time-of-flight expansion, and nonassociated atoms are separated via a magnetic field gradient. After 5 ms, the molecules are dissociated by a reverse field ramp with variable speed (see details in SM [43]). The dissociated molecules continue ballistic expansion as Na and Cs atoms for a variable time of flight until an image of Cs is taken [see Fig. 3(a)]. From the expansion data we obtain the mean kinetic energy via $E_k = (3/2)m_{\text{Cs}}\Delta\sigma^2/\Delta t_{\text{TOF}}^2$ [49], where m_{Cs} is the mass of a Cs atom.

To extract the mean kinetic energy of the molecules prior to dissociation, we take into account that E_k is comprised of the mean kinetic energy of the molecules E_{mol} and an additional kinetic energy from the dissociative reverse ramp




FIG. 3. Measuring the mean kinetic energy of NaCs Feshbach molecules. (a) Cloud size σ vs time of flight t_{TOF} for a dissociation ramp speed of 10 G/ms. σ is obtained from a two-dimensional Gaussian fit to the Cs clouds after dissociation. The solid line shows a linear fit where the slope equals to $\Delta\sigma^2/\Delta t_{\text{TOF}}^2$ (see text). (b) Mean kinetic energy of the resulting Cs cloud vs ramp speed. The solid line shows a fit of $\xi E_{\text{mol}} + \Delta E(\dot{B})$, where E_{mol} is the only fitting parameter. For this data set, the fit yields $T_{\text{mol}} = 124(13)$ nK.

ΔE . The additional kinetic energy arises from nonadiabaticity of the dissociation ramp and depends on the ramp speed and the coupling strength of the Feshbach resonance; the latter is characterized by the background scattering length a_{bg} and the resonance width Δ . This has been previously considered for homonuclear molecules [49,50]. For our case, involving heteronuclear molecules, we find that the additional kinetic energy carried by Cs depends on the ramp speed as [43]

$$\Delta E(\dot{B}) = \frac{m_{\text{Na}}}{m_{\text{Na}} + m_{\text{Cs}}} \Gamma(5/3) \left(\frac{3\hbar^2 \dot{B}}{4|\Delta|a_{\text{bg}}\sqrt{2\mu}} \right)^{2/3}, \quad (1)$$

where Γ is the gamma function, m_{Na} is the mass of a Na atom, \hbar is Planck's constant divided by 2π , $\mu = m_{\text{Na}}m_{\text{Cs}}/(m_{\text{Na}} + m_{\text{Cs}})$ the reduced mass of the molecule, and \dot{B} the ramp speed across the resonance. Using the fitted values for a_{bg} and Δ (see Table I), we quantify ΔE without free parameters using Eq. (1) and find good agreement with our experimental data for different ramp speeds, as shown in Fig. 3(b). From the fit we extract the energy ξE_{mol} at zero ramp speed (full adiabaticity), where E_{mol} is the mean kinetic energy of the molecular gas prior to dissociation and $\xi = m_{\text{Cs}}/(m_{\text{Na}} + m_{\text{Cs}})$.

In Fig. 4, we study the dependence of molecule number, temperature, and phase-space density [43] on the conditions of the initial atomic mixtures. To this end, we associate molecules from thermal gases of Na and Cs, from Na BECs overlapping with a thermal gas of Cs, and from overlapping BECs of Na and Cs. For Na, the initial numbers range from 3.5×10^6 to 5×10^5 , and, for Cs, from 9×10^5 to 4×10^4 atoms. The Na temperatures are typically slightly lower than the Cs temperatures due to a lag in thermalization of Cs during sympathetic cooling with Na [42]. We use the Cs temperature T_{Cs} as a parameter that characterizes the starting conditions at B_{prep} prior to association.

We observe NaCs samples with $1.2(2) \times 10^4$ molecules when associated from thermal mixtures, and $6(1) \times 10^2$ molecules when associated from overlapping BECs [see Fig. 4(a)]. The detected molecule numbers correspond to 4(1)% of the Cs atoms prior to association, independent of the temperature of the initial mixture. While this fraction is similar to previous observations in heteronuclear bosonic molecules [23,27,51], it is likely not a fundamental limit. Using the pairing model of Ref. [52] we expect that 80% of the Cs atoms are associated into molecules, owing to well-matched densities and good cloud overlap [see Fig. 4(d)]. Indeed, we observe that the majority of Cs atoms vanishes during Feshbach association from dual BECs, suggesting that the molecule formation efficiency is high, but collisional loss between molecules and atoms during the relatively slow atom removal procedure (\sim ms) reduces the number of detectable molecules. A fast atom removal procedure via optical clean-out ($\sim \mu$ s) is expected to improve detectable molecule numbers [53].

The temperatures of the molecular clouds range from 2(1) μ K to 68(15) nK [see Fig. 4(b)]. We calculate the temperature via $T_{\text{mol}} = 3E_{\text{mol}}/2k_B$, which serves as a measure of mean kinetic energy as we do not have direct evidence for thermalization of the molecular ensembles. The molecules follow the trend of the Cs temperatures, especially for the association from thermal mixtures and from Na BECs




FIG. 4. Creation of NaCs Feshbach molecules across a wide parameter range. Molecules are associated from overlapping thermal clouds of Na and Cs (right), overlapping Na BECs and Cs thermal clouds (middle), and overlapping Na and Cs BECs (left). (a) Molecule number vs Cs temperature T_{Cs} . The gray-shaded region indicates a fraction of 4(1)% of the initial Cs atom number. (b) Molecule temperature T_{mol} vs T_{Cs} . The solid line indicates $T_{mol} = T_{Cs}$. (c) Phase-space density of the molecular cloud vs T_{Cs} . Numbered data points correspond to Fig. 1(b). (d) Density profiles and cloud overlap of Na (orange) and Cs (red). The shift of the cloud centers reflects the differential gravitational sag between Na and Cs.

overlapping with a Cs thermal cloud. For the association from dual BECs, a marked increase of temperature compared to the initial Cs temperature is observed; we attribute this to heating of the Cs BEC during the ramp from B_{prep} to below B_{res} , where the Cs scattering length varies from moderately repulsive to strongly attractive [see Fig. 2(e)]. Despite the heating, this is one of the lowest temperatures reported for heteronuclear bosonic Feshbach molecules.

Combining the data on molecule numbers and temperatures, we determine phase-space densities that range from $2.4(4) \times 10^{-4}$ up to $4(1) \times 10^{-2}$. We find the highest phase-space densities when molecules are associated from a Na




FIG. 5. Lifetime of NaCs Feshbach molecules. Decay dynamics are recorded for molecules at a binding energy of $h \times 1.3$ MHz. The solid (dashed) line is a fit to two-body (one-body) decay. The inset shows the derived two-body loss coefficients as a function of molecular binding energy for a cloud temperature of $1.8(4) \mu$ K. We attribute the increased decay at $E_b = h \times 50$ kHz to thermal dissociation of Feshbach molecules as the binding energy is comparable to the temperature.

BEC overlapping with a thermal Cs gas at the verge of condensation. The data demonstrate excellent starting conditions for the preparation of ultracold NaCs ground-state molecules. Recently, effective evaporative cooling has been shown for ground-state molecules using resonant collisional shielding [54,55] and microwave shielding [56,57]. For NaCs, resonant shielding is predicted to lead to a superb ratio of elastic to inelastic collisions of 10^6 [55,58] and microwave shielding should be highly effective owing to weak hyperfine interactions [59,60]. This makes NaCs a promising candidate for direct evaporation to a BEC of dipolar ground-state molecules, in particular when starting from a high initial phase-space density.

Finally, we study the lifetime of NaCs molecular ensembles, while held in the vicinity of the Feshbach resonance. Figure 5 shows a decay curve for a gas in the regime of highest phase-space density with a peak density of $n_0 = 8(2) \times 10^{11}$ cm $^{-3}$ and a temperature of $147(9)$ nK. The data are consistent with two-body loss with a decay rate of $\Gamma_{2B} = 5(2) \times 10^{-10}$ cm 3 s $^{-1}$ and a characteristic decay time of $1/(\Gamma_{2B}\bar{n}) = 6(1)$ ms, where \bar{n} is the average density [61]. We attribute the two-body decay to vibrational relaxation when two molecules collide. We do not observe a strong dependence of the decay rate on molecular binding energy (see the inset of Fig. 5). This is expected as the Franck-Condon factor for vibrational relaxation is proportional to the closed-channel fraction, which, as shown above, saturates for small binding energies (see Fig. 2). The observed lifetime is sufficiently long for fast STIRAP transfer ($\sim 100 \mu$ s) of a high phase-space density gas of NaCs molecules into the rovibrational ground state.

In conclusion, we have demonstrated the creation of high phase-space density gases of NaCs Feshbach molecules, which is an ideal stepping stone for the preparation of ultracold gases of NaCs ground-state molecules. In the ground state, evaporative cooling supported by microwave [56,57,59,60] or resonant shielding [54,55] may open the door to a BEC of dipolar NaCs molecules. In addition, mixtures of Na and Cs have favorable properties for the

creation of dual Mott insulators in an optical lattice. This offers a path towards the efficient formation of Feshbach molecules [62,63] and low-entropy lattice gases of strongly dipolar NaCs ground-state molecules.

We thank Tarik Yefsah for fruitful discussions and for helpful comments on the manuscript. We also thank Joseph Lee and Edita Bytyqi for experimental assistance. This work was supported by an NSF CAREER Award (Award No.

1848466), an ONR DURIP Award (Award No. N00014-21-1-2721), and a Lenfest Junior Faculty Development Grant from Columbia University. S.W. acknowledges additional support from the Alfred P. Sloan Foundation. I.S. was supported by the Ernest Kempton Adams Fund. C.W. acknowledges support from the Natural Sciences and Engineering Research Council of Canada (NSERC) and the Chien-Shiung Wu Family Foundation. W. Y. acknowledges support from the Croucher Foundation.

-
- [1] M. Greiner, C. A. Regal, and D. S. Jin, Emergence of a molecular Bose–Einstein condensate from a fermi gas, *Nature (London)* **426**, 537 (2003).
- [2] M. W. Zwierlein, C. A. Stan, C. H. Schunck, S. M. F. Raupach, S. Gupta, Z. Hadzibabic, and W. Ketterle, Observation of Bose-Einstein Condensation of Molecules, *Phys. Rev. Lett.* **91**, 250401 (2003).
- [3] S. Jochim, M. Bartenstein, A. Altmeyer, G. Hendl, S. Riedl, C. Chin, J. H. Denschlag, and R. Grimm, Bose-Einstein condensation of molecules, *Science* **302**, 2101 (2003).
- [4] Z. Zhang, L. Chen, K.-X. Yao, and C. Chin, Transition from an atomic to a molecular Bose–Einstein condensate, *Nature (London)* **592**, 708 (2021).
- [5] C. A. Regal, M. Greiner, and D. S. Jin, Observation of Resonance Condensation of Fermionic Atom Pairs, *Phys. Rev. Lett.* **92**, 040403 (2004).
- [6] M. W. Zwierlein, C. A. Stan, C. H. Schunck, S. M. F. Raupach, A. J. Kerman, and W. Ketterle, Condensation of Pairs of Fermionic Atoms near a Feshbach Resonance, *Phys. Rev. Lett.* **92**, 120403 (2004).
- [7] G. B. Partridge, K. E. Strecker, R. I. Kamar, M. W. Jack, and R. G. Hulet, Molecular Probe of Pairing in the BEC-BCS Crossover, *Phys. Rev. Lett.* **95**, 020404 (2005).
- [8] M. W. Zwierlein, J. R. Abo-Shaer, A. Schirotzek, C. H. Schunck, and W. Ketterle, Vortices and superfluidity in a strongly interacting Fermi gas, *Nature (London)* **435**, 1047 (2005).
- [9] A. Schirotzek, C.-H. Wu, A. Sommer, and M. W. Zwierlein, Observation of Fermi Polarons in a Tunable Fermi Liquid of Ultracold Atoms, *Phys. Rev. Lett.* **102**, 230402 (2009).
- [10] S. Palzer, C. Zipkes, C. Sias, and M. Köhl, Quantum Transport through a Tonks-Girardeau Gas, *Phys. Rev. Lett.* **103**, 150601 (2009).
- [11] S. Nascimbène, N. Navon, K. J. Jiang, L. Tarruell, M. Teichmann, J. McKeever, F. Chevy, and C. Salomon, Collective Oscillations of an Imbalanced Fermi Gas: Axial Compression Modes and Polaron Effective Mass, *Phys. Rev. Lett.* **103**, 170402 (2009).
- [12] K.-K. Ni, S. Ospelkaus, M. De Miranda, A. Pe’Er, B. Neyenhuis, J. Zirbel, S. Kotochigova, P. Julienne, D. Jin, and J. Ye, A high phase-space-density gas of polar molecules, *Science* **322**, 231 (2008).
- [13] J. G. Danzl, M. J. Mark, E. Haller, M. Gustavsson, R. Hart, J. Aldegunde, J. M. Hutson, and H.-C. Nägerl, An ultracold high-density sample of rovibronic ground-state molecules in an optical lattice, *Nat. Phys.* **6**, 265 (2010).
- [14] L. D. Carr, D. DeMille, R. V. Krems, and J. Ye, Cold and ultracold molecules: science, technology and applications, *New J. Phys.* **11**, 055049 (2009).
- [15] T. Lahaye, C. Menotti, L. Santos, M. Lewenstein, and T. Pfau, The physics of dipolar bosonic quantum gases, *Rep. Prog. Phys.* **72**, 126401 (2009).
- [16] M. A. Baranov, M. Dalmonte, G. Pupillo, and P. Zoller, Condensed matter theory of dipolar quantum gases, *Chem. Rev.* **112**, 5012 (2012).
- [17] A. Micheli, G. Brennen, and P. Zoller, A toolbox for lattice-spin models with polar molecules, *Nat. Phys.* **2**, 341 (2006).
- [18] H. P. Büchler, E. Demler, M. Lukin, A. Micheli, N. Prokof’ev, G. Pupillo, and P. Zoller, Strongly Correlated 2D Quantum Phases with Cold Polar Molecules: Controlling the Shape of the Interaction Potential, *Phys. Rev. Lett.* **98**, 060404 (2007).
- [19] B. Capogrosso-Sansone, C. Trefzger, M. Lewenstein, P. Zoller, and G. Pupillo, Quantum Phases of Cold Polar Molecules in 2D Optical Lattices, *Phys. Rev. Lett.* **104**, 125301 (2010).
- [20] D. DeMille, Quantum Computation with Trapped Polar Molecules, *Phys. Rev. Lett.* **88**, 067901 (2002).
- [21] C. Ospelkaus, S. Ospelkaus, L. Humbert, P. Ernst, K. Sengstock, and K. Bongs, Ultracold Heteronuclear Molecules in a 3D Optical Lattice, *Phys. Rev. Lett.* **97**, 120402 (2006).
- [22] C. Weber, G. Barontini, J. Catani, G. Thalhammer, M. Inguscio, and F. Minardi, Association of ultracold double-species bosonic molecules, *Phys. Rev. A* **78**, 061601(R) (2008).
- [23] M. P. Köppinger, D. J. McCarron, D. L. Jenkin, P. K. Molony, H.-W. Cho, S. L. Cornish, C. R. Le Sueur, C. L. Blackley, and J. M. Hutson, Production of optically trapped $^{87}\text{RbCs}$ Feshbach molecules, *Phys. Rev. A* **89**, 033604 (2014).
- [24] T. Takekoshi, L. Reichsöllner, A. Schindewolf, J. M. Hutson, C. R. Le Sueur, O. Dulieu, F. Ferlaino, R. Grimm, and H.-C. Nägerl, Ultracold Dense Samples of Dipolar RbCs Molecules in the Rovibrational and Hyperfine Ground State, *Phys. Rev. Lett.* **113**, 205301 (2014).
- [25] C.-H. Wu, J. W. Park, P. Ahmadi, S. Will, and M. W. Zwierlein, Ultracold Fermionic Feshbach Molecules of $^{23}\text{Na}^{40}\text{K}$, *Phys. Rev. Lett.* **109**, 085301 (2012).
- [26] A.-C. Voigt, M. Taglieber, L. Costa, T. Aoki, W. Wieser, T. W. Hänsch, and K. Dieckmann, Ultracold Heteronuclear Fermi-Fermi Molecules, *Phys. Rev. Lett.* **102**, 020405 (2009).
- [27] F. Wang, X. He, X. Li, B. Zhu, J. Chen, and D. Wang, Formation of ultracold NaRb Feshbach molecules, *New J. Phys.* **17**, 035003 (2015).
- [28] M.-S. Heo, T. T. Wang, C. A. Christensen, T. M. Rvachov, D. A. Cotta, J.-H. Choi, Y.-R. Lee, and W. Ketterle, Formation of

- ultracold fermionic NaLi Feshbach molecules, *Phys. Rev. A* **86**, 021602(R) (2012).
- [29] L. De Marco, G. Valtolina, K. Matsuda, W. G. Tobias, J. P. Covey, and J. Ye, A degenerate Fermi gas of polar molecules, *Science* **363**, 853 (2019).
- [30] M. Duda, X.-Y. Chen, A. Schindewolf, R. Bause, J. von Milczewski, R. Schmidt, I. Bloch, and X.-Y. Luo, Transition from a polaronic condensate to a degenerate Fermi gas of heteronuclear molecules, [arXiv:2111.04301](https://arxiv.org/abs/2111.04301).
- [31] V. A. Yurovsky, A. Ben-Reuven, P. S. Julienne, and C. J. Williams, Atom loss from Bose-Einstein condensates due to Feshbach resonance, *Phys. Rev. A* **60**, R765 (1999).
- [32] D. S. Petrov, C. Salomon, and G. V. Shlyapnikov, Weakly Bound Dimers of Fermionic Atoms, *Phys. Rev. Lett.* **93**, 090404 (2004).
- [33] J. He, X. Ye, J. Lin, M. Guo, G. Quémener, and D. Wang, Observation of resonant dipolar collisions in ultracold $^{23}\text{Na}^{87}\text{Rb}$ rotational mixtures, *Phys. Rev. Research* **3**, 013016 (2021).
- [34] M. Aymar and O. Dulieu, Calculation of accurate permanent dipole moments of the lowest $^1\Sigma^+$ states of heteronuclear alkali dimers using extended basis sets, *J. Chem. Phys.* **122**, 204302 (2005).
- [35] J. P. Shaffer, W. Chalupczak, and N. P. Bigelow, Photoassociative Ionization of Heteronuclear Molecules in a Novel Two-Species Magneto-optical Trap, *Phys. Rev. Lett.* **82**, 1124 (1999).
- [36] J. Kleinert, C. Haimberger, P. J. Zabawa, and N. P. Bigelow, Trapping of Ultracold Polar Molecules with a Thin-Wire Electrostatic Trap, *Phys. Rev. Lett.* **99**, 143002 (2007).
- [37] C. Haimberger, J. Kleinert, P. Zabawa, A. Wakim, and N. Bigelow, Formation of ultracold, highly polar $X^1\Sigma^+$ NaCs molecules, *New J. Phys.* **11**, 055042 (2009).
- [38] P. Zabawa, A. Wakim, M. Haruza, and N. P. Bigelow, Formation of ultracold $X^1\Sigma^+(v''=0)$ NaCs molecules via coupled photoassociation channels, *Phys. Rev. A* **84**, 061401(R) (2011).
- [39] L. Liu, J. Hood, Y. Yu, J. Zhang, N. Hutzler, T. Rosenband, and K.-K. Ni, Building one molecule from a reservoir of two atoms, *Science* **360**, 900 (2018).
- [40] J. T. Zhang, Y. Yu, W. B. Cairncross, K. Wang, L. R. B. Picard, J. D. Hood, Y.-W. Lin, J. M. Hutson, and K.-K. Ni, Forming a Single Molecule by Magnetoassociation in an Optical Tweezer, *Phys. Rev. Lett.* **124**, 253401 (2020).
- [41] W. B. Cairncross, J. T. Zhang, L. R. B. Picard, Y. Yu, K. Wang, and K.-K. Ni, Assembly of a Rovibrational Ground State Molecule in an Optical Tweezer, *Phys. Rev. Lett.* **126**, 123402 (2021).
- [42] C. Warner, A. Z. Lam, N. Bigagli, H. C. Liu, I. Stevenson, and S. Will, Overlapping Bose-Einstein condensates of ^{23}Na and ^{133}Cs , *Phys. Rev. A* **104**, 033302 (2021).
- [43] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevResearch.4.L022019> for details on the coarse search of the Feshbach resonance via trap loss spectroscopy, imaging of Feshbach molecules, measurement of the molecular binding energy, measurement of molecule temperature prior to dissociation, measurement of mean kinetic energy versus ramp speed, and calculation of the molecular phase-space density.
- [44] O. Docenko, M. Tamanis, J. Zaharova, R. Ferber, A. Pashov, H. Knöckel, and E. Tiemann, The coupling of the $X^1\Sigma^+$ and $a^3\Sigma^+$ states of the atom pair Na + Cs and modelling cold collisions, *J. Phys. B: At., Mol. Opt. Phys.* **39**, S929 (2006).
- [45] The magnetic moment of Na (Cs) is $0.5\mu_B$ ($0.75\mu_B$) and the magnetic moment of NaCs Feshbach molecules is $-1.4(1)\mu_B$.
- [46] S. T. Thompson, E. Hodby, and C. E. Wieman, Ultracold Molecule Production via a Resonant Oscillating Magnetic Field, *Phys. Rev. Lett.* **95**, 190404 (2005).
- [47] S. Knoop, T. Schuster, R. Scelle, A. Trautmann, J. Appmeier, M. K. Oberthaler, E. Tiesinga, and E. Tiemann, Feshbach spectroscopy and analysis of the interaction potentials of ultracold sodium, *Phys. Rev. A* **83**, 042704 (2011).
- [48] M. Berninger, A. Zenesini, B. Huang, W. Harm, H.-C. Nägerl, F. Ferlaino, R. Grimm, P. S. Julienne, and J. M. Hutson, Feshbach resonances, weakly bound molecular states, and coupled-channel potentials for cesium at high magnetic fields, *Phys. Rev. A* **87**, 032517 (2013).
- [49] T. Mukaiyama, J. R. Abo-Shaer, K. Xu, J. K. Chin, and W. Ketterle, Dissociation and Decay of Ultracold Sodium Molecules, *Phys. Rev. Lett.* **92**, 180402 (2004).
- [50] S. Dürr, T. Volz, and G. Rempe, Dissociation of ultracold molecules with Feshbach resonances, *Phys. Rev. A* **70**, 031601(R) (2004).
- [51] K. K. Voges, P. Gersema, T. Hartmann, T. A. Schulze, A. Zenesini, and S. Ospelkaus, Formation of ultracold weakly bound dimers of bosonic $^{23}\text{Na}^{39}\text{K}$, *Phys. Rev. A* **101**, 042704 (2020).
- [52] E. Hodby, S. T. Thompson, C. A. Regal, M. Greiner, A. C. Wilson, D. S. Jin, E. A. Cornell, and C. E. Wieman, Production Efficiency of Ultracold Feshbach Molecules in Bosonic and Fermionic Systems, *Phys. Rev. Lett.* **94**, 120402 (2005).
- [53] Recently, we have implemented a faster optical clean-out procedure and observe an enhancement of the detectable molecule number by at least a factor of 2.
- [54] K. Matsuda, L. De Marco, J.-R. Li, W. G. Tobias, G. Valtolina, G. Quémener, and J. Ye, Resonant collisional shielding of reactive molecules using electric fields, *Science* **370**, 1324 (2020).
- [55] J.-R. Li, W. G. Tobias, K. Matsuda, C. Miller, G. Valtolina, L. De Marco, R. R. Wang, L. Lassablière, G. Quémener, J. L. Bohn *et al.*, Tuning of dipolar interactions and evaporative cooling in a three-dimensional molecular quantum gas, *Nat. Phys.* **17**, 1144 (2021).
- [56] L. Anderegg, S. Burchesky, Y. Bao, S. S. Yu, T. Karman, E. Chae, K.-K. Ni, W. Ketterle, and J. M. Doyle, Observation of microwave shielding of ultracold molecules, *Science* **373**, 779 (2021).
- [57] A. Schindewolf, R. Bause, X.-Y. Chen, M. Duda, T. Karman, I. Bloch, and X.-Y. Luo, Evaporation of microwave-shielded polar molecules to quantum degeneracy, [arXiv:2201.05143](https://arxiv.org/abs/2201.05143).
- [58] M. L. González-Martínez, J. L. Bohn, and G. Quémener, Adimensional theory of shielding in ultracold collisions of dipolar rotors, *Phys. Rev. A* **96**, 032718 (2017).
- [59] T. Karman and J. M. Hutson, Microwave Shielding of Ultracold Polar Molecules, *Phys. Rev. Lett.* **121**, 163401 (2018).
- [60] L. Lassablière and G. Quémener, Controlling the Scattering Length of Ultracold Dipolar Molecules, *Phys. Rev. Lett.* **121**, 163402 (2018).

- [61] The fit function for two-body decay has the form $N(t) = N_0/(1 + \Gamma_{2B}\bar{n}t)$, where N_0 is the initial molecule number. The average density \bar{n} of a thermal gas in a harmonic trap is given by $\bar{n} = n_0/2^{3/2}$, where n_0 is the peak density.
- [62] S. A. Moses, J. P. Covey, M. T. Miecnikowski, B. Yan, B. Gadway, J. Ye, and D. S. Jin, Creation of a low-entropy quantum gas of polar molecules in an optical lattice, [Science](#) **350**, 659 (2015).
- [63] L. Reichsöllner, A. Schindewolf, T. Takekoshi, R. Grimm, and H.-C. Nägerl, Quantum Engineering of a Low-Entropy Gas of Heteronuclear Bosonic Molecules in an Optical Lattice, [Phys. Rev. Lett.](#) **118**, 073201 (2017).