




Monitoring-induced entanglement entropy and sampling complexity

Mathias Van Regemortel,^{1,*} Oles Shtanko,² Luis Pedro García-Pintos,¹ Abhinav Deshpande,³ Hossein Dehghani ¹,
Alexey V. Gorshkov ¹ and Mohammad Hafezi ¹

¹*Joint Quantum Institute and Joint Center for Quantum Information and Computer Science, NIST/University of Maryland, College Park, Maryland 20742, USA*

²*IBM Quantum, IBM Research – Almaden, San Jose, California 95120, USA*

³*Institute for Quantum Information and Matter, California Institute of Technology, Pasadena, California 91125, USA*



(Received 1 February 2022; accepted 12 July 2022; published 9 August 2022)

The dynamics of open quantum systems is generally described by a master equation, which describes the loss of information into the environment. By using a simple model of uncoupled emitters, we illustrate how the recovery of this information depends on the monitoring scheme applied to register the decay clicks. The dissipative dynamics, in this case, is described by pure-state stochastic trajectories, and we examine different unravelings of the same master equation. More precisely, we demonstrate how registering the sequence of clicks from spontaneously emitted photons through a linear optical interferometer induces entanglement in the trajectory states. Since this model consists of an array of single-photon emitters, we show a direct equivalence with Fock-state boson sampling and link the hardness of sampling the outcomes of the quantum jumps with the scaling of trajectory entanglement.

DOI: [10.1103/PhysRevResearch.4.L032021](https://doi.org/10.1103/PhysRevResearch.4.L032021)

The coupling of a quantum system to an environment generally leads to decoherence and, under certain conditions, can be modeled by a Markovian master equation that could generically result in a mixed (nonpure) density matrix [1]. An alternative but equivalent approach describes the “unraveling” of the same density matrix in terms of pure-state stochastic wave-function trajectories [2–5]. Interestingly, for a given master equation, the unraveling in terms of stochastic trajectories is not unique. For example, note that a Lindblad master equation,

$$\partial_t \rho = \gamma \sum_j \left(c_j \rho c_j^\dagger - \frac{1}{2} \{c_j^\dagger c_j, \rho\} \right), \quad (1)$$

is invariant under any transformation $c_i \rightarrow \sum_j U_{ij} c_j$, where U is a unitary matrix and γ is the decoherence rate. Here, c_j are the jump operators that describe dissipative coupling to the environment (see Supplemental Material [6]). In particular, this implies that any observable $\langle O \rangle = \text{Tr}(\rho O)$ preserves its expectation value, independent of the choice of U . In the unraveling picture, on the other hand, the unitary U is of direct importance for the stochastic quantum states, as can be understood by evaluating the effect of a quantum jump $c_i |\psi\rangle$. Nevertheless, averaging expectation values over different trajectory states will converge back to the U -independent result

from the master equation, $\mathbb{E}_\psi \langle \psi | O | \psi \rangle = \text{Tr}(\rho O)$, where \mathbb{E}_ψ is the expectation over all individual trajectories $|\psi\rangle$. This is in contrast with the case of nonlinear quantities, such as bipartite entanglement entropy, which may show an unraveling dependence.

Physically, the specific choice of unraveling of a master equation is determined by the physical observable that is monitored in a dissipative process [7–12], e.g., detecting the decay of a two-level system by observing the emitted single photon. Remarkably, such stochastic quantum trajectories were observed in several pioneering experiments in trapped-ion systems [13–16] and circuit quantum electrodynamics (circuit QED) [17]. Moreover, it has been shown that monitoring such trajectories can be used to manipulate stochastic quantum systems [18–22], with potential applications in quantum error correction [23,24].

Furthermore, from a theoretical perspective, monitoring may have a profound impact on the stochastic trajectory states when it competes with coherent processes. Specifically, it was shown in Refs. [25–28] that a scaling transition for averaged trajectory entanglement entropy can occur. In these works, dissipation was studied in the context of a measurement-induced phase transition [29,30], and the master equation associated with the dissipative dynamics was changing across the phase transition. This implies that the effect of the monitoring protocol itself and the corresponding choice of unraveling remain largely unexplored for the scaling of entanglement entropy in the stochastic trajectory states.

In this Research Letter, we consider different monitoring schemes that correspond to different unravelings of the same master equation and analyze the associated impact on stochastic quantum dynamics. We consider an array of uncoupled single-photon emitters whose decay can be monitored by

*mvanrege@umd.edu




FIG. 1. (a) A schematic illustration of the setup, consisting of a chain of N two-level emitters, M of which are initially in the $|\uparrow\rangle$ state, with the remaining $N - M$ in the $|\downarrow\rangle$ state. The quantum jumps from the spontaneous emissions in the chain are monitored through the output ports of a linear optical network represented by an $N \times N$ unitary U , giving new jump operators c_i . (b) The case $N = M = 22$ and U sampled from the $N \times N$ Haar measure: half-chain entropy for some stochastic trajectories (red) and the averaged value (blue). The inset shows the volume-law scaling of the maximal averaged entanglement entropy \bar{S}_{\max} . (c) After registering M clicks, the jump outcome probabilities are given by Fock-state boson sampling from Eq. (5). A comparison is given for $N = 7, M = 4$, giving 210 possible outcomes, and a Haar-random U , sampled with 10 000 quantum trajectories from the associated unraveling.

detected photons. A linear optical network (LON) is positioned between the emitters and the detectors, as shown in Fig. 1(a), so that the new jump operators correspond to a LON-determined linear combination of the decay jump operators. As the sequence of jump clicks is recorded, a buildup and a decay of entanglement entropy are generated in the state of the emitters; see Fig. 1(b). When the LON unitary is Haar random (see, e.g., Ref. [31]), the averaged entanglement entropy reaches a maximum over time that has volume-law scaling, as shown in the inset of Fig. 1(b). Moreover, since a series of single-photon emissions is recorded, we analytically verify a direct equivalence between sampling the outcomes of the decay jumps and the Fock-state boson sampling problem [32], as we also numerically demonstrate in Fig. 1(c). Finally, we illustrate in Fig. 2 that the depth of the LON determines the scaling of maximal trajectory entanglement entropy over time, ranging from area law for constant depth to volume law when the depth is proportional to the number of emitters. Given the connection of our system to Fock-state boson sampling, we relate the scaling of maximal trajectory entanglement entropy to the hardness of classically sampling the jump-outcome probabilities: polynomial vs superpolynomial time, respectively [33,34]. Utilizing the setup described above, we therefore establish clear connections between the invariance properties of the master equation, the scaling of the associated trajectory entanglement entropy, and the sampling complexity of jump outcomes.

The model. Our setup consists of a chain of N two-level systems that emit photons via deexcitation and are monitored through the output arms of a LON, represented by an $N \times N$ unitary U . We start from a state with M two-level systems in the excited state $|\uparrow\rangle$ and $N - M$ in the ground state $|\downarrow\rangle$, i.e., $|\psi_0(M, N)\rangle \equiv |\uparrow_1 \cdots \uparrow_M \downarrow_{M+1} \cdots \downarrow_{M-N}\rangle$, and assume a




FIG. 2. The entanglement generated in the chain of emitters is studied by monitoring the decays through a LON. (a) Schematic of the setup, where N emitters are excited and monitored through a D -layered LON consisting of staggered layers of 2×2 Haar random unitaries from Eq. (3). (b) and (c) A network of constant depth D shows area-law scaling: In (b), increasing N , $\bar{S}_{\max}^{(D)}$ remains stable, and in (c), entanglement profiles $\bar{S}_N^{(D)}(l, k_{\max}(D))$ for $N = 100$, selected after k_{\max} , when the maximal entropy is reached, saturate in the bulk (we find that k_{\max} is independent of l). (d) and (e) Taking D to scale with system size as $D = pN$ gives a volume law for entropy, converging to the result of an $N \times N$ Haar-random unitary for large p . (d) Scaling of $\bar{S}_{\max}^{(D)}$ with system size shows linear growth, and (e) the profiles $\bar{S}_N^{(D)}(l, k_{\max}(p))$ for $N = 22$ show a strong dependence on subsystem size l .

uniform rate γ for the excited emitters to spontaneously emit a photon and relax to the ground state, as depicted in Fig. 1(a).

It is assumed that $\tau_d \ll 1/(M\gamma)$, with τ_d comprising the time for a photon to traverse the LON and the detector dead time. A jump click recorded in output arm i of the LON U now corresponds to applying the jump operator

$$c_i \equiv \sum_{j=1}^N U_{ij} \sigma_j^-, \quad (2)$$

with $\sigma_j^- = (\sigma_j^x - i\sigma_j^y)/2$ being the decay operator of emitter j and $\sigma_j^{x,y,z}$ being the Pauli (x, y, z) operator acting on site j .

As was emphasized earlier and shown in more detail in the Supplemental Material [6], the Lindblad master equation, given by $\partial_t \rho = \gamma \sum_i (\sigma_i^- \rho \sigma_i^+ - \frac{1}{2} \{\sigma_i^+ \sigma_i^-, \rho\})$, is invariant under unitary mixing of the jump operators (2). On the level of the master equation, the dynamics of the (uncoupled) emitters is a simple classically mixed state, for which the single-emitter density matrix entries evolve for each emitter independently as $\rho_{\uparrow\uparrow} = 1 - \rho_{\downarrow\downarrow} = e^{-\gamma t}$, $\rho_{\downarrow\uparrow} = \rho_{\uparrow\downarrow} = 0$ with $\rho_{ij} = |i\rangle\langle j|$.

Stochastic quantum trajectories. A crucial element in this Research Letter is the explicit monitoring and recording of the jumps c_i (2). The stochastic dynamics resulting from registering the photon clicks in the output arms of U can be simulated with pure-state trajectories [2–4]. Given a state $|\psi(t)\rangle$, we evaluate the probability for jump c_i to occur in a short time interval Δt as $p_i(t) = \gamma \Delta t \langle \psi(t) | c_i^\dagger c_i | \psi(t) \rangle$. The probability $p_{\text{jump}}(t) = \sum_i p_i(t)$ determines whether a jump happens at time t or not. If a jump happens, then c_i is selected with probability $\propto p_i(t)$, and we evaluate $|\psi(t + \Delta t)\rangle = c_i |\psi(t)\rangle$. If there is no jump, the system evolves for time Δt under the

effective non-Hermitian Hamiltonian $H_{\text{eff}} = -\frac{i\gamma}{2} \sum_j c_j^\dagger c_j$. In both scenarios, the state is renormalized after each time step. In the limit $\Delta t \rightarrow 0$, averaging $\langle O \rangle$ over sampled trajectory states is equivalent to computing $\langle O \rangle$ via the master equation (1).

Note that H_{eff} only depends on the number of excited emitters $N_{\text{exc}} = \sum_i \sigma_i^+ \sigma_i^- = \sum_j c_j^\dagger c_j$, and that $|\psi(t)\rangle$ is an eigenstate of N_{exc} between jumps if we start from $|\psi_0(N, M)\rangle$. This means that, after renormalization, the evolution between jumps does not change the stochastic state $|\psi(t)\rangle$.

For the rest of this work, we will therefore discard the explicit time dimension and express the evolution in terms of the jump sequence (m_1, \dots, m_M) , with m_k representing the k th click in output arm $1 \leq m_k \leq N$ and $1 \leq k \leq M$. This sequence can be obtained reliably when $\tau_d \ll 1/(M\gamma)$, since the photon clicks are now registered with an accuracy significantly higher than the duration of emission (the temporal extent of the photonic wave packet).

Connection to remote entanglement of two emitters. To intuitively explain the idea and illustrate the underlying correspondence with bosonic statistics, we start with the simple case of two excited emitters and a 2×2 LON ($N = M = 2$) parametrized as

$$U = \begin{pmatrix} a & b \\ -e^{i\phi} b^* & e^{i\phi} a^* \end{pmatrix}, \quad (3)$$

with $|a|^2 + |b|^2 = 1$, quantifying the mixing between the modes, and ϕ being the relative phase shift. Setting $a = b = 1/\sqrt{2}$ and $\phi = \pi$, corresponding to a 50 : 50 beam splitter, gives two new jumps $c_s = \frac{1}{\sqrt{2}}(\sigma_1^- + \sigma_2^-)$ and $c_a = \frac{1}{\sqrt{2}}(\sigma_1^- - \sigma_2^-)$, the symmetric and antisymmetric jump, respectively. In case a symmetric click is observed, the symmetric jump c_s is applied to the initial state $|\uparrow\uparrow\rangle$, giving the symmetric Bell state $|\psi_s\rangle = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle)$. This state can only decay another time with the same symmetric jump c_s , as seen immediately by evaluating the probabilities $P_i \propto \langle \psi_s | c_i^\dagger c_i | \psi_s \rangle$, with $i = (a, s)$. The same story holds for the antisymmetric jump c_a , and therefore, upon monitoring the output arms of the beam splitter, either the jump sequence (m_s, m_s) or (m_a, m_a) is detected, each with probability $\frac{1}{2}$, and never the sequence (m_s, m_a) or (m_a, m_s) . This is equivalent to the celebrated Hong-Ou-Mandel effect for two indistinguishable photons, incident on the two input arms of a 50 : 50 beam splitter [35]. In our case, however, the indistinguishable photonic wave packets are detected after a time much shorter than the duration of emission. As a result, an intermediate maximally entangled (anti)symmetric Bell state between the two emitters is established to convey the interference between the emitted photons. A similar procedure was considered to generate entanglement between cold atoms in a lattice configuration [36] and experimentally implemented to entangle two distant trapped ions [37]. The effect can also be viewed as super-radiant emission [38].

Correspondence with boson sampling. We now generalize the system to N emitters, of which M are excited, and an $N \times N$ unitary U , representing the LON with monitored output arms; see Fig. 1(a). After having registered all M clicks, an observer knows that all emitters have reached the ground state $|\psi\rangle = |\downarrow\downarrow \dots\rangle$. The probability of detecting the

M clicks in the Markovian sequence $\vec{m} \equiv (m_1, m_2, \dots, m_M)$ can be evaluated as (see Supplemental Material [6])

$$\begin{aligned} P(\vec{m}) &= \frac{1}{M!} \langle \psi_0(M, N) | c_{m_1}^\dagger \dots c_{m_M}^\dagger c_{m_M} \dots c_{m_1} | \psi_0(M, N) \rangle \\ &= \frac{1}{M!} \sum_{\vec{k}, \vec{l}} U_{m_1, k_1}^* \dots U_{m_M, k_M}^* U_{m_M, l_M} \dots U_{m_1, l_1} \\ &\quad \times \langle \psi_0(M, N) | \sigma_{k_1}^+ \dots \sigma_{k_M}^+ \sigma_{l_M}^- \dots \sigma_{l_1}^- | \psi_0(M, N) \rangle \\ &= \frac{|\text{Per}(U_T)|^2}{M!}. \end{aligned} \quad (4)$$

Here, $\text{Per}(A) = \sum_{\sigma \in S_M} \prod_{i=1}^M A_{i, \sigma(i)}$ is the permanent of an $M \times M$ matrix A , with S_M being the symmetric group, i.e., the summation is performed over the $M!$ possible permutations of the numbers $1, \dots, M$. U_T is the $M \times M$ matrix constructed from U by taking the first M columns and repeating the i th row n_i times, where n_i is the number of times detector i appears in the sequence \vec{m} . $|\text{Per}(U_T)|^2$ arises from gathering all terms that give unit (nonzero) expectation value in the second line of Eq. (4). Expression (4) can also be obtained with multiboson correlation sampling, i.e., by evaluating the M th-order temporal correlation function of the photonic quantum state at the output ports of the LON [39].

We see that $P(\vec{m})$ is the same for all \vec{m} that give rise to a given $\vec{n} = (n_1, \dots, n_N)$. Therefore the probability of registering clicks \vec{n} with $\sum_i n_i = M$ is obtained simply by multiplying the expression (4) by the number of sequences \vec{m} that give rise to this \vec{n} , so that

$$P(\vec{n}) = \frac{|\text{Per}(U_T)|^2}{\prod_i n_i!}. \quad (5)$$

The jump outcome probabilities $P(\vec{n})$ in Eq. (5) are exactly the ones found for Fock-state (conventional) boson sampling when M indistinguishable photons are sampled after passing through an $N \times N$ interferometer [32,40], as verified in Fig. 1(c). When U is drawn from the Haar measure and $N = O(M^2)$, it has been proven that sampling from the output distribution is classically hard (takes superpolynomial time) unless the polynomial hierarchy collapses to the third level. This follows from the #P hardness of classically computing the output probabilities in Eq. (5).

Experimentally, Fock-state boson sampling has been implemented for small numbers of photons, well within the classically simulable regime [41–43]. Gaussian boson sampling [44], using squeezed states instead of single photons as input, can be scaled up further, leading to one of the first claims of experimental quantum advantage [45]. Interestingly, by engineering long-range interactions, Fock-state boson sampling was also proven to be equivalent to sampling spin measurement outcomes after a short Hamiltonian time evolution [46,47].

Trajectory entanglement entropy. Our primary interest lies in evaluating nonlinear properties of the stochastic trajectory states of the emitters. For this, we focus on the *averaged trajectory entanglement entropy* of a subsystem of size $l < N$, after having registered $0 \leq k \leq M$ clicks in the output arms

of a network U , evaluated as

$$\bar{S}_M^{(U)}(l, k) = \frac{1}{N_s} \sum_{i=1}^{N_s} S(l)[|\psi_M^{(U)}(k)\rangle_i], \quad (6)$$

with N_s being the number of samples taken and $|\psi_M^{(U)}(k)\rangle_i \propto c_{m_k} \cdots c_{m_1} |\psi_0(M, N)\rangle$, i.e., the state after some sequence \vec{m} of k detected jumps c_{m_j} (2). Furthermore, $S(l)[|\psi\rangle] = -\text{Tr}[\rho_{\mathcal{A}} \ln \rho_{\mathcal{A}}]$ is the von Neumann entanglement entropy of state $|\psi\rangle$, with $\rho_{\mathcal{A}} = \text{Tr}_{\mathcal{B}} |\psi\rangle\langle\psi|$ being the reduced density matrix of subsystem \mathcal{A} , containing l adjacent sites starting from the boundary, and \mathcal{B} , containing the remaining $N - l$ sites.

From a photonic perspective, an equivalent state $|\psi_M^{(U)}(k)\rangle_i$ can be obtained by subtracting k single photons from the M -photon wave function at the output ports (m_1, \dots, m_k) from U and sending the remaining $M - k$ photons back through U .

By sampling stochastic trajectories using matrix-product states (MPSs) [48], we show in Fig. 1(b) that when U is drawn from the Haar measure, a volume-law scaling for entanglement entropy is observed, as seen in the inset. In this case, each new jump c_i (2) generally has a nonzero overlap with any σ_j^- and will induce long-range entanglement between all emitters in the chain. Yet, the initial growth of entanglement is upper bounded by $\bar{S}_M^{(U)}(N/2, k=1) \leq \ln 2$, independent of N , which is obtained from the concavity of entanglement entropy [49] (see Supplemental Material [6] for details).

LON and the sampling procedure. In what follows, we restrict ourselves to the case $N = M$, i.e., all M emitters are initialized in the excited state $|\psi_N(k=0)\rangle = |\uparrow\uparrow\cdots\rangle$. The $N \times N$ unitary $U(N, D)$ that encodes the quantum jumps is implemented through a LON that consists of D staggered layers of Haar random 2×2 unitaries, each of which can be written as Eq. (3) [see Fig. 2(a)]. For a sufficiently deep LON, one can show that sampling instances from the LON converge to drawing the $N \times N$ unitaries from the Haar measure [50].

Each instance in the sample set is obtained by (i) sampling a $U(N, D)$ and (ii) sampling a quantum trajectory, thus yielding a jump sequence m_k and the corresponding stochastic series of (pure) states $|\psi_N(k)\rangle$, with $0 \leq k \leq N$ being the number of registered jump clicks. After repeating this procedure N_s times, we obtain a set of sampled trajectories, and the averaged entanglement entropies $\bar{S}_N^{(D)}(l, k)$ for subsystem size l can be evaluated, yielding the entanglement of the trajectories averaged over unitaries $U(N, D)$.

Previously, a number of works have investigated the entanglement entropy of the M -photon wave function for Fock-state boson sampling in an N -mode LON. In the Haar regime, the photonic wave function shows volume-law scaling of entanglement entropy when exiting the LON [51,52]. In this chain of two-level emitters, on the other hand, the spontaneously emitted photons themselves are short-lived (stemming from the Born-Markov approximation of the quantum trajectory approach), and we study the buildup and decay of entanglement entropy between the emitters induced by registering and applying the jumps c_j (2). Additionally, this also marks a significant difference with the measurement-induced phase transition studied in circuit models [29,30] since no projective measurements are performed on the emitters.

Numerical results and scaling of complexity. The stochastic simulations were run with MPSs [48,53], using the C++ package ITENSOR [54].

In Figs. 2(b) and 2(c), we first study the scaling of entanglement entropy by monitoring outputs of a LON with fixed depth D . The largest achieved averaged entanglement entropy $\bar{S}_{N,\max}^{(D)} \equiv \max_{k,l} [\bar{S}_N^{(D)}(k, l)]$ shows an area-law behavior. In Fig. 2(b), it is seen that $\bar{S}_{N,\max}^{(D)}$ does not scale with system size for fixed D . This is further confirmed in Fig. 2(c) for subsystem scaling for the case $N = 100$, where it is seen that $\bar{S}_N^{(D)}(k, l)$ converges to a finite value in the bulk. Note that, for any k , the maximal $\bar{S}_N^{(D)}(k, l)$ is always reached for $l = N/2$.

Intuitively, after detecting a click from a jump c_j when $D = \text{const}$, an observer can pinpoint a subset of adjacent emitters of size $2D$ from which the decay could have originated, independent of N . Therefore registering a click can only generate local entanglement in the chain. LONs of fixed depth $D \ll N$ are represented by a unitary U that is formulated as a banded matrix of width $2D$. Interestingly, there exist polynomial-time algorithms to efficiently evaluate $\text{Per}(U_T)$ of banded matrices, which encode output probabilities of outcomes with few or no collisions via Eq. (5) [33,34,55]. The efficient evaluation of the output probabilities is in line with our result: The area law of entanglement entropy ensures that the output configurations $P(\vec{n})$ can be efficiently sampled using MPSs of fixed maximal bond dimension to represent the quantum state of the emitters after k clicks [48].

As shown in Figs. 2(d) and 2(e), the situation drastically changes when the network depth D scales linearly with system size: $D = pN$. In Fig. 2(d), we show the maximal averaged entanglement entropy $\bar{S}_{N,\max}^{(pN)}$, which now has a clear linear dependence on system size N , thus establishing a *volume law*. The simulation quickly gets out of reach for efficient simulation with MPSs of a given maximal bond dimension χ_{\max} (set to $\chi_{\max} = 700$). Also, the entanglement profiles of subsystem size l , shown in Fig. 2(e), acquire a strong dependence on subsystem size l when p is increased, which we identify as volume law for the scaling for subsystem entanglement entropy. As p increases, the entanglement entropy approaches the value obtained by sampling U from the $N \times N$ Haar measure [black dashed line in Figs. 2(d) and 2(e)].

In order to secure the classical sampling hardness, the original proof for Fock-state boson sampling requires that $N = O(M^2)$ to ensure *collision-free* samples [32]. While we are not in that regime, to our knowledge no efficient classical algorithm is known to sample the jump outcomes if $N = M$ and $D \propto N$. In our unraveling picture, we face a correlation in complexity: The entanglement entropy between the emitters in the trajectory states has volume-law scaling and quickly surpasses the limit of efficient simulation with MPSs.

In contrast, when the trajectory-averaged entanglement entropy scales as an area law, the sample complexity (the number of trajectory states required in order to accurately sample the density matrix) may be expected to increase exponentially. This is captured by the scaling of the (classical) Shannon entropy of the distribution over quantum trajectory states. Hence there is a trade-off between sample complexity of trajectories and the complexity of simulating each trajectory. It might

be possible to practically exploit this trade-off in a classical algorithm; see Supplemental Material [6] for a more detailed explanation.

Conclusions and outlook. It was illustrated that changing the unraveling of a straightforward, uncoupled master equation of emitters may cause drastic changes in both the entanglement of stochastic trajectory states and the sampling hardness of jump outcomes. Moreover, changing the unraveling is immediately related to an observer monitoring the decay clicks in the output arms of a LON, resulting in the unitary mixing of the decay jumps. Sampling the jump outcomes in the established monitoring scheme is equivalent to the problem of Fock-state boson sampling. Finally, a connection was established between the scaling of entanglement entropy between emitters and the classical hardness of sampling the jump outcomes.

While we have reported different scaling behavior for the trajectory entanglement entropy, we have not yet seen a conclusive signature of a scaling transition for the trajectory entanglement entropy across a critical point, such as presented in, e.g., Refs. [26,28]. For example, one can investigate fermionic or Gaussian models to access larger systems for the scaling analysis.

Note added. Recently, we became aware of a recent work, where an entanglement scaling transition was reported in a homodyne monitoring scheme [56].

Acknowledgments. We acknowledge stimulating discussions with Alireza Seif and Dominik Hangleiter. M.V.R., H.D., and M.H. were sponsored by ARO W911NF2010232, AFOSR FA9550-19-1-0399, NSF OMA-2120757, QSA-DOE, and the Simons Foundation. L.P.G.-P. and A.V.G. acknowledge funding by the DOE ASCR Accelerated Research in Quantum Computing program (Award No. DE-SC0020312), DARPA SAVANT ADVENT, NSF QLCI (Award No. OMA-2120757), DOE QSA, ARO MURI, DOE ASCR Quantum Testbed Pathfinder program (Award No. DE-SC0019040), NSF PFCQC program, AFOSR, AFOSR MURI, and U.S. Department of Energy Award No. DE-SC0019449. A.D. acknowledges support from the National Science Foundation (RAISE-TAQS 1839204). The Institute for Quantum Information and Matter is an NSF Physics Frontiers Center (PHY-1733907). This work used the Extreme Science and Engineering Discovery Environment (XSEDE), supported by National Science Foundation Grants No. ACI-1548562 and ACI-1928147, at the Pittsburgh Supercomputing Center (PSC) [57].

-
- [1] C. Gardiner and Zoller, *Quantum Noise: A Handbook of Markovian and Non-Markovian Quantum Stochastic Methods with Applications to Quantum Optics* (Springer, New York, 2004).
- [2] J. Dalibard, Y. Castin, and K. Mølmer, Wave-Function Approach to Dissipative Processes in Quantum Optics, *Phys. Rev. Lett.* **68**, 580 (1992).
- [3] R. Dum, A. S. Parkins, P. Zoller, and C. W. Gardiner, Monte Carlo simulation of master equations in quantum optics for vacuum, thermal, and squeezed reservoirs, *Phys. Rev. A* **46**, 4382 (1992).
- [4] H. J. Carmichael, Quantum Trajectory Theory for Cascaded Open Systems, *Phys. Rev. Lett.* **70**, 2273 (1993).
- [5] H. Weimer, A. Kshetrimayum, and R. Orús, Simulation methods for open quantum many-body systems, *Rev. Mod. Phys.* **93**, 015008 (2021).
- [6] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevResearch.4.L032021> for more details about the master equation invariance and its impact on stochastic trajectory states, the probability of observing a specific quantum jump sequence, a bound on the initial growth of entanglement and, finally, the effect of trajectory entanglement on the statistical fluctuations of sampling the density matrix.
- [7] J. I. Cirac, R. Blatt, A. S. Parkins, and P. Zoller, Quantum collapse and revival in the motion of a single trapped ion, *Phys. Rev. A* **49**, 1202 (1994).
- [8] V. Bühner and C. Tamm, Resonance fluorescence spectrum of a trapped ion undergoing quantum jumps, *Phys. Rev. A* **61**, 061801(R) (2000).
- [9] H. Nha and H. J. Carmichael, Entanglement within the Quantum Trajectory Description of Open Quantum Systems, *Phys. Rev. Lett.* **93**, 120408 (2004).
- [10] P. Kok, W. J. Munro, K. Nemoto, T. C. Ralph, J. P. Dowling, and G. J. Milburn, Linear optical quantum computing with photonic qubits, *Rev. Mod. Phys.* **79**, 135 (2007).
- [11] J. Gambetta, A. Blais, M. Boissonneault, A. A. Houck, D. I. Schuster, and S. M. Girvin, Quantum trajectory approach to circuit QED: Quantum jumps and the Zeno effect, *Phys. Rev. A* **77**, 012112 (2008).
- [12] H. M. Wiseman and J. M. Gambetta, Are Dynamical Quantum Jumps Detector Dependent?, *Phys. Rev. Lett.* **108**, 220402 (2012).
- [13] J. C. Bergquist, R. G. Hulet, W. M. Itano, and D. J. Wineland, Observation of Quantum Jumps in a Single Atom, *Phys. Rev. Lett.* **57**, 1699 (1986).
- [14] W. Nagourney, J. Sandberg, and H. Dehmelt, Shelved Optical Electron Amplifier: Observation of Quantum Jumps, *Phys. Rev. Lett.* **56**, 2797 (1986).
- [15] T. Sauter, W. Neuhauser, R. Blatt, and P. E. Toschek, Observation of Quantum Jumps, *Phys. Rev. Lett.* **57**, 1696 (1986).
- [16] D. Leibfried, R. Blatt, C. Monroe, and D. Wineland, Quantum dynamics of single trapped ions, *Rev. Mod. Phys.* **75**, 281 (2003).
- [17] K. Murch, S. Weber, C. Macklin, and I. Siddiqi, Observing single quantum trajectories of a superconducting quantum bit, *Nature (London)* **502**, 211 (2013).
- [18] S. Weber, A. Chantasri, J. Dressel, A. N. Jordan, K. Murch, and I. Siddiqi, Mapping the optimal route between two quantum states, *Nature (London)* **511**, 570 (2014).
- [19] L. Sun, A. Petrenko, Z. Leghtas, B. Vlastakis, G. Kirchmair, K. Sliwa, A. Narla, M. Hatridge, S. Shankar, J. Blumoff, L. Frunzio, M. Mirrahimi, M. H. Devoret, and R. J. Schoelkopf, Tracking photon jumps with repeated quantum non-demolition parity measurements, *Nature (London)* **511**, 444 (2014).
- [20] S. Hacohen-Gourgy, L. P. García-Pintos, L. S. Martin, J. Dressel, and I. Siddiqi, Incoherent Qubit Control Using the Quantum Zeno Effect, *Phys. Rev. Lett.* **120**, 020505 (2018).
- [21] Q. Ficheux, S. Jezouin, Z. Leghtas, and B. Huard, Dynamics of a qubit while simultaneously monitoring its relaxation and dephasing, *Nat. Commun.* **9**, 1926 (2018).

- [22] E. Flurin, L. S. Martin, S. Hacothen-Gourgy, and I. Siddiqi, Using a Recurrent Neural Network to Reconstruct Quantum Dynamics of a Superconducting Qubit from Physical Observations, *Phys. Rev. X* **10**, 011006 (2020).
- [23] N. Akerman, S. Kotler, Y. Glickman, and R. Ozeri, Reversal of Photon-Scattering Errors in Atomic Qubits, *Phys. Rev. Lett.* **109**, 103601 (2012).
- [24] Z. K. Mineev, S. O. Mundhada, S. Shankar, P. Reinhold, R. Gutiérrez-Jáuregui, R. J. Schoelkopf, M. Mirrahimi, H. J. Carmichael, and M. H. Devoret, To catch and reverse a quantum jump mid-flight, *Nature (London)* **570**, 200 (2019).
- [25] X. Cao, A. Tilloy, and A. De Luca, Entanglement in a fermion chain under continuous monitoring, *SciPost Phys.* **7**, 024 (2019).
- [26] O. Alberton, M. Buchhold, and S. Diehl, Entanglement Transition in a Monitored Free-Fermion Chain: From Extended Criticality to Area Law, *Phys. Rev. Lett.* **126**, 170602 (2021).
- [27] Y. Fuji and Y. Ashida, Measurement-induced quantum criticality under continuous monitoring, *Phys. Rev. B* **102**, 054302 (2020).
- [28] M. Van Regemortel, Z.-P. Cian, A. Seif, H. Dehghani, and M. Hafezi, Entanglement Entropy Scaling Transition under Competing Monitoring Protocols, *Phys. Rev. Lett.* **126**, 123604 (2021).
- [29] A. Nahum, J. Ruhman, S. Vijay, and J. Haah, Quantum Entanglement Growth under Random Unitary Dynamics, *Phys. Rev. X* **7**, 031016 (2017).
- [30] B. Skinner, J. Ruhman, and A. Nahum, Measurement-Induced Phase Transitions in the Dynamics of Entanglement, *Phys. Rev. X* **9**, 031009 (2019).
- [31] B. Collins and P. Śniady, Integration with respect to the Haar measure on unitary, orthogonal and symplectic group, *Commun. Math. Phys.* **264**, 773 (2006).
- [32] S. Aaronson and A. Arkhipov, The computational complexity of linear optics, in *Proceedings of the Forty-Third Annual ACM Symposium on Theory of Computing* (Association for Computing Machinery, New York, 2011), pp. 333–342.
- [33] K. Temme and P. Wocjan, Efficient computation of the permanent of block factorizable matrices, [arXiv:1208.6589](https://arxiv.org/abs/1208.6589).
- [34] P. Lundow and K. Markström, Efficient computation of permanents, with applications to boson sampling and random matrices, *J. Comput. Phys.* **455**, 110990 (2022).
- [35] C.-K. Hong, Z.-Y. Ou, and L. Mandel, Measurement of Subpicosecond Time Intervals Between Two Photons by Interference, *Phys. Rev. Lett.* **59**, 2044 (1987).
- [36] T. J. Elliott, W. Kozłowski, S. F. Caballero-Benitez, and I. B. Mekhov, Multipartite Entangled Spatial Modes of Ultracold Atoms Generated and Controlled by Quantum Measurement, *Phys. Rev. Lett.* **114**, 113604 (2015).
- [37] D. L. Moehring, P. Maunz, S. Olmschenk, K. C. Younge, D. N. Matsukevich, L.-M. Duan, and C. Monroe, Entanglement of single-atom quantum bits at a distance, *Nature (London)* **449**, 68 (2007).
- [38] R. Wiegner, S. Oettel, D. Bhatti, J. von Zanthier, and G. S. Agarwal, Simulating superradiance from higher-order-intensity-correlation measurements: Single atoms, *Phys. Rev. A* **92**, 033832 (2015).
- [39] V. Tamma and S. Laibacher, Multi-boson correlation sampling, *Quantum Inf. Proc.* **15**, 1241 (2016).
- [40] M. A. Broome, A. Fedrizzi, S. Rahimi-Keshari, J. Dove, S. Aaronson, T. C. Ralph, and A. G. White, Photonic boson sampling in a tunable circuit, *Science* **339**, 794 (2013).
- [41] M. Tillmann, B. Dakić, R. Heilmann, S. Nolte, A. Szameit, and P. Walther, Experimental boson sampling, *Nat. Photonics* **7**, 540 (2013).
- [42] J. B. Spring, B. J. Metcalf, P. C. Humphreys, W. S. Kolthammer, X.-M. Jin, M. Barbieri, A. Datta, N. Thomas-Peter, N. K. Langford, D. Kundys, J. C. Gates, B. J. Smith, P. G. R. Smith, and I. A. Walmsley, Boson sampling on a photonic chip, *Science* **339**, 798 (2013).
- [43] N. Spagnolo, C. Vitelli, M. Bentivegna, D. J. Brod, A. Crespi, F. Flamini, S. Giacomini, G. Milani, R. Ramponi, P. Mataloni, R. Osellame, E. F. Galvão, and F. Sciarrino, Experimental validation of photonic boson sampling, *Nat. Photonics* **8**, 615 (2014).
- [44] C. S. Hamilton, R. Kruse, L. Sansoni, S. Barkhofen, C. Silberhorn, and I. Jex, Gaussian Boson Sampling, *Phys. Rev. Lett.* **119**, 170501 (2017).
- [45] H.-S. Zhong, H. Wang, Y.-H. Deng, M.-C. Chen, L.-C. Peng, Y.-H. Luo, J. Qin, D. Wu, X. Ding, Y. Hu, P. Hu, X.-Y. Yang, W.-J. Zhang, H. Li, Y. Li, X. Jiang, L. Gan, G. Yang, L. You, Z. Wang *et al.*, Quantum computational advantage using photons, *Science* **370**, 1460 (2020).
- [46] D. Gonzalez Olivares, B. Peropadre, A. Aspuru-Guzik, and J. J. García-Ripoll, Quantum simulation with a boson sampling circuit, *Phys. Rev. A* **94**, 022319 (2016).
- [47] B. Peropadre, A. Aspuru-Guzik, and J. J. García-Ripoll, Equivalence between spin Hamiltonians and boson sampling, *Phys. Rev. A* **95**, 032327 (2017).
- [48] D. Perez-Garcia, F. Verstraete, M. Wolf, and J. Cirac, Matrix product state representations, *Quantum Inf. Comput.* **7**, 401 (2007).
- [49] D. P. DiVincenzo, C. A. Fuchs, H. Mabuchi, J. A. Smolin, A. Thapliyal, and A. Uhlmann, Entanglement of assistance, in *NASA International Conference on Quantum Computing and Quantum Communications* (Springer, New York, 1998), pp. 247–257.
- [50] J. Emerson, E. Livine, and S. Lloyd, Convergence conditions for random quantum circuits, *Phys. Rev. A* **72**, 060302(R) (2005).
- [51] H.-L. Huang, W.-S. Bao, and C. Guo, Simulating the dynamics of single photons in boson sampling devices with matrix product states, *Phys. Rev. A* **100**, 032305 (2019).
- [52] C. Oh, K. Noh, B. Fefferman, and L. Jiang, Classical simulation of lossy boson sampling using matrix product operators, *Phys. Rev. A* **104**, 022407 (2021).
- [53] A. J. Daley, Quantum trajectories and open many-body quantum systems, *Adv. Phys.* **63**, 77 (2014).
- [54] M. Fishman, S. R. White, and E. M. Stoudenmire, The ITensor software library for tensor network calculations, [arXiv:2007.14822](https://arxiv.org/abs/2007.14822).
- [55] D. Cifuentes and P. A. Parrilo, An efficient tree decomposition method for permanents and mixed discriminants, *Linear Algebra Appl.* **493**, 45 (2016).
- [56] T. Vovk and H. Pichler, Entanglement-Optimal Trajectories of Many-Body Quantum Markov Processes, *Phys. Rev. Lett.* **128**, 243601 (2022).
- [57] J. Towns, T. Cockerill, M. Dahan, I. Foster, K. Gaither, A. Grimshaw, V. Hazlewood, S. Lathrop, D. Lifka, G. D. Peterson, R. Roskies, J. R. Scott, and N. Wilkins-Diehr, XSEDE: Accelerating scientific discovery, *Comput. Sci. Eng.* **16**, 62 (2014).