

# Electron-phonon interactions from first principles

Feliciano Giustino

*Department of Materials, University of Oxford, Parks Road, Oxford OX1 3PH, United Kingdom*

(published 16 February 2017)

This article reviews the theory of electron-phonon interactions in solids from the point of view of *ab initio* calculations. While the electron-phonon interaction has been studied for almost a century, predictive nonempirical calculations have become feasible only during the past two decades. Today it is possible to calculate from first principles many materials properties related to the electron-phonon interaction, including the critical temperature of conventional superconductors, the carrier mobility in semiconductors, the temperature dependence of optical spectra in direct and indirect-gap semiconductors, the relaxation rates of photoexcited carriers, the electron mass renormalization in angle-resolved photoelectron spectra, and the nonadiabatic corrections to phonon dispersion relations. In this article a review of the theoretical and computational framework underlying modern electron-phonon calculations from first principles as well as landmark investigations of the electron-phonon interaction in real materials is given. The first part of the article summarizes the elementary theory of electron-phonon interactions and their calculations based on density-functional theory. The second part discusses a general field-theoretic formulation of the electron-phonon problem and establishes the connection with practical first-principles calculations. The third part reviews a number of recent investigations of electron-phonon interactions in the areas of vibrational spectroscopy, photoelectron spectroscopy, optical spectroscopy, transport, and superconductivity.

DOI: [10.1103/RevModPhys.89.015003](https://doi.org/10.1103/RevModPhys.89.015003)

## CONTENTS

I. Introduction	2	3. Expressions for the phonon self-energy used in <i>ab initio</i> calculations	23
II. Historical Development	3	B. Effects of the electron-phonon interaction on electrons	25
A. Early approaches to the electron-phonon interaction	3	1. Electron self-energy: Fan-Migdal and Debye-Waller terms	25
1. Metals	3	2. Expressions for the electron self-energy used in <i>ab initio</i> calculations	27
2. Semiconductors	4	3. Temperature dependence of electronic band structures	28
3. Ionic crystals	5	4. Carrier lifetimes	30
B. The pseudopotential method	5	5. Kinks and satellites	31
C. <i>Ab initio</i> self-consistent field calculations	6	6. Model Hamiltonians, polarons, and the cumulant expansion	32
III. Electron-phonon Interaction in Density-functional Theory	7	VI. Efficient Calculations of Matrix Elements and Their Integrals	32
A. Lattice vibrations in crystals	7	A. Wannier interpolation	33
B. Electron-phonon coupling Hamiltonian	8	1. Maximally localized Wannier functions	33
1. Kohn-Sham Hamiltonian	8	2. Interpolation of electron-phonon matrix elements	33
2. Electron-phonon coupling Hamiltonian to first and second order in the atomic displacements	9	3. Electron-phonon matrix elements in polar materials	34
3. Calculation of electron-phonon matrix elements using density-functional perturbation theory	10	B. Fermi-surface harmonics	36
4. The dielectric approach	11	VII. Nonadiabatic Vibrational Frequencies and Linewidths	36
5. Connection with early formulations	12	VIII. Electron-phonon Interactions in Photoelectron Spectroscopy	38
IV. Field-theoretic Approach to the Electron-phonon Interaction	13	A. Electron mass enhancement in metals	41
A. Operators and distinguishability	13	IX. Electron-phonon Effects in the Optical Properties of Semiconductors	41
B. Electron Green's function	14	A. Temperature dependence of band gaps and band structures	41
1. Equation of motion and self-energy	14	1. Perturbative calculations based on the Allen-Heine theory	41
2. The screened Coulomb interaction	16	2. Nonperturbative adiabatic calculations	44
3. Nuclear contribution to the screened Coulomb interaction	17	B. Phonon-assisted optical absorption	44
C. Phonon Green's function	18	X. Carrier Dynamics and Transport	45
D. Hedin-Baym equations	20	A. Electron linewidths and lifetimes	45
V. From a Many-body Formalism to Practical Calculations	20		
A. Effects of the electron-phonon interaction on phonons	21		
1. Phonons in the Born-Oppenheimer adiabatic approximation	21		
2. Phonons beyond the adiabatic approximation	22		

B. Phonon-limited mobility	46
XI. Phonon-mediated Superconductors	47
A. McMillan–Allen–Dynes formula	47
B. Anisotropic Migdal–Eliashberg theory	48
C. Density-functional theory for superconductors	49
XII. Electron-phonon Interactions Beyond the Local Density Approximation to DFT	50
XIII. Conclusions	51
Acknowledgments	52
Appendix A: Born–Von Kármán Boundary Conditions	52
Appendix B: Ladder Operators in Extended Systems	52
References	54

## I. INTRODUCTION

The interaction between fermions and bosons is one of the cornerstones of many-particle physics. It is therefore not surprising that, despite being one of the most thoroughly studied chapters of solid state physics, the interaction between electrons and phonons in solids continues to attract unrelenting attention.

Electron-phonon interactions (EPIs) are ubiquitous in condensed matter and materials physics. For example, they underpin the temperature dependence of the electrical resistivity in metals and the carrier mobility in semiconductors, they give rise to conventional superconductivity, and they contribute to optical absorption in indirect-gap semiconductors. In addition, EPIs enable the thermalization of hot carriers, determine the temperature dependence of electron energy bands in solids, and distort band structures and phonon dispersion relations of metals, leading to characteristic kinks and Kohn anomalies in photoemission and Raman and neutron spectra. EPIs also play a role in the areas of spintronics and quantum information, for example, by coupling lattice and spin degrees of freedom in electromagnons, or by modulating the lifetimes of electron spins in color centers.

Given the fundamental and practical importance of electron-phonon interactions, it is perhaps surprising that the majority of theoretical studies in this area still rely on semiempirical model Hamiltonians, especially in times when *ab initio* calculations have become pervasive in every area of condensed matter and materials physics. The reason for this lag can be found in the complexity of electron-phonon calculations: while density-functional theory (DFT) calculations of total energies and structural properties were already well established in the early 1980s (Martin, 2004), systematic *ab initio* calculations of EPIs had to wait for the development of density-functional perturbation theory (DFPT) for lattice dynamics between the late 1980s and the mid 1990s (Baroni, Giannozzi, and Testa, 1987; Gonze, Allan, and Teter, 1992; Savrasov, 1992).

Despite this delayed start, the past two decades have witnessed tremendous progress in this area, and new exciting applications are becoming accessible as first-principles techniques for studying EPIs catch up with more established DFT methods. These advances are driving the evolution from qualitative and descriptive theories of electron-phonon effects in model solids to quantitative and predictive theories of real materials. As the methodology for calculating EPIs from first

principles is rapidly reaching maturity, it appears that the time is ripe for reviewing this vast, complex, and fascinating landscape.

One of the most authoritative reviews on the theory of EPIs is the classic book by Grimvall (1981). This monumental work represents an unmissable reference for the specialist. However, as this book predates the rise of *ab initio* computational methods based on DFT, it inevitably misses the most recent developments in this area. The present article constitutes an attempt at filling this gap by reflecting on what DFT calculations can contribute to the study of electron-phonon physics. In addition, this article is also an opportunity to establish a unified conceptual and mathematical framework in this incredibly diverse landscape, shed light on the key approximations, and identify some of the challenges and opportunities ahead.

As emphasized by the title “Electron-phonon interactions from first principles,” the aim of this article is to review the *ab initio* theory of EPIs and to survey modern advances in *ab initio* calculations of EPIs. The reader interested in the fundamentals of electron-phonon physics or in theoretical developments relating to model Hamiltonians is referred to the outstanding monographs by Ziman (1960), Grimvall (1981), Schrieffer (1983), Mahan (1993), and Alexandrov and Devreese (2010).

Among significant recent advances that are covered in this review we mention the zero-point renormalization and the temperature dependence of electronic band structures, the calculation of phonon-assisted optical absorption spectra, the electron mass renormalization and the kinks in angle-resolved photoemission spectra, the thermalization of hot carriers in semiconductors, the calculation of phonon-limited mobility, the development of efficient computational techniques for calculating EPIs, and efforts to improve the predictive power of EPI calculations by going beyond standard density-functional theory.

The review is organized as follows: Sec. II provides an historical perspective on the development of theories of the EPI, from early semiempirical approaches to modern first-principles calculations. In Sec. III we examine the various components of DFT calculations of EPIs in solids and set the formalism which will be used throughout this article. Section IV provides a synthesis of the most advanced field-theoretic approaches employed to study EPIs, and Sec. V makes the link between the most general formalism and DFT calculations for real materials. In this section the reader will find a number of expressions which are useful for practical implementations. Section VI reviews advanced computational techniques for performing calculations of EPIs efficiently and accurately, such as Wannier interpolation and Fermi-surface harmonics. Here we also discuss recent progress in the study of electron-phonon couplings in polar semiconductors. In Sec. VII we discuss recent calculations of phonons beyond the adiabatic Born-Oppenheimer approximation. Section VIII reviews calculations of EPIs in the context of photoelectron spectroscopy. Section IX focuses on the optical properties of semiconductors and insulators, in particular, the temperature dependence of the band structure and phonon-assisted optical processes. In Sec. X we review calculations on the effects of EPIs on carrier dynamics and transport, including carrier

thermalization rates and mobilities. Section XI discusses EPI calculations in the area of phonon-mediated superconductivity. Attempts at improving the accuracy and predictive power of *ab initio* EPI calculations by using more sophisticated electronic structure methods are discussed in Sec. XII. Finally in Sec. XIII we highlight the most pressing challenges in the study of EPIs from first principles, and we present our conclusions. The Appendixes give some notational remarks and more technical discussions.

## II. HISTORICAL DEVELOPMENT

The notion of “electron-phonon interactions” is as old as the quantum theory of solids. In fact, in the very same work where Bloch (1929) discussed the formal solutions of the Schrödinger equation in periodic potentials, Sec. V begins with the all-telling title: “The interaction of the electrons and the elastic waves of the lattice.” In this work the first quantum theory of the temperature-dependent electrical resistivity of metals was developed. It took only a few years for Bloch’s “elastic waves” to be replaced by the brand-name “phonon” by Frenkel (1932), thus establishing a tradition that continues unaltered almost a century later (Walker and Slack, 1970).

In order to discuss the early approaches to the electron-phonon problem, it is useful to state right from the start the standard form of the Hamiltonian describing a coupled electron-phonon system:

$$\begin{aligned} \hat{H} = & \sum_{nk} \varepsilon_{nk} \hat{c}_{nk}^\dagger \hat{c}_{nk} + \sum_{\mathbf{q}\nu} \hbar \omega_{\mathbf{q}\nu} (\hat{a}_{\mathbf{q}\nu}^\dagger \hat{a}_{\mathbf{q}\nu} + 1/2) \\ & + N_p^{-1/2} \sum_{\substack{\mathbf{k}, \mathbf{q} \\ mn\nu}} g_{mn\nu}(\mathbf{k}, \mathbf{q}) \hat{c}_{m\mathbf{k}+\mathbf{q}}^\dagger \hat{c}_{n\mathbf{k}} (\hat{a}_{\mathbf{q}\nu} + \hat{a}_{-\mathbf{q}\nu}^\dagger) \\ & + \left[ N_p^{-1} \sum_{\substack{\mathbf{k}, \mathbf{q}, \mathbf{q}' \\ mn\nu\nu'}} g_{mn\nu\nu'}^{\text{DW}}(\mathbf{k}, \mathbf{q}, \mathbf{q}') \hat{c}_{m\mathbf{k}+\mathbf{q}+\mathbf{q}'}^\dagger \hat{c}_{n\mathbf{k}} \right. \\ & \left. \times (\hat{a}_{\mathbf{q}\nu} + \hat{a}_{-\mathbf{q}\nu}^\dagger) (\hat{a}_{\mathbf{q}'\nu'} + \hat{a}_{-\mathbf{q}'\nu'}^\dagger) \right]. \end{aligned} \quad (1)$$

In this equation the first line describes the separate electron and phonon subsystems using the usual second-quantized formalism, while the second line specifies the mutual coupling between electrons and phonons to first order in the atomic displacements (Mahan, 1993). Here  $\varepsilon_{nk}$  is the single-particle eigenvalue of an electron with crystal momentum  $\mathbf{k}$  in the band  $n$ ,  $\omega_{\mathbf{q}\nu}$  is the frequency of a lattice vibration with crystal momentum  $\mathbf{q}$  in the branch  $\nu$ , and  $\hat{c}_{nk}^\dagger$  and  $\hat{c}_{nk}$  ( $\hat{a}_{\mathbf{q}\nu}^\dagger$  and  $\hat{a}_{\mathbf{q}\nu}$ ) are the associated fermionic (bosonic) creation and destruction operators.  $N_p$  is the number of unit cells in the Born–von Kármán (BvK) supercell (see Appendix A). The third and fourth lines of Eq. (1) describe the electron-phonon coupling Hamiltonian to second order in the atomic displacements. This contribution is rarely found in the early literature (hence the square brackets), but it plays an important role in the theory of temperature-dependent band structures (Sec. V.B.1). The matrix elements  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$  and  $g_{mn\nu\nu'}^{\text{DW}}(\mathbf{k}, \mathbf{q}, \mathbf{q}')$  measure the strength of the coupling between the electron and the phonon subsystems and have physical dimensions of an

energy. Here the superscript “DW” stands for Debye-Waller and relates to the Debye-Waller self-energy discussed in Sec. V.B.2. Complete details as well as a derivation of Eq. (1) are provided in Sec. III.

The formal simplicity of Eq. (1) conceals some important difficulties that one faces when attempting to use this equation for predictive calculations. For example, the electronic Hamiltonian relies on the assumption that the system under consideration can be described in terms of well-defined quasiparticle excitations. Similarly, the phonon term is meaningful only within the harmonic and the adiabatic approximations. More importantly, Eq. (1) does not provide us with any prescription for determining the numerical parameters  $\varepsilon_{nk}$ ,  $\omega_{\mathbf{q}\nu}$ ,  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$ , and  $g_{mn\nu\nu'}^{\text{DW}}(\mathbf{k}, \mathbf{q}, \mathbf{q}')$ .

In a sense, the history of the study of electron-phonon interactions is really the history of how to calculate the parameters entering Eq. (1) using procedures that can be at once rigorous, reliable, and practical. As will become clear in Sec. IV, despite great progress in this area, some conceptual difficulties still remain.

### A. Early approaches to the electron-phonon interaction

#### 1. Metals

A clear account of the theory of EPIs until the late 1950s is given by Ziman (1960). In the following we highlight only those aspects that are relevant to the subsequent discussion in this article.

Early studies of electron-phonon interactions in solids were motivated by the quest for a quantum theory of the electrical resistivity in metals (Hoddeson and Baym, 1980). The common denominator of most early approaches is that the electronic excitations in Eq. (1) were described using the free electron gas model  $\varepsilon_{nk} = \hbar^2 \mathbf{k}^2 / 2m_e - \varepsilon_F$ , with  $m_e$  being the electron mass and  $\varepsilon_F$  the Fermi energy; the lattice vibrations were described as acoustic waves using the Debye model  $\omega_{\mathbf{q}\nu} = v_s |\mathbf{q}|$ , where  $v_s$  is the speed of sound in the solid. Both approximations were reasonable given that the systems of interest included almost exclusively elemental metals and primarily monovalent alkali and noble metals (Mott and Jones, 1936). While these approximations were fairly straightforward, it was considerably more challenging to determine the EPI matrix elements  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$  using realistic approximations.

The very first expression of the electron-phonon matrix element was derived by Bloch (1929); using contemporary notation it can be written as

$$g_{mn\nu}(\mathbf{k}, \mathbf{q}) = -i \left( \frac{\hbar}{2N_p M_\kappa \omega_{\mathbf{q}\nu}} \right)^{1/2} \mathbf{q} \cdot \mathbf{e}_{\kappa\nu}(\mathbf{q}) V_0. \quad (2)$$

Here  $M_\kappa$  is the mass of the  $\kappa$ th nucleus, and  $\mathbf{e}_{\kappa\nu}(\mathbf{q})$  is the polarization of the acoustic wave corresponding to the wave vector  $\mathbf{q}$  and mode  $\nu$ . The term  $V_0$  represents a unit-cell average of the “effective” potential experienced by the electrons in the crystal. Equation (2) was meant to describe the scattering from an initial electronic state with wave vector  $\mathbf{k}$  to a final state with wave vector  $\mathbf{k} + \mathbf{q}$ , via an acoustic phonon of wave vector  $\mathbf{q}$  and frequency  $\omega_{\mathbf{q}\nu}$ . The formula was

developed for free electron metals and neglects so-called “umklapp” (folding) processes, i.e., scattering events whereby  $\mathbf{k}$  goes into  $\mathbf{k} + \mathbf{q} + \mathbf{G}$  with  $\mathbf{G}$  being a reciprocal lattice vector. A derivation of Eq. (2) is provided in Sec. III.B.5. In order to determine  $V_0$  Bloch (1929) argued that the crystal may be described as a continuous deformable medium. Starting from this assumption he reached the conclusion that the average potential can be approximated as  $V_0 = \hbar^2/(16m_e a_0^2)$  ( $a_0$  is the Bohr radius). Even though Bloch’s matrix element is no longer in use, this model provides helpful insight into the nature of EPIs in monovalent metals. For example the so-called “polarization factor” in Eq. (2),  $\mathbf{q} \cdot \mathbf{e}_{\kappa\nu}(\mathbf{q})$ , shows that (in the absence of umklapp processes) only longitudinal sound waves scatter electrons.

Nordheim (1931) proposed a refinement of Bloch’s model whereby the average potential  $V_0$  in Eq. (2) is replaced by the Fourier component  $V_\kappa(\mathbf{q})$  of the ionic Coulomb potential (see Sec. III.B.5). The key assumption underlying this model is that the effective potential experienced by the electrons is simply the sum of the individual bare ionic potentials of each nucleus. When a nucleus is displaced from its equilibrium position, the corresponding potential also shifts rigidly. This is the so-called “rigid-ion” approximation.

The main difficulty that arises with the rigid-ion model is that the Fourier transform of the Coulomb potential diverges as  $q^{-2}$  for  $q = |\mathbf{q}| \rightarrow 0$ ; this leads to unrealistically strong EPIs. In order to circumvent this difficulty Mott and Jones (1936) proposed to truncate the ionic potential at the boundary of the Wigner-Seitz unit cell of the crystal. This choice represents the first attempt at including the *electronic screening* of the nuclear potential in a rudimentary form. In practice Mott and Jones (1936) calculated the Fourier transform of  $V_\kappa(\mathbf{r})$  by restricting the integration over a Wigner-Seitz cell; the resulting potential is no longer singular at long wavelengths. A detailed discussion of this model can be found in Ziman (1960).

Despite some initial successes in the study of the electrical conductivity of metals, the descriptive power of these early models was undermined by the complete neglect of the electronic response to the ionic displacements. The first attempt at describing the effect of the electronic screening was made by Bardeen (1937). In his model the average potential  $V_0$  in Eq. (2) is replaced by

$$V_0 \rightarrow V_\kappa(\mathbf{q})/\epsilon(q), \quad (3)$$

where  $\epsilon(q)$  is the Lindhard function (Mahan, 1993):

$$\epsilon(q) = 1 + (k_{\text{TF}}/q)^2 F(q/2k_{\text{F}}). \quad (4)$$

Here  $k_{\text{TF}}$  and  $k_{\text{F}}$  are the Thomas-Fermi screening wave vector and the Fermi wave vector, respectively, and  $F(x) = 1/2 + (4x)^{-1}(1-x^2) \log|1+x|/|1-x|$ . A derivation of Bardeen’s model is provided in Sec. III.B.5. Since  $\epsilon(q) \rightarrow (k_{\text{TF}}/q)^2$  for  $q \rightarrow 0$ , the singularity of the electron-nuclei potential is removed in Bardeen’s matrix element. The work of Bardeen (1937) can be considered as a precursor of modern *ab initio* approaches, insofar as the calculation of the matrix element was carried out using a self-consistent field method

within the linearized Hartree theory. This strategy is similar in spirit to modern DFPT calculations.

The key qualitative difference between the approach of Bardeen (1937) and modern techniques lies in the neglect of exchange and correlation effects in the screening. A possible route to overcome this limitation was proposed by Bardeen and Pines (1955). In this work they considered the role of a screened exchange interaction in the electron-phonon problem (see Appendix B of their work); however, the mathematical complexity of the formalism prevented further progress in this direction. Similar efforts were undertaken by Hone (1960), and a more detailed account of the early approximations to exchange and correlation can be found in Grimvall (1981).

The most interesting aspect of the work by Bardeen and Pines (1955), as well as previous work along the same lines by Nakajima (1954), is that for the first time the electron-phonon problem was addressed using a *field-theoretic* approach.

One interesting feature in the theory of Bardeen and Pines is that their field-theoretic formulation naturally leads to a *retarded* electron-phonon vertex: the effective potential experienced by electrons upon the displacement of nuclei depends on how fast this displacement takes place. In this approach the effective potential  $V_0$  in Eq. (2) is replaced by the dynamically screened potential:

$$V_0 \rightarrow V_\kappa(\mathbf{q})/\epsilon(q, \omega_{\mathbf{q}\nu}). \quad (5)$$

Here  $\epsilon(q, \omega)$  is the frequency-dependent Lindhard function (Mahan, 1993), and the effect of electronic screening is evaluated at the phonon frequency  $\omega = \omega_{\mathbf{q}\nu}$ . Somewhat surprisingly, this development was not followed up in the literature on *ab initio* calculations of EPIs.

## 2. Semiconductors

While the investigation of electron-phonon effects was initially restricted to monovalent metals, the formal developments were soon extended to the case of more complex systems such as semiconductors. Carriers in semiconductors are typically confined within a narrow energy range near the band extrema; consequently it is expected that the dominant electron-phonon scattering mechanisms will involve long-wavelength phonons ( $q \rightarrow 0$ ). This concept was formalized by Bardeen and Shockley (1950) and Shockley and Bardeen (1950), laying the foundations of the “deformation-potential” method.

In the deformation-potential approach it is assumed that the atomic displacements can be described by long-wavelength acoustic waves, and these can be related in turn to the elastic strain of the crystal. Using concepts from the effective mass theory, Bardeen and Shockley showed that in this approximation the potential  $V_0$  in Eq. (2) can be replaced by

$$V_0 \rightarrow E_{1,n\mathbf{k}} = \Omega \partial \epsilon_{n\mathbf{k}} / \partial \Omega, \quad (6)$$

where  $\Omega$  represents the volume of the unit cell, and the electron eigenvalues correspond to the valence or conduction band extrema. The derivation of this result can be found in Appendix B of Bardeen and Shockley (1950). The deformation potentials  $E_1$  were obtained empirically; for example,

Bardeen and Shockley determined these values for the band extrema of silicon by fitting mobility data. More complex scenarios such as anisotropic constant-energy surfaces in semiconductors were subsequently addressed by considering the effects of shear deformations (Dumke, 1956). While the concept of deformation potentials has become a classic in semiconductor physics, this method relies on a semiempirical approach and lacks predictive power.

### 3. Ionic crystals

A class of materials that played an important role in the development of the theory of EPIs is that of ionic crystals. The qualitative difference between ionic solids and the systems discussed in Secs. II.A.1 and II.A.2 is that the atomic displacements can generate long-ranged electric fields; these fields provide a new scattering channel for electrons and holes.

The theory of polar electron-phonon coupling started with the investigation of the electron mean free path in ionic crystals in search for a theoretical model of dielectric breakdown in insulators (Fröhlich, 1937; Fröhlich and Mott, 1939). The central idea of these models is that in insulators the density of free carriers is very low; therefore it is sensible to consider a single electron interacting with the polarization field of the ionic lattice.

The Fröhlich model is similar in spirit to the contemporary work of Bardeen (1937) for metals. The main difference is that Fröhlich considered the screening arising from the dielectric polarization of an insulating crystal, while Bardeen considered the screening arising from the response of the Fermi sea.

Fröhlich, Pelzer, and Zienau (1950) showed that in the case of isotropic ionic crystals the effective potential  $V_0$  appearing in Eq. (2) must be replaced by

$$V_0 \rightarrow - \left[ \frac{e^2 M_\kappa \omega_{\mathbf{q}\nu}^2}{\epsilon_0 \Omega} \left( \frac{1}{\epsilon^\infty} - \frac{1}{\epsilon^0} \right) \right]^{1/2} \frac{1}{|\mathbf{q}|^2}. \quad (7)$$

In this expression  $e$  is the electron charge,  $\epsilon_0$  is the dielectric permittivity of a vacuum, and  $\epsilon^0$  and  $\epsilon^\infty$  are the static and the high-frequency relative permittivities, respectively. This result is derived in Sec. VI.A.3. Using Eqs. (2) and (7) we see that when  $\epsilon_0 > \epsilon^\infty$  the matrix element  $g_{m\nu}(\mathbf{k}, \mathbf{q})$  diverges as  $|\mathbf{q}|^{-1}$  at long wavelengths. This singular behavior can lead to very strong EPIs and provides the physical basis for the phenomenon of electron self-trapping in polarons (Pekar, 1946; Emin, 2013). The initial studies in this area were rapidly followed by more refined approaches based on field-theoretic methods (Lee, Low, and Pines, 1953). A comprehensive discussion of the various models can be found in the original review article by Fröhlich (1954).

### B. The pseudopotential method

The approximations underpinning the models discussed in Sec. II.A become inadequate when one tries to study EPIs for elements across the periodic table. This and other limitations stimulated the development of the pseudopotential method, starting in the late 1950s with the work of Phillips and Kleinman (1959). The theory of pseudopotentials is too vast to be summarized in a few lines, and the reader is referred to

Chapter 11 of Martin (2004) for a thorough discussion. Here we highlight only the aspects that are relevant to the calculation of EPIs.

The genesis of the pseudopotential method is linked with the question of how the valence electrons of metals could be described using the electron gas model, even though the orthogonality to the core states imposes rapid fluctuations of the valence wave functions near the atomic cores. In order to address this question, it is useful to go through the key steps of the orthogonalized plane wave method (OPW) (Herring, 1940). In this method one considers plane waves  $|\mathbf{k} + \mathbf{G}\rangle$  for the wave vector  $\mathbf{k} + \mathbf{G}$  and projects out the component belonging to the Hilbert subspace spanned by core electrons. This is done by defining  $|\mathbf{k} + \mathbf{G}\rangle_{\text{OPW}} = |\mathbf{k} + \mathbf{G}\rangle - \sum_c |\phi_c\rangle \langle \phi_c | \mathbf{k} + \mathbf{G}\rangle$ , where  $|\phi_c\rangle$  represents the core states of all atoms in the system. The functions  $|\mathbf{k} + \mathbf{G}\rangle_{\text{OPW}}$  are by construction orthogonal to core states, and therefore they can be used to expand the valence electron wave functions  $|\psi_{n\mathbf{k}}\rangle$  using only a few basis elements:  $|\psi_{n\mathbf{k}}\rangle = \sum_{\mathbf{G}} c_{\mathbf{k}}(\mathbf{G}) |\mathbf{k} + \mathbf{G}\rangle_{\text{OPW}}$ . In the language of pseudopotential theory  $|\psi_{n\mathbf{k}}\rangle$  is referred to as the “all-electron” wave function, while the function  $|\tilde{\psi}_{n\mathbf{k}}\rangle = \sum_{\mathbf{G}} c_{\mathbf{k}}(\mathbf{G}) |\mathbf{k} + \mathbf{G}\rangle$  is referred to as the “pseudo” wave function. The all-electron and the pseudo wave functions are simply related as follows:

$$|\psi_{n\mathbf{k}}\rangle = \hat{T} |\tilde{\psi}_{n\mathbf{k}}\rangle, \quad \text{with} \quad \hat{T} = 1 - \sum_c |\phi_c\rangle \langle \phi_c|. \quad (8)$$

Here we used a modern notation borrowed from the projector-augmented wave (PAW) method of Blöchl (1994). By construction, the pseudo wave function  $|\tilde{\psi}_{n\mathbf{k}}\rangle$  does not exhibit rapid fluctuations near the atomic cores. The projector operator  $\hat{T}$  is now used to rewrite the single-particle Schrödinger equation for the all-electron wave function [e.g., the Kohn-Sham (KS) equations] in terms of the pseudo wave functions. Using  $\hat{H} |\psi_{n\mathbf{k}}\rangle = \epsilon_{n\mathbf{k}} |\psi_{n\mathbf{k}}\rangle$  and Eq. (8) we have

$$\hat{T}^\dagger \hat{H} \hat{T} |\tilde{\psi}_{n\mathbf{k}}\rangle = \epsilon_{n\mathbf{k}} \hat{T}^\dagger \hat{T} |\tilde{\psi}_{n\mathbf{k}}\rangle, \quad (9)$$

which is a generalized eigenvalue problem. By replacing the definition of  $\hat{T}$  given earlier one finds (Phillips and Kleinman, 1959)

$$(\hat{H} + \hat{V}^{\text{rep}}) |\tilde{\psi}_{n\mathbf{k}}\rangle = \epsilon_{n\mathbf{k}} |\tilde{\psi}_{n\mathbf{k}}\rangle, \quad (10)$$

with  $\hat{V}^{\text{rep}} = \sum_c (\epsilon_{n\mathbf{k}} - \epsilon_c) |\phi_c\rangle \langle \phi_c|$  and  $\epsilon_c$  the eigenvalue of a core electron. Clearly the additional potential  $\hat{V}^{\text{rep}}$  is strongly repulsive and is localized near the atomic cores. Cohen and Heine (1961) showed that this extra potential largely cancels the attractive potential of the nuclei. This is the reason why valence electrons in metals behave almost like free electrons.

The practical consequence of these developments is that it is possible to define smooth effective “pseudopotentials” for systematic band structure calculations, whose form factors include only a few Fourier components (Phillips, 1958; Heine and Abarenkov, 1964; Animalu and Heine, 1965; Cohen and Bergstresser, 1966).

The use of pseudopotentials in electron-phonon calculations started with the works of Sham (1961) and Sham and Ziman (1963). Sham (1961) showed that if the pseudopotential can be described by a local function, then the electron-phonon matrix element  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$  can be calculated by replacing the all-electron potentials and wave functions by the corresponding pseudopotentials and pseudo wave functions. In this approach the pseudopotentials move around rigidly with the ionic cores; therefore we are dealing effectively with an improved version of the rigid-ion approximation discussed in Sec. II.A.

The pseudopotential method was employed by Shuey (1965) in order to calculate the electron-phonon matrix elements in germanium. Shortly afterward many calculations of electron-phonon interactions based on the pseudopotential method appeared in the literature, including work on the resistivity of metals (Carbotte and Dynes, 1967; Dynes and Carbotte, 1968; Hayman and Carbotte, 1971; Kaveh and Wisner, 1972), the electron mass enhancement in metals (Ashcroft and Wilkins, 1965; Grimvall, 1969; Allen and Cohen, 1970, 1972; Allen, 1972a), the superconducting transition temperatures within the McMillan formalism (Allen *et al.*, 1968; Allen and Cohen, 1969), the mobility of semiconductors (Ralph, 1970), and the temperature dependence of semiconductor band structures (Allen and Cardona, 1981, 1983). These calculations were mostly based on phonon dispersion relations extracted from neutron scattering data, and the results were in reasonable agreement with experiment. It seems fair to say that the pseudopotential method enabled the evolution from qualitative to quantitative calculations of electron-phonon interactions.

Before proceeding we note that, although Eqs. (8) and (9) were introduced starting from the method of orthogonalized plane waves, there exists considerable freedom in the choice of the operator  $\hat{T}$ . In practice  $\hat{T}$  can be chosen so as to make  $\tilde{\psi}_{n\mathbf{k}}$  as smooth as possible, while retaining information on the all-electron wave functions near the ionic cores. This was achieved by the PAW method of Blöchl (1994). Broadly speaking it is also possible to reinterpret the historical development of the pseudopotential method as the evolution of the projector  $\hat{T}$ . In fact Blöchl showed how the most popular pseudopotential methods (Hamann, Schlüter, and Chiang, 1979; Bachelet, Hamann, and Schlüter, 1982; Vanderbilt, 1990; Troullier and Martins, 1991) can be derived from the PAW method under specific approximations.

### C. *Ab initio* self-consistent field calculations

Predictive calculations of EPIs became possible with the development of *ab initio* DFT techniques. The key advantage of DFT methods is the possibility of calculating electron band structures, phonon dispersion relations, and electron-phonon matrix elements entirely from first principles. Historically, DFT started with the works of Hohenberg and Kohn (1964) and Kohn and Sham (1965). However, its widespread use had to wait for the development of accurate parametrizations of the exchange and correlation energy of the electron gas (Hedin and Lundqvist, 1971; von Barth and Hedin, 1972; Gunnarsson, Lundqvist, and Wilkins, 1974; Ceperley and

Alder, 1980; Perdew and Zunger, 1981). An introduction to DFT techniques can be found in the books by Parr and Yang (1994) (advanced), Martin (2004) (intermediate), and Giustino (2014) (elementary).

The first calculation of electron-phonon interactions using DFT was carried out by Dacorogna, Cohen, and Lam (1985) using a “frozen-phonon” approach (see Sec. III.B.3). In this work they computed electron bands, phonon dispersions, and electron-phonon matrix elements of Al entirely from first principles. Quoting from the original manuscript: “This calculation is *ab initio* since only information about the Al atom, i.e., the atomic number and atomic mass, is used as input.” Dacorogna *et al.* calculated the so-called electron-phonon coupling strength  $\lambda_{\mathbf{q}\nu}$  for several phonon branches  $\nu$  and momenta  $\mathbf{q}$  throughout the Brillouin zone, as well as the phonon linewidths arising from the EPI (see Secs. VII and XI.A). The average coupling strength was found to be in good agreement with that extracted from the superconducting transition temperature. In the approach of Chang *et al.* (1985), Dacorogna, Chang, and Cohen (1985), Dacorogna, Cohen, and Lam (1985), and Lam, Dacorogna, and Cohen (1986) the electron-phonon matrix element was calculated using

$$g_{mn\nu}(\mathbf{k}, \mathbf{q}) = \langle u_{m\mathbf{k}+\mathbf{q}} | \Delta_{\mathbf{q}\nu} v^{\text{KS}} | u_{n\mathbf{k}} \rangle_{\text{uc}}, \quad (11)$$

with  $u_{n\mathbf{k}}$  and  $u_{m\mathbf{k}+\mathbf{q}}$  being the Bloch-periodic components of the Kohn-Sham electron wave functions,  $\Delta_{\mathbf{q}\nu} v^{\text{KS}}$  is the phonon-induced variation of the self-consistent potential experienced by the electrons, and the integral extending over one unit cell. Equation (11) is discussed in Sec. III.B.2. The scattering potential  $\Delta_{\mathbf{q}\nu} v^{\text{KS}}$  was calculated by explicitly taking into account the rearrangement of the electronic charge following a small displacement of the nuclei. The inclusion of the self-consistent response of the electrons constitutes a considerable step forward beyond the rigid-ion approximation of Sec. II.B.

The next and most recent step in the evolution of electron-phonon calculations came with the development of DFPT for lattice dynamics (Baroni, Giannozzi, and Testa, 1987; Gonze, Allan, and Teter, 1992; Savrasov, 1992). In contrast to frozen-phonon calculations, which may require large supercells, DFPT enables the calculations of vibrational frequencies and eigenmodes at arbitrary wave vectors in the Brillouin zone. This innovation was critical in the context of electron-phonon physics, since the calculation of many physical quantities requires the evaluation of nontrivial integrals over the Brillouin zone. The first calculations of EPIs using DFPT were reported by Savrasov, Savrasov, and Andersen (1994), Liu and Quong (1996), Mauri *et al.* (1996), and Bauer *et al.* (1998). They calculated the electrical resistivity, thermal conductivity, mass enhancement, and superconducting critical temperature of a number of elemental metals (e.g., Al, Au, Cu, Mo, Nb, Pb, Pd, Ta, V, and Te), and reported good agreement with experiment.

By the late 1990s most of the basic ingredients required for the *ab initio* calculation of EPIs were available. Subsequent studies focused on using these techniques for calculating a variety of material properties and on improving the efficiency

and accuracy of the methodology. The most recent advances are reviewed in Secs. VI–XII.

### III. ELECTRON-PHONON INTERACTION IN DENSITY-FUNCTIONAL THEORY

In this section we review the basic formalism underlying the calculation of EPIs using DFT, and we establish the link with the Hamiltonian in Eq. (1). We start by introducing the standard formalism for lattice vibrations and the electron-phonon coupling Hamiltonian. Then we briefly summarize established methods of DFPT for calculating electron-phonon matrix elements. For the time being we describe electrons and phonons as separate subsystems; a rigorous theoretical framework for addressing the coupled electron-phonon system will be discussed in Sec. IV.

#### A. Lattice vibrations in crystals

The formalism for studying lattice dynamics in crystals is covered in many excellent textbooks such as Born and Huang (1954), Ziman (1960), Kittel (1963, 1976), and Ashcroft and Mermin (1976). Here we introduce the notation and summarize those aspects which will be useful for subsequent discussions in this section and in Secs. IV and V.

We consider  $M$  nuclei or ions in the unit cell. The position vector and Cartesian coordinates of the nucleus  $\kappa$  in the primitive unit cell are denoted by  $\boldsymbol{\tau}_\kappa$  and  $\tau_{\kappa\alpha}$ , respectively. We describe the infinitely extended solid using BvK boundary conditions. In this approach, periodic boundary conditions are applied to a large supercell which contains  $N_p$  unit cells, identified by the direct lattice vectors  $\mathbf{R}_p$ , with  $p = 1, \dots, N_p$ . The position of the nucleus  $\kappa$  belonging to the unit cell  $p$  is indicated by  $\boldsymbol{\tau}_{\kappa p} = \mathbf{R}_p + \boldsymbol{\tau}_\kappa$ . The Bloch wave vectors  $\mathbf{q}$  are taken to define a uniform grid of  $N_p$  points in one unit cell of the reciprocal lattice, and the vectors of the reciprocal lattice are indicated by  $\mathbf{G}$ . In Appendix A we provide additional details on the notation, and we state the Fourier transforms between the direct and reciprocal lattices.

Using standard DFT techniques it is possible to calculate the total potential energy of electrons and nuclei in the BvK supercell. This quantity is denoted as  $U(\{\boldsymbol{\tau}_{\kappa p}\})$ , where the braces are a short-hand notation for the coordinates of all the ions. The total potential energy refers to electrons in their ground state, with the nuclei being represented as *classical* particles *clamped* at the coordinates  $\boldsymbol{\tau}_{\kappa p}$ . Every DFT software package available today provides the quantity  $U$  as a standard output.

In order to study lattice vibrations, one begins by making the *harmonic* approximation. Accordingly, the total potential energy is expanded to second order in the displacements  $\Delta\boldsymbol{\tau}_{\kappa p}$  of the ions in the BvK supercell away from their equilibrium positions  $\boldsymbol{\tau}_{\kappa p}^0$ :

$$U = U_0 + \frac{1}{2} \sum_{\substack{\kappa\alpha p \\ \kappa'\alpha' p'}} \frac{\partial^2 U}{\partial\tau_{\kappa\alpha p} \partial\tau_{\kappa'\alpha' p'}} \Delta\tau_{\kappa\alpha p} \Delta\tau_{\kappa'\alpha' p'}, \quad (12)$$

where  $U_0$  denotes the total energy calculated for the ions in their equilibrium positions, and the derivatives are evaluated

for the equilibrium structure. The second derivatives of the total energy with respect to the nuclear coordinates define the matrix of “interatomic force constants”:

$$C_{\kappa\alpha p, \kappa'\alpha' p'} = \partial^2 U / \partial\tau_{\kappa\alpha p} \partial\tau_{\kappa'\alpha' p'}. \quad (13)$$

The Fourier transform of the interatomic force constants yields the “dynamical matrix” (Maradudin and Vosko, 1968):

$$D_{\kappa\alpha, \kappa'\alpha'}^{\text{dm}}(\mathbf{q}) = (M_\kappa M_{\kappa'})^{-1/2} \sum_p C_{\kappa\alpha 0, \kappa'\alpha' p} \exp(i\mathbf{q} \cdot \mathbf{R}_p), \quad (14)$$

where  $M_\kappa$  is the mass of the  $\kappa$ th ion. The superscript “dm” is there to distinguish this quantity from the many-body phonon propagators  $D(12)$  and  $D_{\kappa\alpha p, \kappa'\alpha' p'}$  that will be introduced in Sec. IV.B. The dynamical matrix is Hermitian and therefore admits real eigenvalues, which we denote as  $\omega_{\mathbf{q}\nu}^2$ :

$$\sum_{\kappa'\alpha'} D_{\kappa\alpha, \kappa'\alpha'}^{\text{dm}}(\mathbf{q}) e_{\kappa'\alpha', \nu}(\mathbf{q}) = \omega_{\mathbf{q}\nu}^2 e_{\kappa\alpha, \nu}(\mathbf{q}). \quad (15)$$

In classical mechanics, each  $\omega_{\mathbf{q}\nu}$  corresponds to the vibrational frequency of an independent harmonic oscillator. The Hermiticity of the dynamical matrix allows us to choose the eigenvectors  $e_{\kappa\alpha, \nu}(\mathbf{q})$  to be orthonormal for each  $\mathbf{q}$ :

$$\sum_\nu e_{\kappa'\alpha', \nu}^*(\mathbf{q}) e_{\kappa\alpha, \nu}(\mathbf{q}) = \delta_{\kappa\kappa'} \delta_{\alpha\alpha'}, \quad (16)$$

$$\sum_{\kappa\alpha} e_{\kappa\alpha, \nu}^*(\mathbf{q}) e_{\kappa\alpha, \nu'}(\mathbf{q}) = \delta_{\nu\nu'}. \quad (17)$$

Here the index  $\nu$  runs from 1 to  $3M$ . The column vectors  $e_{\kappa\alpha, \nu}(\mathbf{q})$  for a given  $\nu$  are called the “normal modes of vibration” or the “polarization” of the vibration wave. The following relations can be derived from Eq. (14):

$$\omega_{-\mathbf{q}\nu}^2 = \omega_{\mathbf{q}\nu}^2, \quad e_{\kappa\alpha, \nu}(-\mathbf{q}) = e_{\kappa\alpha, \nu}^*(\mathbf{q}). \quad (18)$$

These relations between normal modes carry a degree of arbitrariness in the choice of phases; here we have chosen to follow the same phase convention as Maradudin and Vosko (1968).

Using Eqs. (12) and (13) the Hamiltonian for nuclei considered as quantum particles can be written as

$$\hat{H}_p = \frac{1}{2} \sum_{\substack{\kappa\alpha p \\ \kappa'\alpha' p'}} C_{\kappa\alpha p, \kappa'\alpha' p'} \Delta\tau_{\kappa\alpha p} \Delta\tau_{\kappa'\alpha' p'} - \sum_{\kappa\alpha p} \frac{\hbar^2}{2M_\kappa} \frac{\partial^2}{\partial\tau_{\kappa\alpha p}^2}, \quad (19)$$

where the ground-state energy  $U_0$  has been omitted and the last term is the kinetic energy operator. The Hamiltonian in Eq. (19) corresponds to the energy of an entire BvK supercell. Equation (19) relies on two approximations: (i) the harmonic approximation, which coincides with the truncation of Eq. (12) to second order in the displacements; and (ii) the Born-Oppenheimer *adiabatic* approximation. This latter approximation is made when one calculates the interatomic force constants with the electrons in their ground state. The scope and validity of the adiabatic approximation are

discussed in detail in Sec. V.A.1. We note incidentally that, strictly speaking, the Born-Oppenheimer approximation does not need to be invoked were one to use the generalization of DFT to multicomponent systems introduced by Kreibich and Gross (2001) and Kreibich, van Leeuwen, and Gross (2008).

For practical purposes it is convenient to rewrite Eq. (19) by introducing the quanta of lattice vibrations. This is accomplished by defining the standard creation ( $\hat{a}_{\mathbf{q}\nu}^\dagger$ ) and destruction ( $\hat{a}_{\mathbf{q}\nu}$ ) operators for each phonon of energy  $\hbar\omega_{\mathbf{q}\nu}$  and polarization  $e_{\kappa\alpha,\nu}(\mathbf{q})$ . This operation is not entirely trivial and is described in detail in Appendix B. The formal definition of the ladder operators is given in Eqs. (B13) and (B14). These operators obey the commutation relations  $[\hat{a}_{\mathbf{q}\nu}, \hat{a}_{\mathbf{q}'\nu'}^\dagger] = \delta_{\nu\nu'}\delta_{\mathbf{q}\mathbf{q}'}$  and  $[\hat{a}_{\mathbf{q}\nu}, \hat{a}_{\mathbf{q}'\nu'}] = [\hat{a}_{\mathbf{q}\nu}^\dagger, \hat{a}_{\mathbf{q}'\nu'}^\dagger] = 0$ , where  $\delta$  is the Kronecker symbol. From these relations we know that the quanta of the harmonic oscillations in crystals obey Bose-Einstein statistics. Appendix B shows that the atomic displacements can be expressed in terms of the ladder operators as follows:

$$\Delta\tau_{\kappa\alpha p} = \left(\frac{M_0}{N_p M_\kappa}\right)^{1/2} \sum_{\mathbf{q}} e^{i\mathbf{q}\cdot\mathbf{R}_p} e_{\kappa\alpha,\nu}(\mathbf{q}) l_{\mathbf{q}\nu} (\hat{a}_{\mathbf{q}\nu} + \hat{a}_{-\mathbf{q}\nu}^\dagger), \quad (20)$$

where  $l_{\mathbf{q}\nu}$  is the ‘‘zero-point’’ displacement amplitude:

$$l_{\mathbf{q}\nu} = [\hbar/(2M_0\omega_{\mathbf{q}\nu})]^{1/2}. \quad (21)$$

Here  $M_0$  is an arbitrary reference mass which is introduced to ensure that  $l_{\mathbf{q}\nu}$  has the dimensions of a length and is similar in magnitude to  $\Delta\tau_{\kappa\alpha p}$ . Typically  $M_0$  is chosen to be the proton mass.

Using Eqs. (13)–(21) the nuclear Hamiltonian can be written in terms of  $3MN_p$  independent harmonic oscillators as follows:

$$\hat{H}_p = \sum_{\mathbf{q}\nu} \hbar\omega_{\mathbf{q}\nu} (\hat{a}_{\mathbf{q}\nu}^\dagger \hat{a}_{\mathbf{q}\nu} + 1/2), \quad (22)$$

where the sum is over all wave vectors. The ground-state wave function of this Hamiltonian is a product of Gaussians, and all other states can be generated by acting on the ground state with the operators  $\hat{a}_{\mathbf{q}\nu}^\dagger$ . In the case of  $|\mathbf{q}| = 0$  there are three normal modes for which  $\omega_{\mathbf{q}\nu} = 0$ . For these modes, which correspond to global translations of the crystal, the zero-point displacement  $l_{\mathbf{q}\nu}$  is not defined. Throughout this article it is assumed that these modes are skipped in summations containing zero-point amplitudes. A detailed derivation of Eq. (22) and a discussion of the eigenstates of  $\hat{H}_p$  are provided in Appendix B.

## B. Electron-phonon coupling Hamiltonian

Having outlined the standard formalism for addressing lattice vibrations in crystals, we now proceed to make the connection between DFT calculations and the remaining terms of Eq. (1). The electronic band structure  $\varepsilon_{n\mathbf{k}}$  and

electron-phonon matrix elements  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$  are almost invariably calculated by using the KS Hamiltonian (Hohenberg and Kohn, 1964; Kohn and Sham, 1965). A justification for these choices is provided in Sec. V; for now we limit ourselves to outline the key elements of practical calculations.

### 1. Kohn-Sham Hamiltonian

Let us denote the KS eigenfunctions by  $\psi_{n\mathbf{k}}(\mathbf{r})$  and use  $\mathbf{k}$  to indicate both the wave vector and the spin. We restrict ourselves to systems with collinear spins. The KS eigenfunctions satisfy the equation  $\hat{H}^{\text{KS}}\psi_{n\mathbf{k}}(\mathbf{r}) = \varepsilon_{n\mathbf{k}}\psi_{n\mathbf{k}}(\mathbf{r})$ , with the Hamiltonian given by

$$\hat{H}^{\text{KS}} = -\frac{\hbar^2}{2m_e}\nabla^2 + V^{\text{KS}}(\mathbf{r}; \{\tau_{\kappa\alpha p}\}). \quad (23)$$

Here the potential  $V^{\text{KS}}$  is the sum of the nuclear (or ionic) contribution  $V^{\text{en}}$ , the Hartree electronic screening  $V^{\text{H}}$ , and the exchange and correlation potential  $V^{\text{xc}}$  (Martin, 2004):

$$V^{\text{KS}} = V^{\text{en}} + V^{\text{H}} + V^{\text{xc}}. \quad (24)$$

The potentials appearing in Eq. (24) are defined as follows. The electron-nuclei potential energy is given by

$$V^{\text{en}}(\mathbf{r}; \{\tau_{\kappa\alpha p}\}) = \sum_{\kappa p, \mathbf{T}} V_\kappa(\mathbf{r} - \tau_{\kappa p} - \mathbf{T}), \quad (25)$$

where  $V_\kappa(\mathbf{r})$  is the interaction between an electron and the nucleus  $\kappa$  located at the center of the reference frame, and  $\mathbf{T}$  denotes a lattice vector of the BvK supercell. In the case of all-electron DFT calculations,  $V_\kappa(\mathbf{r})$  is the Coulomb interaction:

$$V_\kappa(\mathbf{r}) = -\frac{e^2}{4\pi\epsilon_0} \frac{Z_\kappa}{|\mathbf{r}|}, \quad (26)$$

where  $Z_\kappa$  is the atomic number of the nucleus  $\kappa$ . In the case of pseudopotential implementations  $V_\kappa$  is a function that goes as in Eq. (26) at large  $|\mathbf{r}|$ , but remains finite at  $|\mathbf{r}| = 0$ . Furthermore the nuclear charge is replaced by the ionic charge, which is the difference between the nuclear charge and the number of core electrons described by the pseudopotential. In all modern pseudopotential implementations  $V_\kappa(\mathbf{r})$  is nonlocal due to the separation of the angular momentum channels (Martin, 2004). However, since this nonlocality is short ranged and is inconsequential in the following discussion, it will be ignored here in order to maintain a light notation. The Hartree term is obtained from the electron density  $n(\mathbf{r}'; \{\tau_{\kappa\alpha p}\})$ :

$$V^{\text{H}}(\mathbf{r}; \{\tau_{\kappa\alpha p}\}) = \frac{e^2}{4\pi\epsilon_0} \sum_{\mathbf{T}} \int_{\text{sc}} \frac{n(\mathbf{r}'; \{\tau_{\kappa\alpha p}\})}{|\mathbf{r} - \mathbf{r}' - \mathbf{T}|} d\mathbf{r}', \quad (27)$$

where the integral extends over the supercell. The exchange and correlation potential is the functional derivative of the exchange and correlation energy with respect to the electron density (Kohn and Sham, 1965):



$$V^{xc}(\mathbf{r}; \{\tau_{\kappa\alpha p}\}) = \delta E^{xc}[n]/\delta n|_{n(\mathbf{r}; \{\tau_{\kappa\alpha p}\})}. \quad (28)$$

The eigenfunctions  $\psi_{n\mathbf{k}}$  of  $\hat{H}^{\text{KS}}$  can be expressed in the following Bloch form:

$$\psi_{n\mathbf{k}}(\mathbf{r}) = N_p^{-1/2} u_{n\mathbf{k}}(\mathbf{r}) e^{i\mathbf{k}\cdot\mathbf{r}}, \quad (29)$$

with  $u_{n\mathbf{k}}$  a lattice-periodic function. The wave function  $\psi_{n\mathbf{k}}$  is taken to be normalized to one in the supercell, while the periodic part  $u_{n\mathbf{k}}(\mathbf{r})$  is normalized to one in the crystal unit cell. The electron density is  $n(\mathbf{r}) = \sum_{v\mathbf{k}} |\psi_{v\mathbf{k}}(\mathbf{r})|^2$ , where  $v$  indicates occupied states. In order to determine  $\psi_{n\mathbf{k}}$  and  $\varepsilon_{n\mathbf{k}}$  the Kohn-Sham equations are solved self-consistently. This requires one to start from a reasonable guess for the electron density (for example, a superposition of atomic electron densities), calculate the potentials in Eq. (24), and determine the solutions of the KS Hamiltonian in Eq. (23). The electron density is recalculated using these solutions, and the cycle is repeated until convergence.

In order to establish the link with Eq. (1), we can regard the KS Hamiltonian as an effective one-body operator and make the transition to a second quantized formalism by using the standard prescription (Merzbacher, 1998):

$$\hat{H}_e = \sum_{n\mathbf{k}, n'\mathbf{k}'} \langle \psi_{n\mathbf{k}} | \hat{H}^{\text{KS}} | \psi_{n'\mathbf{k}'} \rangle \hat{c}_{n\mathbf{k}}^\dagger \hat{c}_{n'\mathbf{k}'} = \sum_{n\mathbf{k}} \varepsilon_{n\mathbf{k}} \hat{c}_{n\mathbf{k}}^\dagger \hat{c}_{n\mathbf{k}}. \quad (30)$$

This expression is useful for performing formal manipulations in the study of coupled electron-phonon systems. However, Eq. (30) implicitly introduces the drastic approximation that the electronic system can be described in terms of sharp quasiparticle excitations. A field-theoretic approach that does not rely on any such approximation is discussed in Sec. IV.

## 2. Electron-phonon coupling Hamiltonian to first and second order in the atomic displacements

Within the DFT Kohn-Sham formalism, the coupling Hamiltonian appearing in the second line of Eq. (1) is obtained by expanding the Kohn-Sham effective potential in terms of the nuclear displacements  $\Delta\tau_{\kappa p}$  from their equilibrium positions  $\tau_{\kappa p}^0$ . The potential to first order in the displacements is

$$V^{\text{KS}}(\{\tau_{\kappa p}\}) = V^{\text{KS}}(\{\tau_{\kappa p}^0\}) + \sum_{\kappa\alpha p} \frac{\partial V^{\text{KS}}}{\partial \tau_{\kappa\alpha p}} \Delta\tau_{\kappa\alpha p}. \quad (31)$$

This expression can be rewritten into normal mode coordinates using Eq. (20):

$$V^{\text{KS}} = V^{\text{KS}}(\{\tau_{\kappa p}^0\}) + N_p^{-1/2} \sum_{\mathbf{q}\nu} \Delta_{\mathbf{q}\nu} V^{\text{KS}}(\hat{a}_{\mathbf{q}\nu} + \hat{a}_{-\mathbf{q}\nu}^\dagger), \quad (32)$$

having defined

$$\Delta_{\mathbf{q}\nu} V^{\text{KS}} = e^{i\mathbf{q}\cdot\mathbf{r}} \Delta_{\mathbf{q}\nu} v^{\text{KS}}, \quad (33)$$

$$\Delta_{\mathbf{q}\nu} v^{\text{KS}} = l_{\mathbf{q}\nu} \sum_{\kappa\alpha} (M_0/M_\kappa)^{1/2} e_{\kappa\alpha,\nu}(\mathbf{q}) \partial_{\kappa\alpha,\nu} v^{\text{KS}}, \quad (34)$$

$$\partial_{\kappa\alpha,\nu} v^{\text{KS}} = \sum_p e^{-i\mathbf{q}\cdot(\mathbf{r}-\mathbf{R}_p)} \frac{\partial V^{\text{KS}}}{\partial \tau_{\kappa\alpha}} \Big|_{\mathbf{r}-\mathbf{R}_p}. \quad (35)$$

From the last expression we see that  $\partial_{\kappa\alpha,\nu} v^{\text{KS}}$  and  $\Delta_{\mathbf{q}\nu} v^{\text{KS}}$  are lattice-periodic functions. The transition to second quantization is performed as in Eq. (30) (Merzbacher, 1998):

$$\hat{H}_{\text{ep}} = \sum_{n\mathbf{k}, n'\mathbf{k}'} \langle \psi_{n\mathbf{k}} | V^{\text{KS}}(\{\tau_{\kappa p}\}) - V^{\text{KS}}(\{\tau_{\kappa p}^0\}) | \psi_{n'\mathbf{k}'} \rangle \hat{c}_{n\mathbf{k}}^\dagger \hat{c}_{n'\mathbf{k}'}, \quad (36)$$

where the bra and ket indicate an integral over the supercell. After using Eqs. (29), (32)–(35), and (A1) we have

$$\hat{H}_{\text{ep}} = N_p^{-1/2} \sum_{\substack{\mathbf{k}, \mathbf{q} \\ m\nu}} g_{m\nu}(\mathbf{k}, \mathbf{q}) \hat{c}_{m\mathbf{k}+\mathbf{q}}^\dagger \hat{c}_{n\mathbf{k}} (\hat{a}_{\mathbf{q}\nu} + \hat{a}_{-\mathbf{q}\nu}^\dagger), \quad (37)$$

where the electron-phonon matrix element is given by

$$g_{m\nu}(\mathbf{k}, \mathbf{q}) = \langle u_{m\mathbf{k}+\mathbf{q}} | \Delta_{\mathbf{q}\nu} v^{\text{KS}} | u_{n\mathbf{k}} \rangle_{\text{uc}}. \quad (38)$$

Here the subscript ‘‘uc’’ indicates that the integral is carried out within one unit cell of the crystal. The coupling Hamiltonian in Eq. (37) yields the energy of an entire supercell. In the case of the three translational modes at  $|\mathbf{q}| = 0$  we set the matrix elements  $g_{m\nu}(\mathbf{k}, \mathbf{q})$  to zero, as a consequence of the *acoustic sum rule* (see discussion in Sec. IX.A.1).

Taken together, Eqs. (22), (30), and (37) constitute the starting point of most first-principles calculations of electron-phonon interactions. It remains to be seen how one calculates the electron-phonon matrix elements  $g_{m\nu}(\mathbf{k}, \mathbf{q})$ ; the most common procedures are described in Sec. III.B.3.

Before proceeding, we briefly discuss the second-order coupling Hamiltonian which appears in the third and fourth lines of Eq. (1). The rationale for incorporating this extra term is that the expansion of the Kohn-Sham potential to first order in the atomic displacements, Eq. (31), is somewhat inconsistent with the choice of expanding the total potential energy in Eq. (12) to second order in the atomic displacements. This aspect was discussed by Allen and Heine (1976) and Allen (1978). In order to obtain an electron-phonon coupling Hamiltonian including terms of second order in the displacements, we must include the second derivatives of the Kohn-Sham potential in Eq. (31) and follow the same steps which led to Eq. (37). By calling the extra term  $\hat{H}_{\text{ep}}^{(2)}$  we have

$$\hat{H}_{\text{ep}}^{(2)} = N_p^{-1} \sum_{\substack{\mathbf{k}, \mathbf{q}, \mathbf{q}' \\ m\nu\nu'}} g_{m\nu\nu'}^{\text{DW}}(\mathbf{k}, \mathbf{q}, \mathbf{q}') \hat{c}_{m\mathbf{k}+\mathbf{q}+\mathbf{q}'}^\dagger \hat{c}_{n\mathbf{k}} \times (\hat{a}_{\mathbf{q}\nu} + \hat{a}_{-\mathbf{q}\nu}^\dagger) (\hat{a}_{\mathbf{q}'\nu'} + \hat{a}_{-\mathbf{q}'\nu'}^\dagger), \quad (39)$$

where

$$g_{m\nu\nu'}^{\text{DW}}(\mathbf{k}, \mathbf{q}, \mathbf{q}') = \frac{1}{2} \langle u_{m\mathbf{k}+\mathbf{q}+\mathbf{q}'} | \Delta_{\mathbf{q}\nu} \Delta_{\mathbf{q}'\nu'} v^{\text{KS}} | u_{n\mathbf{k}} \rangle_{\text{uc}}. \quad (40)$$

The variations  $\Delta_{\mathbf{q}\nu}$  are the same as in Eqs. (33)–(35).

The second-order coupling Hamiltonian in Eq. (39) is considerably more involved than its first-order counterpart. The increased complexity partly explains why in the literature this term has largely been ignored. So far the Hamiltonian  $\hat{H}_{\text{ep}}^{(2)}$  has been described using only an approximation based on first-order perturbation theory (Allen and Heine, 1976). In this special case, the only terms in Eq. (39) that can modify the electron excitation spectrum are those with  $\mathbf{q}' = -\mathbf{q}$ . The corresponding energy shift is  $\Delta\varepsilon_{n\mathbf{k}} = N_p^{-1} \sum_{\mathbf{q}\nu} g_{m\nu\nu}^{\text{DW}}(\mathbf{k}, \mathbf{q}, -\mathbf{q})(2n_{\mathbf{q}\nu} + 1)$ , with  $n_{\mathbf{q}\nu}$  the number of phonons in each mode. We come back to this point in Sec. IX.A.1.

### 3. Calculation of electron-phonon matrix elements using density-functional perturbation theory

In this section we review how the scattering potential  $\Delta_{\mathbf{q}\nu} v^{\text{KS}}$  appearing in Eq. (38) is calculated in first-principles approaches. The most intuitive approach is to evaluate the derivatives appearing in Eq. (35) by using finite atomic displacements in a supercell:

$$\left. \frac{\partial V^{\text{KS}}}{\partial \tau_{\kappa\alpha p}^0} \right|_{\tau_{\kappa p}^0} \simeq [V^{\text{KS}}(\mathbf{r}; \tau_{\kappa\alpha p}^0 + b) - V^{\text{KS}}(\mathbf{r}; \tau_{\kappa\alpha p}^0)]/b. \quad (41)$$

In this expression  $b$  is a small displacement of the order of the zero-point amplitude ( $\sim 0.1 \text{ \AA}$ ), and the atom  $\kappa$  in the unit cell  $p$  is displaced along the direction  $\alpha$ . The first calculations of electron-phonon interactions within DFT employed a variant of this ‘‘supercell approach’’ whereby all atoms are displaced according to a chosen vibrational eigenmode (Chang *et al.*, 1985; Dacorogna, Chang, and Cohen, 1985; Dacorogna, Cohen, and Lam, 1985; Lam, Dacorogna, and Cohen, 1986). This strategy is usually referred to as the frozen-phonon method.

One disadvantage of the frozen-phonon method is that the supercell may become impractically large when evaluating matrix elements corresponding to long-wavelength phonons. This difficulty can be circumvented by using DFPT (Baroni, Giannozzi, and Testa, 1987; Gonze, Allan, and Teter, 1992; Savrasov, 1992). The main strength of DFPT is that the scattering potential  $\Delta_{\mathbf{q}\nu} v^{\text{KS}}$  in Eq. (38) is obtained by performing calculations within a single unit cell. Since the computational workload of standard (not linear-scaling) DFT calculations scales as the cube of the number of electrons, the saving afforded by DFPT over the frozen-phonon method is proportional to  $N_p^2$ , and typically corresponds to a factor  $> 10^3$ .

In the DFPT approach of Baroni *et al.* (2001) one calculates the lattice-periodic scattering potential  $\partial_{\kappa\alpha, \mathbf{q}} v^{\text{KS}}$  defined by Eq. (35). By differentiating Eq. (24) via Eq. (35) this potential is written as

$$\partial_{\kappa\alpha, \mathbf{q}} v^{\text{KS}} = \partial_{\kappa\alpha, \mathbf{q}} v^{\text{en}} + \partial_{\kappa\alpha, \mathbf{q}} v^{\text{H}} + \partial_{\kappa\alpha, \mathbf{q}} v^{\text{xc}}. \quad (42)$$

The variation of the ionic potential is obtained from Eqs. (25) and (35). The result is conveniently expressed in reciprocal space

$$\partial_{\kappa\alpha, \mathbf{q}} v^{\text{en}}(\mathbf{G}) = -i(\mathbf{q} + \mathbf{G})_{\alpha} V_{\kappa}(\mathbf{q} + \mathbf{G}) e^{-i(\mathbf{q} + \mathbf{G}) \cdot \tau_{\kappa}}, \quad (43)$$

where the convention for the Fourier transform is  $f(\mathbf{G}) = \Omega^{-1} \int_{\text{uc}} d\mathbf{r} e^{-i\mathbf{G} \cdot \mathbf{r}} f(\mathbf{r})$ , and  $\Omega$  is the volume of the unit cell. In order to keep the presentation as general as possible we avoid explicitly indicating the nonlocality of  $V_{\kappa}$  which arises in pseudopotential implementations. The adaptation of this equation and the following ones to the case of nonlocal pseudopotentials, ultrasoft pseudopotentials, and the projector-augmented wave method can be found in Giannozzi *et al.* (1991), Dal Corso, Pasquarello, and Baldereschi (1997), and Audouze *et al.* (2006), respectively. The variation of the Hartree and exchange-correlation contributions to the Kohn-Sham potential is obtained from the self-consistent charge density response to the perturbation in Eq. (43). After a few manipulations using Eqs. (27) and (35) one obtains

$$\partial_{\kappa\alpha, \mathbf{q}} v^{\text{H}}(\mathbf{G}) = \Omega v^{\text{C}}(\mathbf{q} + \mathbf{G}) \partial_{\kappa\alpha, \mathbf{q}} n(\mathbf{G}), \quad (44)$$

where  $v^{\text{C}}(\mathbf{q}) = \Omega^{-1} \int d\mathbf{r} e^{-i\mathbf{q} \cdot \mathbf{r}} e^2/4\pi\epsilon_0|\mathbf{r}|$  is the Fourier transform of the Coulomb potential. For the exchange and correlation potential we use Eq. (28) and the Taylor expansion of a functional to find

$$\partial_{\kappa\alpha, \mathbf{q}} v^{\text{xc}}(\mathbf{G}) = \Omega \sum_{\mathbf{G}'} f^{\text{xc}}(\mathbf{q} + \mathbf{G}, \mathbf{q} + \mathbf{G}') \partial_{\kappa\alpha, \mathbf{q}} n(\mathbf{G}'), \quad (45)$$

where  $f^{\text{xc}}$  indicates the standard exchange and correlation kernel, which is the second-order functional derivative of the exchange and correlation energy  $E^{\text{xc}}$  with respect to the electron density (Hohenberg and Kohn, 1964)

$$f^{\text{xc}}(\mathbf{r}, \mathbf{r}') = \left. \frac{\delta^2 E^{\text{xc}}[n]}{\delta n(\mathbf{r}) \delta n(\mathbf{r}')} \right|_{n(\mathbf{r}; \{\tau_{\kappa p}^0\})}. \quad (46)$$

In the case of the local density approximation (LDA) to DFT, the exchange and correlation kernel reduces to a local function (Parr and Yang, 1994), and Eq. (45) is more conveniently evaluated in real space. Today DFPT calculations can be performed using one of several exchange and correlation kernels. The effect of the kernel on the calculation of lattice-dynamical properties of solids has been analyzed in several works; see, for example, Dal Corso (2013) and He *et al.* (2014). The formal structure of the DFPT equations discussed in this section remains unchanged if we replace the DFT kernel in Eq. (46) by more sophisticated versions. For example, both DFPT calculations based on Hubbard-corrected DFT (Floris *et al.*, 2011) and DFPT coupled with dynamical mean-field theory (Savrasov and Kotliar, 2003) have been demonstrated.

It should be noted that in Eqs. (45) and (46) we are implicitly assuming a spin-unpolarized system. The adaptation of these equations as well as the other DFPT equations to the most general case of noncollinear spin systems can be found in Dal Corso (2007, 2008) and Verstraete *et al.* (2008).

From Eqs. (44) and (45) we see that the evaluation of  $g_{m\nu}(\mathbf{k}, \mathbf{q})$  goes through the calculation of the variation of the electron density induced by the perturbation  $\partial_{\kappa\alpha, \mathbf{q}} v^{\text{KS}}(\mathbf{r}) e^{i\mathbf{q} \cdot \mathbf{r}}$ . Within DFPT such a variation is obtained by evaluating the

change of the Kohn-Sham wave functions to first order in perturbation theory. After inspection of the perturbed Hamiltonian it becomes evident that the wave function change must be of the form  $\partial u_{n\mathbf{k},\mathbf{q}} e^{i\mathbf{q}\cdot\mathbf{r}}$ , with  $\partial u_{n\mathbf{k},\mathbf{q}}$  a lattice-periodic function. Using this observation the first-order variation of the Kohn-Sham equations can be written as a Sternheimer equation (Sternheimer, 1954)

$$(\hat{H}_{\mathbf{k}+\mathbf{q}}^{\text{KS}} - \varepsilon_{v\mathbf{k}}) \partial u_{v\mathbf{k},\mathbf{q}} = -\partial_{\kappa\alpha,\mathbf{q}} v^{\text{KS}} u_{v\mathbf{k}}, \quad (47)$$

where  $\hat{H}_{\mathbf{k}+\mathbf{q}}^{\text{KS}} = e^{-i(\mathbf{k}+\mathbf{q})\cdot\mathbf{r}} \hat{H}^{\text{KS}} e^{i(\mathbf{k}+\mathbf{q})\cdot\mathbf{r}}$ . In this equation the index  $v$  indicates an occupied state. For  $|\mathbf{q}| = 0$  one needs also to consider a shift of the energy eigenvalues which introduces an additional term  $\langle u_{v\mathbf{k}} | \partial_{\kappa\alpha,0} v | u_{v\mathbf{k}} \rangle_{\text{uc}} u_{v\mathbf{k}}$  on the right-hand side of Eq. (47). In practice this term is canceled by the use of the projectors described in Eq. (48), unless one is dealing with metallic systems. This aspect is discussed in detail by de Gironcoli (1995) and Baroni *et al.* (2001). The principal advantage of Eq. (47) over standard perturbation theory is that it does not involve unoccupied electronic states.

A practical problem arises when attempting to solve Eq. (47): the linear system on the left-hand side is ill conditioned owing to small eigenvalues corresponding to  $\varepsilon_{v\mathbf{k}} \approx \varepsilon_{v'\mathbf{k}+\mathbf{q}}$ ; furthermore, in the case of accidental degeneracies  $\varepsilon_{v\mathbf{k}} = \varepsilon_{v'\mathbf{k}+\mathbf{q}}$ , the system becomes singular. In order to make the system nonsingular, Giannozzi *et al.* (1991) noted that the variation of the electron density involves only the component of  $\partial u_{v\mathbf{k},\mathbf{q}}$  belonging to the unoccupied manifold of Kohn-Sham states. As a consequence, what is really needed is only  $\partial \tilde{u}_{v\mathbf{k},\mathbf{q}} = (1 - \hat{P}_{\mathbf{k}+\mathbf{q}}^{\text{occ}}) \partial u_{v\mathbf{k},\mathbf{q}}$ , where  $\hat{P}_{\mathbf{k}+\mathbf{q}}^{\text{occ}} = \sum_v |u_{v\mathbf{k}+\mathbf{q}}\rangle \langle u_{v\mathbf{k}+\mathbf{q}}|$  is the projector over the occupied states with wave vector  $\mathbf{k} + \mathbf{q}$ . The equation for this ‘‘trimmed’’ wave function variation is simply obtained by projecting both sides of Eq. (47) onto  $(1 - \hat{P}_{\mathbf{k}+\mathbf{q}}^{\text{occ}})$  and noting that  $\hat{P}_{\mathbf{k}+\mathbf{q}}^{\text{occ}}$  and  $\hat{H}_{\mathbf{k}+\mathbf{q}}^{\text{KS}}$  do commute

$$(\hat{H}_{\mathbf{k}+\mathbf{q}}^{\text{KS}} - \varepsilon_{v\mathbf{k}}) \partial \tilde{u}_{v\mathbf{k},\mathbf{q}} = -(1 - \hat{P}_{\mathbf{k}+\mathbf{q}}^{\text{occ}}) \partial_{\kappa\alpha,\mathbf{q}} v^{\text{KS}} u_{v\mathbf{k}}. \quad (48)$$

At this point it is possible to remove all small or null eigenvalues of the operator on the left-hand side by adding a term  $\alpha \hat{P}_{\mathbf{k}+\mathbf{q}}^{\text{occ}}$  to the Hamiltonian. This term has no effect on the wave function variation, since  $\hat{P}_{\mathbf{k}+\mathbf{q}}^{\text{occ}} \partial \tilde{u}_{v\mathbf{k},\mathbf{q}} = 0$  by construction. The operator is made nonsingular by choosing the parameter  $\alpha$  larger than the valence bandwidth (Baroni *et al.*, 2001). From the wave function variation obtained by solving Eq. (48), it is now possible to construct the following density response associated with the wave vector  $\mathbf{q}$ :

$$\partial n_{\kappa\alpha,\mathbf{q}}(\mathbf{r}) = 2N_p^{-1} \sum_{v\mathbf{k}} u_{v\mathbf{k}}^* \partial \tilde{u}_{v\mathbf{k},\mathbf{q}}. \quad (49)$$

For simplicity a spin-degenerate system has been assumed (a factor of 2 is implicitly included in the sum over  $\mathbf{k}$ ), and time-reversal symmetry has been used in order to make the expression more compact (yielding the factor of 2 on the right-hand side).

In practical DFPT calculations, Eq. (48) is solved using an iterative procedure which is similar to standard DFT total

energy calculations. One sets the starting perturbation  $\partial_{\kappa\alpha,\mathbf{q}} v^{\text{KS}}$  to be equal to the electron-nuclei potential in Eq. (43). By solving Eq. (48) for each occupied state  $v$  and each wave vector  $\mathbf{k}$  using standard linear algebra techniques, one obtains the induced density in Eq. (49). The new density is now used to construct the variations of the Hartree and exchange-correlation potentials in Eqs. (44) and (45). These induced potentials are added to the electron-nuclei potential, yielding a ‘‘screened’’ perturbation  $\partial_{\kappa\alpha,\mathbf{q}} v^{\text{KS}}$  in Eq. (48). The cycle is repeated until the change of  $\partial n_{\kappa\alpha,\mathbf{q}}$  between two successive cycles is smaller than a set tolerance.

It can be shown that the screened perturbation  $\partial_{\kappa\alpha,\mathbf{q}} v^{\text{KS}}$  described in this section is also the key ingredient required for calculating the interatomic force constants in Eq. (13) (Baroni *et al.*, 2001). As a practical consequence, every software implementation that supports DFPT calculations already contains all the information necessary for evaluating the electron-phonon matrix elements  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$ .

All the quantities introduced in this section can equivalently be calculated using an alternative, variational formulation of density-functional perturbation theory (Gonze, Allan, and Teter, 1992; Gonze, 1995a, 1997; Gonze and Lee, 1997). A thorough discussion of the connection between the Sternheimer approach and the variational approach to DFPT is provided by Gonze (1995b).

The second-order matrix elements  $g_{mn\nu}^{\text{DW}}(\mathbf{k}, \mathbf{q}, \mathbf{q}')$  given by Eq. (40) involve the second derivative of the Kohn-Sham potential with respect to the nuclear displacements. The evaluation of these quantities requires the solution of second-order Sternheimer equations for the second variations of the Kohn-Sham wave functions. The general structure of second-order Sternheimer equations can be found in Sec. IV.H of Gonze (1995b). Since these calculations are rather involved, most practical implementations employ an approximation whereby the Debye-Waller matrix elements are expressed in terms of products of the standard matrix elements  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$ . Such an alternative formulation was developed by Allen and Heine (1976) and Allen and Cardona (1981) and is discussed in Sec. IX.A.1. All recent *ab initio* calculations of electron-phonon interactions based on DFPT employed this latter approach.<sup>1</sup>

#### 4. The dielectric approach

Besides the DFPT method described in the previous section, it is also possible to calculate the screened perturbation  $\partial_{\kappa\alpha,\mathbf{q}} v^{\text{KS}}$  using the so-called ‘‘dielectric approach’’ (Pick, Cohen, and Martin, 1970; Quong and Klein, 1992). This latter approach did not find as widespread an application as those of Baroni, Giannozzi, and Testa (1987), Gonze, Allan, and Teter (1992), and Savrasov (1992), but it is useful to establish a link between DFT calculations of electron-phonon matrix elements and the field-theoretic formulation discussed in Sec. IV.

<sup>1</sup>See, for example, Marini (2008), Giustino, Louie, and Cohen (2010), Gonze, Boulanger, and Côté (2011), Cannuccia and Marini (2013), Antonius *et al.* (2014), Kawai *et al.* (2014), Poncé, Antonius, Boulanger *et al.* (2014), Poncé, Antonius, Gillet *et al.* (2014), and Poncé *et al.* (2015).

For consistency with Sec. III.B.3, we derive the key expressions of the dielectric approach starting from DFPT. To this aim we expand the variation of the wave function  $\partial \tilde{u}_{v\mathbf{k},\mathbf{q}}$  using the complete set of states  $u_{n\mathbf{k}+\mathbf{q}}$  (with  $n$  referring to both occupied and empty Kohn-Sham states). Then we replace this expansion inside Eq. (48), project onto an arbitrary conduction state, and insert the result into Eq. (49). After taking into account time-reversal symmetry, these steps lead to the following result:

$$\partial_{\kappa\alpha,\mathbf{q}} n(\mathbf{r}) = \int_{\text{uc}} d\mathbf{r}' \chi_{\mathbf{q}}^0(\mathbf{r}, \mathbf{r}') \partial_{\kappa\alpha,\mathbf{q}} v^{\text{KS}}(\mathbf{r}'), \quad (50)$$

having defined

$$\begin{aligned} \chi_{\mathbf{q}}^0(\mathbf{r}, \mathbf{r}') &= N_p^{-1} \sum_{m\mathbf{k}} \frac{f_{n\mathbf{k}} - f_{m\mathbf{k}+\mathbf{q}}}{\varepsilon_{n\mathbf{k}} - \varepsilon_{m\mathbf{k}+\mathbf{q}}} \\ &\times u_{n\mathbf{k}}^*(\mathbf{r}) u_{m\mathbf{k}+\mathbf{q}}(\mathbf{r}) u_{m\mathbf{k}+\mathbf{q}}^*(\mathbf{r}') u_{n\mathbf{k}}(\mathbf{r}'). \end{aligned} \quad (51)$$

In this expression  $f_{n\mathbf{k}}$  and  $f_{m\mathbf{k}+\mathbf{q}}$  are the occupations of each state, and the indices run over all bands. A factor of 2 for the spin degeneracy is implicitly included in the sum over  $\mathbf{k}$ . The quantity  $\chi_{\mathbf{q}}^0$  in Eq. (51) is the lattice-periodic component for the wave vector  $\mathbf{q}$  of the “independent-electron polarizability” (Adler, 1962; Wiser, 1963; Pick, Cohen, and Martin, 1970; Quong and Klein, 1992).

For ease of notation we write Eq. (50) in symbolic form as  $\partial n = \chi^0 \partial v^{\text{KS}}$ . Using the same symbolic notation it is also possible to formally rewrite Eqs. (42), (44), and (45) as follows:

$$\partial v^{\text{KS}} = \partial v^{\text{en}} + (v^{\text{C}} + f^{xc}) \chi^0 \partial v^{\text{KS}}, \quad (52)$$

from which one obtains

$$\partial v^{\text{KS}} = (e^{\text{Hxc}})^{-1} \partial v^{\text{en}}, \quad (53)$$

having defined the dielectric matrix

$$e^{\text{Hxc}} = 1 - (v^{\text{C}} + f^{xc}) \chi^0. \quad (54)$$

The superscript “Hxc” refers to the Hartree and exchange and correlation components of the screening. In the language of many-body perturbation theory  $e^{\text{Hxc}}$  is referred to as the “test electron” dielectric matrix, hinting at the fact that the electron density redistribution in response to a perturbation arises both from classical electrostatics (the Hartree term  $v^{\text{C}} \chi^0$ ) and from quantum effects (the exchange and correlation terms  $f^{xc} \chi^0$ ). If we neglect the kernel  $f^{xc}$  in this expression, then we obtain the “test charge” dielectric matrix, which is most commonly known as the dielectric matrix in the random-phase approximation (RPA) (Pines and Bohm, 1952)

$$e^{\text{H}} = 1 - v^{\text{C}} \chi_0. \quad (55)$$

The symbolic expressions outlined here remain almost unchanged when using a reciprocal space representation. As an example, Eq. (55) becomes simply

$$e_{\mathbf{G}\mathbf{G}'}^{\text{H}}(\mathbf{q}) = \delta_{\mathbf{G}\mathbf{G}'} - \Omega^2 \sum_{\mathbf{G}''} \chi_{\mathbf{G}''\mathbf{G}'}^0(\mathbf{q}) v^{\text{C}}(\mathbf{q} + \mathbf{G}) \delta_{\mathbf{G}\mathbf{G}''}. \quad (56)$$

Taken together Eqs. (38) and (53) show that the calculation of electron-phonon matrix elements using DFPT is equivalent to screening the bare electron-nucleus interaction using  $e^{\text{Hxc}}$ . In this case we say that the screening is described at the “RPA + xc” level of approximation.

At this point it is worth pointing out that so far we considered only the screening of *static* perturbations: in fact  $\partial v^{\text{en}}$  was implicitly taken to be frequency independent. Physically this choice corresponds to describing phonons as quasistatic perturbations, so that at each set of instantaneous atomic positions during a vibration cycle, the electrons have enough time to readjust and reach their ground state. This is a statement of the adiabatic approximation (Born and Oppenheimer, 1927). The importance of *retardation* effects in the electron-phonon problem was already recognized in the early work of Bardeen and Pines (1955), but the first *ab initio* calculations of these effects appeared much later [see Lazzeri and Mauri (2006)]. The formal framework required to incorporate retardation in the study of EPIs is presented in Sec. IV.

## 5. Connection with early formulations

For completeness, we illustrate the link between electron-phonon matrix elements obtained within DFPT (Sec. III.B.3) and the early approaches of Bloch (1929) and Bardeen (1937) (Sec. II.A).

The Bloch matrix element can be derived as follows. We assume that the scattering potential is unscreened and corresponds to the bare pseudopotentials  $V_{\kappa}$  in Eq. (43), that there is only one atom at the origin of the unit cell, and the Kohn-Sham wave functions can be described by free electrons  $u_{n\mathbf{k}}(\mathbf{r}) = \Omega^{-1/2} \exp(i\mathbf{G}_n \cdot \mathbf{r})$ . In the last expression, the subscript in  $\mathbf{G}_n$  is used in order to stress the one-to-one correspondence between the reciprocal lattice vectors and the energy bands of the free electron gas in the reduced-zone scheme. Using these approximations in Eqs. (21), (34), (38), and (43), we find

$$\begin{aligned} g_{m\nu}(\mathbf{k}, \mathbf{q}) &= -i[\hbar/(2N_p M_{\kappa} \omega_{\mathbf{q}\nu})]^{1/2} V_{\kappa}(\mathbf{q} + \mathbf{G}_m - \mathbf{G}_n) \\ &\times (\mathbf{q} + \mathbf{G}_m - \mathbf{G}_n) \cdot \mathbf{e}_{\kappa,\nu}(\mathbf{q}). \end{aligned} \quad (57)$$

By further neglecting umklapp processes ( $\mathbf{G}_m \neq \mathbf{G}_n$ ) the previous result becomes (Grimvall, 1981, Sec. 3.4)

$$g_{m\nu}(\mathbf{k}, \mathbf{q}) = -i[\hbar/(2N_p M_{\kappa} \omega_{\mathbf{q}\nu})]^{1/2} \mathbf{q} \cdot \mathbf{e}_{\kappa,\nu}(\mathbf{q}) V_{\kappa}(\mathbf{q}). \quad (58)$$

The expression obtained by Bloch (1929) and reproduced in Eq. (2) is simply obtained by replacing  $V_{\kappa}(\mathbf{q})$  with the effective potential  $V_0$ .

The Bardeen matrix element is more elaborate and can be derived as follows. We describe the screening of the bare ionic potential within the RPA approximation and determine the dielectric matrix by replacing the Kohn-Sham wave functions by free electrons. Using  $u_{n\mathbf{k}}(\mathbf{r}) = \Omega^{-1/2} \exp(i\mathbf{G}_n \cdot \mathbf{r})$  and

$\varepsilon_{n\mathbf{k}} = \hbar^2(\mathbf{k} + \mathbf{G}_n)^2/2m_e - \varepsilon_F$  in Eq. (51), the polarizability reduces to

$$\chi_{\mathbf{G}\mathbf{G}'}^0(\mathbf{q}) = -\frac{m_e k_F}{\pi^2 \hbar^2 \Omega} F(|\mathbf{q} + \mathbf{G}|/2k_F) \delta_{\mathbf{G}\mathbf{G}'}, \quad (59)$$

where  $F$  is the function defined below Eq. (4). The derivation of this result requires making the transition from the first Brillouin zone to the extended-zone scheme. If we use Eq. (59) inside Eq. (56), neglect the exchange and correlation kernel and use the Fourier transform of the Coulomb potential, we find

$$\varepsilon_{\mathbf{G}\mathbf{G}'}(\mathbf{q}) = \delta_{\mathbf{G}\mathbf{G}'} [1 + (k_{\text{TF}}^2/|\mathbf{q} + \mathbf{G}|^2) F(|\mathbf{q} + \mathbf{G}|/2k_F)], \quad (60)$$

where the Thomas-Fermi screening length is given by  $k_{\text{TF}} = [4me^2k_F/(4\pi\varepsilon_0\pi\hbar^2)]^{1/2}$ . Equation (60) is the well-known Lindhard dielectric matrix, and the diagonal matrix elements are the same as in Eq. (4) [see Mahan (1993) and Giuliani and Vignale (2005) for in-depth discussions of the Lindhard function]. By following the same steps that led to Eq. (58), replacing the bare ionic potential by its screened counterpart, and using Eq. (53) with  $\varepsilon$  instead of  $\varepsilon^{\text{Hxc}}$ , we obtain

$$g_{mn\nu}(\mathbf{k}, \mathbf{q}) = -i[\hbar/(2N_p M_\kappa \omega_{\mathbf{q}\nu})]^{1/2} \mathbf{q} \cdot \mathbf{e}_{\kappa,\nu}(\mathbf{q}) V_\kappa(\mathbf{q})/\varepsilon(\mathbf{q}). \quad (61)$$

Here we considered only one atom at the center of the unit cell, and we neglected umklapp processes. This is essentially the result derived by Bardeen (1937) and reproduced in Eq. (3).

#### IV. FIELD-THEORETIC APPROACH TO THE ELECTRON-PHONON INTERACTION

In Sec. III we discussed how the material parameters entering the electron-phonon Hamiltonian in Eq. (1), namely,  $\varepsilon_{n\mathbf{k}}$ ,  $\omega_{\mathbf{q}\nu}$ , and  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$  can be calculated from first principles using DFT and DFPT. Today the formalism and techniques described in Sec. III constitute *de facto* the standard tool in quantitative studies of electron-phonon interactions in solids (see Secs. VII–XII).

However, note that the DFT approach to EPIs does not rest on strong theoretical foundations. For one, the evaluation of the EPI matrix elements via Eq. (38) relies on the assumption that the interaction between electrons and nuclei is governed by the effective Kohn-Sham potential. Therefore we can expect the matrix elements to be sensitive to the exchange and correlation functional (see Sec. XII). Furthermore, the very definition of phonons starting from Eq. (12) relies on the Born-Oppenheimer approximation, and one might ask whether this choice is accurate enough in metals and narrow-gap semiconductors (see Sec. VII). Finally, if one were to go beyond the Born-Oppenheimer approximation, then it would seem sensible to also incorporate retardation effects in the calculation of the EPI matrix elements.

On top of these practical points, and at a more fundamental level, we expect that the electron-phonon interaction will

modify both the electronic structure and the lattice dynamics of a solid, and these modifications will in turn affect the coupling between electrons and phonons. It is therefore clear that a complete theory of interacting electrons and phonons must be self-consistent. In order to address these issues it is necessary to formulate the electron-phonon problem using a rigorous and general theory of interacting electrons and phonons in solids.

The most systematic and elegant approach is based on quantum field theory (Schwinger, 1951) and is tightly connected to the development of the *GW* method (Hedin, 1965). The first attempts in this direction were from Nakajima (1954), Bardeen and Pines (1955), Migdal (1958), and Engelsberg and Schrieffer (1963). However, from the point of view of the present article, these works are of limited usefulness since they were mostly developed around the homogeneous electron gas.

A completely general formulation of the problem, which seamlessly applies to metals, semiconductors, and insulators, was first provided by Baym (1961) and subsequently by Hedin and Lundqvist (1969). The formalism developed in these articles constitutes today the most complete theory of the electron-phonon problem. In fact, many aspects of this formalism are yet to be explored within the context of *ab initio* calculations. After these seminal works several others contributed to clarifying various aspects of the many-body theory of the coupled electron-phonon system, including Sjölander and Johnson (1965), Keating (1968), Gillis (1970), Maksimov (1976), and Vogl (1976), and more recently van Leeuwen (2004) and Marini, Poncé, and Gonze (2015). In particular, van Leeuwen focused on the issues of translational and rotational invariance of the resulting theory, while Marini *et al.* analyzed the connection between many-body perturbation theory approaches and DFT calculations.

Since the mathematical notation of the original articles is obsolete and rather difficult to follow, in Secs. IV.A–IV.D we cover the theory in some detail using contemporary notation. The following derivations can be found across the works of Kato, Kobayashi, and Namiki (1960), Baym (1961), Hedin and Lundqvist (1969), and Maksimov (1976). Here we provide a synthesis of these contributions using a unified notation, and we fill the gaps wherever it is necessary. The presentation requires some familiarity with field operators [see, for example, Merzbacher (1998) for a succinct introduction].

##### A. Operators and distinguishability

The starting point for studying EPIs using a field-theoretic approach is to define the Fock space and the field operators for electrons and nuclei. In the case of electrons the choice is unambiguous, since any many-body state can be represented as a linear combination of Slater determinants constructed using a basis of single-particle wave functions. In the case of nuclei the situation is slightly more ambiguous: in principle we might proceed in a very general way by choosing to focus on the nuclei as our quantum particles, as opposed to their displacements from equilibrium. In practice this choice leads to a dead end for two reasons. First, the quantum statistics of nuclei would be dependent on their spin, and therefore we

would end up with an unwieldy mix of fermions and bosons depending on the solid. Second, the notion of indistinguishable particles, which is central to second quantization, does not apply to nuclei in solids (at least in thermodynamic equilibrium and far from a solid-liquid phase transition). In fact, in many cases we can directly label the nuclei, for example, by means of experimental probes such as scanning tunneling microscopy and electron diffraction. In order to avoid these issues, it is best to study the electron-phonon problem by considering (i) *indistinguishable* electrons, for which it is convenient to use second-quantized operators; (ii) *distinguishable* nuclei, for which it is best to use first quantization in the displacements; and (iii) *indistinguishable* phonons, resulting from the quantization of the nuclear displacements. In this latter case the distinction between first and second quantization is irrelevant. These aspects are briefly mentioned by Baym (1961) and Maksimov (1976).

With these choices, the dynamical variables of the problem are the electronic field operators  $\hat{\psi}$  (discussed later) and the nuclear displacements from equilibrium  $\Delta\hat{\tau}$  (discussed in Sec. IV.C). In this theory the equilibrium coordinates of the nuclei are regarded as *external* parameters and are to be obtained, for example, from crystallography or DFT calculations. Throughout this section, we limit ourselves to consider *equilibrium* Green's functions at zero temperature. As a result, all expectation values will be evaluated for the electron-nuclei ground state  $|0\rangle$ . The extension of the main results to finite temperature is presented in Sec. V. We do not specify how to obtain the ground state, since the following discussion is independent of the precise shape of this state. In order to derive expressions that are useful for first-principles calculations, at the very end the ground state will be approximated using standard DFT wave functions and phonons (see Sec. V).

The electronic field creation and destruction operators are denoted by  $\hat{\psi}^\dagger(\mathbf{x})$  and  $\hat{\psi}(\mathbf{x})$ , where the variable  $\mathbf{x}$  indicates both the position  $\mathbf{r}$  and the spin label  $\sigma$ . These operators obey the anticommutation relations (Merzbacher, 1998)  $\{\hat{\psi}(\mathbf{x}), \hat{\psi}(\mathbf{x}')\} = \{\hat{\psi}^\dagger(\mathbf{x}), \hat{\psi}^\dagger(\mathbf{x}')\} = 0$  and  $\{\hat{\psi}(\mathbf{x}), \hat{\psi}^\dagger(\mathbf{x}')\} = \delta(\mathbf{x} - \mathbf{x}')$ . The most general nonrelativistic Hamiltonian for a system of coupled electrons and nuclei can be written as

$$\hat{H} = \hat{T}_e + \hat{T}_n + \hat{U}_{ee} + \hat{U}_{nn} + \hat{U}_{en}, \quad (62)$$

where each term is introduced hereafter. The electron kinetic energy is

$$\hat{T}_e = -\frac{\hbar^2}{2m_e} \int d\mathbf{x} \hat{\psi}^\dagger(\mathbf{x}) \nabla^2 \hat{\psi}(\mathbf{x}), \quad (63)$$

with  $m_e$  the electron mass, and the integrals  $\int d\mathbf{x}$  denoting the sum over spin and the integration over space  $\sum_\sigma \int d\mathbf{r}$ . The electron-electron interaction is

$$\hat{U}_{ee} = \frac{1}{2} \int d\mathbf{r} \int d\mathbf{r}' \hat{n}_e(\mathbf{r}) [\hat{n}_e(\mathbf{r}') - \delta(\mathbf{r} - \mathbf{r}')] v(\mathbf{r}, \mathbf{r}'), \quad (64)$$

where the electron particle density operator is given by  $\hat{n}_e(\mathbf{r}) = \sum_\sigma \hat{\psi}^\dagger(\mathbf{x}) \hat{\psi}(\mathbf{x})$ , and  $v(\mathbf{r}, \mathbf{r}') = e^2 / (4\pi\epsilon_0 |\mathbf{r} - \mathbf{r}'|)$  is the Coulomb interaction between two particles of charge  $e$ .

In Eqs. (63) and (64) the integrals are over the entire crystal. This corresponds to considering a supercell of infinite size (therefore the lattice vectors  $\mathbf{T}$  of the supercell drop out) and a dense sampling of wave vectors  $\mathbf{q}$  in the Brillouin zone. This choice is useful in order to maintain the formalism as light as possible. Accordingly, all sums over  $\mathbf{q}$  are replaced using  $N_p^{-1} \sum_{\mathbf{q}} \rightarrow \Omega_{\text{BZ}}^{-1} \int d\mathbf{q}$ , where the integral is over the Brillouin zone of volume  $\Omega_{\text{BZ}}$ . Similarly the closure relations in Eq. (A1) are replaced by  $\int d\mathbf{q} \exp(i\mathbf{q} \cdot \mathbf{R}_p) = \Omega_{\text{BZ}} \delta_{p0}$  and  $\sum_p \exp(i\mathbf{q} \cdot \mathbf{R}_p) = \Omega_{\text{BZ}} \delta(\mathbf{q})$ . The nuclear kinetic energy operator is the same as the last term in Eq. (19). Using the same notation as in Sec. III the nucleus-nucleus interaction energy is

$$\hat{U}_{nn} = \frac{1}{2} \sum_{\kappa' p' \neq \kappa p} Z_\kappa Z_{\kappa'} v(\boldsymbol{\tau}_{\kappa p}^0 + \Delta\hat{\boldsymbol{\tau}}_{\kappa p}, \boldsymbol{\tau}_{\kappa' p'}^0 + \Delta\hat{\boldsymbol{\tau}}_{\kappa' p'}). \quad (65)$$

Here  $\boldsymbol{\tau}_{\kappa p}^0$  denotes the classical equilibrium position of each nucleus, and the displacement operators  $\Delta\hat{\boldsymbol{\tau}}_{\kappa p}$  will later be expressed in terms of the ladder operators from Appendix B. The electron-nucleus interaction energy is

$$\hat{U}_{en} = \int d\mathbf{r} \int d\mathbf{r}' \hat{n}_e(\mathbf{r}) \hat{n}_n(\mathbf{r}') v(\mathbf{r}, \mathbf{r}'), \quad (66)$$

where the nuclear charge density operator is given by

$$\hat{n}_n(\mathbf{r}) = -\sum_{\kappa p} Z_\kappa \delta(\mathbf{r} - \boldsymbol{\tau}_{\kappa p}^0 - \Delta\hat{\boldsymbol{\tau}}_{\kappa p}). \quad (67)$$

Here the density operators are expressed in units of the electron charge, so that the expectation value of the total charge density is  $-e\langle 0 | \hat{n}(\mathbf{r}) | 0 \rangle$  with  $\hat{n}(\mathbf{r}) = \hat{n}_e(\mathbf{r}) + \hat{n}_n(\mathbf{r})$ .

We underline the asymmetry between Eqs. (64) and (65): in the case of electrons one considers the electrostatic energy of a continuous distribution of charge, and the unphysical self-interaction is removed by the Dirac delta. Whereas in the case of nuclei, the particles are distinguishable and therefore one has to take into account all pairwise interactions individually.

## B. Electron Green's function

### 1. Equation of motion and self-energy

In this section we focus on the electrons. By combining Eqs. (62)–(66) and using the anticommutation relations for the field operators one finds the following standard expression:

$$\hat{H} = \hat{T}_n + \hat{U}_{nn} + \int d\mathbf{x} \hat{\psi}^\dagger(\mathbf{x}) \left[ -\frac{\hbar^2}{2m_e} \nabla^2 + \hat{V}_n(\mathbf{r}) \right] \hat{\psi}(\mathbf{x}) + \frac{1}{2} \int d\mathbf{x} d\mathbf{x}' v(\mathbf{r}, \mathbf{r}') \hat{\psi}^\dagger(\mathbf{x}) \hat{\psi}^\dagger(\mathbf{x}') \hat{\psi}(\mathbf{x}') \hat{\psi}(\mathbf{x}), \quad (68)$$

where the nuclear potential  $\hat{V}_n$  is given by

$$\hat{V}_n(\mathbf{r}) = \int d\mathbf{r}' v(\mathbf{r}, \mathbf{r}') \hat{n}_n(\mathbf{r}'). \quad (69)$$

In order to study the excitation spectrum of the many-body Hamiltonian  $\hat{H}$  at equilibrium we need to determine the time-ordered one-electron Green's function (Kato, Kobayashi, and Namiki, 1960; Fetter and Walecka, 2003). At zero temperature this function is defined as

$$G(\mathbf{x}t, \mathbf{x}'t') = -\frac{i}{\hbar} \langle 0 | \hat{T} \psi(\mathbf{x}t) \psi^\dagger(\mathbf{x}'t') | 0 \rangle, \quad (70)$$

where  $\hat{T}$  is Wick's time-ordering operator for fermions and ensures that the times of the subsequent operators increase toward the left. The formal definition of the Wick operator is  $\hat{T} \psi(\mathbf{x}t) \psi^\dagger(\mathbf{x}'t') = \theta(t - t') \psi(\mathbf{x}t) \psi^\dagger(\mathbf{x}'t') - \theta(t' - t) \psi^\dagger(\mathbf{x}'t') \psi(\mathbf{x}t)$ , where  $\theta$  is the Heaviside function. Based on this definition we see that for  $t > t'$  the Green's function in Eq. (70) corresponds to the scalar product between the initial state  $\psi^\dagger(\mathbf{x}'t') | 0 \rangle$  and the final state  $\psi(\mathbf{x}t) | 0 \rangle$ . This product is precisely the probability amplitude for finding an electron in the position  $\mathbf{x}$  at the time  $t$ , after having introduced an electron in  $\mathbf{x}'$  at an earlier time  $t'$ . In the case  $t < t'$  the situation is reversed and the Green's function describes the propagation of a hole created in the system at the time  $t$ .

In order to determine  $G(\mathbf{x}t, \mathbf{x}'t')$  we need to establish an equation of motion for the field operators. This can be done by describing the time dependence of the operators within the Heisenberg picture:

$$\hat{\psi}(\mathbf{x}t) = e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar}, \quad (71)$$

where  $\hat{H}$  was defined in Eq. (68). From this definition it follows immediately that

$$i\hbar \frac{\partial}{\partial t} \hat{\psi}(\mathbf{x}t) = [\hat{\psi}(\mathbf{x}t), \hat{H}]. \quad (72)$$

By combining Eqs. (68) and (72) and using the anticommutation relations for the field operators one obtains

$$i\hbar \frac{\partial}{\partial t} \hat{\psi}(\mathbf{x}t) = \left[ -\frac{\hbar^2}{2m_e} \nabla^2 + \int d\mathbf{r}' v(\mathbf{r}, \mathbf{r}') \hat{n}(\mathbf{r}') \right] \hat{\psi}(\mathbf{x}t), \quad (73)$$

where the time dependence in  $\hat{n}(\mathbf{r}'t)$  is to be understood in the Heisenberg sense, as in Eq. (71). This equation of motion allows us to write the corresponding equation for the electron Green's function in Eq. (70):

$$\begin{aligned} & \left[ i\hbar \frac{\partial}{\partial t} + \frac{\hbar^2}{2m_e} \nabla^2 - \varphi(\mathbf{r}t) \right] G(\mathbf{x}t, \mathbf{x}'t') \\ &= \delta(\mathbf{x}t, \mathbf{x}'t') - \frac{i}{\hbar} \int d\mathbf{r}'' dt'' v(\mathbf{r}t, \mathbf{r}''t'') \\ & \quad \times \langle \hat{T} \hat{n}(\mathbf{r}''t'') \psi(\mathbf{x}t) \psi^\dagger(\mathbf{x}'t') \rangle. \end{aligned} \quad (74)$$

Here  $v(\mathbf{r}t, \mathbf{r}''t'') = v(\mathbf{r}, \mathbf{r}'') \delta(t - t'')$ , the bra and ket  $\langle \cdots \rangle$  are a short-hand notation for  $\langle 0 | \cdots | 0 \rangle$ , and the additional term  $\varphi$  is discussed later. In order to obtain Eq. (74) we used once again the anticommutation relations, and we noted that the derivative of the Heaviside function is a Dirac delta.

The new term  $\varphi(\mathbf{r}t)$  which appeared in Eq. (74) is a scalar electric potential which couples to both electronic and nuclear charges. This potential has been introduced in order to perturb the system via the additional Hamiltonian  $\hat{H}_1(t) = \int d\mathbf{r} \hat{n}(\mathbf{r}t) \varphi(\mathbf{r}t)$ . The physical idea behind this choice is to use  $\varphi(\mathbf{r}t)$  in order to induce forced oscillations in the system. When the system resonates with the perturbation we know that the resonant frequency must correspond to a free oscillation, i.e., a many-body eigenmode. From a formal point of view, the potential  $\varphi(\mathbf{r}t)$  is introduced in order to exploit Schwinger's functional derivative technique (Kato, Kobayashi, and Namiki, 1960, Appendix II) and is set to zero at the end of the derivation.

One complication arising from the introduction of  $\varphi(\mathbf{r}t)$  in Eq. (74) is that the time evolution in Eq. (71) is no longer valid, since the perturbed Hamiltonian now depends on the time variable. The way around this complication is to switch from the Heisenberg picture to the interaction picture. This change amounts to replacing the exponentials in Eq. (71) by the time-ordered Dyson series  $\hat{U}(t) = \hat{T} \exp[-i\hbar^{-1} \int_0^t \hat{H}(t') dt']$  (Fetter and Walecka, 2003). Since this would lead to an overlong derivation, we prefer to leave this aspect aside and refer the interested reader to Aryasetiawan and Gunnarsson (1998) for a more comprehensive discussion.

In order to write Eq. (74) in a manageable form, we use the identity (Kato, Kobayashi, and Namiki, 1960)

$$\frac{\delta \langle \hat{T} \hat{a}(t_1) \hat{b}(t_2) \rangle}{\delta \varphi(\mathbf{r}''t'')} = -\frac{i}{\hbar} \langle \hat{T} [\hat{n}(\mathbf{r}''t'') - \langle \hat{n}(\mathbf{r}''t'') \rangle] \hat{a}(t_1) \hat{b}(t_2) \rangle. \quad (75)$$

In this and the following expressions  $\delta/\delta\varphi(\mathbf{r}''t'')$  denotes the *functional* derivative with respect to  $\varphi(\mathbf{r}''t'')$  and should not be confused with the Dirac delta functions  $\delta(\mathbf{x}t, \mathbf{x}'t')$ . Equation (75) is proven by Kato, Kobayashi, and Namiki (1960), Appendix II, and by Hedin and Lundqvist (1969), Appendix B.a. After identifying  $\hat{a}$  and  $\hat{b}$  with  $\hat{\psi}(\mathbf{x}t)$  and  $\hat{\psi}^\dagger(\mathbf{x}'t')$ , respectively, Eq. (74) becomes

$$\begin{aligned} & \left[ i\hbar \frac{\partial}{\partial t} + \frac{\hbar^2}{2m_e} \nabla^2 - V_{\text{tot}}(\mathbf{r}t) - i\hbar \int d\mathbf{r}'' dt'' v(\mathbf{r}t + \eta, \mathbf{r}''t'') \right. \\ & \quad \left. \times \frac{\delta}{\delta \varphi(\mathbf{r}''t'')} \right] G(\mathbf{x}t, \mathbf{x}'t') = \delta(\mathbf{x}t, \mathbf{x}'t'), \end{aligned} \quad (76)$$

where  $\eta$  is a positive infinitesimal arising from the time ordering, and  $V_{\text{tot}}(\mathbf{r}t)$  is the total potential acting on the electronic and nuclear charges, averaged over the many-body quantum state  $|0\rangle$

$$V_{\text{tot}}(\mathbf{r}t) = \int d\mathbf{r}' v(\mathbf{r}, \mathbf{r}') \langle \hat{n}(\mathbf{r}'t) \rangle + \varphi(\mathbf{r}t). \quad (77)$$

Equation (76) was first derived by Kato, Kobayashi, and Namiki (1960). In order to avoid a proliferation of variables, it is common practice to replace the letters by integer numbers, using the convention  $(\mathbf{x}t)$  or  $(\mathbf{r}t) \rightarrow 1$ ,  $(\mathbf{x}'t')$  or  $(\mathbf{r}'t') \rightarrow 2$ ,

$(\mathbf{r}t + \eta) \rightarrow 1^+$ , and so on. Using this convention the last two equations become

$$\left[ i\hbar \frac{\partial}{\partial t_1} + \frac{\hbar^2}{2m_e} \nabla^2(1) - V_{\text{tot}}(1) - i\hbar \int d3v(1+3) \frac{\delta}{\delta\varphi(3)} \right] G(12) = \delta(12) \quad (78)$$

and

$$V_{\text{tot}}(1) = \int d2v(12) \langle \hat{n}(2) \rangle + \varphi(1). \quad (79)$$

In these expressions the spin labels are implied for the Green's function and for the Dirac delta.

At this point, a set of self-consistent equations for coupled electrons and phonons can be generated by eliminating the functional derivative in Eq. (78). For this purpose one first relates the total screened electrostatic potential  $V_{\text{tot}}$  to the external potential  $\varphi$  by introducing the inverse dielectric matrix  $\epsilon^{-1}$  as a functional derivative:

$$\epsilon^{-1}(12) = \delta V_{\text{tot}}(1) / \delta\varphi(2). \quad (80)$$

The function  $\epsilon^{-1}(12)$  is the many-body counterpart of the dielectric matrix discussed in Sec. III.B.4. The form given by Eq. (80) is the most general field-theoretic formulation for a system of interacting electrons and nuclei.

The next step is to rewrite  $\delta G / \delta\varphi$  inside Eq. (78) in terms of the inverse Green's function  $G^{-1}$ . By using the fact that  $\delta \int d2G(12)G^{-1}(23) = 0$  and the rule for the functional derivative of a product (Kadanoff and Baym, 1962) one obtains

$$\frac{\delta G(12)}{\delta\varphi(3)} = - \int d(45)G(14) \frac{\delta G^{-1}(45)}{\delta\varphi(3)} G(52). \quad (81)$$

In order to eliminate any explicit reference to  $\varphi$  we can express the functional derivative on the right-hand side using the chain rule for functional differentiation (Kadanoff and Baym, 1962):

$$\frac{\delta G^{-1}(45)}{\delta\varphi(3)} = \int d6 \frac{\delta G^{-1}(45)}{\delta V_{\text{tot}}(6)} \frac{\delta V_{\text{tot}}(6)}{\delta\varphi(3)}. \quad (82)$$

It is customary to call ‘‘vertex’’ the three-point quantity defined by

$$\Gamma(123) = -\delta G^{-1}(12) / \delta V_{\text{tot}}(3). \quad (83)$$

By combining Eqs. (78) and (80)–(83) one finds

$$\left[ i\hbar \frac{\partial}{\partial t_1} + \frac{\hbar^2}{2m_e} \nabla^2(1) - V_{\text{tot}}(1) \right] G(12) - \int d3 \Sigma(13)G(32) = \delta(12), \quad (84)$$

having introduced the following so-called ‘‘electron self-energy’’  $\Sigma$ :

$$\Sigma(12) = i\hbar \int d(34)G(13)\Gamma(324)W(41^+), \quad (85)$$

which in turn contains the ‘‘screened Coulomb interaction’’  $W$  defined as

$$W(12) = \int d3\epsilon^{-1}(13)v(32) = \int d(3)v(13)\epsilon^{-1}(23). \quad (86)$$

The last equality can be obtained by observing that  $\delta \langle \hat{n}(1) \rangle / \delta\varphi(2) = \delta \langle \hat{n}(2) \rangle / \delta\varphi(1)$  after Eq. (75); therefore  $W(12) = W(21)$ .

Now, by inverting Eq. (84) and using Eq. (83), we can express the vertex  $\Gamma$  in terms of  $\Sigma$  and  $G$ :

$$\Gamma(123) = \delta(12)\delta(13) + \int d(4567) \frac{\delta \Sigma(12)}{\delta G(45)} G(46)G(75)\Gamma(673). \quad (87)$$

The derivation of this result is rather lengthy: it requires the use of the chain rule, in symbols  $\delta / \delta V_{\text{tot}} = (\delta G / \delta V_{\text{tot}}) \delta / \delta G$ , as well as Eq. (81) with  $\Sigma$  and  $V_{\text{tot}}$  instead of  $G$  and  $\varphi$ , respectively.

Equations (84)–(87) form a nonlinear system of equations for the electron Green's function  $G$ , the electron self-energy  $\Sigma$ , the *total* screened Coulomb interaction  $W$ , and the vertex  $\Gamma$ . In order to close the loop it remains to specify the relation between  $W$  and the other quantities. The next section is devoted to this aspect.

## 2. The screened Coulomb interaction

The following equation for the screened Coulomb interaction can be found by combining Eqs. (79), (80), and (86):

$$W(12) = v(12) + \int d(34)v(13) \frac{\delta \langle \hat{n}(3) \rangle}{\delta V_{\text{tot}}(4)} W(42). \quad (88)$$

By defining the ‘‘polarization propagator’’ as

$$P(12) = \frac{\delta \langle \hat{n}(1) \rangle}{\delta V_{\text{tot}}(2)}, \quad (89)$$

the previous expression takes the usual form (Hedin, 1965)

$$W(12) = v(12) + \int d(34)v(13)P(34)W(42). \quad (90)$$

This result can be combined with Eq. (86) in order to express the dielectric matrix in terms of the polarization:

$$\epsilon(12) = \delta(12) - \int d(3)v(13)P(32). \quad (91)$$

We now consider the special case whereby the nuclei are regarded as classical point charges clamped to their equilibrium positions. In this situation, the variation of the charge density  $\delta \langle \hat{n} \rangle$  in Eq. (89) corresponds to the redistribution of the electronic charge in response to the perturbation  $\delta V_{\text{tot}}$ . In order to describe this special case it is convenient to introduce



a new polarization propagator  $P_e$  associated with the electronic response only:

$$P_e(12) = \frac{\delta\langle\hat{n}_e(1)\rangle}{\delta V_{\text{tot}}(2)} = -i\hbar \sum_{\sigma_1} \int d(34)G(13)G(41^+)\Gamma(342). \quad (92)$$

The last equality in this expression is obtained by using Eq. (81) with  $V_{\text{tot}}$  instead of  $\varphi$ , together with Eq. (83), and by considering that the electron density is related to the Green's function via

$$\langle\hat{n}_e(1)\rangle = -i\hbar \sum_{\sigma_1} G(11^+). \quad (93)$$

In conjunction with  $P_e$  it is natural to define the Coulomb interaction screened by the electronic polarization only:

$$W_e(12) = v(12) + \int d(34)v(13)P_e(34)W_e(42), \quad (94)$$

as well as the associated dielectric matrix, in analogy with Eq. (91):

$$\epsilon_e(12) = \delta(12) - \int d(34)v(13)P_e(32). \quad (95)$$

Taken together, Eqs. (84)–(87) with  $W$  replaced by  $W_e$  constitute the well-known Hedin's equations for a system of interacting electrons when the nuclei are clamped to their equilibrium positions (Hedin, 1965).

In order to go back to the most general case whereby the nuclei are not clamped to their equilibrium positions, one has to describe the readjustment of both electronic and nuclear charges. To this aim we combine Eqs. (80), (86), (88), (92), (94), and (95). The result is

$$W(12) = W_e(12) + \int d(34)W_e(13)\frac{\delta\langle\hat{n}_n(3)\rangle}{\delta\varphi(4)}v(42). \quad (96)$$

An explicit expression for  $\delta\langle\hat{n}_n\rangle/\delta\varphi$  can be obtained using the following reasoning. We go into the details since this is a delicate passage. Equation (75) provides a recipe for evaluating the variation of any operator with respect to a potential  $\varphi(\mathbf{r}t)$  which couples to the total charge density operator  $\hat{n}(\mathbf{r}t)$  via  $\hat{H}_1(t) = \int d\mathbf{r}\hat{n}(\mathbf{r}t)\varphi(\mathbf{r}t)$ . Therefore we can replace  $\hat{a}\hat{b}$  in Eq. (75) by  $\hat{n}_n$  to obtain

$$\frac{\delta\langle\hat{n}_n(1)\rangle}{\delta\varphi(2)} = -\frac{i}{\hbar}\langle\hat{T}[\hat{n}(2) - \langle\hat{n}(2)\rangle][\hat{n}_n(1) - \langle\hat{n}_n(1)\rangle]\rangle. \quad (97)$$

In addition, if we introduce a second perturbation  $\hat{H}_2(t) = \int d\mathbf{r}\hat{n}_n(\mathbf{r}t)J(\mathbf{r}t)$ , which couples only to the nuclear charges, we can repeat the same reasoning as in Eq. (97) after replacing  $\varphi$  by  $J$  and  $\hat{n}$  by  $\hat{n}_n$ :

$$\frac{\delta\langle\hat{n}(1)\rangle}{\delta J(2)} = -\frac{i}{\hbar}\langle\hat{T}[\hat{n}_n(2) - \langle\hat{n}_n(2)\rangle][\hat{n}(1) - \langle\hat{n}(1)\rangle]\rangle. \quad (98)$$

The comparison between Eqs. (97) and (98) yields

$$\frac{\delta\langle\hat{n}_n(1)\rangle}{\delta\varphi(2)} = \frac{\delta\langle\hat{n}(2)\rangle}{\delta J(1)}. \quad (99)$$

This can be restated by using the chain rule  $\delta\langle\hat{n}_e\rangle/\delta J = \delta\langle\hat{n}_e\rangle/\delta V_{\text{tot}} \times \delta V_{\text{tot}}/\delta\langle n\rangle \times \delta\langle n\rangle/\delta J$ :

$$\frac{\delta\langle\hat{n}(1)\rangle}{\delta J(2)} = \int d(34)\epsilon_e^{-1}(13)\frac{\delta\langle\hat{n}_n(3)\rangle}{\delta J(2)}. \quad (100)$$

The variation  $\delta\langle\hat{n}_n\rangle/\delta J$  on the right-hand side can be expressed as in Eq. (98):

$$\frac{\delta\langle\hat{n}_n(1)\rangle}{\delta J(2)} = -\frac{i}{\hbar}\langle\hat{T}[\hat{n}_n(2) - \langle\hat{n}_n(2)\rangle]\hat{n}_n(1)\rangle, \quad (101)$$

and since  $\langle\hat{n}_n - \langle\hat{n}_n\rangle\rangle = 0$  this can also be rewritten as

$$\delta\langle\hat{n}_n(1)\rangle/\delta J(2) = D(21), \quad (102)$$

having defined

$$D(12) = -\frac{i}{\hbar}\langle\hat{T}[\hat{n}_n(1) - \langle\hat{n}_n(1)\rangle][\hat{n}_n(2) - \langle\hat{n}_n(2)\rangle]\rangle. \quad (103)$$

This quantity is called the ‘‘density-density correlation function’’ for the nuclei. Finally, we combine Eqs. (96), (99), (100), and (103) to obtain

$$W(12) = W_e(12) + W_{\text{ph}}(12), \quad (104)$$

where  $W_{\text{ph}}$  is the *nuclear* contribution to the screened Coulomb interaction and is given by

$$W_{\text{ph}}(12) = \int d(34)W_e(13)D(34)W_e(42). \quad (105)$$

This important result was first derived by Hedin and Lundqvist (1969).

### 3. Nuclear contribution to the screened Coulomb interaction

In view of the forthcoming discussion, it is useful to derive a more explicit expression for the screened interaction  $W_{\text{ph}}$  in Eq. (105). Here we follow Baym (1961) and Maksimov (1976). The Taylor expansion of the Dirac delta to second order in the displacement  $\mathbf{u}$  reads

$$\delta(\mathbf{r} - \mathbf{u}) = \delta(\mathbf{r}) - \mathbf{u} \cdot \nabla\delta(\mathbf{r}) + \frac{1}{2}\mathbf{u} \cdot \nabla\nabla\delta(\mathbf{r}) \cdot \mathbf{u}, \quad (106)$$

where  $\mathbf{u} \cdot \nabla\nabla \cdot \mathbf{u}$  is a short-hand notation for the second-order derivative  $\sum_{\alpha\alpha'}u_\alpha u_{\alpha'}\nabla_\alpha\nabla_{\alpha'}$ . Equation (106) derives from the Fourier representation of the Dirac delta. Using Eq. (106) inside Eq. (67) we deduce

$$\begin{aligned}\hat{n}_n(\mathbf{r}) &= n_n^0(\mathbf{r}) + \sum_{\kappa p} Z_\kappa \Delta \hat{\boldsymbol{\tau}}_{\kappa p} \cdot \nabla \delta(\mathbf{r} - \boldsymbol{\tau}_{\kappa p}^0) \\ &\quad - \frac{1}{2} \sum_{\kappa p} Z_\kappa \Delta \hat{\boldsymbol{\tau}}_{\kappa p} \cdot \nabla \nabla \delta(\mathbf{r} - \boldsymbol{\tau}_{\kappa p}^0) \cdot \Delta \hat{\boldsymbol{\tau}}_{\kappa p},\end{aligned}\quad (107)$$

where  $n_n^0(\mathbf{r})$  is the density of nuclear point charges at the classical equilibrium positions  $\boldsymbol{\tau}_{\kappa p}^0$ . After taking into account the time evolution in the Heisenberg picture as in Eq. (71), we can replace this expansion inside Eq. (103) to obtain

$$\begin{aligned}D(12) &= \sum_{\substack{\kappa p \\ \kappa' \alpha' p'}} Z_\kappa \nabla_{1,\alpha} \delta(\mathbf{r}_1 - \boldsymbol{\tau}_{\kappa p}^0) D_{\kappa \alpha p, \kappa' \alpha' p'}(t_1 t_2) \\ &\quad \times Z_{\kappa'} \nabla_{2,\alpha'} \delta(\mathbf{r}_2 - \boldsymbol{\tau}_{\kappa' p'}^0).\end{aligned}\quad (108)$$

On the right-hand side we introduced the ‘‘displacement-displacement correlation function’’:

$$D_{\kappa \alpha p, \kappa' \alpha' p'}(tt') = -\frac{i}{\hbar} \langle \hat{T} \Delta \hat{\boldsymbol{\tau}}_{\kappa \alpha p}(t) \Delta \hat{\boldsymbol{\tau}}_{\kappa' \alpha' p'}(t') \rangle.\quad (109)$$

If we insert the last two equations into Eq. (105) we find

$$\begin{aligned}W_{\text{ph}}(12) &= \sum_{\substack{\kappa p \\ \kappa' \alpha' p'}} \int d(34) \epsilon_e^{-1}(13) \nabla_{3,\alpha} V_\kappa(\mathbf{r}_3 - \boldsymbol{\tau}_{\kappa p}^0) \\ &\quad \times D_{\kappa \alpha p, \kappa' \alpha' p'}(t_3 t_4) \epsilon_e^{-1}(24) \nabla_{4,\alpha'} V_{\kappa'}(\mathbf{r}_4 - \boldsymbol{\tau}_{\kappa' p'}^0).\end{aligned}\quad (110)$$

In this expression  $V_\kappa$  is the bare Coulomb potential of a nucleus or its ionic pseudopotential.

At this point in the derivation, Hedin and Lundqvist introduced the approximation in which the electronic dielectric matrix in Eq. (110) can be replaced by its static counterpart. This choice implies the Born-Oppenheimer adiabatic approximation. In view of keeping the formalism as general as possible, we prefer to retain retardation effects, following the earlier works by Bardeen and Pines (1955) and Baym (1961). We come back to this aspect in Secs. V.A.3 and V.B.2.

We stress the fact that the sole approximation used until this point is to truncate the density operator for the nuclei to the second order in the atomic displacements. This is the standard harmonic approximation. Apart from this approximation, which is useful to express  $W_{\text{ph}}$  in a tractable form, no other assumptions are made. Gillis (1970) proposed a generalization of the results by Baym (1961) which does not use the harmonic approximation. However, the resulting formalism is exceedingly complex and has not been followed up.

### C. Phonon Green’s function

In order to complete the set of self-consistent many-body equations for the coupled electron-phonon system, it remains to specify a prescription for calculating the displacement-displacement correlation function  $D_{\kappa \alpha p, \kappa' \alpha' p'}(tt')$ . This function is seldom referred to as the ‘‘phonon Green’s function,’’ even though strictly speaking this name should be reserved for the quantity  $-(i/\hbar) \langle \hat{T} \hat{a}_{\mathbf{q}\nu}(t) \hat{a}_{\mathbf{q}'\nu'}^\dagger(t') \rangle$  discussed in Sec. V.A.1.

In the following we describe the procedure originally devised by Baym (1961), and subsequently analyzed by Keating (1968), Hedin and Lundqvist (1969), Gillis (1970), and Maksimov (1976).

The starting point is the equation of motion for the displacement operators  $\Delta \hat{\boldsymbol{\tau}}_{\kappa p}(t)$ . In analogy with Eq. (72) we have  $i\hbar \partial/\partial t \Delta \hat{\boldsymbol{\tau}}_{\kappa p}(t) = [\Delta \hat{\boldsymbol{\tau}}_{\kappa p}(t), \hat{H}]$ . Since we are considering the harmonic approximation and we expect the nuclei to oscillate around their equilibrium positions, it is convenient to aim for an expression resembling Newton’s equation. This can be done by taking the time derivative of the equation of motion

$$M_\kappa \frac{\partial^2}{\partial t^2} \Delta \hat{\boldsymbol{\tau}}_{\kappa p} = -\frac{M_\kappa}{\hbar^2} [[\Delta \hat{\boldsymbol{\tau}}_{\kappa p}, \hat{H}], \hat{H}].\quad (111)$$

After evaluating the commutators using Eqs. (62)–(66) and performing the derivatives with respect to the nuclear displacements by means of Eq. (106), we obtain (the steps are laborious but straightforward)

$$\begin{aligned}M_\kappa \frac{\partial^2}{\partial t^2} \Delta \hat{\boldsymbol{\tau}}_{\kappa p}(t) &= Z_\kappa \int d\mathbf{r} d\mathbf{r}' \hat{n}^{(\kappa p)}(\mathbf{r}t) v(\mathbf{r}, \mathbf{r}') \\ &\quad \times \{-\nabla' \delta(\mathbf{r}' - \boldsymbol{\tau}_{\kappa p}^0) \\ &\quad + \nabla' [\nabla' \delta(\mathbf{r}' - \boldsymbol{\tau}_{\kappa p}^0) \cdot \Delta \hat{\boldsymbol{\tau}}_{\kappa p}(t)]\}.\end{aligned}\quad (112)$$

Here  $\hat{n}^{(\kappa p)}(\mathbf{r})$  is the total charge density of electrons and nuclei, after removing the contribution of the nucleus  $\kappa$  in the unit cell  $p$ . In the second line  $\nabla'$  indicates that the derivatives are taken with respect to the variable  $\mathbf{r}'$ . At this point, we can use the functional derivative technique as in Sec. IV.B in order to determine an expression involving the displacement-displacement correlation function from Eq. (109). Here, instead of using  $J(\mathbf{r})$  as in Sec. IV.B.2 for the nuclear density, it is convenient to work with the individual displacements and introduce a third perturbation  $\hat{H}_3(t) = \sum_{\kappa p} \mathbf{F}_{\kappa p}(t) \cdot \Delta \hat{\boldsymbol{\tau}}_{\kappa p}(t)$ . The extra terms  $\mathbf{F}_{\kappa p}(t)$  have the meaning of external forces acting on the nuclei. Using this perturbation, it is possible to write the displacement-displacement correlation function in a manner similar to Eq. (101)

$$\frac{\delta \langle \Delta \hat{\boldsymbol{\tau}}_{\kappa \alpha p}(t) \rangle}{\delta \mathbf{F}_{\kappa' \alpha' p'}(t')} = D_{\kappa \alpha p, \kappa' \alpha' p'}(tt').\quad (113)$$

This result was derived by Baym (1961) using a finite-temperature formalism. As in the case of the electron Green’s function in Sec. IV.B, it can be obtained only by working in the interaction picture and by taking into account the explicit time dependence of the Hamiltonian  $\hat{H} + \hat{H}_3(t)$ . Also in the present case, we omit these details for conciseness.

If we take the expectation value of Eq. (112) in the ground state, after having added the new force term  $-\mathbf{F}_{\kappa p}(t)$ , and carry out the functional derivative with respect to  $\mathbf{F}_{\kappa' p'}(t')$ , we obtain

$$\begin{aligned}
 M_\kappa \frac{\partial^2}{\partial t^2} D_{\kappa\alpha p, \kappa' \alpha' p'}(tt') &= -\delta_{\kappa\alpha p, \kappa' \alpha' p'} \delta(tt') + Z_\kappa \int d\mathbf{r} d\mathbf{r}' \\
 &\times \left[ -\frac{\delta \langle \hat{n}^{(\kappa p)}(\mathbf{r}t) \rangle}{\delta F_{\kappa' \alpha' p'}(t')} v(\mathbf{r}, \mathbf{r}') \nabla'_\alpha \delta(\mathbf{r}' - \boldsymbol{\tau}_{\kappa p}^0) \right. \\
 &\left. + \langle \hat{n}^{(\kappa p)}(\mathbf{r}t) \rangle v(\mathbf{r}, \mathbf{r}') \nabla'_\alpha \nabla'_\gamma \delta(\mathbf{r}' - \boldsymbol{\tau}_{\kappa p}^0) D_{\kappa\gamma p, \kappa' \alpha' p'}(tt') \right], \quad (114)
 \end{aligned}$$

where the sum over the Cartesian directions  $\gamma$  is implied. The derivation of this result is rather cumbersome and involves the following considerations: (i) the Dirac deltas in the first line come from the force terms  $-\mathbf{F}_{\kappa p}$  added to the Hamiltonian; (ii) within the harmonic approximation the expectation value  $\langle \hat{n}^{(\kappa p)} \Delta \hat{\boldsymbol{\tau}}_{\kappa p} \rangle$  can be replaced by  $\langle \hat{n}^{(\kappa p)} \rangle \langle \Delta \hat{\boldsymbol{\tau}}_{\kappa p} \rangle$  (Baym, 1961); and (iii) the expectation value  $\langle \Delta \hat{\boldsymbol{\tau}}_{\kappa p} \rangle$  can be set to zero, because at the end of the derivation one sets  $|\mathbf{F}_{\kappa p}| = 0$ , hence the expectation values of the displacements vanish. We note that Hedin and Lundqvist (1969) omitted the last line of Eq. (114) in their derivation, but this term was correctly included by Baym (1961) and Maksimov (1976).

The remaining functional derivative in Eq. (114) can be expressed in terms of the nuclear charge density using the same strategy which led to Eq. (100). The result is

$$\begin{aligned}
 \frac{\delta \langle \hat{n}^{(\kappa p)}(\mathbf{r}t) \rangle}{\delta F_{\kappa' \alpha' p'}(t')} &= \int d\mathbf{r}'' dt'' e^{-1}(\mathbf{r}t, \mathbf{r}''t'') \frac{\delta \langle \hat{n}_n(\mathbf{r}''t'') \rangle}{\delta F_{\kappa' \alpha' p'}(t')} \\
 &- \sum_\gamma Z_\kappa D_{\kappa\gamma p, \kappa' \alpha' p'}(tt') \nabla_\gamma \delta(\mathbf{r} - \boldsymbol{\tau}_{\kappa p}^0). \quad (115)
 \end{aligned}$$

By inserting this result inside Eq. (114) and using the expansion in Eq. (107), we finally obtain the equation of motion for the displacement-displacement correlation function:

$$\begin{aligned}
 M_\kappa \frac{\partial^2}{\partial t^2} D_{\kappa\alpha p, \kappa' \alpha' p'}(tt') &= -\delta_{\kappa\alpha p, \kappa' \alpha' p'} \delta(tt') - \sum_{\kappa'' \alpha'' p''} \int dt'' \Pi_{\kappa\alpha p, \kappa'' \alpha'' p''}(tt'') \\
 &\times D_{\kappa'' \alpha'' p'', \kappa' \alpha' p'}(t''t'). \quad (116)
 \end{aligned}$$

The quantity  $\Pi_{\kappa p \alpha, \kappa' p' \alpha'}(tt')$  in this expression is called the ‘‘phonon self-energy’’ and is given by

$$\begin{aligned}
 \Pi_{\kappa\alpha p, \kappa' \alpha' p'}(tt') &= \int d\mathbf{r} d\mathbf{r}' [Z_\kappa \nabla_\alpha \delta(\mathbf{r} - \boldsymbol{\tau}_{\kappa p}^0) W_e(\mathbf{r}t, \mathbf{r}'t') Z_{\kappa'} \nabla'_\alpha \delta(\mathbf{r}' - \boldsymbol{\tau}_{\kappa' p'}^0) \\
 &+ \delta_{\kappa p, \kappa' p'} \delta(tt') \nabla_\alpha \langle \hat{n}(\mathbf{r}) \rangle v(\mathbf{r}, \mathbf{r}') Z_{\kappa'} \nabla'_\alpha \delta(\mathbf{r}' - \boldsymbol{\tau}_{\kappa' p'}^0)]. \quad (117)
 \end{aligned}$$

The derivation of Eq. (117) is nontrivial and is not found consistently in the literature. It requires converting the derivatives with respect to the position variables  $\mathbf{r}, \mathbf{r}'$  into derivatives with respect to the nuclear coordinates; integrating by parts in order to rearrange the derivatives with respect to  $\mathbf{r}$  and  $\mathbf{r}'$ ; invoking the harmonic approximation; and considering that, after setting the forces  $\mathbf{F}_{\kappa p}(t) = 0$  and the field  $\varphi(\mathbf{r}t) = 0$  at the

end, the expectation value  $\langle \hat{n}(\mathbf{r}t) \rangle$  does not depend on time. The term in the third line of Eq. (117) is what Baym (1961) called the ‘‘static force,’’ since it arises from the forces experienced by the nuclei in their equilibrium configuration.

In order to simplify Eq. (117) it is convenient to move from the time to the frequency domain. We use the following convention for the Fourier transform of a function  $f(t)$ :  $f(\omega) = \int_{-\infty}^{\infty} dt f(t) e^{i\omega t}$ . Since we are considering equilibrium properties in the absence of time-dependent external potentials, the time variables enter in these quantities only as differences (Abrikosov, Gor'kov, and Dzyaloshinski, 1975); for example,  $W_e(\mathbf{r}t, \mathbf{r}'t') = W_e(\mathbf{r}, \mathbf{r}', t - t')$ . As a consequence, Eq. (116) is rewritten as

$$\begin{aligned}
 \sum_{\kappa'' \alpha'' p''} [M_\kappa \omega^2 \delta_{\kappa\alpha p, \kappa'' \alpha'' p''} - \Pi_{\kappa\alpha p, \kappa'' \alpha'' p''}(\omega)] \\
 \times D_{\kappa'' \alpha'' p'', \kappa' \alpha' p'}(\omega) = \delta_{\kappa\alpha p, \kappa' \alpha' p'}, \quad (118)
 \end{aligned}$$

whereas the phonon self-energy in the frequency domain at equilibrium reads

$$\begin{aligned}
 \Pi_{\kappa\alpha p, \kappa' \alpha' p'}(\omega) &= \int d\mathbf{r} d\mathbf{r}' [Z_\kappa \nabla_\alpha \delta(\mathbf{r} - \boldsymbol{\tau}_{\kappa p}^0) W_e(\mathbf{r}, \mathbf{r}', \omega) \\
 &+ \delta_{\kappa p, \kappa' p'} \nabla_\alpha \langle \hat{n}(\mathbf{r}) \rangle v(\mathbf{r}, \mathbf{r}') Z_{\kappa'} \nabla'_\alpha \delta(\mathbf{r}' - \boldsymbol{\tau}_{\kappa' p'}^0)]. \quad (119)
 \end{aligned}$$

The second line in this expression is conveniently rewritten by making use of the acoustic sum rule. This sum rule is well known in the theory of lattice dynamics of crystals (Born and Huang, 1954) and can be generalized to the case of many-body Green's function approaches as follows (Hedin and Lundqvist, 1969):

$$\sum_{\kappa' p'} \Pi_{\kappa\alpha p, \kappa' \alpha' p'}(\omega = 0) = 0 \quad \text{for any } \alpha, \alpha'. \quad (120)$$

This relation was first derived by Baym (1961) by imposing the condition that the nuclei in the crystal must remain near their equilibrium positions due to fictitious restoring forces. Physically this condition corresponds to considering a crystal which is held fixed in the laboratory reference frame. In this approach, the crystal cannot translate or rotate as a whole. Similar relations were derived by Sjölander and Johnson (1965) and Gillis (1970).

If we combine Eqs. (119) and (120), perform integrations by parts, and carry out the integrations in  $\mathbf{r}$  and  $\mathbf{r}'$  we obtain

$$\begin{aligned}
 \Pi_{\kappa\alpha p, \kappa' \alpha' p'}(\omega) &= \sum_{\kappa'' p''} Z_\kappa Z_{\kappa''} \frac{\partial^2}{\partial r_\alpha \partial r'_\alpha} \Big|_{\mathbf{r}=\boldsymbol{\tau}_{\kappa p}^0, \mathbf{r}'=\boldsymbol{\tau}_{\kappa'' p''}^0} \\
 &[\delta_{\kappa' p', \kappa'' p''} W_e(\mathbf{r}, \mathbf{r}', \omega) - \delta_{\kappa p, \kappa' p'} W_e(\mathbf{r}, \mathbf{r}', 0)], \quad (121)
 \end{aligned}$$

which fulfills the sum rule in Eq. (120).

Equations (118) and (121) completely define the nuclear dynamics in the harmonic approximation. After obtaining the displacement-displacement correlation function  $D_{\kappa\alpha p, \kappa' \alpha' p'}(tt')$  by solving this set of equations, it is possible to construct the expectation value of the nuclear density using Eqs. (107) and (109)

TABLE I. Self-consistent Hedin-Baym equations for the coupled electron-phonon system in the harmonic approximation.

Eq.	Description	Expression
(93)	Electronic charge density	$\langle \hat{n}_e(1) \rangle = -i\hbar \sum_{\sigma_1} G(11^+)$
(122)	Nuclear charge density	$\langle \hat{n}_n(\mathbf{r}t) \rangle = n_n^0(\mathbf{r}) - (i\hbar/2) \sum_{\kappa p, \alpha \alpha'} Z_\kappa \partial^2 \delta(\mathbf{r} - \boldsymbol{\tau}_{\kappa p}^0) / \partial r_\alpha \partial r_{\alpha'} D_{\kappa \alpha p, \kappa \alpha' p}(t^+ t)$
(79)	Total electrostatic potential	$V_{\text{tot}}(1) = \int d2 v(12) [\langle \hat{n}_e(2) \rangle + \langle \hat{n}_n(2) \rangle]$
(84)	Equation of motion, electrons	$[i\hbar \partial / \partial t_1 + (\hbar^2 / 2m_e) \nabla^2(1) - V_{\text{tot}}(1)] G(12) - \int d3 \Sigma(13) G(32) = \delta(12)$
(118)	Equation of motion, nuclei	$\sum_{\kappa' \alpha' p'} [M_\kappa \omega^2 \delta_{\kappa \alpha p, \kappa' \alpha' p'} - \Pi_{\kappa \alpha p, \kappa' \alpha' p'}(\omega)] D_{\kappa' \alpha' p', \kappa \alpha p}(\omega) = \delta_{\kappa \alpha p, \kappa' \alpha' p'}$
(85)	Electron self-energy	$\Sigma(12) = i\hbar \int d(34) G(13) \Gamma(324) [W_e(41^+) + W_{\text{ph}}(41^+)]$
(94)	Screened Coulomb, electrons	$W_e(12) = v(12) + \int d(34) v(13) P_e(34) W_e(42)$
(92)	Electronic polarization	$P_e(12) = -i\hbar \sum_{\sigma_1} \int d(34) G(13) G(41^+) \Gamma(342)$
(95)	Electronic dielectric matrix	$\epsilon_e(12) = \delta(12) - \int d(3) v(13) P_e(32)$
(87)	Vertex	$\Gamma(123) = \delta(12) \delta(13) + \int d(4567) [\delta \Sigma(12) / \delta G(45)] G(46) G(75) \Gamma(673)$
(110)	Screened Coulomb, nuclei	$W_{\text{ph}}(12) = \sum_{\kappa \alpha p, \kappa' \alpha' p'} \int d(34) \epsilon_e^{-1}(13) \nabla_{3,\alpha} V_\kappa(\mathbf{r}_3 - \boldsymbol{\tau}_{\kappa p}^0) \times D_{\kappa \alpha p, \kappa' \alpha' p'}(t_3 t_4) \epsilon_e^{-1}(24) \nabla_{4,\alpha'} V_{\kappa'}(\mathbf{r}_4 - \boldsymbol{\tau}_{\kappa' p'}^0)$
(121)	Phonon self-energy	$\Pi_{\kappa \alpha p, \kappa' \alpha' p'}(\omega) = \sum_{\kappa'' p''} Z_\kappa Z_{\kappa''} (\partial^2 / \partial r_\alpha \partial r_{\alpha'}) \times [\delta_{\kappa' p', \kappa'' p''} W_e(\mathbf{r}, \mathbf{r}', \omega) - \delta_{\kappa p, \kappa' p'} W_e(\mathbf{r}, \mathbf{r}', 0)]_{\mathbf{r}=\boldsymbol{\tau}_{\kappa p}^0, \mathbf{r}'=\boldsymbol{\tau}_{\kappa' p'}^0}$

$$\langle \hat{n}_n(\mathbf{r}t) \rangle = n_n^0(\mathbf{r}) - \frac{i\hbar}{2} \sum_{\kappa p, \alpha \alpha'} Z_\kappa \frac{\partial^2 \delta(\mathbf{r} - \boldsymbol{\tau}_{\kappa p}^0)}{\partial r_\alpha \partial r_{\alpha'}} D_{\kappa \alpha p, \kappa \alpha' p}(t^+ t). \quad (122)$$

We emphasize that, according to Eq. (121), the coupling of the nuclear displacements to the electrons is completely defined by the electronic dielectric matrix through  $W_e$ . Similarly, the nuclei affect the electronic structure via the dielectric matrix which enters  $W_{\text{ph}}$  in Eq. (110) and via the nuclear density inside  $V_{\text{tot}}$  in Eq. (79). From these considerations it should be clear that the *electronic* dielectric matrix  $\epsilon_e(\mathbf{r}, \mathbf{r}', \omega)$  plays an absolutely central role in the field-theoretic approach to the electron-phonon problem.

#### D. Hedin-Baym equations

Apart from making use of the harmonic approximation, the set of equations given by Eqs. (79), (84), (85), (87), (92), (93), (94), (95), (110), (118), (121), and (122) describe the coupled electron-phonon system entirely from *first principles*. This set of equations can be regarded as the most sophisticated description of interacting electrons and phonons available today. Since the self-consistent equations for the electrons were originally derived by Hedin (1965), and those for the nuclei were derived first by Baym (1961), we refer to the complete set as the *Hedin-Baym equations*. Given the importance of these relations, we summarized them schematically in Table I. The standard Hedin's equations for interacting electrons in the potential of clamped nuclei (Hedin, 1965) are immediately recovered from the Hedin-Baym equations by setting to zero the displacement-displacement correlation function of the nuclei  $D_{\kappa \alpha p, \kappa' \alpha' p'} = 0$ .

Table I provides a closed set of self-consistent equations whose solution yields the Green's functions of a fully interacting electron-phonon system, within the *harmonic* approximation. We stress that these relations are

fundamentally different from diagrammatic approaches. In fact, here the coupled electron-phonon system is not addressed using Feynman-Dyson perturbation theory as was done, for example, by Keating (1968). Instead, in Table I, electrons and phonons are described nonperturbatively by means of a coupled set of nonlinear equations for the exact propagators. In particular, we emphasize that this approach does not require the Born-Oppenheimer adiabatic approximation, and therefore it encompasses insulators, intrinsic as well as doped semiconductors, metals, and superconductors.

Almost every property related to electron-phonon interactions in solids that can be calculated today from first principles can be derived from these equations. Examples to be discussed in Secs. V–X include the renormalization of the Fermi velocity, the band gap renormalization in semiconductors and insulators, the nonadiabatic corrections to vibrational frequencies, the Frölich interaction, and the lifetimes of electrons and phonons. The generalization of these results to the case of finite temperature should also be able to describe phonon-mediated superconductivity, although this phenomenon is best addressed by directly studying the propagation of Cooper pairs (see Sec. XI).

Baym's theory can in principle be extended to go beyond the harmonic approximation (Gillis, 1970). However, the mathematical complexity of the resulting formalism is formidable, due to the appearance of many additional terms which are neglected in the harmonic approximation.

#### V. FROM A MANY-BODY FORMALISM TO PRACTICAL CALCULATIONS

The Hedin-Baym equations summarized in Table I define a rigorous formalism for studying interacting electrons and phonons in metals, semiconductors, and insulators entirely from first principles. However, a direct numerical solution of these equations for real materials is currently out of reach, and approximations are needed for practical calculations.

The following sections establish the connection between the Hedin-Baym equations and standard expressions which are currently in use in *ab initio* calculations of electron-phonon interactions.

## A. Effects of the electron-phonon interaction on phonons

### 1. Phonons in the Born-Oppenheimer adiabatic approximation

The vibrational eigenmodes of the nuclei can be identified with the resonances of the displacement-displacement correlation function  $D_{\kappa p \alpha, \kappa' p' \alpha'}(t t')$  in the frequency domain. If we denote by  $\mathbf{M}$  the diagonal matrix having the nuclear masses  $M_\kappa$  along its diagonal, then the formal solution of Eq. (118) can be written as

$$\mathbf{D}(\omega) = [\mathbf{M}\omega^2 - \Pi(\omega)]^{-1}, \quad (123)$$

where  $\mathbf{D}$  is the matrix with elements  $D_{\kappa p \alpha, \kappa' p' \alpha'}$ . The resonant frequencies of the system correspond to the solutions of the nonlinear equations

$$\Omega_\nu(\omega) - \omega = 0, \quad \text{with } \nu = 1, \dots, 3M, \quad (124)$$

where  $\Omega_\nu^2(\omega)$  is an eigenvalue of  $\mathbf{M}^{-1/2}\Pi(\omega)\mathbf{M}^{-1/2}$ , parametric in the variable  $\omega$ .

As expected, the study of lattice vibrations within a field-theoretic framework resembles the standard eigenvalue problem reviewed in Sec. III.A. In particular, the matrix  $\Pi(\omega)$  represents the many-body counterpart of the matrix of interatomic force constants  $C_{\kappa p \alpha, \kappa' p' \alpha'}$  introduced in Eq. (13). However, despite its formal simplicity, Eq. (124) conceals the full wealth of information associated with the many-body electronic screening  $\epsilon_e(\mathbf{r}, \mathbf{r}', \omega)$  via Eq. (121). In fact, the phonon self-energy is generally *complex* and *frequency dependent*. Therefore we can expect to find roots of Eq. (124) outside of the real frequency axis, as well as multiple roots for the same ‘‘eigenmode.’’

The link between Eq. (124) and phonon calculations by means of DFT is established by noting that DFT relies on the Born-Oppenheimer adiabatic approximation. In the adiabatic approximation the nuclei are considered immobile during characteristic electronic time scales. Formally, this approximation is introduced by setting  $\omega = 0$  in Eq. (121) (Keating, 1968). In practice, this assumption corresponds to stating that  $\epsilon_e(\mathbf{r}, \mathbf{r}', \omega)$  can be replaced by  $\epsilon_e(\mathbf{r}, \mathbf{r}', 0)$  in the frequency range of the vibrational excitations. Obviously this is not always the case, and important exceptions are discussed in Sec. V.A.2.

In order to see more clearly the connection with the formalism discussed in Sec. III.A, we partition the phonon self-energy into ‘‘adiabatic’’ and ‘‘nonadiabatic’’ contributions:

$$\Pi(\omega) = \Pi^A + \Pi^{\text{NA}}(\omega), \quad (125)$$

with  $\Pi^A = \Pi(\omega = 0)$ . As we will see, the adiabatic term  $\Pi^A$  will be taken to describe ‘‘noninteracting’’ phonons, and the nonadiabatic self-energy  $\Pi^{\text{NA}}$  will be used to describe the effects of electron-phonon interactions.

In the early literature it is common to find a different partitioning, whereby the noninteracting system is defined by the bare interatomic force constants, corresponding to nuclei in the absence of electrons (Grimvall, 1981). This alternative choice is not useful in modern calculations, because the resulting noninteracting phonon dispersions are very different from the fully interacting dispersions. The present choice of using instead adiabatic phonons as the noninteracting system is more convenient in the context of modern *ab initio* techniques, since calculations of adiabatic phonon spectra are routinely performed within DFPT.

In the remainder of this section we concentrate on the adiabatic term, and we defer the discussion of the nonadiabatic self-energy to Sec. V.A.2. Using Eq. (121), we can rewrite the adiabatic self-energy as follows:

$$\begin{aligned} \Pi_{\kappa p \alpha, \kappa' p' \alpha'}^A &= \sum_{\kappa'' p''} (\delta_{\kappa' p' \alpha' \kappa'' p''} - \delta_{\kappa p \alpha, \kappa' p' \alpha'}) \\ &\times \left[ \int d\mathbf{r} \frac{\partial \langle \hat{n}_e(\mathbf{r}) \rangle}{\partial \tau_{\kappa' \alpha' p''}} \frac{\partial V^{\text{en}}(\mathbf{r})}{\partial \tau_{\kappa p \alpha}} + \frac{\partial^2 U_{\text{nn}}}{\partial \tau_{\kappa p \alpha} \partial \tau_{\kappa' \alpha' p''}} \right]. \end{aligned} \quad (126)$$

In this expression  $U_{\text{nn}}$  is the nucleus-nucleus interaction energy from Eq. (65),  $V^{\text{en}}$  is the electron-nuclei interaction from Eq. (25), and all the derivatives are taken at the equilibrium coordinates. The derivation of Eq. (126) requires the use of the identity

$$\frac{\partial \langle \hat{n}_e(\mathbf{r}) \rangle}{\partial \tau_{\kappa p \alpha}^0} = -Z_\kappa \int d\mathbf{r}' [\epsilon_e^{-1}(\mathbf{r}, \mathbf{r}'; 0) - \delta(\mathbf{r}, \mathbf{r}')] \nabla'_\alpha \delta(\mathbf{r}' - \tau_{\kappa p}^0). \quad (127)$$

This identity follows from the same reasoning leading to Eq. (100), after considering an external potential which modifies the position of the nucleus  $\kappa$  in the cell  $p$ . Equation (126) can be recast in a familiar form by exploiting the acoustic sum rule in Eq. (120). Indeed after a few tedious but straightforward manipulations we obtain

$$\begin{aligned} \Pi_{\kappa p \alpha, \kappa' p' \alpha'}^A &= \int d\mathbf{r} \frac{\partial \langle \hat{n}_e(\mathbf{r}) \rangle}{\partial \tau_{\kappa' \alpha' p'}} \frac{\partial V^{\text{en}}(\mathbf{r})}{\partial \tau_{\kappa p \alpha}} + \int d\mathbf{r} \langle \hat{n}_e(\mathbf{r}) \rangle \frac{\partial^2 V^{\text{en}}(\mathbf{r})}{\partial \tau_{\kappa p \alpha} \partial \tau_{\kappa' \alpha' p'}} \\ &+ \frac{\partial^2 U_{\text{nn}}}{\partial \tau_{\kappa p \alpha} \partial \tau_{\kappa' \alpha' p'}}. \end{aligned} \quad (128)$$

In this form, one can see that the adiabatic self-energy gives precisely the interatomic force constants that we would obtain using the Born-Oppenheimer approximation and the Hellman-Feynman theorem; compare Eq. (128), for example, with Eq. (10) of Baroni *et al.* (2001).

The difference between  $\Pi_{\kappa p \alpha, \kappa' p' \alpha'}^A$  in Eq. (128) and  $C_{\kappa p \alpha, \kappa' p' \alpha'}$  in Eq. (13) is that, in the former case, the electron density response to atomic displacements is governed by the exact many-body dielectric matrix  $\epsilon_e(\mathbf{r}, \mathbf{r}', 0)$  and electron density  $\langle \hat{n}_e(\mathbf{r}) \rangle$ , as shown by Eqs. (127) and (128). As a result,  $\Pi_{\kappa p \alpha, \kappa' p' \alpha'}^A$  corresponds to force constants and electron density *dressed* by all many-body interactions of the system (both electron-electron and electron-phonon interactions). In

contrast, when the force constants in Eq. (13) are calculated using DFT, the electron density response to an atomic displacement is evaluated using the RPA +  $xc$  screening, that is,  $\epsilon^{\text{Hxc}}(\mathbf{r}, \mathbf{r}')$  from Sec. III.B.4, and the ground-state electron density is calculated at clamped nuclei.

The use of the adiabatic approximation in the study of phonons carries the important advantage that the many-body force constants  $\Pi^{\text{A}}$  form a real and symmetric matrix. This can be seen by rewriting Eq. (121) for  $\omega = 0$ , and using  $W_{\mathbf{e}}(\mathbf{r}, \mathbf{r}', \omega) = W_{\mathbf{e}}(\mathbf{r}', \mathbf{r}, -\omega)$  which follows from the property  $W(12) = W(21)$  (see Sec. IV.B). Since  $\Pi^{\text{A}}$  is real and symmetric, all its eigenvalues are guaranteed to be real. In this approximation, the excitations of the lattice correspond to sharp resonances in the displacement-displacement correlation function  $\mathbf{D}(\omega)$ , and it is meaningful to talk about phonons as long-lived excitations of the system. In fact these excitations are infinitely long lived in the harmonic approximation. In practical calculations, the many-body  $\Pi^{\text{A}}$  is invariably replaced by the DFT interatomic force constants, and in this case the agreement of the calculated phonon frequencies with experiment is excellent in most cases. Illustrative examples can be found among others in Yin and Cohen (1982), Giannozzi *et al.* (1991), Dal Corso *et al.* (1993), Lee, Ghosez, and Gonze (1994), Kresse, Furthmüller, and Hafner (1995), Bungaro, Rapcewicz, and Bernholc (2000), Dal Corso and de Gironcoli (2000), Karki *et al.* (2000), Baroni *et al.* (2001), Díaz-Sánchez, Romero, and Gonze (2007), and Dal Corso (2013).

The most obvious criticism to the adiabatic approximation is that, in the case of *metals*, the assumption  $\epsilon_{\mathbf{e}}(\mathbf{r}, \mathbf{r}', \omega) \simeq \epsilon_{\mathbf{e}}(\mathbf{r}, \mathbf{r}', 0)$  is inadequate. This can intuitively be understood by recalling that the dielectric function of the homogeneous electron gas diverges when  $\omega, \mathbf{q} \rightarrow 0$  (Mahan, 1993). In practical calculations, this divergence is connected with vanishing denominators in Eq. (51) for  $\mathbf{q} \rightarrow 0$ . An approximate, yet successful strategy for overcoming this problem is to replace the occupation numbers in Eq. (51) by smoothing functions such as the Fermi-Dirac distribution and to describe the singular terms analytically (de Gironcoli, 1995). Most first-principles calculations of phonon dispersion relations in metals have been carried out using this strategy. Improvements to this strategy are discussed in Sec. V.A.2.

The adiabatic approximation leads naturally to the definition of an adiabatic propagator  $\mathbf{D}^{\text{A}}(\omega)$ , which can be obtained from Eq. (123) after replacing the phonon self-energy by its static limit

$$\mathbf{D}^{\text{A}}(\omega) = [\mathbf{M}\omega^2 - \Pi^{\text{A}}]^{-1}. \quad (129)$$

Now, if we identify  $\Pi_{\kappa\alpha p, \kappa'\alpha' p'}^{\text{A}}$  with the interatomic force constant  $C_{\kappa\alpha p, \kappa'\alpha' p'}$  in Eq. (13), we can obtain an explicit expression for the adiabatic phonon propagator in terms of the eigenmodes  $e_{\kappa\alpha, \nu}(\mathbf{q})$  and eigenfrequencies  $\omega_{\mathbf{q}\nu}$  introduced in Sec. III.A. To this end we invert Eq. (129) using Eqs. (14)–(17) and recall that the dynamical matrix is Hermitian and obeys  $D_{\kappa\alpha, \kappa'\alpha'}^{\text{dm},*}(\mathbf{q}) = D_{\kappa\alpha, \kappa'\alpha'}^{\text{dm}}(-\mathbf{q})$  (Maradudin and Vosko, 1968). After tedious but straightforward steps we find

$$D_{\kappa\alpha p, \kappa'\alpha' p'}^{\text{A}}(\omega) = \sum_{\nu} \int \frac{d\mathbf{q}}{\Omega_{\text{BZ}}} S_{\mathbf{q}\nu, \kappa\alpha p}^* S_{\mathbf{q}\nu, \kappa'\alpha' p'} \frac{2\omega_{\mathbf{q}\nu}}{\omega^2 - \omega_{\mathbf{q}\nu}^2}, \quad (130)$$

with the definition

$$S_{\mathbf{q}\nu, \kappa\alpha p} = e^{i\mathbf{q}\cdot\mathbf{R}_p} (2M_{\kappa}\omega_{\mathbf{q}\nu})^{-1/2} e_{\kappa\alpha, \nu}(\mathbf{q}). \quad (131)$$

This result suggests that, as expected, the propagator should take a simple form in the eigenmodes representation. In fact, by using the inverse transform of Eq. (131) we have  $D_{\mathbf{q}\nu, \mathbf{q}'\nu'}^{\text{A}}(\omega) = \Omega_{\text{BZ}} \delta(\mathbf{q} - \mathbf{q}') D_{\mathbf{q}\nu\nu'}^{\text{A}}(\omega)$ , with

$$D_{\mathbf{q}\nu\nu'}^{\text{A}}(\omega) = 2\omega_{\mathbf{q}\nu} / (\omega^2 - \omega_{\mathbf{q}\nu}^2) \delta_{\nu\nu'}. \quad (132)$$

This result can alternatively be obtained starting from the ladder operators of Appendix B. In fact, after using Eqs. (20), (109), and (131) we find

$$D_{\mathbf{q}\nu\nu'}^{\text{A}}(t t') = -i \langle \hat{T} [\hat{a}_{\mathbf{q}\nu}^{\dagger}(t) \hat{a}_{\mathbf{q}\nu}(t') + \hat{a}_{-\mathbf{q}\nu}(t) \hat{a}_{-\mathbf{q}\nu}^{\dagger}(t')] \rangle \delta_{\nu\nu'}. \quad (133)$$

An explicit evaluation of the right-hand side using the Heisenberg time evolution generated by the phonon Hamiltonian in Eq. (22) yields precisely Eq. (132), with the added advantage that it is easier to keep track of the time ordering. The result is

$$D_{\mathbf{q}\nu\nu'}^{\text{A}}(\omega) = \frac{1}{\omega - \omega_{\mathbf{q}\nu} + i\eta} - \frac{1}{\omega + \omega_{\mathbf{q}\nu} - i\eta}, \quad (134)$$

with  $\eta$  a positive real infinitesimal. This alternative approach is very common in textbooks; see, for example, Schrieffer (1983) and Schäfer and Wegener (2002). However, it does not carry general validity in a field-theoretic framework since it rests on the adiabatic approximation.

## 2. Phonons beyond the adiabatic approximation

In order to go beyond the adiabatic approximation, it is necessary to determine the complete propagator  $\mathbf{D}(\omega)$  in Eq. (123). Formally this can be done by combining Eqs. (123) and (125) to obtain the following Dyson-like equation:

$$\mathbf{D}(\omega) = \mathbf{D}^{\text{A}}(\omega) + \mathbf{D}^{\text{A}}(\omega) \Pi^{\text{NA}}(\omega) \mathbf{D}(\omega). \quad (135)$$

In this form it is apparent that the nonadiabatic phonon self-energy  $\Pi^{\text{NA}}(\omega)$  “dresses” the noninteracting phonons obtained within the adiabatic approximation, as shown schematically in Fig. 1(a). It is convenient to rewrite the Dyson equation in such a way as to show more clearly the poles of the propagator. Using Eqs. (131) and (132) we find

$$D_{\mathbf{q}\nu\nu'}^{-1}(\omega) = \frac{1}{2\omega_{\mathbf{q}\nu}} [\delta_{\nu\nu'} (\omega^2 - \omega_{\mathbf{q}\nu}^2) - 2\omega_{\mathbf{q}\nu} \Pi_{\mathbf{q}\nu\nu'}^{\text{NA}}(\omega)], \quad (136)$$

where  $\Pi_{\mathbf{q}\nu\nu'}^{\text{NA}}$  and  $D_{\mathbf{q}\nu\nu'}^{-1}$  are obtained using the transform of Eq. (131) and its inverse, respectively.

From Eq. (136) we see that the nonadiabatic self-energy  $\Pi^{\text{NA}}$  modifies the adiabatic phonon spectrum in four distinct ways: (i) the real part of the diagonal elements  $\Pi_{\mathbf{q}\nu}^{\text{NA}}$  shifts the adiabatic frequencies; (ii) the imaginary part introduces spectral broadening; (iii) the off-diagonal elements of  $\Pi_{\mathbf{q}\nu}^{\text{NA}}$  introduce a coupling between the adiabatic vibrational eigenmodes; and (iv) the frequency dependence of  $\Pi_{\mathbf{q}\nu}^{\text{NA}}(\omega)$  might lead to multiple poles for the same mode  $\nu$ , thereby introducing new structures in the phonon spectrum.

Today it is relatively common to calculate phonon linewidths arising from electron-phonon interactions (Allen, 1972b; Bauer *et al.*, 1998). Recently it has also become possible to study the frequency renormalization due to non-adiabatic effects (Saitta *et al.*, 2008; Calandra, Profeta, and Mauri, 2010).

The possibility of observing new features in vibrational spectra arising from the EPI has not been studied from first principles, but the underlying phenomenology should be similar to that of plasmon satellites in photoelectron spectra (see Sec. V.B.6). Generally speaking we expect satellites whenever  $\epsilon_e(\mathbf{r}, \mathbf{r}', \omega)$  exhibits dynamical structure close to vibrational frequencies. This can happen, for example, in the case of degenerate polar semiconductors, when phonon and plasmon energies are in resonance. In these cases, phonons and plasmons can combine into “coupled plasmon-phonon modes” (Richter, 1984), which are the electronic analog of photon polaritons (Yu and Cardona, 2010). This phenomenon was predicted theoretically (Varga, 1965) and subsequently confirmed by Raman measurements on GaAs (Mooradian and Wright, 1966). We speculate that it should be possible to obtain coupled plasmon-phonon modes from the frequency dependence of the phonon self-energy in Eq. (136); it would be interesting to perform first-principles calculations in order to shed light on these aspects.

In practical calculations the nonadiabatic corrections to the adiabatic phonon spectrum are evaluated from Eq. (136) using first-order perturbation theory, by retaining only the diagonal elements of  $\Pi^{\text{NA}}$ . If we denote the complex zeros of  $D_{\mathbf{q}\nu}^{-1}(\omega)$  by  $\tilde{\Omega}_{\mathbf{q}\nu} = \Omega_{\mathbf{q}\nu} - i\gamma_{\mathbf{q}\nu}$ , in the case of nondegenerate eigenmodes Eq. (136) gives

$$\tilde{\Omega}_{\mathbf{q}\nu}^2 = \omega_{\mathbf{q}\nu}^2 + 2\omega_{\mathbf{q}\nu}\Pi_{\mathbf{q}\nu}^{\text{NA}}(\tilde{\Omega}_{\mathbf{q}\nu}), \quad (137)$$

therefore

$$\gamma_{\mathbf{q}\nu} = -\frac{\omega_{\mathbf{q}\nu}}{\Omega_{\mathbf{q}\nu}} \text{Im}\Pi_{\mathbf{q}\nu}^{\text{NA}}(\Omega_{\mathbf{q}\nu} - i\gamma_{\mathbf{q}\nu}), \quad (138)$$

$$\Omega_{\mathbf{q}\nu}^2 = \omega_{\mathbf{q}\nu}^2 + \gamma_{\mathbf{q}\nu}^2 + 2\omega_{\mathbf{q}\nu}\text{Re}\Pi_{\mathbf{q}\nu}^{\text{NA}}(\Omega_{\mathbf{q}\nu} - i\gamma_{\mathbf{q}\nu}). \quad (139)$$

Apart from the small  $\gamma_{\mathbf{q}\nu}^2$  term in Eq. (139), these expressions are identical to those provided by Allen (1972b) and Grimvall (1981). Since nonadiabatic corrections are usually small as compared to the adiabatic phonon frequencies, Eqs. (138) and (139) are often simplified further by using the additional approximations  $|\Omega_{\mathbf{q}\nu} - \omega_{\mathbf{q}\nu}| \ll \omega_{\mathbf{q}\nu}$  and  $|\gamma_{\mathbf{q}\nu}| \ll \omega_{\mathbf{q}\nu}$ , leading to  $\gamma_{\mathbf{q}\nu} \approx -\text{Im}\Pi_{\mathbf{q}\nu}^{\text{NA}}(\omega_{\mathbf{q}\nu})$  and  $\Omega_{\mathbf{q}\nu} \approx \omega_{\mathbf{q}\nu} + \text{Re}\Pi_{\mathbf{q}\nu}^{\text{NA}}(\omega_{\mathbf{q}\nu})$ . In these forms it becomes evident that the real part of the

self-energy shifts the adiabatic phonon frequencies, and the imaginary part is responsible for the spectral broadening of the resonances. Using these expressions in Eq. (136) and going back to the time domain, it is seen that, as a result of the EPI, phonons acquire a finite lifetime given by  $\tau_{\mathbf{q}\nu}^{\text{ph}} = (2\gamma_{\mathbf{q}\nu})^{-1}$ .

### 3. Expressions for the phonon self-energy used in *ab initio* calculations

In the literature on electron-phonon interactions, the phonon self-energy  $\Pi$  is almost invariably expressed in terms of an electron-phonon vertex  $g$  and the electron Green’s function  $G$  as  $\Pi = |g|^2 GG$  in symbolic notation; see, for example, Grimvall (1981). While this has become common practice also in *ab initio* calculations, the origin of this choice is not entirely transparent. One could derive this expression directly from Eq. (1), using standard Green’s function techniques. However, this procedure does not answer the key question of how to calculate the electron-phonon matrix elements  $g$ .

A closer inspection of the theory reveals that this is a rather nontrivial point. In fact, on the one hand, a straightforward expansion of the second-quantized Hamiltonian of Eq. (68) in terms of the nuclear coordinates leads to “bare” electron-phonon matrix elements  $g^{\text{b}}$ , which contain the bare Coulomb interaction between electrons and nuclei. On the other hand, if we go back to Sec. III.B.2, we see that the electron-phonon matrix elements in DFT are dressed by the self-consistent response of the electrons. The difference between the bare and the dressed vertex is not only quantitative, but also qualitative: for example, in metals the bare vertex is long ranged, while the screened vertex is short ranged.

The relation between bare and dressed electron-phonon vertices and the derivation of explicit expressions for the phonon self-energy have been discussed by many; see, for example, Scalapino (1969) and Rickayzen (1980). In short, the argument is that the lowest-order Feynman diagram starting and ending with a phonon line must contain precisely two bare electron-phonon vertices, as shown in Fig. 1(b). By construction this diagram corresponds to having  $\Pi = |g^{\text{b}}|^2 GG$ . In order to make the transition from the bare vertex to the dressed vertex it is necessary to collect together all the proper electronic polarization diagrams around the vertex. However, these steps have been carried out only for the homogeneous electron gas (Scalapino, 1969; Rickayzen, 1980). In the following we show how the dressed electron-phonon vertex emerges from a nonperturbative analysis based on the Hedin-Baym equations.

The nonadiabatic phonon self-energy  $\Pi^{\text{NA}}$  introduced in Sec. V.A.2 can be written explicitly by combining Eqs. (119) and (125):

$$\begin{aligned} \Pi_{\kappa\alpha p, \kappa'\alpha' p'}^{\text{NA}}(\omega) &= \int d\mathbf{r}d\mathbf{r}' Z_{\kappa} \nabla_{\alpha} \delta(\mathbf{r} - \boldsymbol{\tau}_{\kappa p}^0) \\ &\quad \times [W_e(\mathbf{r}, \mathbf{r}', \omega) - W_e(\mathbf{r}, \mathbf{r}', 0)] Z_{\kappa'} \nabla_{\alpha'} \delta(\mathbf{r}' - \boldsymbol{\tau}_{\kappa' p'}^0). \end{aligned} \quad (140)$$

Using the Dyson equation for the screened Coulomb interaction, it can be seen that this expression does indeed contain electron-phonon matrix elements. In fact, by inserting Eq. (94)




FIG. 1. Diagrammatic representation of the phonon Green's function and self-energy. (a) Dyson equation for the phonon propagator, Eq. (135). The thick wavy line represents the fully interacting, nonadiabatic propagator; the thin wavy line is the adiabatic propagator; the disk is the nonadiabatic self-energy. (b) Lowest-order diagrammatic expansion of the phonon self-energy in terms of the bare electron-phonon vertices and the RPA electronic polarization. The small dots are the bare electron-phonon coupling functions, and the thin lines are the non-interacting (for example, Kohn-Sham) electron Green's functions. This diagram is the simplest possible term which begins and ends with a phonon line. (c) Nonperturbative representation of the phonon self-energy in terms of the bare coupling, the dressed coupling (large gray disk), the fully interacting electron Green's functions (thick lines), and the vertex  $\Gamma$  from Eq. (83). This diagram was proposed by Keating (1968) and describes the first line of Eq. (141). (d) Schematic representation of the relation between the dressed electron-phonon coupling  $g$  and the bare coupling  $g^b$ , from Eq. (144). Vogl (1976) reports a similar diagram, although with the bare coupling function on the far right; the difference stems from the present choice of using the irreducible polarization  $P = G\Gamma G$  instead of the reducible polarization employed by Vogl.

into Eq. (140) we find terms like  $vP_e W_e$ , and the electron-phonon matrix elements will arise from taking the gradients of  $v$  and  $W_e$ . By working in the eigenmodes representation via Eq. (131), after lengthy manipulations this procedure yields

$$\begin{aligned} \hbar\Pi_{\mathbf{q}\nu,\mathbf{q}'\nu'}^{\text{NA}}(\omega) &= \int d\mathbf{r}d\mathbf{r}'g_{\mathbf{q}\nu}^b(\mathbf{r})P_e(\mathbf{r},\mathbf{r}',\omega)g_{\mathbf{q}'\nu'}^{\text{cc}}(\mathbf{r}',\omega) \\ &\quad - \int d\mathbf{r}d\mathbf{r}'g_{\mathbf{q}\nu}^b(\mathbf{r})P_e(\mathbf{r},\mathbf{r}',0)g_{\mathbf{q}'\nu'}^*(\mathbf{r}',0), \end{aligned} \quad (141)$$

where we introduced electron-phonon ‘‘coupling functions’’ as follows. The bare coupling  $g^b$  is defined as

$$g_{\mathbf{q}\nu}^b(\mathbf{r}) = \Delta_{\mathbf{q}\nu}V^{\text{en}}(\mathbf{r}), \quad (142)$$

where  $V^{\text{en}}$  is the potential of the nuclei from Eq. (25); in practical calculations this quantity is replaced by the usual ionic pseudopotentials. The meaning of the variation  $\Delta_{\mathbf{q}\nu}$  is the same as in Eqs. (33)–(35). The dressed couplings  $g$  and  $g^{\text{cc}}$  are defined as (Hedin and Lundqvist, 1969)

$$g_{\mathbf{q}\nu}(\mathbf{r},\omega) = \int d\mathbf{r}'\epsilon_e^{-1}(\mathbf{r},\mathbf{r}',\omega)g_{\mathbf{q}\nu}^b(\mathbf{r}'), \quad (143)$$

$$g_{\mathbf{q}\nu}^{\text{cc}}(\mathbf{r},\omega) = \int d\mathbf{r}'\epsilon_e^{-1}(\mathbf{r},\mathbf{r}',\omega)g_{\mathbf{q}\nu}^{b*}(\mathbf{r}'). \quad (144)$$

Since the dielectric matrix is real at  $\omega = 0$ , we have the simple relation  $g_{\mathbf{q}\nu}^{\text{cc}}(\mathbf{r},0) = g_{\mathbf{q}\nu}^*(\mathbf{r},0)$ . In order to derive Eq. (141) it is best to carry out the algebra in the time domain. We emphasize that the result expressed by Eq. (141) is *nonperturbative* and relies solely on the harmonic approximation.

Equation (141) is in agreement with the standard result for the homogeneous electron gas (Scalapino, 1969). The same expression was also obtained by Keating (1968) using a detailed diagrammatic analysis. Keating's diagrammatic representation of the self-energy is shown in Fig. 1(c) and can be obtained from Eq. (141) by noting that, in symbolic notation,  $P_e = G\Gamma G$  from Eq. (92); therefore  $\Pi = g^b G\Gamma G g$ . For completeness we also show in Fig. 1(d) a diagrammatic representation of the dressed electron-phonon coupling function  $g$  as given by Eq. (144). This representation is obtained by observing that  $g = \epsilon^{-1}g^b$ ,  $\epsilon = 1 - vP$ , and  $P = G\Gamma G$ ; therefore  $g = g^b + vG\Gamma G g$ .

In view of practical first-principles calculations it is useful to have a simplified expression for the nonadiabatic phonon self-energy in Eq. (141). To this aim we make the following approximations:

- (i) The vertex function in Eq. (92) is set to  $\Gamma(123) = \delta(12)\delta(13)$ . This is the same approximation at the center of the GW method (Hedin, 1965; Hybertsen and Louie, 1986; Onida, Reining, and Rubio, 2002).
- (ii) The fully interacting electron Green's function  $G$  is replaced by its noninteracting counterpart, using the Kohn-Sham eigenstates and eigenvalues evaluated with the nuclei held in their equilibrium positions.
- (iii) The fully interacting dielectric matrix in Eq. (144) is replaced by the RPA +  $xc$  response obtained from a DFT calculation, as discussed in Sec. III.B.2.
- (iv) The frequency dependence of the screened electron-phonon coupling defined in Eq. (144) is neglected:  $g_{\mathbf{q}\nu}^{\text{cc}}(\mathbf{r},\omega) \simeq g_{\mathbf{q}\nu}^{\text{cc}}(\mathbf{r},0) = g_{\mathbf{q}\nu}^*(\mathbf{r},0)$ . This approximation is ubiquitous in the literature but it is never mentioned explicitly.
- (v) For notational simplicity, we consider a spin-degenerate system with time-reversal symmetry; this simplification is easily removed.

Using these assumptions we can rewrite the component of Eq. (141) for  $\mathbf{q} = \mathbf{q}'$  as



$$\begin{aligned} \hbar\Pi_{\mathbf{q}\nu'}^{\text{NA}}(\omega) &= 2\sum_{mn} \int \frac{d\mathbf{k}}{\Omega_{\text{BZ}}} g_{mn\nu}^{\text{b}}(\mathbf{k}, \mathbf{q}) g_{mn\nu'}^*(\mathbf{k}, \mathbf{q}) \\ &\times \left[ \frac{f_{m\mathbf{k}+\mathbf{q}} - f_{n\mathbf{k}}}{\varepsilon_{m\mathbf{k}+\mathbf{q}} - \varepsilon_{n\mathbf{k}} - \hbar(\omega + i\eta)} - \frac{f_{m\mathbf{k}} - f_{n\mathbf{k}+\mathbf{q}}}{\varepsilon_{m\mathbf{k}} - \varepsilon_{n\mathbf{k}+\mathbf{q}} - \hbar(\omega + i\eta)} \right]. \end{aligned} \quad (145)$$

We note that the components of the phonon self-energy for  $\mathbf{q} \neq \mathbf{q}'$  vanish due to the periodicity of the crystalline lattice. In Eq. (145) the sums run over all the Kohn-Sham states, with occupations  $f_{n\mathbf{k}}$ , and  $\eta$  is a real positive infinitesimal. In this case we indicate explicitly the factor of 2 arising from the spin degeneracy. The matrix element  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$  is the same as in Eq. (38), and it is precisely the quantity calculated by most linear-response codes. The matrix element  $g_{mn\nu}^{\text{b}}(\mathbf{k}, \mathbf{q})$  is obtained from  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$  by replacing the variation of the Kohn-Sham potential by the corresponding variation of the ionic (pseudo)potentials. The field-theoretic phonon self-energy given by Eq. (145) is in agreement with the expression derived by Calandra, Profeta, and Mauri (2010) starting from time-dependent density-functional perturbation theory.

The presence of both the bare electron-phonon matrix element and the screened matrix element in Eq. (145) has not been fully appreciated in the literature, and most *ab initio* calculations employ an approximate self-energy whereby  $g^{\text{b}}$  is replaced by  $g$ . The replacement of the bare matrix elements by their screened counterparts in the phonon self-energy goes a long way back and can be found already in the seminal work by Allen (1972b). As a result many investigators (including myself) calculated phonon lifetimes using the following expression (Grimvall, 1981):

$$\begin{aligned} \frac{1}{\tau_{\mathbf{q}\nu}^{\text{ph}}} &= \frac{2\pi}{\hbar} 2\sum_{mn} \int \frac{d\mathbf{k}}{\Omega_{\text{BZ}}} |g_{mn\nu}(\mathbf{k}, \mathbf{q})|^2 (f_{n\mathbf{k}} - f_{m\mathbf{k}+\mathbf{q}}) \\ &\times \delta(\varepsilon_{m\mathbf{k}+\mathbf{q}} - \varepsilon_{n\mathbf{k}} - \hbar\omega_{\mathbf{q}\nu}). \end{aligned} \quad (146)$$

This is obtained from Eq. (145) by taking the imaginary part and by making the replacement  $g^{\text{b}} \rightarrow g$ . While Eq. (146) can be derived from the Fermi golden rule in an independent-particle approximation [see Albers *et al.* (1976), Appendix B], the choice of the electron-phonon matrix elements is somewhat arbitrary. In future calculations of the phonon self-energy it will be important to assess the significance of using the correct vertex structure, that is, replacing  $|g_{mn\nu}(\mathbf{k}, \mathbf{q})|^2$  by  $g_{mn\nu}^{\text{b}}(\mathbf{k}, \mathbf{q})g_{mn\nu}^*(\mathbf{k}, \mathbf{q})$  in Eq. (146).

In general, the effects of the nonadiabatic self-energy on the phonon spectrum are expected to be significant only in the case of metals and small-gap semiconductors. In fact, by combining Eqs. (138), (139), and (145) it is seen that  $\Pi^{\text{NA}}$  can be large only when occupied and empty single-particle states are separated by an energy of the order of the characteristic phonon energy. In such a case, we can expect a shift of the adiabatic phonon frequencies and a concomitant broadening of the lines. A clear illustration of these effects was provided by Maksimov and Shulga (1996), who analyzed a simplified model of a metal with linear bands near the Fermi level.

Calculations of phonon linewidths based on Eq. (146) have been reported by several<sup>2</sup> and have become commonplace in first-principles studies of electron-phonon physics. On the other hand, calculations of the nonadiabatic phonon frequencies using Eq. (145) have been reported only by Lazzeri and Mauri (2006), Caudal *et al.* (2007), Piscanec *et al.* (2007), Saitta *et al.* (2008), and Calandra, Profeta, and Mauri (2010), using the approximation that the bare vertex  $g^{\text{b}}$  can be replaced by the screened vertex  $g$ . Examples of such calculations are reviewed in Sec. VII.

Equation (145) suggests several avenues worth exploring in the future: first, the use of the bare vertex should not pose a challenge in practical calculations, since this quantity is already being calculated in linear-response DFT codes. Testing the impact of the bare vertex on phonon linewidths and frequency renormalizations will be important. Second, Eq. (145) contains off-diagonal couplings, which are usually ignored. It will be interesting to check the effect of using the complete matrix self-energy. Third, the dynamical structure of the self-energy may contain interesting information such as, for example, spectral satellites and coupled phonon-plasmon modes. Last, the move from Eq. (141) to Eq. (145) involves the approximation that the frequency dependence of the electron-phonon matrix elements can be neglected. The validity of this approximation is uncertain, and there are no reference *ab initio* calculations on this. However, we note that frequency-dependent electron-phonon matrix elements have been employed systematically in theoretical models of doped semiconductors (Mahan, 1993, Sec. 6.3).

Before closing this section we note that the formalism discussed here is based on zero-temperature Green's functions. In order to extend the present results to finite temperature it is necessary to repeat all derivations using the Matsubara representation and then perform the analytic continuation of the self-energy to the real frequency axis. Detailed derivations can be found in Baym (1961) and Gillis (1970) and will not be repeated here. Fortunately it turns out that Eq. (145) can be extended to finite temperature by simply replacing the occupation factors  $f_{n\mathbf{k}}$  and  $f_{m\mathbf{k}+\mathbf{q}}$  by the corresponding Fermi-Dirac distributions.<sup>3</sup>

## B. Effects of the electron-phonon interaction on electrons

### 1. Electron self-energy: Fan-Migdal and Debye-Waller terms

In Sec. V.A we discussed the link between the Hedin-Baym equations summarized in Table I and *ab initio* calculations of phonons. We first identified a Hermitian eigenvalue problem for the vibrational frequencies via the adiabatic approximation, and then we improved upon this description by means of

<sup>2</sup>See, for example, Butler, Pinski, and Allen (1979), Bauer *et al.* (1998), Shukla *et al.* (2003), Lazzeri *et al.* (2006), Giustino, Cohen, and Louie (2007), Park, Giustino, Cohen, and Louie (2008), and Heid *et al.* (2010).

<sup>3</sup>Throughout this article, when  $f_{n\mathbf{k}}$  and  $n_{\mathbf{q}\nu}$  have the meaning of Fermi-Dirac and Bose-Einstein distributions, respectively, they are defined as follows:  $f_{n\mathbf{k}} = f[(\varepsilon_{n\mathbf{k}} - \varepsilon_{\text{F}})/k_{\text{B}}T]$  with  $f(x) = 1/(e^x + 1)$ , where  $\varepsilon_{\text{F}}$  is the Fermi energy;  $n_{\mathbf{q}\nu} = n(\hbar\omega_{\mathbf{q}\nu}/k_{\text{B}}T)$  with  $n(x) = 1/(e^x - 1)$ .

a nonadiabatic self-energy. In this section, we adopt a similar strategy in order to discuss electronic excitation energies: first we identify an approximation to the Hedin-Baym equations which does not include any electron-phonon interactions, and then we introduce an electron self-energy to incorporate such interactions.

The single most common approximation in first-principles electronic structure calculations is to describe nuclei as classical particles clamped in their equilibrium positions. Within this approximation the expectation value of the nuclear charge density operator in Eq. (67),  $\langle \hat{n}_n(\mathbf{r}) \rangle$ , is replaced by the first term in Eq. (107),  $n_n^0(\mathbf{r})$ . From Eq. (122) we see that this approximation formally corresponds to setting to zero the displacement-displacement correlation function of the nuclei. This observation suggests that, in order to unambiguously single out electron-phonon interactions in the Hedin-Baym equations, we need to define a noninteracting problem by setting  $D_{\kappa\alpha p, \kappa'd'p'} = 0$  and identify the electron-phonon interaction with the remainder. In the following, we write an equation of motion for the electrons analogous to Eq. (84), except with the nuclei clamped in their equilibrium positions; then we use a Dyson-like equation to recover the fully interacting electron Green's function.

The equation of motion for the electron Green's function at *clamped nuclei*, which we denote as  $G^{\text{cn}}$ , reads

$$\left[ i\hbar \frac{\partial}{\partial t_1} + \frac{\hbar^2}{2m_e} \nabla^2(1) - V_{\text{tot}}^{\text{cn}}(1) \right] G^{\text{cn}}(12) - \int d3 \Sigma_e^{\text{cn}}(13) G^{\text{cn}}(32) = \delta(12). \quad (147)$$

Here the potential  $V_{\text{tot}}^{\text{cn}}$  differs from its counterpart  $V_{\text{tot}}$  of Eq. (79) in that the total density of electrons and nuclei  $\langle \hat{n} \rangle$  is replaced by the density calculated at clamped nuclei  $\langle \hat{n}^{\text{cn}} \rangle$ :

$$V_{\text{tot}}^{\text{cn}}(1) = \int d2 v(1, 2) \langle \hat{n}^{\text{cn}}(2) \rangle, \quad (148)$$

where

$$\langle \hat{n}^{\text{cn}}(1) \rangle = -i\hbar \sum_{\sigma_1} G^{\text{cn}}(11^+) + n_n^0(\mathbf{r}_1). \quad (149)$$

The term  $\Sigma_e^{\text{cn}}$  in Eq. (147) represents the electronic part of Hedin's self-energy in Eq. (85), evaluated at clamped nuclei:

$$\Sigma_e^{\text{cn}}(12) = i\hbar \int d(34) G^{\text{cn}}(13) \Gamma^{\text{cn}}(324) W_e^{\text{cn}}(41^+). \quad (150)$$

In this expression, the vertex  $\Gamma^{\text{cn}}$  and the screened Coulomb interaction  $W_e^{\text{cn}}$  are both evaluated via the Hedin-Baym equations at clamped nuclei. Equations (147)–(150) lead directly to the well-known Hedin's equations (Hedin, 1965). Hedin's equations and the associated *GW* method at clamped nuclei are addressed in a number of excellent reviews (Hedin and Lundqvist, 1969; Hybertsen and Louie, 1986; Aryasetiawan and Gunnarsson, 1998; Onida, Reining, and Rubio, 2002); hence they will not be discussed here.

In order to recover the complete Hedin-Baym equation of motion, Eq. (84), starting from Eqs. (147)–(150), it is sufficient to introduce the Dyson equation

$$G(12) = G^{\text{cn}}(12) + \int d(34) G^{\text{cn}}(13) \Sigma^{\text{ep}}(34) G(42), \quad (151)$$

together with the electron self-energy  $\Sigma^{\text{ep}}$  arising from electron-phonon interactions

$$\Sigma^{\text{ep}} = \Sigma^{\text{FM}} + \Sigma^{\text{DW}} + \Sigma^{\text{dGW}}, \quad (152)$$

where

$$\Sigma^{\text{FM}}(12) = i\hbar \int d(34) G(13) \Gamma(324) W_{\text{ph}}(41^+), \quad (153)$$

$$\Sigma^{\text{DW}}(12) = \int d3 v(13) [\langle \hat{n}(3) \rangle - \langle \hat{n}^{\text{cn}}(3) \rangle] \delta(12), \quad (154)$$

$$\Sigma^{\text{dGW}}(12) = \Sigma_e(12) - \Sigma_e^{\text{cn}}(12). \quad (155)$$

We emphasize that Eqs. (147)–(155) are just an alternative formulation of the Hedin-Baym equations in Table I. The advantage of this formulation is that it better reflects standard practice, whereby the DFT equations and the *GW* quasiparticle corrections are evaluated at clamped nuclei. Equations (147)–(155) are formally exact within the harmonic approximation.

A schematic representation of the Dyson equation for the electron Green's function and the decomposition of the electron self-energy are given in Fig. 2. The self-energy contribution  $\Sigma^{\text{FM}}$  in Eq. (153) is a dynamic correction to the electronic excitation energies and is analogous to the *GW* self-energy at clamped nuclei. Indeed, in the same way as the correlation part of the standard *GW* self-energy describes the effect of the dynamic electronic polarization upon the addition of electrons or holes to the system, the term  $GW_{\text{ph}}$  in Eq. (153) describes the effect of the dynamic polarization of the lattice.

In the semiconductor community, the self-energy obtained from Eq. (153) by setting  $\Gamma(123) = \delta(13)\delta(23)$  is commonly referred to as the Fan self-energy (Fan, 1951; Allen and Heine, 1976; Allen and Cardona, 1981; Cardona, 2001). In the metal and superconductor communities, the same term is traditionally referred to as the self-energy in the Migdal approximation (Migdal, 1958; Engelsberg and Schrieffer, 1963; Scalapino, 1969; Schrieffer, 1983). By extension we refer to the self-energy  $\Sigma^{\text{FM}}$  in Eq. (153) as the ‘‘Fan-Migdal’’ (FM) self-energy.

The static term  $\Sigma^{\text{DW}}$  in Eq. (154) corresponds to the difference between the self-consistent potential  $V_{\text{tot}}$  calculated for the fully interacting system and the same potential evaluated with the nuclei clamped in their equilibrium positions  $V_{\text{tot}}^{\text{cn}}$ . Intuitively this term corresponds to a time-independent correction to the ‘‘crystal potential’’ that arises from the fuzziness of the nuclear charge density around the equilibrium nuclear positions. This term is similar to the one appearing in the study of the temperature dependence of x-ray




FIG. 2. Diagrammatic representation of the electron Green's function and electron-phonon self-energy. (a) Dyson equation for the electron Green's function, Eq. (151). The thick straight line represents the fully dressed electron propagator, the thin straight line is the propagator calculated at clamped nuclei, and the disk is the electron-phonon self-energy. (b) Decomposition of the electron-phonon self-energy into Fan-Migdal self-energy, Eq. (153), Debye-Waller contribution, Eq. (154), and the remainder given by Eq. (155). (c) Fan-Migdal electron-phonon self-energy expressed in terms of the dressed electron-phonon coupling function (dark gray disk as in Fig. 1), the fully interacting electron Green's functions (thick straight line), the fully interacting phonon propagator (thick wavy line), and the vertex  $\Gamma$  from Eq. (83). (d) Debye-Waller contribution resulting from the fully interacting phonon propagator (thick wavy line) and the matrix element in Eq. (40) (hatched disk). (e) Correction to Hedin's  $GW$  self-energy arising from the modification of the electronic structure induced by the electron-phonon interaction.  $W_e$  is the screened Coulomb interaction of Eq. (94) (bold dashed double line).  $W_e^{\text{cn}}$  is the screened Coulomb interaction evaluated at clamped nuclei (thin dashed double line).  $\Gamma^{\text{cn}}$  is the vertex of Eq. (83), but evaluated at clamped nuclei.

diffraction and neutron diffraction spectra (Mermin, 1966; Ashcroft and Mermin, 1976) and is commonly referred to as the Debye-Waller term (Antončík, 1955; Walter *et al.*, 1970; Cardona and Thewalt, 2005). Hedin and Lundqvist (1969) did not include this term in their classic work; however, this contribution was discussed by Allen and Heine (1976) [see also Allen (1978)].

The last term Eq. (152),  $\Sigma^{\text{dGW}}$ , is the correction to the standard Hedin self-energy arising from the fact that the fully interacting electron Green's function and density are slightly different from those evaluated at clamped nuclei, owing to the electron-phonon interaction. The magnitude of this term

corresponds to a fraction of the  $GW$  quasiparticle corrections at clamped nuclei. Since  $\Sigma^{\text{dGW}}$  has never been investigated so far, we will not discuss this term further.

## 2. Expressions for the electron self-energy used in *ab initio* calculations

A complete self-consistent solution of Eqs. (147)–(155) from first principles is not possible at present, and one has to replace the various entries of Eq. (152) by the best approximations available. In practice, one resorts either to DFT or to  $GW$  calculations; recent progress is reviewed in Secs. VIII and IX.

Using Eqs. (110), (130), (131), and (144) we can rewrite the Fan-Migdal self-energy as follows:

$$\begin{aligned} \Sigma^{\text{FM}}(12) &= i \sum_{\nu\nu'} \int \frac{d\omega}{2\pi} \frac{d\mathbf{q}}{\Omega_{\text{BZ}}} d(34) e^{-i\omega(t_4-t_1^+)} \\ &\times G(13) \Gamma(324) g_{\mathbf{q}\nu}^{\text{cc}}(\mathbf{r}_4, \omega) \\ &\times D_{\mathbf{q}\nu\nu'}(\omega) g_{\mathbf{q}\nu'}(\mathbf{r}_1, \omega). \end{aligned} \quad (156)$$

This shows that the Fan-Migdal self-energy is, in symbolic notation, of the type  $\Sigma = g^2 D G \Gamma$ ; a graphical representation of this term is given in Fig. 2(c). In order to make Eq. (156) amenable to *ab initio* calculations, it is common to make the following approximations, which are similar to those introduced earlier for the phonon self-energy:

- (i) The vertex  $\Gamma(123)$  is set to  $\delta(13)\delta(23)$ .
- (ii) The fully interacting electron Green's function is replaced by the Kohn-Sham Green's function evaluated at clamped nuclei.
- (iii) The fully interacting phonon propagator  $D_{\mathbf{q}\nu\nu'}(\omega)$  is replaced by the adiabatic propagator  $D_{\mathbf{q}\nu\nu'}^{\text{A}}(\omega)$  from Eq. (134).
- (iv) The screened electron-phonon vertex is evaluated using the RPA +  $xc$  electronic screening from a DFT calculation.
- (v) The frequency dependence of the electron-phonon coupling is neglected,  $g_{\mathbf{q}\nu}(\mathbf{r}, \omega) \approx g_{\mathbf{q}\nu}(\mathbf{r}, 0)$ .

After using these approximations in Eq. (156), we obtain the following result for the  $\mathbf{k} = \mathbf{k}'$  matrix elements of the Fan-Migdal self-energy in the basis of Kohn-Sham states:

$$\begin{aligned} \Sigma_{m\mathbf{k}'\mathbf{k}}^{\text{FM}}(\omega) &= \frac{1}{\hbar} \sum_{m\nu} \int \frac{d\mathbf{q}}{\Omega_{\text{BZ}}} g_{m\nu}^*(\mathbf{k}, \mathbf{q}) g_{m\nu'}(\mathbf{k}, \mathbf{q}) \\ &\times \left[ \frac{1 - f_{m\mathbf{k}+\mathbf{q}}}{\omega - \varepsilon_{m\mathbf{k}+\mathbf{q}}/\hbar - \omega_{\mathbf{q}\nu} + i\eta} \right. \\ &\left. + \frac{f_{m\mathbf{k}+\mathbf{q}}}{\omega - \varepsilon_{m\mathbf{k}+\mathbf{q}}/\hbar + \omega_{\mathbf{q}\nu} - i\eta} \right]. \end{aligned} \quad (157)$$

Here  $f_{m\mathbf{k}+\mathbf{q}} = 1$  for occupied Kohn-Sham states and 0 otherwise, and the matrix element  $g_{m\nu}(\mathbf{k}, \mathbf{q})$  is obtained from Eq. (38). As for the phonon self-energy in Eq. (145), in this case the components of the electron self-energy for  $\mathbf{k} \neq \mathbf{k}'$  also vanish due to the periodicity of the lattice. The result in Eq. (157) is obtained by closing the contour of the frequency integration in the upper complex plane, owing to

the  $t_1^+$  in the exponential of Eq. (156). The infinitesimals inside the electron and phonon propagators, which reflect the time ordering, are crucial to obtain the correct result (Schrieffer, 1983). The spin label is omitted in Eq. (157) since this contribution to the self-energy is diagonal in the spin indices.

This result is valid only at zero temperature. The extension to finite temperature requires going through the Matsubara representation and then continuing the self-energy from the imaginary axis to the real axis. The procedure is described in many textbooks; see, for example, Sec. 3.5 of Mahan (1993). The result is that at finite temperature the square brackets of Eq. (157) are to be modified as follows:

$$\left[ \frac{1 - f_{m\mathbf{k}+\mathbf{q}}}{\dots + i\eta} + \frac{f_{m\mathbf{k}+\mathbf{q}}}{\dots - i\eta} \right] \rightarrow \left[ \frac{1 - f_{m\mathbf{k}+\mathbf{q}} + n_{q\nu}}{\dots + i\eta} + \frac{f_{m\mathbf{k}+\mathbf{q}} + n_{q\nu}}{\dots + i\eta} \right], \quad (158)$$

where  $f_{m\mathbf{k}+\mathbf{q}}$  and  $n_{q\nu}$  are now Fermi-Dirac and Bose-Einstein distribution functions, respectively (see footnote 3). The change of sign in the imaginary infinitesimal on the second fraction has to do with the fact that in the Matsubara formalism the analytic continuation from the imaginary frequency axis to the real axis through the upper complex plane leads to the so-called *retarded* self-energy, that is, a self-energy with all poles below the real axis (Abrikosov, Gor'kov, and Dzyaloshinski, 1975; Mahan, 1993).

The Debye-Waller term in Eq. (154) can also be written in a form which is convenient for practical calculations by expanding the total density operator  $\hat{n}(3)$  to second order in the atomic displacements. Using Eqs. (79) and (109) we find

$$\Sigma^{\text{DW}}(12) = \delta(12) \frac{i\hbar}{2} \sum_{\substack{\kappa\alpha p \\ \kappa'\alpha' p'}} \frac{\partial^2 V_{\text{tot}}(1)}{\partial \tau_{\kappa\alpha p}^0 \partial \tau_{\kappa'\alpha' p'}^0} D_{\kappa\alpha p, \kappa'\alpha' p'}(t_1^+, t_1). \quad (159)$$

In order to arrive at this result, it is necessary to make the additional approximation that the electronic field operators and the operators for the nuclear displacements are uncorrelated, that is  $\langle \hat{n} \Delta \hat{\tau}_{\kappa\alpha p} \rangle = \langle \hat{n} \rangle \langle \Delta \hat{\tau}_{\kappa\alpha p} \rangle$ , and similarly for the second power of the displacements. This requirement was noted by Gillis (1970) and is trivially satisfied if we describe phonons within the adiabatic approximation of Sec. V.A.1. Equation (159) motivates the diagrammatic representation of the Debye-Waller self-energy shown in Fig. 2(d), whereby the phonon line begins and ends at the same time point. Note that Eq. (159) involves the variation of the screened potential  $V_{\text{tot}}$ ; this result, which we derived here starting from Schwinger's functional derivative technique, is also obtained when starting from a perturbative diagrammatic analysis (Marini, Ponc e, and Gonze, 2015). The Debye-Waller self-energy can be simplified further if we make use of the following approximations, in the same spirit as for the Fan-Migdal self-energy:

- (vi) The fully interacting phonon propagator is replaced by the adiabatic propagator  $D_{q\nu}^A(\omega)$  from Eq. (134).

- (vii) The total many-body potential  $V_{\text{tot}}$  of Eq. (79) is replaced by the Kohn-Sham potential  $V^{\text{KS}}(\mathbf{r})$  evaluated at clamped nuclei. Strictly speaking, the Kohn-Sham effective potential also includes contributions from exchange and correlation, which in the Hedin-Baym equations are all contained in the electron self-energy. However, the present discussion is unchanged if we add any local and frequency-independent potential to  $V_{\text{tot}}$  in Eq. (84), while removing the same potential from the self-energy (Keating, 1968).

Using these simplifications together with Eqs. (130) and (131), we can write the  $\Sigma^{\text{DW}}$  in the basis of Kohn-Sham eigenstates as follows:

$$\Sigma_{nn'\mathbf{k}}^{\text{DW}} = \sum_{\nu} \int \frac{d\mathbf{q}}{\Omega_{\text{BZ}}} g_{nn'\nu}^{\text{DW}}(\mathbf{k}, \mathbf{q}, -\mathbf{q}), \quad (160)$$

where the Debye-Waller matrix element  $g^{\text{DW}}$  is obtained from Eq. (40), and the presence of only the diagonal terms  $\nu = \nu'$  is a result of the Kronecker delta in Eq. (132). In going from Eq. (159) to Eq. (160) the frequency integration is performed by using Eq. (134), after closing the contour in the lower half plane. The resulting expression is diagonal in the spin indices.

The expression for the Debye-Waller term in Eq. (160) is valid only at zero temperature. In this case the extension to finite temperature is immediate since the self-energy does not involve the electron Green's function; hence we need to evaluate only the canonical average of Eq. (133) at equal times. The result is that Eq. (160) is simply to be modified as follows:

$$g_{nn'\nu}^{\text{DW}}(\mathbf{k}, \mathbf{q}, -\mathbf{q}) \rightarrow g_{nn'\nu}^{\text{DW}}(\mathbf{k}, \mathbf{q}, -\mathbf{q})(2n_{q\nu} + 1), \quad (161)$$

with  $n_{q\nu}$  the Bose-Einstein occupations (see footnote 3).

The Debye-Waller contribution to the electron self-energy is almost invariably ignored in the literature on metals and superconductors, but it is well known in the theory of temperature-dependent band structures of semiconductors (Allen and Heine, 1976; Allen and Cardona, 1981; Marini, 2008; Giustino, Louie, and Cohen, 2010; Faber *et al.*, 2011). Neglecting  $\Sigma^{\text{DW}}$  in metals is partly justified by the fact that this term is frequency independent; therefore it is expected to be a slowly varying function over each Fermi-surface sheet. A detailed first-principles analysis of this aspect is currently lacking.

### 3. Temperature dependence of electronic band structures

Once the electron self-energy has been determined as in Sec. V.B.2, it is possible to study the modification of the electronic structure induced by the EPI. To this aim it is convenient to rewrite Eq. (151) in the basis of Kohn-Sham eigenstates as follows:

$$G_{nn'\mathbf{k}}^{-1}(\omega) = G_{nn'\mathbf{k}}^{\text{cn},-1}(\omega) - \Sigma_{nn'\mathbf{k}}^{\text{ep}}(\omega). \quad (162)$$

Assuming that the electronic structure problem at clamped nuclei has been solved using DFT or DFT + *GW* calculations,

the Green's function  $G^{\text{cn}}$  can be written in terms of simple poles at the Kohn-Sham or quasiparticle eigenvalues  $\varepsilon_{n\mathbf{k}}$  (Hedin and Lundqvist, 1969). In this case Eq. (162) reduces to

$$G_{nn'\mathbf{k}}^{-1}(\omega) = (\hbar\omega - \tilde{\varepsilon}_{n\mathbf{k}})\delta_{nn'} - \Sigma_{nn'\mathbf{k}}^{\text{ep}}(\omega), \quad (163)$$

where  $\tilde{\varepsilon}_{n\mathbf{k}} = \varepsilon_{n\mathbf{k}} \pm i\hbar\eta$  with the upper and lower signs corresponding to occupied and unoccupied states, respectively. The spin indices are omitted since these self-energy contributions do not mix states with opposite spin.

In order to gain insight into the effects of the electron-phonon interaction, we start from the approximation that  $\Sigma^{\text{ep}}$  leads only to a small shift of the quasiparticle poles, from the noninteracting energies  $\varepsilon_{n\mathbf{k}}$  to the renormalized energies  $\tilde{E}_{n\mathbf{k}} = E_{n\mathbf{k}} + i\Gamma_{n\mathbf{k}}$ . In this approximation, the fully interacting Green's function is expressed as a sum of simple poles, given by the zeros of Eq. (163):

$$E_{n\mathbf{k}} = \varepsilon_{n\mathbf{k}} + \text{Re}\Sigma_{nn\mathbf{k}}^{\text{ep}}(\tilde{E}_{n\mathbf{k}}/\hbar), \quad (164)$$

$$\Gamma_{n\mathbf{k}} = \text{Im}\Sigma_{nn\mathbf{k}}^{\text{ep}}(\tilde{E}_{n\mathbf{k}}/\hbar). \quad (165)$$

As in the case of vibrational frequencies in Eq. (137), we are considering for simplicity nondegenerate electronic states and making the assumption that the off-diagonal elements of the self-energy  $\Sigma_{nn'\mathbf{k}}^{\text{ep}}$  with  $n \neq n'$  can be neglected. In more general situations the right-hand side of Eq. (163) needs to be diagonalized, or alternatively the off-diagonal terms  $\Sigma_{nn'\mathbf{k}}^{\text{ep}}$  need to be treated perturbatively. The energies  $E_{n\mathbf{k}}$  obtained from Eq. (164) yield the band structure renormalized by the EPIs, to be discussed later. The imaginary part  $\Gamma_{n\mathbf{k}}$  in Eq. (165) is connected with the quasiparticle lifetimes and is discussed in Sec. V.B.4.

Equation (164) is to be solved self-consistently for  $E_{n\mathbf{k}}$  and  $\Gamma_{n\mathbf{k}}$ . When Eq. (164) is used in combination with the standard approximations to the Fan-Migdal and Debye-Waller self-energies given by Eqs. (157) and (160), the result that one obtains is equivalent to describing electron-phonon couplings to second order in the Brillouin-Wigner perturbation theory (Mahan, 1993). Similarly one recovers the more basic Rayleigh-Schrödinger perturbation theory by making the replacements  $E_{n\mathbf{k}} \rightarrow \varepsilon_{n\mathbf{k}}$  and  $\Gamma_{n\mathbf{k}} \rightarrow 0$  in Eq. (164).

By combining Eqs. (157) and (158), (160) and (161), and (164), we obtain the temperature-dependent “band structure renormalization” arising from the EPI:

$$\begin{aligned} E_{n\mathbf{k}} = & \varepsilon_{n\mathbf{k}} + \sum_{\nu} \int \frac{d\mathbf{q}}{\Omega_{\text{BZ}}} \sum_m |g_{m\nu}(\mathbf{k}, \mathbf{q})|^2 \\ & \times \text{Re} \left[ \frac{1 - f_{m\mathbf{k}+\mathbf{q}} + n_{\mathbf{q}\nu}}{E_{n\mathbf{k}} - \varepsilon_{m\mathbf{k}+\mathbf{q}} - \hbar\omega_{\mathbf{q}\nu} + i\Gamma_{n\mathbf{k}}} \right. \\ & \left. + \frac{f_{m\mathbf{k}+\mathbf{q}} + n_{\mathbf{q}\nu}}{E_{n\mathbf{k}} - \varepsilon_{m\mathbf{k}+\mathbf{q}} + \hbar\omega_{\mathbf{q}\nu} + i\Gamma_{n\mathbf{k}}} \right] \\ & + \sum_{\nu} \int \frac{d\mathbf{q}}{\Omega_{\text{BZ}}} g_{n\nu\nu}^{\text{DW}}(\mathbf{k}, \mathbf{q}, -\mathbf{q})(2n_{\mathbf{q}\nu} + 1). \quad (166) \end{aligned}$$

For practical calculations it is important to bear in mind that this result rests on the approximations (i)–(vii) introduced previously, as well as the harmonic approximation.

The theory of temperature-dependent band structures developed by Allen and Heine (1976) makes two additional approximations in addition to Eq. (166): the Brillouin-Wigner perturbation theory is replaced by the Rayleigh-Schrödinger perturbation theory, and the phonon energies in the denominators are neglected. Using these additional approximations Eq. (166) becomes

$$\begin{aligned} E_{n\mathbf{k}} = & \varepsilon_{n\mathbf{k}} + \sum_{\nu} \int \frac{d\mathbf{q}}{\Omega_{\text{BZ}}} \left[ \sum_m \frac{|g_{m\nu}(\mathbf{k}, \mathbf{q})|^2}{\varepsilon_{n\mathbf{k}} - \varepsilon_{m\mathbf{k}+\mathbf{q}}} \right. \\ & \left. + g_{n\nu\nu}^{\text{DW}}(\mathbf{k}, \mathbf{q}, -\mathbf{q}) \right] (2n_{\mathbf{q}\nu} + 1), \quad (167) \end{aligned}$$

which is referred to as the “adiabatic Allen-Heine formula.” By setting  $T = 0$  the Bose-Einstein factors  $n_{\mathbf{q}\nu}$  vanish and we have the so-called “zero-point renormalization” of the energy bands  $\Delta E_{n\mathbf{k}}^{\text{ZP}} = E_{n\mathbf{k}}(T = 0) - \varepsilon_{n\mathbf{k}}$ . This is the modification of the electronic energies evaluated at clamped nuclei, which arises from the zero-point fluctuations of the atoms around their equilibrium sites.

An expression that is essentially identical to Eq. (167) can also be obtained directly from Eq. (1) using second-order Rayleigh-Schrödinger perturbation theory in Fock space, following the same lines as in Kittel (1963). A detailed derivation of the formalism starting from Eq. (1) was given by Chakraborty and Allen (1978).

Historically, the Allen-Heine theory (Allen and Heine, 1976) was developed by starting from a straightforward perturbation theory expansion of the electron energies in terms of the atomic displacements within the adiabatic approximation, followed by a canonical average of the displacements using Bose-Einstein statistics. It is reassuring that, after making a few well-defined approximations, a field-theoretic method leads to the same result.

Equation (167) was employed in many semiempirical calculations.<sup>4</sup> More recently, this expression was used in the context of first-principles DFT calculations by Marini (2008) and Giustino, Louie, and Cohen (2010). DFT calculations of band structure renormalization based on Eqs. (166) or (167) are becoming increasingly popular, and the latest developments are reviewed in Sec. IX.A.1.

The nature of the band structure renormalization by electron-phonon interactions can be understood at a qualitative level by considering a drastically simplified model, consisting of a semiconductor with parabolic and nondegenerate valence and conduction bands, with the band extrema coupled to all other states by a dispersionless phonon mode of frequency  $\omega_0$ . If the Debye-Waller matrix elements are much

<sup>4</sup>See, for example, Allen and Cardona (1981, 1983), Lautenschlager, Allen, and Cardona (1985), Gopalan, Lautenschlager, and Cardona (1987), Zollner, Cardona, and Gopalan (1992), and Olguin, Cardona, and Cantarero (2002). Detailed reviews of early calculations and comparison to experiments can be found in Cardona (2001, 2005) and Cardona and Thewalt (2005).




FIG. 3. Temperature-dependent band gap and lifetimes in an idealized semiconductor or insulator. (a) Temperature dependence of the band gap according to Eq. (168) (thick solid blue line). The straight thin black line is the asymptotic expansion at high temperature; this line intercepts the vertical axis at the band gap calculated with clamped nuclei  $E_g^{\text{cn}}$ . The difference between the latter value and the band gap at  $T = 0$  including the EPI gives the zero-point renormalization  $\Delta E_g^{\text{ZP}}$ . (b) Temperature dependence of the electron linewidth (solid blue line) and lifetimes (dashed red line) using the same model as in (a). The zero-point broadening is  $\Gamma_{nk}^{\text{ZP}}$ . This simplified trend is valid only when the electron energy is at least one phonon energy away from a band extremum, so that both phonon emission and phonon absorption processes are allowed. The parameters of the model are  $E_g^{\text{cn}} = 1$  eV,  $\Delta E_g^{\text{ZP}} = 100$  meV,  $\hbar\omega_0 = 100$  meV, and  $\Gamma_{nk}^{\text{ZP}} = 50$  meV. These values are representative of common semiconductors.

smaller than the Fan-Migdal matrix elements, then the dominant contributions to Eq. (167) arise from denominators such as  $\varepsilon_{n\mathbf{k}} - \varepsilon_{n\mathbf{k}+\mathbf{q}} \approx \pm \hbar^2 |\mathbf{q}|^2 / 2m_n^*$ , where the upper and lower signs are for the valence and conduction bands, respectively, and  $m_n^*$  are the effective masses. As a result the temperature dependence of the band gap takes the form

$$E_g(T) = E_g^{\text{cn}} - |\Delta E_g^{\text{ZP}}| [1 + 2n(\hbar\omega_0/k_B T)], \quad (168)$$

where  $E_g^{\text{cn}}$  is the gap at clamped nuclei and  $|\Delta E_g^{\text{ZP}}|$  is the zero-point correction. The negative sign in Eq. (168) arises from the curvatures of the valence and conduction bands. In this example the band gap decreases with temperature: this is a well-known effect in semiconductor physics and is often referred to as the ‘‘Varshni effect’’ (Varshni, 1967). The first measurements of such effects were performed by Becker and Fan (1949) and stimulated the development of the first theory of temperature-dependent band gaps (Fan, 1951). A schematic illustration of this qualitative model is provided in Fig. 3(a). The redshift of the gap as a function of temperature is seen in many, albeit not all, semiconductors. For example, copper halides (Göbel *et al.*, 1998) and lead halide perovskites (D’Innocenzo *et al.*, 2014) exhibit an ‘‘inverse Varshni’’ effect, that is, a blueshift of the gap with temperature; in addition, some chalcopyrites exhibit a nonmonotonic temperature dependence of the band gap (Bhosale *et al.*, 2012). We also point out that the qualitative model shown in Fig. 3(a) does not take into account the subtle temperature dependence of the

band gap renormalization at very low temperature. These effects were recently investigated by Allen and Nery (2017).

#### 4. Carrier lifetimes

While the real part of the poles in Eq. (164) describes the energy level renormalization induced by the electron-phonon coupling, the imaginary part  $\Gamma_{nk}$  in Eq. (165) carries information on the spectral broadening, discussed in Sec. V.B.5, and on quasiparticle lifetimes, which we discuss next.

After transforming  $G_{nn'\mathbf{k}}(\omega)$  from Eq. (163) into the time domain it is seen that, for an electron or hole added to the system at time  $t$  in the state  $|n\mathbf{k}\rangle$ , the probability amplitude to persist in the same state decreases as  $\exp[\Gamma_{nk}(t' - t)/\hbar]$ . Using Eqs. (157) and (165) it can be seen that  $\Gamma_{nk} < 0$  for an electron added to the system and  $\Gamma_{nk} > 0$  for a hole. Therefore the average time spent by the particle in the state  $|n\mathbf{k}\rangle$  is  $\tau_{nk} = \hbar/(2|\Gamma_{nk}|)$ .

A popular expression for the electron and hole lifetimes is obtained by making the replacement  $\tilde{E}_{n\mathbf{k}} \rightarrow \varepsilon_{n\mathbf{k}}$  in Eq. (157) and by taking the absolute value of the imaginary part. We find

$$\begin{aligned} \frac{1}{\tau_{nk}} = & \frac{2\pi}{\hbar} \sum_{m\nu} \int \frac{d\mathbf{q}}{\Omega_{\text{BZ}}} |g_{nm\nu}(\mathbf{k}, \mathbf{q})|^2 \\ & \times |(1 - f_{m\mathbf{k}+\mathbf{q}})\delta(\varepsilon_{n\mathbf{k}} - \hbar\omega_{\mathbf{q}\nu} - \varepsilon_{m\mathbf{k}+\mathbf{q}}) \\ & - f_{m\mathbf{k}+\mathbf{q}}\delta(\varepsilon_{n\mathbf{k}} + \hbar\omega_{\mathbf{q}\nu} - \varepsilon_{m\mathbf{k}+\mathbf{q}})|. \end{aligned} \quad (169)$$

A more accurate expression is discussed after Eq. (174). The extension of this result to finite temperature is obtained by taking the absolute value of the imaginary part of Eq. (158):

$$\begin{aligned} \frac{1}{\tau_{nk}} = & \frac{2\pi}{\hbar} \sum_{m\nu} \int \frac{d\mathbf{q}}{\Omega_{\text{BZ}}} |g_{nm\nu}(\mathbf{k}, \mathbf{q})|^2 \\ & \times [(1 - f_{m\mathbf{k}+\mathbf{q}} + n_{\mathbf{q}\nu})\delta(\varepsilon_{n\mathbf{k}} - \hbar\omega_{\mathbf{q}\nu} - \varepsilon_{m\mathbf{k}+\mathbf{q}}) \\ & + (f_{m\mathbf{k}+\mathbf{q}} + n_{\mathbf{q}\nu})\delta(\varepsilon_{n\mathbf{k}} + \hbar\omega_{\mathbf{q}\nu} - \varepsilon_{m\mathbf{k}+\mathbf{q}})]. \end{aligned} \quad (170)$$

We emphasize that the change of sign in the third line results from the analytic continuation to the retarded self-energy. Equation (170) coincides with the expression that one would obtain by using the standard Fermi golden rule (Grimvall, 1981). The intuitive interpretation of this result is that the quasiparticle lifetime is reduced by processes of phonon emission and absorption, corresponding to the second and third lines of Eq. (170), respectively. We note that in deriving Eq. (170) we did not consider the Debye-Waller self-energy. This is because the diagonal matrix elements of  $\Sigma^{\text{DW}}$  are purely real; hence they do not contribute to the quasiparticle widths (Lautenschlager, Allen, and Cardona, 1986). *Ab initio* calculations of carrier lifetimes using Eq. (170) were first reported by Eiguren *et al.* (2002) and Eiguren, de Gironcoli *et al.* (2003). These applications and more recent developments are reviewed in Sec. X.A.

If we evaluate Eq. (170) for the same simplified model introduced for the temperature renormalization, and we neglect the phonon energy in the Dirac delta functions, we obtain  $\Gamma_{nk}(T) = \Gamma_{nk}^{\text{ZP}}[1 + 2n(\hbar\omega_0/k_B T)]$ , where  $\Gamma_{nk}^{\text{ZP}}$  is the linewidth at  $T = 0$ . The dependence of the linewidth and the corresponding lifetime on temperature for this model are

shown in Fig. 3(b). This trend is typical in semiconductors (Lautenschlager, Garriga, Logothetidis, and Cardona, 1987; Lautenschlager, Garriga, Vina, and Cardona, 1987).

### 5. Kinks and satellites

In many cases of interest, the use of the Brillouin-Wigner perturbation theory as given by Eqs. (164) and (165) is not sufficient to provide an adequate description of EPIs, and it becomes necessary to go back to the complete Dyson equation (163). Generally speaking a direct solution of the Dyson equation is important in all those cases where the electronic energy scales are comparable to phonon energies, namely, in metals (including superconductors), narrow-gap semiconductors, and doped semiconductors. In order to study these systems, it is convenient to introduce an auxiliary function called the “spectral density function” or simply spectral function.

In its simplest version the spectral function is defined as (Abrikosov, Gor’kov, and Dzyaloshinski, 1975; Mahan, 1993)

$$A(\mathbf{k}, \omega) = -\frac{1}{\pi} \sum_n \text{Im} G_{n\mathbf{k}}^{\text{ret}}(\omega), \quad (171)$$

where the superscript “ret” stands for “retarded” and simply indicates that all poles of the Green’s function  $G_{n\mathbf{k}}(\omega)$  in the upper complex plane must be replaced by their complex conjugate. The spectral function is positive definite and carries the meaning of a many-body momentum-resolved density of states (Abrikosov, Gor’kov, and Dzyaloshinski, 1975). This is precisely the function that is probed by angle-resolved photoelectron spectroscopy experiments (ARPES) (Damascelli, Hussain, and Shen, 2003). Using Eq. (163) the spectral function can be rewritten as

$$A(\mathbf{k}, \omega) = \sum_n \frac{-(1/\pi) \text{Im} \Sigma_{n\mathbf{k}}^{\text{ep}}(\omega)}{[\hbar\omega - \varepsilon_{n\mathbf{k}} - \text{Re} \Sigma_{n\mathbf{k}}^{\text{ep}}(\omega)]^2 + [\text{Im} \Sigma_{n\mathbf{k}}^{\text{ep}}(\omega)]^2}. \quad (172)$$

In order to obtain the correct spectral function, it is important to use the retarded self-energy. This is done by using Eq. (158) for the Fan-Migdal term, while the static Debye-Waller term remains unchanged.

It is often convenient to approximate the spectral function as a sum of quasiparticle peaks. To this aim, one performs a linear expansion of Eq. (172) around each quasiparticle energy  $E_{n\mathbf{k}}$  to obtain

$$A(\mathbf{k}, \omega) = \sum_n Z_{n\mathbf{k}} \frac{-(1/\pi) Z_{n\mathbf{k}} \text{Im} \Sigma_{n\mathbf{k}}^{\text{ep}}(E_{n\mathbf{k}}/\hbar)}{[\hbar\omega - E_{n\mathbf{k}}]^2 + [Z_{n\mathbf{k}} \text{Im} \Sigma_{n\mathbf{k}}^{\text{ep}}(E_{n\mathbf{k}}/\hbar)]^2}. \quad (173)$$

This is a sum of Lorentzians with strength  $Z_{n\mathbf{k}}$  and width  $Z_{n\mathbf{k}} \text{Im} \Sigma_{n\mathbf{k}}^{\text{ep}}(E_{n\mathbf{k}}/\hbar)$ . Here the “quasiparticle strength” is defined as the homonymous quantity appearing in  $GW$  calculations (Hedin and Lundqvist, 1969):

$$Z_{n\mathbf{k}} = [1 - \hbar^{-1} \partial \text{Re} \Sigma_{n\mathbf{k}}^{\text{ep}}(\omega) / \partial \omega |_{\omega=E_{n\mathbf{k}}/\hbar}]^{-1}. \quad (174)$$

The result expressed by Eq. (173) shows that, in a rigorous field-theoretic approach, the quasiparticle broadening and lifetime given by Eqs. (169) and (170) should be renormalized by  $Z_{n\mathbf{k}}$  and  $Z_{n\mathbf{k}}^{-1}$ , respectively, and should be evaluated using the quasiparticle energy  $E_{n\mathbf{k}}$  instead of  $\varepsilon_{n\mathbf{k}}$ . This result can also be derived from Eq. (165) by performing a Taylor expansion of the self-energy along the imaginary axis and using the Cauchy-Riemann conditions.

In order to illustrate the typical features of the spectral function, we consider a model system characterized by one parabolic conduction band. The occupied electronic states couple to all states within an energy cutoff via a dispersionless phonon mode and a constant electron-phonon matrix element. A simplified version of this model was discussed by Engelsberg and Schrieffer (1963) by considering a constant density of electronic states. By evaluating the spectral function in Eq. (172) using the Fan-Migdal self-energy and neglecting the Debye-Waller term, we obtain the results shown in Fig. 4 for two sets of parameters.

In Fig. 4(a) the Fermi energy is much larger than the characteristic phonon energy. This case is representative of a metallic system with electron bands nearly linear around the Fermi level. Here the electron-phonon interaction leads to (i) a reduction of the band velocity in proximity of the Fermi level, and (ii) a broadening of the spectral function beyond the phonon energy  $\hbar\omega_0$ . A detailed analysis of these features for a slightly simpler model system, including a discussion of the analytic properties of the Green’s function, can be found in the work by Engelsberg and Schrieffer.




FIG. 4. Two-dimensional maps of the electron spectral function  $A(\mathbf{k}, \omega)$  for electrons coupled to a dispersionless phonon of frequency  $\omega_0$ . The noninteracting bands are given by  $\varepsilon(\mathbf{k}) = -\varepsilon_F + \hbar^2 |\mathbf{k}|^2 / 2m^*$ , and the Fermi level coincides with the top of the energy window. The matrix element is  $|g|^2 = \hbar\omega_0 / N_F$  when the electron energies differ by less than the cutoff  $\varepsilon_{\text{max}}$  and zero otherwise ( $N_F$  is the density of states at the Fermi level). (a) Spectral function for the case  $\varepsilon_F = 10\hbar\omega_0$  (white on blue, black), noninteracting band structure (solid line, yellow, light gray), and fully interacting band structure within the Brillouin-Wigner perturbation theory (solid line, red, dark gray). (b) Spectral function for the case  $\varepsilon_F = 2\hbar\omega_0$ . The model parameters are  $m^* = 0.1m_e$ ,  $\hbar\omega_0 = 100$  meV,  $\eta = 20$  meV, and  $\varepsilon_c = 5$  eV. For clarity the calculated spectral functions are cut off at the value  $3 \text{ eV}^{-1}$  and normalized. The self-energy is shifted by a constant so as to have  $\Sigma^{\text{ep}}(0) = 0$ . This correction guarantees the fulfillment of Luttinger’s theorem about the volume enclosed by the Fermi surface (Luttinger, 1960).

The solid line (red, dark gray) in Fig. 4(a) shows the renormalized band structure obtained from the Brillouin-Wigner perturbation theory, Eq. (164). We see that these solutions track the maxima of the spectral function  $A(\mathbf{k}, \omega)$ . The renormalized bands exhibit a characteristic S shape near the Fermi level, corresponding to multiple solutions of Eq. (164) for the same wave vector  $\mathbf{k}$ . Starting from the late 1990s such S-shaped energy-momentum dispersion curves have been observed in a number of ARPES experiments and have become known in the literature as the “photoemission kink” (Valla, Fedorov, Johnson, and Hulbert, 1999). First-principles calculations of kinks were first reported by Eiguren, de Gironcoli *et al.* (2003) and Giustino, Cohen, and Louie (2008) and are reviewed in Sec. VIII.

In Fig. 4(b) the Fermi energy is comparable to the characteristic phonon energy. This case is representative of a degenerately doped semiconductor close to a conduction band minimum. Here the electron-phonon interaction leads to two distinct spectral features: (i) a parabolic band with a heavier mass, which is well described by the Brillouin-Wigner solutions (solid line, red, dark gray), and (ii) a polaron satellite that is visible farther down. In this example, it is clear that Eq. (164) is unable to describe the satellite and that the spectral function carries qualitatively new information about the system. Polaron satellites resembling Fig. 4(b) have been observed in ARPES experiments on doped oxides (Moser *et al.*, 2013; Chen *et al.*, 2015; Cancellieri *et al.*, 2016; Wang *et al.*, 2016) and were recently calculated from first principles (Verdi, Caruso, and Giustino, 2017).

## 6. Model Hamiltonians, polarons, and the cumulant expansion

At the end of this section it is worth mentioning complementary *non-first-principles* approaches for studying the effects of EPIs on the electronic properties of solids. Model EPI Hamiltonians can be derived from Eq. (1) by choosing *a priori* explicit expressions for the electron band energies, the vibrational frequencies, and the coupling matrix elements. Examples of model Hamiltonians are those of Fröhlich (1954), Holstein (1959), and Su, Schrieffer, and Heeger (1979), the Hubbard-Holstein model (Berger, Valášek, and von der Linden, 1995), the Peierls-Hubbard model (Campbell, DeGrand, and Mazumdar, 1984), the *t*-*J* Holstein model (Rösch and Gunnarsson, 2004), and the Su-Schrieffer-Heeger-Holstein model (Perroni *et al.*, 2004). These models involve the tight-binding approximation, the Einstein phonon spectrum, and electron-phonon couplings to first order in the atomic displacements. Using these model Hamiltonians it is possible to go beyond the approximations introduced in Sec. V.B.2 and obtain nonperturbative solutions by means of canonical Lang-Firsov transformations, path-integral methods, exact diagonalization, and variational or quantum Monte Carlo techniques (Alexandrov, 2008; Alexandrov and Devreese, 2010). These models have been used extensively to explore many aspects of polaron physics, for example, the ground-state energy of polarons (weak or strong coupling), their spatial extent (large or small polarons), and transport properties (bandlike or hoppinglike).

Given the considerable body of literature on model EPI Hamiltonians, it is natural to ask whether one could bring

*ab initio* calculations of EPIs to a similar level of sophistication. The main limitation of current first-principles approaches is that, given the complexity of the calculations, the electron self-energies are evaluated using the bare propagators, as in Eq. (157). As a consequence, higher-order interaction diagrams beyond the Migdal approximation (Migdal, 1958) are omitted altogether.

A promising avenue for going beyond the Migdal approximation consists of introducing higher-order diagrams via the “cumulant expansion” approach (Hedin and Lundqvist, 1969; Langreth, 1970; Aryasetiawan, Hedin, and Karlsson, 1996). In the cumulant expansion method, instead of calculating the electron Green’s function via a Dyson equation, one evaluates the time evolution of the Green’s function by formulating the problem in the interaction picture, in symbols  $G_{n\mathbf{k}}(t) = (i/\hbar) \exp[-i(\varepsilon_{n\mathbf{k}}/\hbar)t + C_{n\mathbf{k}}(t)]$  (Aryasetiawan, Hedin, and Karlsson, 1996). The distinctive advantage of this approach is that the cumulant  $C_{n\mathbf{k}}(t)$  can be obtained from a low-order self-energy, for example, the Fan-Migdal self-energy in Eq. (157), and the exponential “resummation” automatically generates higher-order diagrams (Mahan, 1993). Detailed discussions of the cumulant expansion formalism are given by Zhou *et al.* (2015) and Gumhalter *et al.* (2016).

The cumulant method provides an interesting point of contact between *ab initio* and model Hamiltonian approaches. In fact, the cumulant expansion is closely related to the “momentum average approximation” introduced by Berciú (2006) for studying the Green’s function of the Holstein polaron.

The cumulant expansion has proven successful in *ab initio* calculations of electron-electron interactions, in particular, valence band satellites in semiconductors (Kheifets *et al.*, 2003; Guzzo *et al.*, 2011, 2012, 2014; Lischner, Vigil-Fowler, and Louie, 2013; Kas, Rehr, and Reining, 2014; Caruso and Giustino, 2015; Caruso, Lambert, and Giustino, 2015). In the context of EPIs, the *ab initio* cumulant expansion method was applied to elemental metals by Story *et al.* (2014), and to the ARPES spectra of *n*-doped TiO<sub>2</sub> by Verdi, Caruso, and Giustino (2017). In the latter work the cumulant method correctly reproduced the polaron satellites observed in the experiments of Moser *et al.* (2013).

The study of polarons using *ab initio* many-body techniques is yet to begin. However, a first calculation of the spectral function of Fröhlich polarons and an approximate polaron wave function were recently reported by Verdi, Caruso, and Giustino (2017).

## VI. EFFICIENT CALCULATIONS OF MATRIX ELEMENTS AND THEIR INTEGRALS

The study of EPIs from first principles requires evaluating Brillouin-zone integrals of functions that exhibit strong fluctuations. This requirement can be appreciated by inspecting Eqs. (145) and (157): there the denominators become large whenever the difference between two electronic eigenvalues approaches a phonon energy. As a result, while in DFT total energy calculations the Brillouin zone is typically discretized using meshes of the order of  $10 \times 10 \times 10$  points, the numerical convergence of EPI calculations requires much finer grids, sometimes with as many as  $10^6$  wave vectors



(Giustino, Cohen, and Louie, 2007; Poncé *et al.*, 2015). Determining vibrational frequencies  $\omega_{\mathbf{q}\nu}$  and perturbations  $\Delta_{\mathbf{q}\nu} v^{\text{KS}}(\mathbf{r})$  for such a large number of wave vectors is a prohibitive task, since every calculation is roughly as expensive as one total energy minimization.

These difficulties stimulated the development of specialized numerical techniques for making calculations of EPIs affordable. In the following sections two such techniques are reviewed: electron-phonon Wannier interpolation and Fermi-surface harmonics.

## A. Wannier interpolation

### 1. Maximally localized Wannier functions

In addition to the standard description of electrons in solids in terms of Bloch waves, as in Eq. (29), it is possible to adopt an alternative point of view whereby electrons are described as linear combinations of localized orbitals called ‘‘Wannier functions’’ (Wannier, 1937). The most general relation between Wannier functions and Bloch waves can be written as follows. One considers electron bands  $\varepsilon_{n\mathbf{k}}$  with eigenfunctions  $\psi_{n\mathbf{k}}$ , where the index  $n$  is restricted to a set of bands that are separated from all other bands by finite energy gaps above and below. These bands are referred to as ‘‘composite energy bands’’ (Marzari and Vanderbilt, 1997). Wannier functions are defined as

$$w_{mp}(\mathbf{r}) = N_p^{-1} \sum_{n\mathbf{k}} e^{i\mathbf{k}\cdot(\mathbf{r}-\mathbf{R}_p)} U_{nm\mathbf{k}} u_{n\mathbf{k}}(\mathbf{r}), \quad (175)$$

where  $U_{nm\mathbf{k}}$  is a unitary matrix in the indices  $m$  and  $n$ . From this definition and Eq. (A1) it follows that Wannier functions are normalized in the supercell  $\langle w_{mp} | w_{m'p'} \rangle_{\text{sc}} = \delta_{mp, m'p'}$ . Furthermore, since  $u_{n\mathbf{k}}$  is lattice periodic, Wannier functions have the property  $w_{mp}(\mathbf{r}) = w_{m0}(\mathbf{r} - \mathbf{R}_p)$ . The inverse transformation of Eq. (175) is obtained by using the unitary character of  $U_{nm\mathbf{k}}$  together with Eq. (A1):

$$u_{n\mathbf{k}}(\mathbf{r}) = \sum_{mp} e^{-i\mathbf{k}\cdot(\mathbf{r}-\mathbf{R}_p)} U_{mn\mathbf{k}}^\dagger w_{mp}(\mathbf{r}). \quad (176)$$

The unitary matrix  $U_{nm\mathbf{k}}$  is completely arbitrary, and therefore there exists considerable freedom in the construction of Wannier functions. For example, by requiring that  $U_{nm, -\mathbf{k}} = U_{nm\mathbf{k}}^*$  one can make Wannier functions real valued. Marzari and Vanderbilt exploited this degree of freedom to construct Wannier functions that are maximally localized.

A comprehensive and up-to-date review of the theory and applications of maximally localized Wannier functions (MLWFs) is given by Marzari *et al.* (2012). Here we recall only that, in order to minimize the spatial extent of a function in a periodic solid, one needs to use a modified definition of the position operator, since the standard position operator is unbounded in an infinite crystal. This procedure is now well established and it is linked to the development of the modern theory of dielectric polarization (King-Smith and Vanderbilt, 1993; Resta, 1994). Nowadays it is possible to determine MLWFs routinely (Mostofi *et al.*, 2008). The original algorithm of Marzari and Vanderbilt (1997) was also extended to

deal with situations where a composite set of bands cannot be identified. This happens notably in metals for electronic states near the Fermi energy. For these cases, Souza, Marzari, and Vanderbilt (2001) developed a band ‘‘disentanglement’’ procedure, which extracts a subset of composite bands out of a larger set of states.

For the purposes of this article, the most important property of MLWFs is that they are exponentially localized in insulators in the sense that  $|w_{m0}(\mathbf{r})| \sim |\mathbf{r}|^{-\alpha} \exp(-h|\mathbf{r}|)$  for large  $|\mathbf{r}|$ , with  $\alpha, h > 0$  real parameters. This property was demonstrated in one spatial dimension by Kohn (1959a) and He and Vanderbilt (2001), and in two and three dimensions by Brouder *et al.* (2007), under the condition that the system exhibits time-reversal symmetry. In the case of metallic systems, no exponential localization is expected. However, the Wannier functions obtained in metals using the disentanglement procedure of Souza, Marzari, and Vanderbilt (2001) are typically highly localized.

MLWFs are usually comparable in size to atomic orbitals, and this makes them ideally suited for the Slater-Koster interpolation of band structures as shown by Souza, Marzari, and Vanderbilt (2001). This concept was successfully employed in a number of applications requiring accurate calculations of band velocities, effective masses, density of states, Brillouin-zone integrals, and transport coefficients (Wang *et al.*, 2006, 2007; Yates *et al.*, 2007; Pizzi *et al.*, 2014).

### 2. Interpolation of electron-phonon matrix elements

Wannier functions were introduced in the study of EPIs by Giustino, Cohen, and Louie (2007) and Giustino *et al.* (2007). The starting point is the following definition of the electron-phonon matrix element in the Wannier representation<sup>5</sup>:

$$g_{mn\kappa\alpha}(\mathbf{R}_p, \mathbf{R}_{p'}) = \langle w_{m0}(\mathbf{r}) | \frac{\partial V^{\text{KS}}}{\partial \tau_{\kappa\alpha}}(\mathbf{r} - \mathbf{R}_{p'}) | w_{n0}(\mathbf{r} - \mathbf{R}_{p'}) \rangle_{\text{sc}}, \quad (177)$$

where the subscript ‘‘sc’’ indicates that the integral is over the BvK supercell. The relation between these quantities and the standard EPI matrix elements  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$  is found by replacing Eq. (176) inside Eq. (38) and using Eqs. (34) and (35) (Giustino, Cohen, and Louie, 2007):

$$g_{mn\nu}(\mathbf{k}, \mathbf{q}) = \sum_{pp'} e^{i(\mathbf{k}\cdot\mathbf{R}_p + \mathbf{q}\cdot\mathbf{R}_{p'})} \times \sum_{m'n'\kappa\alpha} U_{mm'\mathbf{k}+\mathbf{q}} g_{m'n'\kappa\alpha}(\mathbf{R}_p, \mathbf{R}_{p'}) U_{n'n\mathbf{k}}^\dagger u_{\kappa\alpha, \mathbf{q}\nu}, \quad (178)$$

where  $u_{\kappa\alpha, \mathbf{q}\nu} = (\hbar/2M_\kappa \omega_{\mathbf{q}\nu})^{1/2} e_{\kappa\alpha, \nu}(\mathbf{q})$  and  $e_{\kappa\alpha, \nu}(\mathbf{q})$  are the vibrational eigenmodes of Eq. (15). The inverse relation is

<sup>5</sup>We note that  $g_{mn\kappa\alpha}(\mathbf{R}_p, \mathbf{R}_{p'})$  has dimensions of energy by length, at variance with Eq. (38). For consistency here we use a definition that differs from that given in Giustino, Cohen, and Louie (2007) by a factor  $N_p$ ; this factor is inconsequential.

$$\begin{aligned}
 g_{mn\kappa\alpha}(\mathbf{R}_p, \mathbf{R}_{p'}) &= \frac{1}{N_p N_{p'}} \sum_{\mathbf{k}, \mathbf{q}} e^{-i(\mathbf{k} \cdot \mathbf{R}_p + \mathbf{q} \cdot \mathbf{R}_{p'})} \\
 &\times \sum_{m'n'\nu} u_{\kappa\alpha, \mathbf{q}\nu}^{-1} U_{mm'\mathbf{k}+\mathbf{q}}^\dagger g_{m'n'\nu}(\mathbf{k}, \mathbf{q}) U_{n'n\mathbf{k}},
 \end{aligned}
 \tag{179}$$

with  $u_{\kappa\alpha, \mathbf{q}\nu}^{-1} = (\hbar/2M_\kappa\omega_{\mathbf{q}\nu})^{-1/2} e_{\kappa\alpha, \nu}^*(\mathbf{q})$ . The last two equations define a generalized Fourier transform of the electron-phonon matrix elements between reciprocal space and real space. In Eq. (179) we have  $N_p$  and  $N_{p'}$  to indicate that the BvK supercells for electronic band structures and phonon dispersions may not coincide.

If the quantity  $g_{mn\kappa\alpha}(\mathbf{R}_p, \mathbf{R}_{p'})$  decays rapidly as a function of  $|\mathbf{R}_p|$  and  $|\mathbf{R}_{p'}|$ , then only a small number of matrix elements in the Wannier representation will be sufficient to generate  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$  anywhere in the Brillouin zone by means of Eq. (178). The dependence of the matrix elements on  $\mathbf{R}_p$  and  $\mathbf{R}_{p'}$  can be analyzed by considering the following bound:

$$\begin{aligned}
 |g_{mn\kappa\alpha}(\mathbf{R}_p, \mathbf{R}_{p'})| &\leq \int_{sc} d\mathbf{r} |w_{m0}^*(\mathbf{r}) w_{n0}(\mathbf{r} - \mathbf{R}_p)| \\
 &\times \int_{sc} d\mathbf{r} |\partial V^{\text{KS}} / \partial \tau_{\kappa\alpha}(\mathbf{r} - \mathbf{R}_{p'})|.
 \end{aligned}$$

The first term guarantees that the matrix element decays in the variable  $\mathbf{R}_p$  at least as fast as MLWFs. As a result the worst case scenario corresponds to the choice  $\mathbf{R}_p = 0$ . In this case, the matrix element  $|g_{mn\kappa\alpha}(0, \mathbf{R}_{p'})|$  decays with the variable  $\mathbf{R}_{p'}$  at the same rate as the *screened* electric dipole potential generated by the atomic displacement  $\Delta\tau_{\kappa\alpha}$ . In nonpolar semiconductors and insulators, owing to the analytical properties of the dielectric matrix (Pick, Cohen, and Martin, 1970), this potential decays at least as fast as a quadrupole, that is,  $|\mathbf{R}_{p'}|^{-3}$ . As a result, all matrix elements in reciprocal space are finite for  $\mathbf{q} \rightarrow 0$  (Vogl, 1976) and hence amenable to interpolation. In the case of metals the asymptotic trend of  $\partial V^{\text{KS}} / \partial \tau_{\kappa\alpha}$  is dictated by Fermi-surface nesting, leading to Friedel oscillations that decay as  $|\mathbf{R}_{p'}|^{-4}$  (Fetter and Walecka, 2003). These oscillations are connected to the Kohn anomalies in the phonon dispersion relations (Kohn, 1959b). In practical calculations, Friedel oscillations are usually not an issue since they are suppressed by the numerical smearing of the Fermi-Dirac occupations, and a Yukawa-type exponential decay is recovered. The case of polar materials is more subtle and is discussed in Sec. VI.A.3. Figure 5 illustrates the spatial decay of  $|g_{mn\kappa\alpha}(\mathbf{R}_p, \mathbf{R}_{p'})|$  as a function of  $\mathbf{R}_p$  and  $\mathbf{R}_{p'}$  for the prototypical case of diamond.

The interpolation strategy is entirely analogous to standard techniques for generating phonon dispersion relations using the interatomic force constants (Gonze and Lee, 1997): one first determines matrix elements in the Bloch representation using DFPT on a coarse grid in the Brillouin zone, as in Sec. III.B.3. Then MLWFs are determined using the procedures of Marzari and Vanderbilt (1997) and Souza, Marzari, and Vanderbilt (2001). This yields the rotation matrices  $U_{mn\mathbf{k}}$  to be used in Eq. (179). The Fourier transform to real space is performed via Eq. (179). At this point, one assumes that




FIG. 5. Spatial decay of the electron-phonon matrix elements of diamond in the Wannier representation: (a)  $\max |g_{mn\kappa\alpha}(\mathbf{R}_p, 0)|$  vs  $|\mathbf{R}_p|$ , and (b)  $\max |g_{mn\kappa\alpha}(0, \mathbf{R}_{p'})|$  vs  $|\mathbf{R}_{p'}|$ . The maximum values are taken over all subscript indices, and the data are normalized to the largest value. The insets show the same quantities in logarithmic scale. The calculations were performed using the local density approximation to DFT. From Giustino, Cohen, and Louie, 2007.

matrix elements outside of the Wigner-Seitz supercell defined by the coarse Brillouin-zone grid can be neglected and uses Eq. (178) in order to obtain the matrix elements  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$  on very fine grids. The last step requires the knowledge of the rotation matrices  $U_{mn\mathbf{k}}$  also on the fine grids; these matrices are obtained from the Wannier interpolation of the band structures as described by Souza, Marzari, and Vanderbilt (2001). The operation is computationally inexpensive and enables the calculation of millions of electron-phonon matrix elements. The procedure can now be applied routinely (Noffsinger *et al.*, 2010; Poncé *et al.*, 2016). Figure 6 shows the matrix elements obtained using this method as compared to explicit DFPT calculations.

Wannier interpolation of electron-phonon matrix elements was successfully employed in a number of applications, ranging from metal and superconductors to semiconductors and nanoscale systems.<sup>6</sup>

### 3. Electron-phonon matrix elements in polar materials

In the case of polar materials, that is, systems exhibiting nonzero Born effective charges (Pick, Cohen, and Martin,

<sup>6</sup>See, for example, Park *et al.* (2007, 2014), Giustino, Cohen, and Louie (2008), Park, Giustino, McChesney *et al.* (2008), Noffsinger *et al.* (2009, 2010, 2012), Calandra, Profeta, and Mauri (2010), Giustino, Louie, and Cohen (2010), Vukmirović, Bruder, and Stojanović (2012), Margine and Giustino (2013, 2014), Bernardi *et al.* (2014, 2015), Sjakste *et al.* (2015), and Verdi and Giustino (2015).




FIG. 6. Comparison between Wannier-interpolated electron-phonon matrix elements and explicit DFPT calculations for diamond. The interpolated matrix elements were calculated starting from a coarse  $4^3$  Brillouin-zone grid (dotted line, black), a  $6^3$  grid (dashed line, blue), and a  $8^3$  grid (solid line, red). The dots indicate explicit DFPT calculations. In this example  $|n\mathbf{k}\rangle$  is set to the valence band top at  $\Gamma$ ;  $|m\mathbf{k} + \mathbf{q}\rangle$  spans  $\Lambda_3$ ,  $\Delta_5$ , and  $\Sigma_2$  bands, and the phonon is set to the highest optical branch. From Giustino, Cohen, and Louie, 2007.

1970), the interpolation scheme discussed in Sec. VI.A.2 breaks down. In fact, in these systems the dominant contribution to the potential  $\partial V^{\text{KS}}/\partial \tau_{\kappa\alpha}$  in Eq. (177) is a dipole, which decays as  $|\mathbf{R}_p|^{-2}$ . As a consequence some of the matrix elements in reciprocal space diverge as  $|\mathbf{q}|^{-1}$  for  $\mathbf{q} \rightarrow 0$  and cannot be interpolated straightforwardly from a coarse grid to a fine grid. Physically this singularity corresponds to the ‘‘Fröhlich electron-phonon coupling’’ (Fröhlich, 1954).

The adaptation of the Wannier interpolation method to the case of polar materials was recently given by Sjakste *et al.* (2015) and Verdi and Giustino (2015). In both works the basic idea is to separate the matrix elements into a short-range contribution  $g_{mn\nu}^{\text{S}}(\mathbf{k}, \mathbf{q})$ , which is amenable to standard Wannier interpolation, and a long-range contribution  $g_{mn\nu}^{\text{L}}(\mathbf{k}, \mathbf{q})$ , which is singular and is dealt with analytically. The strategy is analogous to that in use for calculating the splitting between longitudinal-optical (LO) and transverse-optical (TO) phonons in polar materials (Gonze and Lee, 1997). The starting point is to define the long-range component of the matrix elements by considering the potential generated by the Born charges of all the atoms, when displaced according to a given vibrational eigenmode. The derivation relies on standard electrostatics and can be found in (Verdi and Giustino, 2015)

$$g_{mn\nu}^{\text{L}}(\mathbf{k}, \mathbf{q}) = i \frac{4\pi}{\Omega} \frac{e^2}{4\pi\epsilon_0} \sum_{\kappa} \left( \frac{\hbar}{2N_p M_{\kappa} \omega_{\text{qv}}} \right)^{1/2} \times \sum_{\mathbf{G} \neq -\mathbf{q}} \frac{(\mathbf{q} + \mathbf{G}) \cdot \mathbf{Z}_{\kappa}^* \cdot \mathbf{e}_{\kappa\nu}(\mathbf{q})}{(\mathbf{q} + \mathbf{G}) \cdot \epsilon^{\infty} \cdot (\mathbf{q} + \mathbf{G})} \times \langle \psi_{m\mathbf{k}+\mathbf{q}} | e^{i(\mathbf{q}+\mathbf{G}) \cdot (\mathbf{r}-\tau_{\kappa})} | \psi_{n\mathbf{k}} \rangle_{\text{sc}}. \quad (180)$$

In this expression  $\mathbf{Z}_{\kappa}^*$  and  $\epsilon^{\infty}$  denote the Born effective charge tensors and the electronic permittivity tensor (that is, the permittivity evaluated at clamped nuclei). Equation (180) is the generalization of Fröhlich’s model to the case of anisotropic crystalline lattices and multiple phonon modes (Fröhlich, 1954). The result can be derived alternatively using the analytical




FIG. 7. Wannier interpolation of electron-phonon matrix elements for anatase  $\text{TiO}_2$ . The initial state  $|n\mathbf{k}\rangle$  is set to the bottom of the conduction band at  $\Gamma$ , the final state  $|m\mathbf{k} + \mathbf{q}\rangle$  spans the bottom of the conduction band along high-symmetry lines, and the phonon is the highest LO mode. The dots correspond to explicit DFPT calculations. The red dashed line is the short-range component of the matrix elements  $g^{\text{S}}$ . The solid blue curve represents the matrix elements  $g^{\text{S}} + g^{\text{L}}$ , as obtained from the modified Wannier interpolation of Sec. VI.A.3. The interpolation was performed starting from a coarse  $4 \times 4 \times 4$  unshifted grid. From Verdi and Giustino, 2015.

properties of the dielectric matrix (Pick, Cohen, and Martin, 1970) as discussed by Vogl (1976).

In order to perform Wannier interpolation, one subtracts Eq. (180) from the matrix elements computed on a coarse grid, interpolates the remaining short-range part, and then adds back Eq. (180) on the fine grid. This process requires the interpolation of the bra and ket  $\langle \dots \rangle_{\text{sc}}$  in the second line of Eq. (180). Verdi and Giustino showed that, for small  $\mathbf{q} + \mathbf{G}$ , this bra and ket can be interpolated via  $\langle \psi_{m\mathbf{k}+\mathbf{q}} | e^{i(\mathbf{q}+\mathbf{G}) \cdot \mathbf{r}} | \psi_{n\mathbf{k}} \rangle_{\text{sc}} = [U_{\mathbf{k}+\mathbf{q}} U_{\mathbf{k}}^{\dagger}]_{mn}$ , where the rotation matrices  $U_{mn\mathbf{k}}$  are obtained as usual from the procedure of Marzari and Vanderbilt (1997) and Souza, Marzari, and Vanderbilt (2001).<sup>7</sup> Figure 7 shows an example of Wannier interpolation for the prototypical polar semiconductor  $\text{TiO}_2$ : it is seen that the singularity is correctly captured by the modified interpolation method.

At the end of this section, we mention that other interpolation schemes are equally possible (Eiguren and Ambrosch-Draxl, 2008b; Prendergast and Louie, 2009; Agapito *et al.*, 2013; Gunst *et al.*, 2016). For example, Eiguren and Ambrosch-Draxl (2008b) proposed to interpolate only the local component of  $\Delta V_{\text{qv}}^{\text{KS}}$ , while explicitly calculating the nonlocal part of the perturbation as well as the Kohn-Sham wave functions in the Bloch representation. Furthermore, Eq. (178) remains unchanged if MLWFs are replaced by a basis of localized atomic orbitals, and all concepts discussed in this section remain valid. An

<sup>7</sup>Equation (4) of Verdi and Giustino (2015) is missing a factor  $e^{-i(\mathbf{q}+\mathbf{G}) \cdot \tau_{\kappa}}$ ; this factor needs to be retained in order to correctly describe the acoustic modes near  $\mathbf{q} = 0$ . In practical calculations the  $\mathbf{G}$ -vector sum in Eq. (180) is restricted to small  $|\mathbf{q} + \mathbf{G}|$  via the cutoff function  $e^{-a|\mathbf{q}+\mathbf{G}|^2}$ . The results are independent of the choice of the cutoff parameter  $a$ .

interpolation scheme using local orbitals was recently demonstrated by [Gunst \*et al.\* \(2016\)](#).

## B. Fermi-surface harmonics

In the study of *metallic* systems, one is often interested in describing EPIs only for electronic states in the vicinity of the Fermi surface. In these cases, besides the Wannier interpolation discussed in Secs. VI.A.2–VI.A.3, it is possible to perform efficient calculations using “Fermi-surface harmonics” (FSHs) introduced by [Allen \(1976\)](#) and recently revisited by [Eiguren and Gurtubay \(2014\)](#).

The basic idea underlying FSHs is to replace expensive three-dimensional Brillouin-zone integrals by inexpensive one-dimensional integrals in the energy variable. To this aim, [Allen \(1976\)](#) proposed to expand functions of the band index  $n$  and wave vector  $\mathbf{k}$ , say  $A_{n\mathbf{k}}$ , in products of pairs of functions, one depending on the energy  $A_L(\varepsilon)$  and one depending on the wave vector  $\Phi_L(\mathbf{k})$ :

$$A_{n\mathbf{k}} = \sum_L A_L(\varepsilon_{n\mathbf{k}}) \Phi_L(\mathbf{k}). \quad (181)$$

In this expression, the Fermi-surface harmonics  $\Phi_L(\mathbf{k})$  (to be defined) are constructed so as to obey the following orthogonality condition:

$$N_p^{-1} \sum_{n\mathbf{k}} \delta(\varepsilon_{n\mathbf{k}} - \varepsilon) \Phi_L(\mathbf{k}) \Phi_{L'}(\mathbf{k}) = N(\varepsilon) \delta_{LL'}, \quad (182)$$

where  $N(\varepsilon) = N_p^{-1} \sum_{n\mathbf{k}} \delta(\varepsilon_{n\mathbf{k}} - \varepsilon)$  is the density of states. Using Eqs. (181) and (182) one finds

$$A_L(\varepsilon) = N(\varepsilon)^{-1} N_p^{-1} \sum_{n\mathbf{k}} \delta(\varepsilon_{n\mathbf{k}} - \varepsilon) \Phi_L(\mathbf{k}) A_{n\mathbf{k}}. \quad (183)$$

Allen showed that in the FSHs representation a linear system of the kind  $A_{n\mathbf{k}} = N_p^{-1} \sum_{n'\mathbf{k}'} M_{n\mathbf{k}, n'\mathbf{k}'} B_{n'\mathbf{k}'}$  transforms into  $A_L(\varepsilon) = \sum_{L'} \int d\varepsilon' N(\varepsilon') M_{LL'}(\varepsilon, \varepsilon') B_{L'}(\varepsilon')$ . Linear systems of this kind are common in the solution of the Boltzmann transport equation (Sec. X) and the Eliashberg equations for the superconducting gap (Sec. XI.B). If one could perform the expansion using only a few harmonics, then the transformation would be advantageous, since the integrals over the wave vectors would have been absorbed in the expansion coefficients.

In the original proposal by Allen, the harmonics  $\Phi_L(\mathbf{k})$  were defined as polynomials in the band velocities; however, the completeness of the basis set was not established. Recently [Eiguren and Gurtubay \(2014\)](#) proposed to construct these functions as eigenstates of a modified Helmholtz equation

$$|\mathbf{v}_{\mathbf{k}}| \nabla_{\mathbf{k}}^2 \Phi_L(\mathbf{k}) = \omega_L \Phi_L(\mathbf{k}), \quad (184)$$

where  $\mathbf{v}_{\mathbf{k}} = \hbar^{-1} \nabla_{\mathbf{k}} \varepsilon_{n\mathbf{k}}$  is the band velocity for states at the Fermi surface, and  $\omega_L$  is the eigenvalue for the harmonic  $\Phi_L$ . The new definition in Eq. (184) maintains the properties of the original FSHs and carries the added advantage that the basis set is complete. In this case the subscript  $L$  in  $\Phi_L(\mathbf{k})$  labels the

eigenstates of the Helmholtz equation. [Eiguren and Gurtubay](#) demonstrated the construction of “Helmholtz FSHs” for prototypical metals such as Cu, Li, and MgB<sub>2</sub>.

Recent examples of the application of Fermi-surface harmonics to first-principles calculations of EPIs include work on the photoemission kink of YBa<sub>2</sub>Cu<sub>3</sub>O<sub>7</sub> ([Heid \*et al.\*, 2008](#)) and on the Seebeck coefficient of Li ([Xu and Verstraete, 2014](#)).

## VII. NONADIABATIC VIBRATIONAL FREQUENCIES AND LINEWIDTHS

As discussed in Secs. V.A.2 and V.A.3, the electron-phonon interaction can lead to a renormalization of the adiabatic vibrational frequencies and to a broadening of the spectral lines.

The first *ab initio* investigations of the effects of the nonadiabatic renormalization of phonon frequencies were reported by [Lazzeri and Mauri \(2006\)](#) and [Pisana \*et al.\* \(2007\)](#). In these works they concentrated on the  $E_{2g}$  phonon of graphene, which is found at the wave number  $\omega/2\pi c = 1585 \text{ cm}^{-1}$  at room temperature ( $c$  is the speed of light). This phonon corresponds to an in-plane C-C stretching vibration with  $\mathbf{q} = 0$  and has been studied extensively via Raman spectroscopy. In the graphene literature this mode is referred to as the “Raman  $G$  band.” Figure 8 shows a comparison between calculated and measured  $E_{2g}$  phonon frequencies, as a function of doping, from [Pisana \*et al.\* \(2007\)](#). The calculations were performed (i) within the adiabatic approximation and (ii) by including the nonadiabatic frequency renormalization using Eq. (145).<sup>8</sup> From Fig. 8 we see that the adiabatic theory is unable to reproduce the experimental data. On the contrary, the calculations including nonadiabatic effects nicely follow the measured Raman shift. This is a clear example of the limits of the adiabatic Born-Oppenheimer approximation and a demonstration of the importance of the phonon self-energy in Eq. (145).

The fact that the adiabatic approximation is inadequate for the  $E_{2g}$  phonon of doped graphene should have been expected from the discussion in Sec. V.A.3. In fact, graphene is a zero-gap semiconductor; therefore electrons residing in the vicinity of the Dirac points can make “virtual” transitions with  $|\mathbf{q}| = 0$  and energies comparable to that of the  $E_{2g}$  mode. As a result, the condition underlying the adiabatic approximation  $|\varepsilon_{m\mathbf{k}+\mathbf{q}} - \varepsilon_{n\mathbf{k}}| \gg \hbar\omega_{\mathbf{q}\nu}$  does not hold in this case.

The importance of nonadiabatic effects was also confirmed in the case of metallic single-walled carbon nanotubes ([Caudal \*et al.\*, 2007](#); [Piscanec \*et al.\*, 2007](#)). In these works, they studied the phonon dispersion relations in the vicinity of  $|\mathbf{q}| = 0$  and found that the difference between adiabatic and nonadiabatic dispersions is concentrated around the zone center. This finding is consistent with earlier models of nonadiabatic effects. In fact, [Maksimov and Shulga \(1996\)](#) showed that, for metals with linear electron bands crossing the

<sup>8</sup>In the works reviewed in this section they used Eq. (145) with the bare matrix elements  $g_{mn\nu}^b(\mathbf{k}, \mathbf{q})$  replaced by the screened matrix elements  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$ . All calculations were performed within DFT, using either the LDA or gradient-corrected DFT functionals.




FIG. 8. Frequency of the Raman  $G$  band of graphene vs carrier concentration. The black filled disks are from Raman measurements of gated graphene on a silicon substrate at 295 K. The thick horizontal dashed line (red) shows the variation of the  $E_{2g}$  mode frequency with doping, within the adiabatic approximation. The solid blue line shows the variation of the frequency calculated by including nonadiabatic frequency renormalization. From [Pisana \*et al.\*, 2007](#).

Fermi level,  $\Pi^{\text{NA}}$  is significant only for wave vectors  $|\mathbf{q}| \sim \omega/v_F$ , where  $\omega$  is the phonon energy and  $v_F$  is the Fermi velocity. This result can be derived from Eq. (145).

In the previous examples, the nonadiabatic renormalization of the vibrational frequencies is measurable but very small, typically of the order of 1% of the corresponding adiabatic frequencies. [Saitta \*et al.\* \(2008\)](#) considered the question as to whether one could find materials exhibiting large nonadiabatic renormalizations and considered several graphite intercalation compounds, namely,  $\text{LiC}_6$ ,  $\text{LiC}_{12}$ ,  $\text{KC}_8$ ,  $\text{KC}_{24}$ ,  $\text{RbC}_8$ ,  $\text{CaC}_6$ ,  $\text{SrC}_6$ , and  $\text{BaC}_6$ , as well as other metallic systems such as  $\text{MgB}_2$ ,  $\text{Mg}$ , and  $\text{Ti}$ . In order to calculate the nonadiabatic renormalization at a reduced computational cost, the real part of the phonon self-energy was approximated as follows:  $\hbar \text{Re} \Pi_{\mathbf{q}=0,\nu\nu}^{\text{NA}} \approx N_F \langle |g_{nm\nu}(\mathbf{k}, \mathbf{q} = 0)|^2 \rangle_{\text{BZ}}$ , where  $\langle \dots \rangle_{\text{BZ}}$  stands for the average taken over the wave vectors  $\mathbf{k}$  in the Brillouin zone, and  $N_F$  is the density of states at the Fermi level. This expression can be derived from Eq. (145) by replacing the bare matrix elements by their screened counterparts, and by neglecting the “interband” contributions  $m \neq n$  in the sum. Figure 9 shows a comparison between vibrational frequencies from experiment and those calculated with or without including the nonadiabatic self-energy. It is clear that the nonadiabatic frequencies are in much better agreement with experiment than the corresponding adiabatic calculations. Furthermore, in these compounds the renormalization can reach values as large as  $\sim 300 \text{ cm}^{-1}$ . In contrast to this, the renormalization in  $\text{Mg}$  and  $\text{Ti}$  was found to be of only a few wave numbers in  $\text{cm}^{-1}$ .




FIG. 9. Comparison between measured ( $\omega_{\text{exp}}$ ) and calculated ( $\omega_{\text{th}}$ ) vibrational frequencies of the  $E_{2g}$  mode of graphite intercalation compounds. Open symbols are adiabatic DFPT calculations, and filled symbols are calculations including the nonadiabatic corrections. The line corresponds to  $\omega_{\text{th}} = \omega_{\text{exp}}$ . From [Saitta \*et al.\*, 2008](#).

The case of  $\text{MgB}_2$  proved more puzzling: here the calculated nonadiabatic frequency is  $761 \text{ cm}^{-1}$ , while experiments reported  $600 \text{ cm}^{-1}$ . In order to explain this discrepancy, [Saitta \*et al.\*](#) reasoned that a more accurate calculation would require taking into account the relaxation time of the electrons, as pointed out by [Maksimov and Shulga \(1996\)](#). This would act so as to partly wash out nonadiabatic effects. In the field-theoretic language of Sec. V, this observation corresponds to stating that when one approximates the dressed electron propagator  $G$  in Fig. 2(c) using the noninteracting Kohn-Sham Green’s function, one should include at the very least the effects of finite electron lifetimes (due to electron-electron, electron-impurity, and electron-phonon scattering), for example, via the imaginary part of Eq. (157).

The calculations discussed so far in this section addressed the nonadiabatic renormalization of zone-center phonons. The generalization to calculations of complete phonon dispersions was made by [Calandra, Profeta, and Mauri \(2010\)](#). In their work, [Calandra \*et al.\*](#) employed Wannier interpolation (see Sec. VI) in order to calculate the nonadiabatic phonon self-energy of Eq. (145) throughout the Brillouin zone. Figure 10 shows a comparison between the standard DFPT phonon dispersion relations of  $\text{CaC}_6$  and the dispersions obtained after incorporating the nonadiabatic self-energy. It is seen that also in this case nonadiabatic effects are most pronounced at small  $\mathbf{q}$  and can be as large as 7% of the adiabatic frequency.

In their work [Calandra \*et al.\*](#) approximated the bare matrix element  $g_{mn\nu}^b(\mathbf{k}, \mathbf{q})$  appearing in Eq. (145) by the screened matrix element  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$ . This replacement was justified by reasoning that the error is of second order in the induced electron density, hence it should be negligible.

The broadening of vibrational spectra arising from the electron-phonon interaction is almost invariably calculated from first principles using Eq. (146). Since the integration of the Dirac delta is computationally costly, it is common to rewrite that equation by neglecting the phonon energy in the delta function and by taking the low-temperature limit, as proposed by [Allen \(1972b\)](#):




FIG. 10. Phonon dispersion relations of  $\text{CaC}_6$  calculated using Wannier interpolation. The dashed lines (black) and the solid lines (red) represent the standard adiabatic calculation and the nonadiabatic phonon dispersions, respectively. From Calandra, Profeta, and Mauri, 2010.

$$\frac{\gamma_{\mathbf{q}\nu}}{\pi\omega_{\mathbf{q}\nu}} \approx 2 \sum_{mn} \int \frac{d\mathbf{k}}{\Omega_{\text{BZ}}} |g_{m\nu}(\mathbf{k}, \mathbf{q})|^2 \delta(\varepsilon_{n\mathbf{k}} - \varepsilon_{\text{F}}) \delta(\varepsilon_{m\mathbf{k}+\mathbf{q}} - \varepsilon_{\text{F}}), \quad (185)$$

where  $\varepsilon_{\text{F}}$  is the Fermi energy. Oddly enough, this is a sort of adiabatic approximation to the nonadiabatic theory. The main advantage is that Eq. (185) is positive definite, hence easier to converge numerically as compared to the complete expression in Eq. (146). The disadvantages are that the temperature dependence is lost, and that one cannot resolve fine features on the scale of a phonon energy. There exists a vast literature on first-principles calculations of phonon linewidths using Eq. (185), mostly in connection with electron-phonon superconductors.<sup>9</sup> Equation (185) is now implemented in several large software projects (Giannozzi *et al.*, 2009; Gonze *et al.*, 2009), and it is used routinely.

Phonon linewidths range from very small values such as  $\sim 1$  meV in Nb (Bauer *et al.*, 1998) to large values such as  $\sim 30$  meV in  $\text{MgB}_2$  (Shukla *et al.*, 2003). The agreement between calculations and neutron scattering or Raman measurements is usually reasonable.

Calculations of phonon linewidths using the more accurate expression in Eq. (146) are computationally more demanding and have been reported less extensively in the literature.<sup>10</sup>

<sup>9</sup>See, for example, early frozen-phonon calculations (Dacorogna, Cohen, and Lam, 1985; Chang and Cohen, 1986; Lam, Dacorogna, and Cohen, 1986) and more recent DFPT calculations (Savrasov, Savrasov, and Andersen, 1994; Bauer *et al.*, 1998; Shukla *et al.*, 2003; Heid *et al.*, 2010). Earlier calculations not based on DFT were reviewed by Grimvall (1981).

<sup>10</sup>See, for example, Lazzeri *et al.* (2006), Bonini *et al.* (2007), Giustino, Cohen, and Louie (2007), and Park, Giustino, Cohen, and Louie (2008).

So far we considered the effect of EPIs on the frequencies and lifetimes of vibrational excitations in solids. Another important phenomenon which modifies frequencies and lifetimes is *anharmonicity*. Anharmonic effects result from third- and higher-order terms in the Taylor expansion of the total potential energy  $U$  in the atomic displacements (Sec. III.A). These effects can be interpreted as additional interactions that couple the oscillators of the harmonic lattice; for example, third-order anharmonic effects reduce phonon lifetimes via energy upconversion or downconversion processes involving three phonons.

Anharmonic effects can be described using a many-body field-theoretic formalism (Cowley, 1963), in complete analogy with the theory of EPIs discussed in Sec. IV. Calculation of anharmonic effects from first principles goes through the evaluation of third- and fourth-order derivatives of the total potential energy  $U$  in the adiabatic approximation. Third-order coefficients are routinely computed using DFPT (Debernardi, Baroni, and Molinari, 1995; Deinzer, Birner, and Strauch, 2003; Lazzeri, Calandra, and Mauri, 2003; Bonini *et al.*, 2007; Broido *et al.*, 2007). In those cases where the harmonic approximation fails completely, “self-consistent phonon” techniques can be employed (Hooton, 1955; Koehler, 1966). Recent implementations and calculations can be found in Errea, Rousseau, and Bergara (2011), Hellman, Abrikosov, and Simak (2011), Hellman *et al.* (2013), Monserrat, Drummond, and Needs (2013), and Errea, Calandra, and Mauri (2014).

## VIII. ELECTRON-PHONON INTERACTIONS IN PHOTOELECTRON SPECTROSCOPY

In Sec. V.B.5 we have seen how the electron-phonon interaction in metals can lead to band structure “kinks” near the Fermi energy. This is illustrated by the model calculation in Fig. 4(a). The experimental investigation of these features started in the late 1990s, following the development of high-resolution ARPES. Since in ARPES only the component of the photoelectron momentum parallel to the sample surface is conserved (Damascelli, Hussain, and Shen, 2003), complete energy versus wave vector dispersion relations can be measured directly only for 2D or quasi-2D materials. Accordingly, the first observations of kinks were reported for the surface states of elemental metals<sup>11</sup> and for the  $\text{CuO}_2$  planes of copper oxide superconductors.<sup>12</sup> *Ab initio* calculations of ARPES kinks can be performed by using the diagonal part of the Fan-Migdal self-energy (the Debye-Waller self-energy will be discussed at the end of this section). To this aim, it is common to rewrite Eqs. (157) and (158) at finite temperature using a spectral representation:

<sup>11</sup>See, for example, Hengsberger *et al.* (1999) and Valla, Fedorov, Johnson, and Hulbert (1999).

<sup>12</sup>See, for example, Valla *et al.* (1999), Johnson *et al.* (2001), and Lanzara *et al.* (2001).

$$\Sigma_{n\mathbf{k}}^{\text{FM}}(\omega, T) = \int_{-\infty}^{+\infty} d\varepsilon \int_0^{\infty} d\varepsilon' \alpha^2 F_{n\mathbf{k}}(\varepsilon, \varepsilon') \times \left[ \frac{1 - f(\varepsilon/k_B T) + n(\varepsilon'/k_B T)}{\hbar\omega - \varepsilon - \varepsilon' + i\hbar\eta} + \frac{f(\varepsilon/k_B T) + n(\varepsilon'/k_B T)}{\hbar\omega - \varepsilon + \varepsilon' + i\hbar\eta} \right]. \quad (186)$$

Here the function  $\alpha^2 F$  is the so-called ‘‘Eliashberg function’’ and is defined as

$$\alpha^2 F_{n\mathbf{k}}(\varepsilon, \varepsilon') = \sum_{m\nu} \int \frac{d\mathbf{q}}{\Omega_{\text{BZ}}} |g_{nm\nu}(\mathbf{k}, \mathbf{q})|^2 \times \delta(\varepsilon - \varepsilon_{m\mathbf{k}+\mathbf{q}}) \delta(\varepsilon' - \hbar\omega_{\mathbf{q}\nu}). \quad (187)$$

This quantity is positive, temperature independent, and contains all the materials-specific parameters. Physically it is proportional to the scattering rate of an electron in the state  $|n\mathbf{k}\rangle$  into any electronic state at the energy  $\varepsilon$ , by emitting or absorbing any one phonon of energy  $\varepsilon'$ . One complication related to the Eliashberg function is that in the literature many variants of Eq. (187) can be found, each stemming from specific approximations; some of these expressions are summarized by Grimvall (1981).

The first *ab initio* calculations of the phonon-induced electron self-energies and photoemission kinks were reported by Eiguren, de Gironcoli *et al.* (2003) for the surface state of the Be(0001) surface. Since the evaluation of Eqs. (186) and (187) is computationally demanding, Eiguren *et al.* employed simplified expressions which involve the following three approximations: (i) the Eliashberg function is replaced by its isotropic average  $\alpha^2 F_n(\varepsilon, \varepsilon') = \int d\mathbf{k} \delta(\varepsilon_{n\mathbf{k}} - \varepsilon) \times \alpha^2 F_{n\mathbf{k}}(\varepsilon, \varepsilon') / \int d\mathbf{k} \delta(\varepsilon_{n\mathbf{k}} - \varepsilon)$ ; (ii) phonon energies are neglected next to electron energies (as in the adiabatic approximation), and (iii) particle-hole symmetry is assumed. Using these approximations the imaginary part of the Fan-Migdal self-energy becomes

$$|\text{Im}\Sigma_n^{\text{FM}}(\omega, T)| = \pi \int_0^{\infty} d\varepsilon' \alpha^2 F_n(\hbar\omega, \varepsilon') \{1 + 2n(\varepsilon'/k_B T) + f[(\hbar\omega + \varepsilon')/k_B T] - f[(\hbar\omega - \varepsilon')/k_B T]\}, \quad (188)$$

where the average of the self-energy is defined as for the Eliashberg function. The real part of the self-energy can be found starting from the same approximations and is given by Grimvall (1981).

Figure 11 shows the self-energy of the surface state at the Be(0001) surface calculated by Eiguren, de Gironcoli *et al.* (2003) using Eq. (188). The imaginary part resembles a step function, with an onset around the energy threshold for phonon emission by a hole (40–80 meV in this case). At a qualitative level, this trend can be rationalized by replacing the Eliashberg function in Eq. (188) by a Dirac delta at a characteristic phonon energy  $\hbar\omega_{\text{ph}}$ . In this case, the hole self-energy becomes proportional to  $f[(\hbar\omega + \hbar\omega_{\text{ph}})/k_B T]$ . At  $T = 0$  this is precisely a step function with onset at  $-\hbar\omega_{\text{ph}}$ . The real part of the self-energy vanishes for  $|\omega| \gg \omega_{\text{max}}$ , with




FIG. 11. Calculated Fan-Migdal self-energy of the surface state at the Be(0001) surface. (a) Imaginary part of the self-energy, obtained from Eq. (188). The dashed (black) line is the self-energy evaluated using the DFT-LDA bands; the solid lines (color, gray scale) correspond to the self-energy calculated by taking into account the renormalization of the DFT band structure by the electron-phonon interaction. (b) Real part of the self-energy, using the same color, gray scale code as in (a). The inset in (a) compares the renormalized band structure (color) with the ‘‘bare’’ DFT band (black dashed line). The inset of (b) shows the renormalization of the band velocity induced by the electron-phonon interaction. From Eiguren, de Gironcoli *et al.*, 2003.

$\omega_{\text{max}}$  the largest phonon frequency. This can be seen in Fig. 11(b) and is a consequence of the approximation of particle-hole symmetry. Eiguren *et al.* also determined the renormalization of the surface state band structure arising from electron-phonon interactions, using Eq. (164); this is shown in the inset of Fig. 11(a). Overall the calculations of Eiguren, de Gironcoli *et al.* (2003) showed good agreement with photoelectron spectroscopy experiments, in both the shape and magnitude of the self-energy (LaShell, Jensen, and Balasubramanian, 2000).

In addition to these calculations, several studies of the electron-phonon self-energy at metal surfaces were reported, namely, for the Cu(111) and Ag(111) surfaces (Eiguren *et al.*, 2002), the Al(100), Ag(111), Cu(111), and Au(111) surfaces (Eiguren, Hellsing *et al.*, 2003), and the W(110) surface (Eiguren and Ambrosch-Draxl, 2008a). Building on these studies, Eiguren, Ambrosch-Draxl, and Echenique (2009) performed a detailed analysis of the *self-consistent* solutions of the complex Dyson equation for the quasiparticle energies, Eqs. (164) and (165), and illustrated the key concepts in the cases of the W(110) surface and for the phonon-mediated superconductor MgB<sub>2</sub>.

Equation (188) or closely related approximations were also employed in the study of electron and hole lifetimes in bulk Be (Sklyadneva *et al.*, 2005), Pb (Sklyadneva *et al.*, 2006),

and Mg (Leonardo *et al.*, 2007); the photoemission kink in  $\text{YBa}_2\text{Cu}_3\text{O}_7$  (Heid *et al.*, 2008); and the spectral function of Ca-intercalated graphite (Sanna *et al.*, 2012).

In the case of complex systems the validity of the approximations (i)–(iii) leading to Eq. (188) is not warranted, and a direct calculation of the Fan-Migdal self-energy using Eqs. (186) and (187) is necessary. Calculations of the complete self-energy were reported by Park *et al.* (2007) for graphene, by Giustino, Cohen, and Louie (2008) for the high-temperature superconductor  $\text{La}_{1-x}\text{Sr}_x\text{Cu}_2\text{O}_4$ , and by Margine, Lambert, and Giustino (2016) for Ca-intercalated bilayer graphene. Figure 12 shows the calculated self-energy and spectral function of graphene calculated by Park *et al.* (2007, 2009). The structure of the self-energy is similar to that shown in Fig. 11, with one important exception:  $\Sigma_{mnk}^{\text{FM}}(\omega)$  does not vanish a few phonon energies away from the Fermi level, but tends instead toward a linear asymptote. Calandra and Mauri (2007) performed a combined *ab initio* and analytical study of the effects of the electron-phonon interaction on the electron bands of graphene and obtained very similar results. A linear asymptote in the real part of the self-energy is a general feature of systems which do not exhibit particle-hole symmetry. For another example see the work on copper oxides by Giustino, Cohen, and Louie (2008).

Using the Fan-Migdal self-energy, it is possible to calculate the renormalization of the band velocity induced by the electron-phonon interaction. Let us denote by  $\mathbf{v}_{nk} = \hbar^{-1}\nabla_{\mathbf{k}}\varepsilon_{nk}$  the DFT band velocity and  $\mathbf{V}_{nk} = \hbar^{-1}\nabla_{\mathbf{k}}E_{nk}$  the band velocity after including electron-phonon interactions. Using Eq. (164) with  $\Gamma_{nk} = 0$  we find that these two quantities are simply related by  $\mathbf{V}_{nk} = Z_{nk}\mathbf{v}_{nk} = \mathbf{v}_{nk}/(1 + \lambda_{nk})$ , where  $Z_{nk}$  is the quasiparticle strength of Eq. (174), and  $\lambda_{nk}$  is the “mass enhancement parameter” or “electron-phonon coupling strength” (Grimvall, 1981):

$$\lambda_{nk} = Z_{nk}^{-1} - 1 = -\hbar^{-1}\partial\text{Re}\Sigma_{mnk}(\omega)/\partial\omega|_{\omega=E_{nk}/\hbar}. \quad (189)$$

In the study of EPs in metals, the coupling strength  $\lambda_{nk}$  is of significant interest since it is related to the superconducting transition temperature of phonon-mediated superconductors (see Secs. VIII.A and XI).

The velocity renormalization in graphene calculated using Eq. (189) is shown in Figs. 12(c) and 12(d), while a calculation of the complete spectral function  $A(\mathbf{k}, \omega)$  is shown in Fig. 12(e). Here the characteristic photoemission kink is visible between 150 and 200 meV but it is not very pronounced, since in this case  $\lambda_{nk} \sim 0.1$  (Park *et al.*, 2009). These results are in good agreement with measured photoelectron spectra (Bostwick *et al.*, 2007; Zhou *et al.*, 2009).

Incidentally, we remark that in the analysis of ARPES data it is common to extract the coupling strength  $\lambda_{nk}$  directly from the ratio of the band velocities above and below the electron-phonon kink. However, this procedure is subject to a significant uncertainty, since the bare velocity is not known and must be approximated by fitting the fully interacting dispersions using *ad hoc* models. For example, in the vicinity of van Hove singularities this procedure leads to a significant overestimation of the electron-phonon coupling strength  $\lambda_{nk}$  (Park, Giustino, McChesney *et al.*, 2008; Bianchi *et al.*, 2010).




FIG. 12. (a), (b) Calculated real part of the Fan-Migdal self-energy in pristine and  $n$ -doped graphene, respectively (solid black lines). The doping level is  $4 \times 10^{13} \text{ cm}^{-2}$ . The dashed lines correspond to a simplified analytical model where particle-hole symmetry is assumed. (c), (d) Electron band velocity renormalization resulting from the self-energies in (a) and (b). All calculations in (a)–(d) were performed using DFT-LDA. From Park *et al.*, 2007. (e) Calculated spectral function of  $n$ -doped graphene for one of the branches of the Dirac cone.  $\omega_{\text{ph}}$  indicates the characteristic phonon energy leading to the photoemission kink, and  $E_{\text{D}}$  denotes the energy of the Dirac point. The calculations include *GW* quasiparticle corrections. From Park *et al.*, 2009.

In addition to photoemission kinks, recent ARPES measurements revealed the existence of polaron satellites in doped oxides, namely,  $\text{TiO}_2$  (Moser *et al.*, 2013) and  $\text{SrTiO}_3$  (Chen *et al.*, 2015; Cancellieri *et al.*, 2016; Wang *et al.*, 2016). The phenomenology is similar to what was discussed in relation to Fig. 4(b). The first theoretical studies along this direction were reported by Story *et al.* (2014), who applied the cumulant expansion approach to the case of the electron-phonon self-energy; by Antonius *et al.* (2015), who identified satellites in the spectral functions of LiF and MgO; and by Verdi, Caruso, and Giustino (2017), who calculated the ARPES spectra of  $n$ -doped  $\text{TiO}_2$ .



At the end of this section, it is worth returning to the subject of the Debye-Waller self-energy. So far we only discussed the Fan-Migdal self-energy, starting from Eq. (186), and we ignored the Debye-Waller self-energy appearing in Eq. (152). This omission reflects the fact that, in the literature on electron-phonon interactions in metals, the DW term has always been disregarded. In order to rationalize this approximation, we rewrite the DW self-energy as follows, by combining Eqs. (160) and (40):

$$\Sigma_{n\mathbf{k}}^{\text{DW}} = \langle u_{n\mathbf{k}} | V_{\text{DW}} | u_{n\mathbf{k}} \rangle_{\text{uc}}, \quad (190)$$

with  $V_{\text{DW}}(\mathbf{r}) = \Omega_{\text{BZ}}^{-1} \sum_{\nu} \int d\mathbf{q} (n_{\mathbf{q}\nu} + 1/2) \Delta_{\mathbf{q}\nu} \Delta_{-\mathbf{q}\nu} v^{\text{KS}}(\mathbf{r})$ . The subscript ‘‘uc’’ indicates that the integral is over one unit cell. From Eq. (190) we see that  $V_{\text{DW}}$  acts like a static local *potential*; indeed the first calculations including DW effects were performed by directly modifying the ionic pseudopotentials (Antončík, 1955; Walter *et al.*, 1970). From Eq. (190) we also see that the only variation in the DW self-energy comes from the Bloch amplitudes  $u_{n\mathbf{k}}$ . Let us consider the limiting situation of the homogeneous electron gas. In this case  $|u_{n\mathbf{k}}(\mathbf{r})|^2 = 1/\Omega$  (Sec. III.B.5); therefore  $\Sigma_{n\mathbf{k}}^{\text{DW}}$  is a constant, independent of  $\mathbf{k}$ . This behavior should be contrasted with the Fan-Migdal self-energy, which exhibits significant structure near the Fermi energy as can be seen in Fig. 4.

In more realistic situations, such as doped semiconductors,  $\mathbf{k} \cdot \mathbf{p}$  perturbation theory (Kittel, 1963) can be used to show that  $\Sigma_{n\mathbf{k}}^{\text{DW}}$  varies smoothly as a function of  $\mathbf{k}$  within the same band. In contrast to this scenario,  $\Sigma_{n\mathbf{k}}^{\text{DW}}$  exhibits large variations across *different* bands. This carries important consequences for the calculation of temperature-dependent band gaps (Sec. IX.A.1).

### A. Electron mass enhancement in metals

We now come back to the mass enhancement parameter  $\lambda_{n\mathbf{k}}$  introduced in Eq. (189), since this quantity played a central role in the development of the theory of EPIs in metals.

The notion of mass enhancement becomes clear when we consider a parabolic band as in the model calculations of Fig. 4. Near the Fermi surface the noninteracting dispersions are given by  $\varepsilon_{n\mathbf{k}} = \hbar \mathbf{k}_{\text{F}} \cdot \hbar(\mathbf{k} - \mathbf{k}_{\text{F}})/m^*$ , where  $\mathbf{k}_{\text{F}}$  is a wave vector on the Fermi surface, and the electron velocity is  $\mathbf{v}_{n\mathbf{k}} = \hbar \mathbf{k}_{\text{F}}/m^*$ . After taking into account the EPI, the electron velocity is renormalized to  $\mathbf{V}_{n\mathbf{k}} = \mathbf{v}_{n\mathbf{k}}/(1 + \lambda_{n\mathbf{k}})$ . Since the magnitude of the Fermi momentum is unchanged (see the caption of Fig. 4), this renormalization can be interpreted as an effective increase of the band mass from  $m^*$  to  $m_{\text{ep}} = m^*(1 + \lambda_{n\mathbf{k}})$ . This reasoning holds for metals with parabolic bands and for doped semiconductors near band extrema and does not take into account the Debye-Waller self-energy.

The electron mass enhancement is reflected in the increase of the heat capacity of metals at low temperature. In fact, below the Debye temperature the electronic contribution to the heat capacity dominates over the lattice contribution (Kittel, 1976). Since the heat capacity is proportional to the density of states at the Fermi level, and the density of states is inversely

proportional to the band velocity, it follows that the heat capacity is directly proportional to the electron mass. This property can be used as a means to determine the strength of the electron-phonon coupling in simple metals from specific heat measurements (Grimvall, 1975).

The general theory of the effects of electron-phonon interactions on the heat capacity and other thermodynamic functions was developed by Eliashberg (1963), Prange and Kadanoff (1964), and Grimvall (1969). A field-theoretic analysis of the effect of EPIs on thermodynamic functions was developed by Eliashberg (1963) starting from the identities of Luttinger and Ward (1960) in the zero-temperature limit. Eliashberg’s analysis was subsequently extended to all temperatures by Grimvall (1969). Here we quote only Grimvall’s result relating the electronic entropy to the Fan-Migdal self-energy of Sec. V.B.1:

$$S_e = \frac{N_{\text{F}} k_{\text{B}} \hbar^3}{(k_{\text{B}} T)^2} \int_0^{\infty} \frac{\omega [\omega - \hbar^{-1} \text{Re} \Sigma_{n\mathbf{k}}^{\text{FM}}(\omega, T)]}{\cosh^2(\hbar\omega/2k_{\text{B}}T)} d\omega. \quad (191)$$

In order to derive this relation, Grimvall started by expressing the thermodynamic potential of the coupled electron-phonon system in terms of the electron and phonon propagators and self-energies and identified the electronic contribution by neglecting terms of the order of  $(m_e/M_0)^{1/2}$  as well as electron-electron interactions (Grimvall, 1969, Appendix).

Below the Debye temperature, an explicit expression for the entropy in Eq. (191) can be obtained by noting that the function  $\cosh^{-2}(\hbar\omega/2k_{\text{B}}T)$  is nonvanishing only for  $\hbar\omega \lesssim 2k_{\text{B}}T$ . In this range Eq. (189) yields  $\Sigma_{n\mathbf{k}}^{\text{FM}}(\omega, T) \simeq -\lambda_{n\mathbf{k}} \hbar\omega$ , and therefore the integration in Eq. (191) can be carried out explicitly. As a result, the low-temperature heat capacity can be written as

$$C_e = T \frac{\partial S_e}{\partial T} = \frac{2}{3} \pi^2 k_{\text{B}}^2 N_{\text{F}} (1 + \lambda_{n\mathbf{k}}) T. \quad (192)$$

If we ignore the EPI by setting  $\lambda_{n\mathbf{k}} = 0$ , this expression reduces to the standard textbook result for the free electron gas (Kittel, 1976). At high temperature Eq. (192) is no longer valid, and one has to evaluate the integral in Eq. (191) using the complete frequency-dependent FM self-energy. The main result of this procedure is that at high temperature the electronic heat capacity is no longer renormalized by EPIs. A detailed discussion of this aspect is provided by Grimvall (1969, 1981).

Early examples of DFT calculations of mass enhancement parameters and comparisons with specific heat measurements in simple metals can be found in Dacorogna, Cohen, and Lam (1985), Savrasov, Savrasov, and Andersen (1994), and Liu and Quong (1996).

## IX. ELECTRON-PHONON EFFECTS IN THE OPTICAL PROPERTIES OF SEMICONDUCTORS

### A. Temperature dependence of band gaps and band structures

#### 1. Perturbative calculations based on the Allen-Heine theory

In Sec. V.B.3 we discussed how the electron-phonon interaction induces a ‘‘renormalization’’ of the electronic

energy levels and thereby gives rise to “temperature-dependent band structures.” These effects have been studied in detail using the Fan-Migdal and the Debye-Waller self-energies, either via the Raleigh-Schrödinger approximation to Eq. (166) or via its adiabatic counterpart given by Eq. (167).

Equation (167) was first employed in a number of calculations based on empirical pseudopotentials (see footnote 4), following the seminal work of Allen and Heine (1976). Allen and Cardona (1981) offered a clear introduction to the basic theory, a discussion of the computational methodology, as well as an historical perspective on earlier calculations.

The evaluation of the Debye-Waller contribution to the self-energy requires the calculation of the second-order variations of the Kohn-Sham potential with respect to the ionic displacements, Eq. (40). From a computational standpoint, this is challenging because one would have to use second-order DFPT, as discussed at the end of Sec. III.B.3. In order to avoid this complication, it is common practice to recast all second-order derivatives as products of first-order derivatives. This strategy was introduced by Allen and Heine in the case of monatomic crystals and extended to polyatomic unit cells by Allen and Cardona. The key observation behind this approach is that one can impose translational invariance of the theory to second order in the nuclear displacements. Specifically, the variation of the Kohn-Sham eigenvalues ensuing an arbitrary displacement of the nuclei should not change if all nuclei are *further* displaced by the same amount. Using time-independent perturbation theory, this condition yields the following two sum rules:

$$\sum_{\kappa p} \langle \psi_{n\mathbf{k}} | \partial_{\kappa p} V^{\text{KS}} | \psi_{n\mathbf{k}} \rangle_{\text{sc}} = 0, \quad (193)$$

$$\begin{aligned} \sum_{\kappa' p'} \langle \psi_{n\mathbf{k}} | \partial_{\kappa p, \kappa' p'}^2 V^{\text{KS}} | \psi_{n\mathbf{k}} \rangle_{\text{sc}} &= -2\text{Re} \sum_{\kappa' p'} \sum_{m\mathbf{q}} \\ &\times \frac{\langle \psi_{n\mathbf{k}} | \partial_{\kappa p} V^{\text{KS}} | \psi_{m\mathbf{k}+\mathbf{q}} \rangle_{\text{sc}} \langle \psi_{m\mathbf{k}+\mathbf{q}} | \partial_{\kappa' p'} V^{\text{KS}} | \psi_{n\mathbf{k}} \rangle_{\text{sc}}}{\varepsilon_{n\mathbf{k}} - \varepsilon_{m\mathbf{k}+\mathbf{q}}}. \end{aligned} \quad (194)$$

Here  $\partial_{\kappa p} V^{\text{KS}}$  is a short-hand notation for  $\partial V^{\text{KS}} / \partial \tau_{\kappa p}$  and similarly for the second derivative; the primed summation indicates that eigenstates  $\psi_{m\mathbf{k}+\mathbf{q}}$  such that  $\varepsilon_{n\mathbf{k}} = \varepsilon_{m\mathbf{k}+\mathbf{q}}$  are skipped. The first sum rule is equivalent to stating that the electron-phonon matrix elements  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$  associated with the three translational modes at  $|\mathbf{q}| = 0$  must vanish; this is an alternative formulation of the “acoustic sum rule.” The second sum rule, Eq. (194), suggests to express the matrix elements of the second derivatives of the potential in terms of first-order derivatives. However, Eq. (194) cannot be used as it stands, since it involves a sum of matrix elements on the left-hand side. In order to proceed, Allen and Heine employed the rigid-ion approximation, whereby  $V^{\text{KS}}$  is written as a sum of atom-centered contributions (see Sec. II.A.1). Under this approximation all the terms  $\kappa p \neq \kappa' p'$  on the left-hand side of Eq. (194) are neglected, and an explicit expression for  $\langle \psi_{n\mathbf{k}} | \partial_{\kappa p, \kappa p}^2 V^{\text{KS}} | \psi_{n\mathbf{k}} \rangle_{\text{sc}}$  is obtained.

In view of practical DFT implementations, Giustino, Louie, and Cohen (2010) used the sum rule in Eq. (194) in order to rewrite the Debye-Waller self-energy as follows:

$$\Sigma_{mn\mathbf{k}}^{\text{DW}} = - \sum_{\nu m} \int \frac{d\mathbf{q}}{\Omega_{\text{BZ}}} \frac{g_{mn\nu}^{2, \text{DW}}(\mathbf{k}, \mathbf{q})}{\varepsilon_{n\mathbf{k}} - \varepsilon_{m\mathbf{k}}} (2n_{q\nu} + 1). \quad (195)$$

Here  $g_{mn\nu}^{\text{DW}}(\mathbf{k}, \mathbf{q})$  is an effective matrix element, and it is obtained from the standard DFPT matrix elements by means of inexpensive matrix multiplications:

$$g_{mn\nu}^{2, \text{DW}}(\mathbf{k}, \mathbf{q}) = \sum_{\substack{\kappa\alpha, \kappa'\alpha' \\ \nu}} \frac{t_{\kappa\alpha, \kappa'\alpha'}^{\nu}(\mathbf{q})}{2\omega_{q\nu}} h_{mn, \kappa\alpha}^*(\mathbf{k}) h_{mn, \kappa'\alpha'}(\mathbf{k}), \quad (196)$$

$$t_{\kappa\alpha, \kappa'\alpha'}^{\nu}(\mathbf{q}) = \frac{e_{\kappa\alpha\nu}(\mathbf{q}) e_{\kappa'\alpha'\nu}^*(\mathbf{q})}{M_{\kappa}} + \frac{e_{\kappa'\alpha'\nu}(\mathbf{q}) e_{\kappa\alpha\nu}^*(\mathbf{q})}{M_{\kappa'}}, \quad (197)$$

$$h_{mn, \kappa\alpha}(\mathbf{k}) = \sum_{\nu} (M_{\kappa} \omega_{0\nu})^{1/2} e_{\kappa\alpha\nu}(0) g_{mn\nu}(\mathbf{k}, 0). \quad (198)$$

In the case of the three translational modes at  $|\mathbf{q}| = 0$ , these definitions are replaced by  $g_{mn\nu}^{\text{DW}}(\mathbf{k}, \mathbf{q}) = 0$ ; see the discussion in Sec. XIII. The derivation of Eqs. (195)–(198) requires using Eqs. (20), (33)–(35), and (38), as well as taking the canonical average over the adiabatic nuclear quantum states.

Equation (195) involves a summation over unoccupied Kohn-Sham states and so does the Fan-Migdal self-energy in Eq. (167). The numerical convergence of these sums is challenging, since one needs to evaluate a very large number of unoccupied electronic states. To address this issue, Gonze, Boulanger, and Côté (2011) developed a procedure whereby only a subset of unoccupied states is required, along the lines of the DFPT equations of Sec. III.B.3.

The first *ab initio* calculations using the formalism of Allen and Heine were reported by Marini (2008), who investigated the temperature dependence of the optical absorption spectrum of silicon and boron nitride. In this work Marini included excitonic effects by combining the Bethe-Salpeter formalism (Onida, Reining, and Rubio, 2002) with the Allen-Heine theory and obtained good agreement with experiments by calculating the direct absorption peaks using DFT-LDA phonons and matrix elements (indirect optical absorption is discussed in Sec. IX.B).

The second application of the Allen-Heine theory using DFT and LDA was reported by Giustino, Louie, and Cohen (2010) for the case of diamond. They investigated the temperature dependence of the direct band gap of diamond and found that the Fan-Migdal and the Debye-Waller self-energies are of comparable magnitude. The calculations captured the characteristic Varshni effect (Fig. 3) and were able to reproduce the measured redshift of the band gap in the temperature range 80–800 K. These calculations were based on the adiabatic version of the Allen-Heine theory and employed Eqs. (196)–(198) for the Debye-Waller self-energy. The calculations by Giustino *et al.* confirmed the large ( $> 0.5$  eV) zero-point renormalization of the direct gap of diamond predicted by Zollner, Cardona, and Gopalan (1992) using the empirical pseudopotential method.

The unusually large zero-point correction to the electronic structure of diamond stimulated further work on this system: [Cannuccia and Marini \(2011, 2012\)](#) calculated the gap renormalization in diamond by employing both the adiabatic version of the Allen-Heine theory and the nonadiabatic Green's function approach, as described in Sec. V.B.5. Their calculations confirmed the large zero-point renormalization and showed that the adiabatic theory underestimates the effect to some extent. [Cannuccia and Marini \(2012, 2013\)](#) also analyzed the quasiparticle renormalization and the spectral function.

[Antonius et al. \(2014\)](#) revisited the electron-phonon interaction in diamond by assessing the reliability of the rigid-ion approximation and the importance of many-body *GW* quasiparticle corrections to the DFT-LDA band structure. The main findings were that the rigid-ion approximation introduces a very small error in diamond, of the order of  $\sim 10$  meV, while *GW* quasiparticle corrections can increase the electron-phonon renormalization of the band gap by as much as  $\sim 200$  meV. The temperature dependence of the band gap of diamond calculated by [Antonius et al.](#) is shown in Fig. 13.

Further work on diamond was reported by [Poncé, Antonius, Gillet et al. \(2014\)](#), who compared *ab initio* calculations based on the Allen-Heine theory with explicit frozen-phonon calculations (Sec. IX.A.2). The corrections to the rigid-ion approximation were found to be smaller than 4 meV in all cases considered. [Poncé, Antonius, Boulanger et al. \(2014\)](#) and [Poncé, Antonius, Gillet et al. \(2014\)](#) also reported a detailed assessment of the accuracy of the various levels of approximation in the calculation of the zero-point renormalization of energy levels, as well as a thorough comparison between the results of different first-principles implementations.

The electron-phonon renormalization of band structures was also investigated in a number of other systems. For example, [Kawai et al. \(2014\)](#) studied zinc-blende GaN by combining the Allen-Heine theory with the Bethe-Salpeter approach. [Poncé et al. \(2015\)](#) investigated silicon, diamond, BN,  $\alpha$ -AlN, and  $\beta$ -AlN using both the adiabatic version of the

Allen-Heine theory and the nonadiabatic Green's function method of Eqs. (157) and (158). [Friedrich et al. \(2015\)](#) investigated the zero-point renormalization in LiNbO<sub>3</sub> using the adiabatic Allen-Heine theory. [Villegas, Rocha, and Marini \(2016\)](#) studied the anomalous temperature dependence of the band gap of black phosphorous. [Antonius et al. \(2015\)](#) investigated diamond, BN, LiF, and MgO, focusing on the dynamical aspects and the spectral function (see Sec. VIII). [Antonius et al. \(2015\)](#) and [Poncé et al. \(2015\)](#) were the first to report complete band structures at finite temperature.

[Poncé et al. \(2015\)](#) paid special attention to the numerical convergence of the self-energy integrals with respect to the limit  $\eta \rightarrow 0$  of the broadening parameter. They noted that in the case of polar crystals the adiabatic correction to the electron energies of band extrema, as given by Eq. (167), diverges in the limit of dense Brillouin-zone sampling. This behavior stems from the polar singularity in the electron-phonon matrix elements, Eq. (180). In fact, near band extrema the bands are approximately parabolic, and the integrand in the adiabatic Fan-Migdal self-energy goes as  $q^{-4}$  for  $q \rightarrow 0$ , while the volume element goes only as  $d\mathbf{q} = 4\pi q^2 dq$ . This problem can be avoided by first performing the integration over  $\mathbf{q}$  in principal value, without neglecting phonon frequencies, and then taking the limit  $\omega_{\mathbf{q}\nu} \rightarrow 0$  so as to recover the adiabatic approximation ([Fan, 1951](#), Sec. IV). In this way the adiabatic approximation can still be employed without incurring into a singularity in the calculations. A practical strategy to correctly perform the principal value integration in first-principles calculations was recently proposed by [Nery and Allen \(2016\)](#); they treated the singularity via an explicitly analytic integration near  $\mathbf{q} = 0$ . The complications arising in polar materials can also be avoided at once by directly using the more accurate expression in Eq. (166) based on Brillouin-Wigner perturbation theory, or even better by calculating the spectral functions as in [Kawai et al. \(2014\)](#) and [Antonius et al. \(2015\)](#). In particular, Eq. (166) clearly shows that in more accurate approaches the infinitesimal  $\eta$  should be replaced by the finite physical linewidth  $\Gamma_{nk}$ .

Although temperature-dependent band structures of polar materials were recently reported ([Kawai et al., 2014](#); [Antonius et al., 2015](#); [Poncé et al., 2015](#)), the specific role of the Fröhlich coupling discussed in Sec. VI.A.3 has received little attention so far. The only *ab initio* investigations which specifically addressed the role of polar couplings in this context are from [Botti and Marques \(2013\)](#) and [Nery and Allen \(2016\)](#). In order to understand the strategy of Botti and Marques, we consider the Hedin-Baym equations in Sec. IV.B.1. Botti and Marques proposed that, instead of splitting the screened Coulomb interaction  $W$  into electronic and nuclear contributions as in Eq. (104), one could try to directly calculate the screened Coulomb interaction  $W$  including the lattice screening, as in Eqs. (85) and (86). In order to make the calculations tractable, Botti and Marques evaluated the total dielectric matrices using a simplified model based on the Lyddane-Sachs-Teller relations. The resulting formalism combines *GW* calculations and experimentally measured LO-TO splittings. The zero-point renormalization of the band gaps calculated by Botti and Marques for LiF, LiCl, NaCl, and MgO were all  $> 1$  eV. This is an interesting result and




FIG. 13. Temperature dependence of the direct band gap of diamond calculated using the Allen-Heine theory. The upper curve shows the results obtained within DFPT at the LDA level. The lower curve was obtained via *GW* calculations in the frozen-phonon approach. Triangles are experimental data. The zero-point renormalization calculated by including *GW* quasiparticle corrections is 628 meV. From [Antonius et al., 2014](#).

deserves further investigation. We note incidentally that the Allen-Heine theory and that of Botti and Marques can both be derived from the Hedin-Baym equations. Therefore the approach of Botti and Marques should effectively correspond to calculating the Fan-Migdal self-energy by retaining only the long-range part of the polar electron-phonon matrix elements, i.e., Eq. (180). In the case of the work by Nery and Allen (2016), they reported a Fröhlich contribution to the zero-point renormalization of the band gap of GaN of 45 meV to be compared with the total renormalization arising from all modes of 150 meV.

## 2. Nonperturbative adiabatic calculations

An alternative approach to the calculation of temperature-dependent band structures consists of avoiding perturbation theory and electron-phonon matrix elements altogether and replacing the entire methodology discussed in Sec. IX.A.1 by straightforward finite-difference calculations. To see how this alternative strategy works we perform a Taylor expansion of the Kohn-Sham eigenvalues  $\varepsilon_{nk}$  to second order in the atomic displacements  $\Delta\tau_{\kappa\alpha p}$  and then average the result on a nuclear wave function identified by the quantum numbers  $\{n_{q\nu}\}$ . After using Eq. (20) one obtains

$$\langle \varepsilon_{nk} \rangle_{\{n_{q\nu}\}} = \varepsilon_{nk} + \sum_{\nu} \int \frac{d\mathbf{q}}{\Omega_{\text{BZ}}} (n_{q\nu} + 1/2) \frac{\partial \varepsilon_{nk}}{\partial n_{q\nu}}, \quad (199)$$

where we used the formal definition  $\partial/\partial n_{q\nu} = \Delta_{q\nu} \Delta_{-q\nu}$ , and the variations  $\Delta_{q\nu}$  are the same as in Eqs. (33)–(35). The nuclear wave functions are obtained from the ground state in Eq. (B10) by applying the ladder operators, as discussed in Appendix B. Equation (199) can be generalized to finite temperature by considering a canonical average over all possible nuclear states. The result maintains the same form as in Eq. (199), except that we now have the Bose-Einstein occupations  $n_{q\nu}(T)$  (see footnote 3). Equation (199) is precisely the conceptual starting point of the Allen-Heine theory of temperature-dependent band structures and appeared for the first time in Allen and Cardona (1981). If the variations  $\Delta_{q\nu} \Delta_{-q\nu} \varepsilon_{nk}$  are calculated in second-order perturbation theory, one obtains precisely the formalism of Allen and Heine.

It was proposed that the coefficients  $\partial \varepsilon_{nk} / \partial n_{q\nu}$  could alternatively be obtained from the derivatives of the vibrational frequencies with respect to the electronic occupations  $\hbar \partial \omega_{q\nu} / \partial f_{nk}$  (Allen and Hui, 1980; King-Smith *et al.*, 1989; Poncé, Antonius, Gillet *et al.*, 2014). A formal derivation of the link between these alternative approaches can be found in Allen and Hui (1980), Appendix; they refer to this as Brooks' theorem. Incidentally, the first *ab initio* calculation of temperature-dependent band gaps relied on this approach (King-Smith *et al.*, 1989).

Most commonly, the coefficients  $\partial \varepsilon_{nk} / \partial n_{q\nu}$  in Eq. (199) are evaluated using frozen-phonon supercell calculations, via the second derivative of the eigenvalue  $\varepsilon_{nk}$  with respect to collective atomic displacements along the vibrational eigenmodes  $e_{\kappa\alpha\nu}(\mathbf{q})$ . This approach was employed by Capaz *et al.* (2005) to study the temperature dependence of the band gaps in carbon nanotubes (within a tight-binding model),

and by Han and Bester (2013) to obtain the zero-point renormalization and temperature dependence of the gaps of silicon and diamond quantum dots. Recent examples include works on diamond, silicon, SiC (Monserat and Needs, 2014), as well as CsSnI<sub>3</sub> (Patrick, Jacobsen, and Thygesen, 2015).

Frozen-phonon supercell calculations based on Eq. (199) carry the advantage that the rigid-ion approximation, which is necessary to obtain Eqs. (195)–(198), is no longer required. Therefore this approach is more accurate in principle. In practice, however, the calculations are challenging as they require large supercells, and the derivatives must be evaluated for every vibrational mode of the supercell. Several computational strategies were developed to tackle this challenge. Patrick and Giustino (2013) proposed to perform the averages leading to Eq. (199) via importance-sampling Monte Carlo integration. Monserat (2016b) described a constrained Monte Carlo scheme which improves the variance of the Monte Carlo estimator. Recently Zacharias and Giustino (2016) showed that it is possible to perform these calculations more efficiently by replacing the stochastic sampling by a suitable choice of an “optimum” configuration; the result becomes exact in the thermodynamic limit of large supercells. In order to reduce the computational cost associated with the use of large supercells, Lloyd-Williams and Monserat (2015) introduced nondiagonal supercells, which allow one to access phonon wave vectors belonging to a uniform grid of  $N_p$  points using supercells containing only  $N_p^{1/3}$  unit cells.

The merit of these nonperturbative approaches is that they explicitly treat the nuclear wave functions and enable exploring effects which go beyond the Allen-Heine theory. For example, Monserat, Drummond, and Needs (2013), Monserat *et al.* (2014), Engel, Monserat, and Needs (2015), and Monserat, Engel, and Needs (2015) were able to investigate effects beyond the harmonic approximations in several systems, such as LiH, LiD, high-pressure He, molecular crystals of CH<sub>4</sub>, NH<sub>3</sub>, H<sub>2</sub>O, HF, as well as ice. In all these cases they found large zero-point effects on the band gaps.

Finally, we mention that the calculation of electronic properties at finite temperature via the Allen-Heine theory and its variants is closely related to what one would obtain using path-integral molecular dynamics simulations (Della Sala *et al.*, 2004; Ramírez, Herrero, and Hernández, 2006; Ramírez *et al.*, 2008), or even classical molecular dynamics simulations at high enough temperatures (Franceschetti, 2007).

## B. Phonon-assisted optical absorption

In addition to modifying the electron energy levels in solids, the electron-phonon interaction plays an important role in the optical properties of semiconductors and insulators, as it is responsible for phonon-assisted optical transitions. Phonon-assisted processes could be analyzed by considering the many-body electronic screening function  $\epsilon_c(12)$  in Eq. (95), by using the electron Green's function  $G$  dressed by the electron-phonon self-energy  $\Sigma^{\text{ep}}$  as in Eq. (151). Since this requires a lengthy detour, here we simply reproduce the standard result of second-order time-dependent perturbation theory (Bassani and Parravicini, 1975; Ridley, 1993):

$$\begin{aligned}
 \alpha(\omega) = & \frac{\pi e^2}{\epsilon_0 c \Omega \omega n_r(\omega)} \int \frac{d\mathbf{k} d\mathbf{q}}{\Omega_{\text{BZ}}^2} \sum_{m\nu} \sum_{s=\pm 1} (f_{n\mathbf{k}} - f_{m\mathbf{k}+\mathbf{q}}) \\
 & \times \left| \mathbf{e} \cdot \sum_p \left[ \frac{\mathbf{v}_{np}(\mathbf{k}) g_{p\nu}(\mathbf{k}, \mathbf{q})}{\epsilon_{p\mathbf{k}} - \epsilon_{n\mathbf{k}} - \hbar\omega} + \frac{g_{np\nu}(\mathbf{k}, \mathbf{q}) \mathbf{v}_{pm}(\mathbf{k} + \mathbf{q})}{\epsilon_{p\mathbf{k}+\mathbf{q}} - \epsilon_{n\mathbf{k}} + s\hbar\omega_{q\nu}} \right] \right|^2 \\
 & \times (n_{q\nu} + 1/2 + s/2) \delta(\epsilon_{m\mathbf{k}+\mathbf{q}} - \epsilon_{n\mathbf{k}} - \hbar\omega + s\hbar\omega_{q\nu}).
 \end{aligned} \quad (200)$$

In this expression  $\alpha(\omega)$  is the absorption coefficient for visible light,  $\mathbf{e}$  is the photon polarization,  $\mathbf{v}_{mn}$  are the matrix elements of the electron velocity operator, and  $n_r(\omega)$  is the real part of the refractive index. The two denominators in the second line correspond to indirect processes whereby a photon is absorbed and a phonon is absorbed or emitted (left), and processes whereby a phonon is absorbed or emitted, and subsequently a photon is absorbed (right). Equation (200) relies on the electric dipole approximation and is therefore valid for photon energies up to a few electron volts. The theory leading to Eq. (200) was originally developed by Hall, Bardeen, and Blatt (1954).

The first *ab initio* calculation employing Eq. (200) was reported by Noffsinger *et al.* (2012) for the prototypical case of silicon. They employed DFT for computing phonons and electron-phonon matrix elements and the *GW* method for the quasiparticle band structures. The sampling of the Brillouin zone was achieved by means of the interpolation strategy described in Sec. VI. Figure 14 shows that the calculations by Noffsinger *et al.* are in very good agreement with experiment throughout the energy range of indirect absorption.

Further work along similar lines was reported by Kioupakis *et al.* (2010), who calculated the indirect optical absorption by free carriers in GaN, and Peelaers, Kioupakis, and Van de Walle (2015), who studied the indirect absorption by free carriers in the transparent conducting oxide SnO<sub>2</sub>. Recently, the *ab initio* theory of phonon-assisted absorption was also




FIG. 14. Phonon-assisted optical absorption in silicon: comparison between first-principles calculations (solid lines, orange) and experiment (circles, blue). The calculations were performed using the theory of Hall, Bardeen, and Blatt (1954), as given by Eq. (200). Spectra calculated at different temperatures were shifted horizontally so as to match the experimental onsets. From Noffsinger *et al.*, 2012.

extended to the case of indirect Auger recombination by Kioupakis *et al.* (2015).

One limitation of the theory by Hall *et al.* is that the indirect absorption onset is independent of temperature. This is seen by noting that the Dirac delta functions in Eq. (200) contain the band structure energies at *clamped* nuclei. The generalization to incorporate temperature-dependent band structures as discussed in Sec. IX.A.1 is nontrivial. Patrick and Giustino (2014) and Zacharias and Giustino (2016) showed that the electron-phonon renormalization of the band structure modifies the energies of *real* transitions but leaves unchanged the energies of *virtual* transitions. In other words the Allen-Heine renormalization should be incorporated only in the Dirac delta functions and in the first denominator in Eq. (200). In order to avoid these complications, Zacharias, Patrick, and Giustino (2015) developed an alternative approach which relies on the semiclassical approximation of Williams (1951) and Lax (1952). In this approximation, the initial states in the optical transitions are described quantum mechanically, and the final states are replaced by a quasiclassical continuum. In the formulation of Zacharias *et al.* the imaginary part of the temperature-dependent dielectric function takes the following form:

$$\epsilon_2(\omega; T) = \frac{1}{Z} \sum_{\{n_{q\nu}\}} e^{-E_{\{n_{q\nu}\}}/k_B T} \langle \epsilon_2(\omega) \rangle_{\{n_{q\nu}\}}, \quad (201)$$

where  $\epsilon_2(\omega)$  denotes the imaginary part of the dielectric function at clamped nuclei, and the expectation values have the same meaning as in Eq. (199).  $E_{\{n_{q\nu}\}}$  is the energy of the quantum nuclear state specified by the quantum number  $\{n_{q\nu}\}$  and  $Z$  is the canonical partition function. Zacharias *et al.* demonstrated that this approach provides an *adiabatic* approximation to Eq. (200) and seamlessly includes the temperature dependence of the electronic structure within the Allen-Heine theory. Using techniques similar to those of Sec. IX.A.2, they calculated the indirect optical absorption line shape of silicon at various temperatures and obtained very good agreement with experiment. These results were recently extended to the temperature-dependent optical spectra of diamond and gallium arsenide by Zacharias and Giustino (2016).

## X. CARRIER DYNAMICS AND TRANSPORT

### A. Electron linewidths and lifetimes

In Sec. V.B.4 we have seen how the Fan-Migdal self-energy can be used in order to evaluate the quasiparticle lifetimes (or equivalently linewidths) resulting from the electron-phonon interaction. The first *ab initio* calculations of this kind were reported by Figuren *et al.* (2002) and Figuren, de Gironcoli *et al.* (2003) in the study of the decay of metal surface states, and by Sklyadneva *et al.* (2005, 2006) and Leonardo *et al.* (2007) in the study of electron lifetimes of elemental metals. Some of these calculations and the underlying approximations were reviewed in Sec. VIII. Calculations of quasiparticle linewidths were also employed to study the temperature-dependent broadening of the optical spectra in

semiconductors. For example, [Giustino, Louie, and Cohen \(2010\)](#) and [Poncé \*et al.\* \(2015\)](#) investigated the broadening of the direct absorption edge of diamond and silicon, respectively. In both cases good agreement with experiment was obtained. More recently, the same approach was employed to study the broadening of photoluminescence peaks in lead-iodide perovskites ([Wright \*et al.\*, 2016](#)). In this case, it was found that the standard Fermi golden rule expression, Eq. (170), significantly overestimates the experimental data. The agreement with experiment is restored by taking into account the quasiparticle renormalization  $Z_{nk}$ ; see the discussion following Eq. (174).

While these works were primarily concerned with the broadening of the spectral lines in photoemission or optical experiments, Eq. (170) can also be used to study carrier lifetimes in time-resolved experiments. The first *ab initio* study in this direction was reported by [Sjakste, Vast, and Tyuterev \(2007\)](#), who investigated the thermalization of hot electrons in GaAs and the exciton lifetimes in GaP. In the case of GaAs, [Sjakste \*et al.\*](#) found thermalization rates in quantitative agreement with time-resolved luminescence and transient optical absorption measurements. Work along similar lines was also reported for the intervalley scattering times in Ge ([Tyuterev \*et al.\*, 2011](#)). Recently, the thermalization rates of hot electrons in GaAs were revisited by [Bernardi \*et al.\* \(2015\)](#). They employed Eq. (170) and the Wannier interpolation technique described in Sec. VI in order to finely sample the electron-phonon scattering processes near the bottom of the conduction band; see Fig. 15. Based on these calculations they were able to interpret transient absorption measurements in terms of the carrier lifetimes within each valley. Another interesting application of Eq. (170) was reported by [Bernardi \*et al.\* \(2014\)](#), who investigated the rate of hot carrier thermalization in silicon, within the context of photovoltaics applications.

Recently [Sangalli and Marini \(2015a, 2015b\)](#) employed the lifetimes calculated using Eq. (170) in order to study carrier dynamics in silicon in real time. Strictly speaking, these developments lie outside of the scope of equilibrium Green's functions discussed in Sec. IV and require concepts based on nonequilibrium Green's functions ([Kadanoff and Baym, 1962](#)). However, the basic ingredients of the electron-phonon calculations remain unchanged.




FIG. 15. Electron relaxation times in GaAs resulting from electron-phonon scattering. (a) Calculated relaxation times as a function of electron energy with respect to the conduction band bottom. The color code (gray shades) of the data points identifies the valley where each electronic state belongs. (b) Schematic representation of the conduction band valleys in GaAs. From [Bernardi \*et al.\*, 2015](#).

In all calculations discussed in this section, the electron-phonon matrix elements were obtained within DFT. However, in order to accurately describe electron-phonon scattering near band extrema [Bernardi \*et al.\* \(2014, 2015\)](#) and [Wright \*et al.\* \(2016\)](#) employed *GW* quasiparticle band structures. This is important in order to obtain accurate band effective masses, which affect the carrier lifetimes via the density of states.

## B. Phonon-limited mobility

The carrier lifetimes  $\tau_{nk}$  of Eq. (170) are also useful in the calculation of electrical mobility, conductivity, and resistivity, within the context of the semiclassical model of electron dynamics in solids ([Ashcroft and Mermin, 1976](#)). In the semiclassical model, one describes the electronic response to an external perturbation by taking the fermionic occupations  $f_{nk}$  to represent the probability density function in the phase space defined by the unperturbed band structure. The probabilities  $f_{nk}$  are then determined using a standard Boltzmann equation. A comprehensive discussion of these methods can be found in the classic book of [Ziman \(1960\)](#).

Here we touch only upon the key result which is needed in *ab initio* calculations of electrical conductivity. In the semiclassical model, the electrical current is calculated as  $\mathbf{J} = -2e\Omega_{\text{BZ}}^{-1} \sum_n \int d\mathbf{k} \mathbf{v}_{nk} f_{nk}$ . In the absence of external fields and thermal gradients, the occupations  $f_{nk}$  reduce to the standard Fermi-Dirac occupations at equilibrium  $f_{nk}^0$ , and the current vanishes identically. Upon introducing an electric field  $\mathbf{E}$ , the electrons respond by adjusting their occupations. In this model it is assumed that the variation  $f_{nk} - f_{nk}^0$  is so small that the electronic density is essentially the same as in the unperturbed system. The modified occupations can be calculated using the linearized Boltzmann transport equation ([Ziman, 1960](#))

$$\frac{\partial f_{nk}^0}{\partial \varepsilon_{nk}} \mathbf{v}_{nk} \cdot (-e)\mathbf{E} = - \sum_{\nu} \int \frac{d\mathbf{q}}{\Omega_{\text{BZ}}} \Gamma_{m\nu}(\mathbf{k}, \mathbf{q}) \times [(f_{nk} - f_{nk}^0) - (f_{m\mathbf{k}+\mathbf{q}} - f_{m\mathbf{k}+\mathbf{q}}^0)], \quad (202)$$

where the kernel  $\Gamma_{m\nu}(\mathbf{k}, \mathbf{q})$  is defined as

$$\Gamma_{m\nu}(\mathbf{k}, \mathbf{q}) = \sum_{s=\pm 1} \frac{2\pi}{\hbar} |g_{m\nu}(\mathbf{k}, \mathbf{q})|^2 f_{nk}^0 (1 - f_{m\mathbf{k}+\mathbf{q}}^0) \times (n_{\mathbf{q}\nu} + 1/2 - s/2) \delta(\varepsilon_{nk} + s\hbar\omega_{\mathbf{q}\nu} - \varepsilon_{m\mathbf{k}+\mathbf{q}}). \quad (203)$$

The left-hand side of Eq. (202) represents the collisionless term of the Boltzmann equation, that is, the change in occupations due to the particle drift under the electric field. The right-hand side represents the change of occupations resulting from electrons scattered in or out of the state  $|n\mathbf{k}\rangle$  by phonon emission or absorption. The rates in Eq. (203) are simply derived from Fermi's golden rule ([Grimvall, 1981](#)). By solving Eq. (202) self-consistently for all  $f_{nk}$  it is possible to calculate the current and from there the conductivity. The connection with the carrier lifetimes  $\tau_{nk}$  of Eq. (170) is

obtained within the so-called “energy relaxation time approximation.” In this approximation the incoming electrons are neglected in Eq. (202); that is, the last term ( $f_{m\mathbf{k}+\mathbf{q}} - f_{m\mathbf{k}+\mathbf{q}}^0$ ) in the second line is ignored. As a result the entire right-hand side of the equation simplifies to  $-(f_{n\mathbf{k}} - f_{n\mathbf{k}}^0)/\tau_{n\mathbf{k}}$ .

The direct solution of Eq. (202) is computationally challenging, and fully *ab initio* calculations were reported only recently by Li (2015) for Si, MoS<sub>2</sub>, and Al, and by Fiorentini and Bonini (2016) for *n*-doped Si. Figure 16 shows that the mobility of *n*-type silicon calculated by Li is in good agreement with experiment. The theory overestimates the measured values to some extent, and this might have to do with the limitations of the DFT matrix elements (see Sec. XII). In addition to the carrier mobility, Fiorentini and Bonini employed the *ab initio* Boltzmann formalism to calculate thermoelectric properties, such as the Lorenz number and the Seebeck coefficient.

The first *ab initio* calculation of mobility was reported by Restrepo, Varga, and Pantelides (2009) for the case of silicon, within the energy relaxation time approximation. Other recent calculations using various approximations to Eq. (202) focused on silicon (Wang *et al.*, 2011; Liao *et al.*, 2015), graphene (Borysenko *et al.*, 2010; Park *et al.*, 2014; Restrepo *et al.*, 2014; Sohler *et al.*, 2014; Gunst *et al.*, 2016; Kim, Park, and Marzari, 2016), MoS<sub>2</sub> (Kaasbjerg, Thygesen, and Jacobsen, 2012; Li *et al.*, 2013; Restrepo *et al.*, 2014; Gunst *et al.*, 2016), silicene (Li *et al.*, 2013; Gunst *et al.*, 2016), and SrTiO<sub>3</sub> and KTiO<sub>3</sub> (Himmetoglu *et al.*, 2014; Himmetoglu and Janotti, 2016).

*Ab initio* calculations of the resistivity of metals are less challenging than for semiconductors and started appearing already from the work of Bauer *et al.* (1998). Most calculations on metals are based on Ziman’s resistivity formula; see Grimvall (1981). An interesting recent example can be found in the work by Xu and Verstraete (2014) on the transport coefficients of lithium. We also highlight related work on phonon-limited transport in organic crystals (Hannewald and Bobbert, 2004a, 2004b; Ortmann, Bechstedt, and Hannewald, 2009; Vukmirović, Bruder, and Stojanović, 2012).




FIG. 16. Temperature-dependent mobility of *n*-type silicon. The solid line (red) indicates the mobility calculated using the linearized Boltzmann transport equation, Eq. (202), and the dashed line (blue) corresponds to the energy relaxation time approximation. The triangles and diamonds are experimental data points. From Li, 2015.

## XI. PHONON-MEDIATED SUPERCONDUCTORS

The last application of *ab initio* calculations of EPIs that we consider here is the study of phonon-mediated superconductivity (Schrieffer, 1983). This research field is so vast that any attempt at covering it in a few pages would not do justice to the subject. For this reason, it was decided to limit the discussion to those novel theoretical and methodological developments which are aiming at fully *predictive* calculations, namely, the “anisotropic Migdal-Eliashberg theory” (Sec. XI.B) and the “density-functional theory for superconductors” (Sec. XI.C). For completeness, in Sec. XI.A we also summarized the most popular equations employed in the study of phonon-mediated superconductors. All calculations described in this section were performed at the DFT level.

### A. McMillan–Allen-Dynes formula

Most *ab initio* calculations on phonon-mediated superconductors rely on a semiempirical expression for the critical temperature, first introduced by McMillan (1968) and then refined by Allen and Dynes (1975):

$$k_B T_c = \frac{\hbar \omega_{\log}}{1.2} \exp \left[ -\frac{1.04(1 + \lambda)}{\lambda - \mu^*(1 + 0.62\lambda)} \right]. \quad (204)$$

Here  $T_c$  is the superconducting critical temperature,  $\omega_{\log}$  is a logarithmic average of the phonon frequencies (Allen and Dynes, 1975),  $\lambda$  is the electron-phonon “coupling strength,” and  $\mu^*$  is a parameter describing the Coulomb repulsion. The functional form of Eq. (204) was derived by McMillan by determining an approximate solution of the Eliashberg gap equations (see Sec. XI.B).  $\lambda$  and  $\omega_{\log}$  are calculated from the isotropic version of the Eliashberg function in Eq. (187) as follows (McMillan, 1968; Allen and Dynes, 1975; Grimvall, 1981; Allen and Mitrović, 1982):

$$\alpha^2 F(\omega) = \frac{1}{N_F} \int \frac{d\mathbf{k}d\mathbf{q}}{\Omega_{\text{BZ}}^2} \sum_{m\nu} |g_{m\nu}(\mathbf{k}, \mathbf{q})|^2 \delta(\varepsilon_{n\mathbf{k}} - \varepsilon_F) \times \delta(\varepsilon_{m\mathbf{k}+\mathbf{q}} - \varepsilon_F) \delta(\hbar\omega - \hbar\omega_{\mathbf{q}}), \quad (205)$$

$$\lambda = 2 \int_0^\infty \frac{\alpha^2 F(\omega)}{\omega} d\omega, \quad (206)$$

$$\omega_{\log} = \exp \left[ \frac{2}{\lambda} \int_0^\infty d\omega \frac{\alpha^2 F(\omega)}{\omega} \log \omega \right], \quad (207)$$

where  $N_F$  is the density of states at the Fermi level and the matrix elements  $g_{m\nu}(\mathbf{k}, \mathbf{q})$  are the same as in Eq. (38). The remaining parameter  $\mu^*$  (Morel and Anderson, 1962) is obtained as  $1/\mu^* = 1/\mu + \log(\omega_p/\omega_{\text{ph}})$ , where  $\hbar\omega_p$  is the characteristic plasma energy of the system,  $\hbar\omega_{\text{ph}}$  the largest phonon energy, and  $\mu$  is the average electron-electron Coulomb repulsion across the Fermi surface. More specifically,  $\mu = N_F \langle \langle V_{n\mathbf{k},n'\mathbf{k}'} \rangle \rangle_{\text{FS}}$ , where  $\langle \langle \dots \rangle \rangle_{\text{FS}}$  denotes a double average over the Fermi surface, and  $V_{n\mathbf{k},n'\mathbf{k}'} = \langle \mathbf{k}'n', -\mathbf{k}'n' | W | \mathbf{k}n, -\mathbf{k}n \rangle$ , with  $W$  the screened Coulomb interaction of Sec. IV.B.2 (Lee, Chang, and Cohen, 1995; Lee and Chang, 1996).

The coupling strength  $\lambda$  is related to the mass enhancement parameter  $\lambda_{n\mathbf{k}}$  discussed in Sec. VIII.A. The main difference between  $\lambda$  and  $\lambda_{n\mathbf{k}}$  is that the former represents an average over the Fermi surface, while the latter refers to the Fermi velocity renormalization of a specific electron band. While these quantities are related, they do not coincide and hence cannot be used interchangeably.

Equations (204)–(207) involve a number of approximations. For example, it is assumed that the superconductor is isotropic and exhibits a single superconducting gap. Furthermore, almost invariably the effective Coulomb potential  $\mu^*$  is treated as an adjustable parameter on the grounds that it should be in the range  $\mu^* = 0.1$ – $0.2$ . This procedure introduces a large uncertainty in the determination of  $T_c$ , especially at moderate coupling strengths.

### B. Anisotropic Migdal-Eliashberg theory

A first-principles approach to the calculation of the superconducting critical temperature is provided by the anisotropic Migdal-Eliashberg theory (Migdal, 1958; Eliashberg, 1960). This is a field-theoretic approach to the superconducting pairing, formulated in the language of finite-temperature Green's functions. At variance with the Hedin-Baym equations of Table I, the Migdal-Eliashberg theory is best developed within the Nambu-Gor'kov formalism (Gor'kov, 1958; Nambu, 1960), which enables describing the propagation of electron quasiparticles and of superconducting Cooper pairs on the same footing (Scalapino, 1969; Schrieffer, 1983). A comprehensive presentation of the Migdal-Eliashberg theory is provided by Allen and Mitrovic (1982). Their article served as the starting point of current first-principles implementations of the theory.

In the Migdal-Eliashberg theory, one solves the following two coupled equations:

$$Z_{n\mathbf{k}}(i\omega_j) = 1 + \frac{\pi k_B T}{N_F} \sum_{n'\mathbf{k}'j'} \frac{\omega_{j'}/\omega_j}{\sqrt{\hbar^2 \omega_{j'}^2 + \Delta_{n'\mathbf{k}'}^2(i\omega_{j'})}} \times \lambda_{n\mathbf{k},n'\mathbf{k}'}(i\omega_j - i\omega_{j'}) \delta(\varepsilon_{n'\mathbf{k}'} - \varepsilon_F), \quad (208)$$

$$Z_{n\mathbf{k}}(i\omega_j) \Delta_{n\mathbf{k}}(i\omega_j) = \frac{\pi k_B T}{N_F} \sum_{n'\mathbf{k}'j'} \frac{\Delta_{n'\mathbf{k}'}(i\omega_{j'})}{\sqrt{\hbar^2 \omega_{j'}^2 + \Delta_{n'\mathbf{k}'}^2(i\omega_{j'})}} \times [\lambda_{n\mathbf{k},n'\mathbf{k}'}(i\omega_j - i\omega_{j'}) - N_F V_{n\mathbf{k},n'\mathbf{k}'}] \times \delta(\varepsilon_{n'\mathbf{k}'} - \varepsilon_F), \quad (209)$$

where  $\sum_{\mathbf{k}'}$  stands for  $\Omega_{\text{BZ}}^{-1} \int d\mathbf{k}'$ . In these equations,  $T$  is the absolute temperature, and  $Z_{n\mathbf{k}}(i\omega_j)$  is the quasiparticle renormalization function and is analogous to  $Z_{n\mathbf{k}}$  in Eq. (189).  $\Delta_{n\mathbf{k}}(i\omega_j)$  is the superconducting gap function. The functions  $Z_{n\mathbf{k}}(i\omega_j)$  and  $\Delta_{n\mathbf{k}}(i\omega_j)$  are determined along the imaginary frequency axis at the fermion Matsubara frequencies  $i\omega_j = i(2j+1)\pi k_B T/\hbar$  with  $j$  an integer. The anisotropic and frequency-dependent generalization of Eq. (206) to be used in the Migdal-Eliashberg equations is

$$\lambda_{n\mathbf{k},n'\mathbf{k}'}(i\omega) = \frac{N_F}{\hbar} \sum_{\nu} \frac{2\omega_{\mathbf{q}\nu}}{\omega_{\mathbf{q}\nu}^2 + \omega^2} |g_{nn'\nu}(\mathbf{k}, \mathbf{q})|^2, \quad (210)$$

with  $\mathbf{q} = \mathbf{k}' - \mathbf{k}$ . Equations (208) and (209) are to be solved self-consistently for each temperature  $T$ . The superconducting critical temperature is then obtained as the highest temperature for which a nontrivial solution is obtained, that is, a solution with  $\Delta_{n\mathbf{k}}(i\omega_j) \neq 0$ . From the superconducting gap along the imaginary axis it is then possible to obtain the gap at real frequencies by analytic continuation (Marsiglio, Schossmann, and Carbotte, 1988), and from there one can compute various thermodynamic functions.

The first *ab initio* implementation of the anisotropic Migdal-Eliashberg theory was reported by Choi *et al.* (2002a, 2002b) and Choi, Cohen, and Louie (2003) in a study of the superconducting properties of MgB<sub>2</sub>. They succeeded in explaining the anomalous heat capacity of MgB<sub>2</sub> in terms of two distinct superconducting gaps and obtained a  $T_c$  in good agreement with experiment. These calculations were later extended to MgB<sub>2</sub> under pressure (Choi, Louie, and Cohen, 2009a) and other hypothetical borides (Choi, Louie, and Cohen, 2009b). Margine and Giustino (2013) demonstrated an implementation of the Migdal-Eliashberg theory based on the Wannier interpolation scheme of Sec. VI and reported applications to Pb and MgB<sub>2</sub>. The superconducting gap and superconducting density of states of MgB<sub>2</sub> calculated by Margine and Giustino are shown in Fig. 17. In all these calculations, the Coulomb repulsion




FIG. 17. (a) Energy distribution of the superconducting gap function of MgB<sub>2</sub> as a function of temperature, calculated using the anisotropic Migdal-Eliashberg theory. The gap vanishes at the critical temperature (in this calculation  $T_c = 50$  K). Two distinct superconducting gaps can be seen at each temperature. (b) Density of electronic states in the superconducting state of MgB<sub>2</sub> at various temperatures calculated within the Migdal-Eliashberg theory. From Margine and Giustino, 2013.



was described empirically via  $\mu^*$ , and this partly accounts for the slight discrepancy between the calculated  $T_c$  of 50 K and the experimental  $T_c$  of 39 K (Nagamatsu *et al.*, 2001). Additional calculations based on the anisotropic Migdal-Eliashberg theory included a study of doped graphene (Margine and Giustino, 2014), as well as investigations of Li-decorated monolayer graphene (Zheng and Margine, 2016) and Ca-intercalated bilayer graphene (Margine, Lambert, and Giustino, 2016). In this latter work they incorporated Coulomb interactions from first principles, after calculating  $\mu^*$  via the screened Coulomb interaction in the random-phase approximation. The calculated  $T_c = 7\text{--}8$  K was in reasonable agreement with the experimental value of 4 K (Ichinokura *et al.*, 2016). The Migdal-Eliashberg theory was also extended to describe the superconducting state as a function of applied magnetic field. A complete *ab initio* implementation was successfully demonstrated with an application to  $\text{MgB}_2$  (Aperis, Maldonado, and Oppeneer, 2015). Recently Sano *et al.* (2016) performed *ab initio* Migdal-Eliashberg calculations including retardation effects on high-pressure sulfur hydrides, obtaining good agreement with experiment. Interestingly in this work they also checked the effect of the zero-point renormalization of the electron bands within the Allen-Heine theory and found that it accounts for a change in  $T_c$  of up to 20 K.

### C. Density-functional theory for superconductors

Another promising *ab initio* approach to the calculation of the superconducting critical temperature is the density-functional theory for superconductors (Lüders *et al.*, 2005; Marques *et al.*, 2005). The starting point of this approach is a generalization of the Hohenberg-Kohn theorem (Hohenberg and Kohn, 1964) to a system described by three densities: the electron density in the normal state, the density of superconducting pairs, and the nuclear density. Based on this premise, Lüders *et al.* mapped the fully interacting system into an equivalent Kohn-Sham system (Kohn and Sham, 1965) of noninteracting nuclei and noninteracting, yet superconducting, electrons. The resulting Kohn-Sham equations for the electrons take the form of Bogoliubov–de Gennes equations (Bogoliubov, 1958), whereby electrons are paired by an effective gap function  $\Delta(\mathbf{r}, \mathbf{r}')$ .

In its simplest formulation, the density-functional theory for superconductors determines the expectation value of the pairing field over Kohn-Sham eigenstates  $\Delta_{n\mathbf{k}} = \langle u_{n\mathbf{k}}(\mathbf{r}) | \Delta(\mathbf{r}, \mathbf{r}') | u_{n\mathbf{k}}(\mathbf{r}') \rangle$ , using the following gap equation:

$$\Delta_{n\mathbf{k}} = -\mathcal{Z}_{n\mathbf{k}} \Delta_{n\mathbf{k}} - \sum_{n'\mathbf{k}'} \frac{\mathcal{K}_{n\mathbf{k}, n'\mathbf{k}'}}{2E_{n'\mathbf{k}'}} \tanh\left(\frac{E_{n'\mathbf{k}'}}{2k_B T}\right), \quad (211)$$

where  $E_{n\mathbf{k}}^2 = \varepsilon_{n\mathbf{k}}^2 + |\Delta_{n\mathbf{k}}|^2$ . In this expression, the kernel  $\mathcal{K}$  contains information about the phonon-mediated pairing interaction and the Coulomb repulsion between electrons  $\mathcal{K} = \mathcal{K}^{\text{ep}} + \mathcal{K}^{\text{ee}}$ , and  $\mathcal{Z}$  contains information about the electron-phonon interaction. More specifically,  $\mathcal{K}^{\text{ep}}$  and  $\mathcal{Z}$  are evaluated starting from the electron-phonon matrix elements  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$  and the DFT electron band structure and phonon dispersions, as in the Migdal-Eliashberg theory.  $\mathcal{K}^{\text{ee}}$  is

approximated using the screened Coulomb interaction  $V_{n\mathbf{k}, n'\mathbf{k}'}$  introduced below Eq. (207). Complete expressions for  $\mathcal{Z}$  and  $\mathcal{K}$  can be found in Marques *et al.* (2005).

Equation (211) is reminiscent of the gap equation in the Bardeen-Cooper-Schrieffer (BCS) theory (Schrieffer, 1983), with the difference that the *ab initio* kernel  $\mathcal{K}$  replaces the model interaction of the BCS theory, and the function  $\mathcal{Z}$  introduces quasiparticle renormalization as in the Migdal-Eliashberg theory; see Eq. (208). At variance with the Migdal-Eliashberg theory, the gap function in the density-functional theory for superconductors does not carry an explicit frequency dependence. Nevertheless, retardation effects are fully included through the dependence of the kernels  $\mathcal{Z}$  and  $\mathcal{K}$  on the electron bands and the phonon dispersions. An important advantage of this theory is that the Coulomb potential  $\mu^*$  is not required, since the electron-electron repulsion is seamlessly taken into account by means of the kernel  $\mathcal{K}^{\text{ee}}$ .

The density-functional theory for superconductors was successfully employed to study the superconducting properties of  $\text{MgB}_2$  (Floris *et al.*, 2005), Li, K, and Al under pressure (Profeta *et al.*, 2006; Sanna *et al.*, 2006), Pb (Floris *et al.*, 2007), Ca-intercalated graphite (Sanna *et al.*, 2007), high-pressure hydrogen (Cudazzo *et al.*, 2008, 2010), CaBeSi (Bersier *et al.*, 2009), layered nitrides (Akashi *et al.*, 2012), alkali-doped fullerides (Akashi and Arita, 2013b), compressed sulfur hydrides (Akashi *et al.*, 2015), and intercalated layered carbides, silicides, and germanides (Flores-Livas and Sanna, 2015).

An interesting recent development of the theory was the determination of the superconducting order parameter in real space  $\chi(\mathbf{r}, \mathbf{r}') = \langle \hat{\psi}_\uparrow(\mathbf{r}) \hat{\psi}_\downarrow(\mathbf{r}') \rangle$  (Linscheid, Sanna, Floris, and Gross, 2015). In the density-functional theory for superconductors, the order parameter is obtained from the superconducting gap using  $\chi_{n\mathbf{k}} = (\Delta_{n\mathbf{k}}/2|E_{n\mathbf{k}}|) \tanh[E_{n\mathbf{k}}/(2k_B T)]$ . Figure 18 shows the order parameter calculated by Linscheid *et al.* for both  $\text{MgB}_2$  and hole-doped graphane (Savini, Ferrari, and Giustino, 2010). The plots show Friedel-like oscillations of the superconducting density as a function of the relative coordinates between two paired electrons.




FIG. 18. Superconducting order parameter  $\chi(\mathbf{r}, \mathbf{r}')$  in real space, calculated for (a)  $\text{MgB}_2$  and (b) hole-doped graphane. The plots show a top view (top) and a side view (bottom) of the hexagonal layers in each case. The variable  $\mathbf{s} = \mathbf{r} - \mathbf{r}'$  is the relative coordinate in the order parameter, while the center-of-mass coordinate is placed in the middle of a B-B bond or a C-C bond. From Linscheid, Sanna, Floris, and Gross, 2015.

Further developments of the superconducting density-functional theory include the study of nonphononic pairing mechanisms, such as plasmon-assisted superconductivity (Akashi and Arita, 2013a), and the extension to magnetic systems (Linscheid, Sanna, Essenerger, and Gross, 2015; Linscheid, Sanna, and Gross, 2015).

## XII. ELECTRON-PHONON INTERACTIONS BEYOND THE LOCAL DENSITY APPROXIMATION TO DFT

The calculations of electron-phonon interactions reviewed in Secs. VII–XI have in common the fact that most investigations used the local density approximation to DFT or a generalized gradient approximation (GGA) such as the PBE functional (Perdew, Burke, and Ernzerhof, 1996). Although the LDA and the GGA represent the workhorse of electron-phonon calculations from first principles, there is growing evidence that these choices can lead to an underestimation of the electron-phonon coupling strength. At a conceptual level we understand this point by rewriting the electron-phonon matrix element after combining Eqs. (38), (142), and (143):

$$g_{mn\nu}(\mathbf{k}, \mathbf{q}) = \langle u_{m\mathbf{k}+\mathbf{q}} | \int d\mathbf{r}' \epsilon_e^{-1}(\mathbf{r}, \mathbf{r}', \omega) \Delta_{q\nu} v^{\text{en}}(\mathbf{r}') | u_{n\mathbf{k}} \rangle_{\text{uc}}. \quad (212)$$

In DFT the many-body dielectric matrix  $\epsilon_e$  appearing in this expression is replaced by the RPA +  $xc$  screening  $\epsilon^{\text{Hxc}}$  from Eq. (54). Given the DFT band gap problem, we expect  $\epsilon^{\text{Hxc}}$  to overestimate the screening, thereby leading to matrix elements  $g_{mn\nu}(\mathbf{k}, \mathbf{q})$  which are underestimated to some extent.

Several groups investigated this point on quantitative grounds. Zhang, Louie, and Cohen (2007) studied the electron-phonon coupling in a model copper oxide superconductor  $\text{CaCuO}_2$ . By calculating the vibrational frequencies of the half-breathing Cu-O stretching mode, they established that the local spin-density approximation (LSDA) yields phonons which are too soft (65.3 meV) as compared to experiment (80.1 meV). In contrast, the introduction of Hubbard corrections in a LSDA +  $U$  scheme restored agreement with experiment (80.9 meV). Since the electron-phonon matrix elements are connected to the phonon frequencies via the phonon self-energy, Eq. (145), a corresponding underestimation of the matrix elements can be expected. These results were supported by the work of Floris *et al.* (2011), who developed DFPT within LSDA +  $U$  and applied their formalism to the phonon dispersions of antiferromagnetic MnO and NiO. They found that the DFT underestimates measured LO energies by as much as 15 meV in MnO, while the use of LSDA +  $U$  leads to good agreement with experiment. Related work was reported by Hong *et al.* (2012), who investigated the multi-ferroic perovskites  $\text{CaMnO}_3$ ,  $\text{SrMnO}_3$ ,  $\text{BaMnO}_3$ ,  $\text{LaCrO}_3$ ,  $\text{LaFeO}_3$ , and the double perovskite  $\text{La}_2\text{CrFeO}_6$ . They calculated the variation of the vibrational frequencies between the ferromagnetic and antiferromagnetic phases of these compounds as a function of the Hubbard  $U$  parameter and compared DFT +  $U$  calculations with hybrid-functional calculations.

Lazzeri *et al.* (2008) investigated the effect of quasiparticle  $GW$  corrections on the electron-phonon coupling of graphene and graphite, for the  $A_1'$  phonon at  $K$  and the  $E_{2g}$  phonon at  $\Gamma$ . They evaluated the intraband electron-phonon matrix elements using a frozen-phonon approach, noting that  $g_{mn\nu}(\mathbf{k}, \mathbf{q} = 0)$  precisely represents the shift of the Kohn-Sham energy  $\epsilon_{n\mathbf{k}}$  upon displacing the atoms according to the  $\nu$ th phonon eigenmode at  $\mathbf{q} = 0$ . Lazzeri *et al.* found that the matrix elements increase by almost 40% from DFT to  $GW$ . The  $GW$  values led to slopes in the phonon dispersions near  $K$  in very good agreement with inelastic x-ray scattering data (Grüneis *et al.*, 2009b). Similar results, albeit less dramatic, were obtained by Grüneis *et al.* (2009a) for the potassium-intercalated graphite  $\text{KC}_8$ .

Laflamme Janssen *et al.* (2010) studied the electron-phonon coupling in the  $\text{C}_{60}$  molecule as a model for superconducting alkali-doped fullerides. They employed the PBE0 hybrid functional (Perdew, Ernzerhof, and Burke, 1996) with a fraction of exact exchange  $\alpha = 30\%$  and obtained an enhancement of the total coupling strength  $\lambda$  of 42% as compared to PBE. This work was followed up by Faber *et al.* (2011), who used the  $GW$  approximation and obtained a similar enhancement of 48%. We also point out an earlier work by Saito (2002) based on the B3LYP functional, reporting similar results.

Yin, Kutepov, and Kotliar (2013) investigated the effects of using the  $GW$  approximation and the HSE hybrid functional (Heyd, Scuseria, and Ernzerhof, 2003) on the electron-phonon coupling in the superconducting bismuthates  $\text{Ba}_{1-x}\text{K}_x\text{BiO}_3$  and chloronitrides  $\beta\text{-ZrNCl}$ , as well as  $\text{MgB}_2$ . In the case of  $\text{Ba}_{1-x}\text{K}_x\text{BiO}_3$  they obtained a threefold increase in the coupling strength  $\lambda$  from PBE to HSE. This enhancement brought the critical temperature calculated using Eq. (204) to 31 K, very close to the experimental value of 32 K. Similarly, in the case of  $\beta\text{-ZrNCl}$ , Yin *et al.* obtained a 50% increase of  $\lambda$ , bringing the calculated critical temperature of 18 K close to the experimental value of 16 K. Instead, in the case of  $\text{MgB}_2$ , they noticed only a slight increase of the electron-phonon coupling as compared to the standard LDA.

Another application of hybrid functionals to the study of EPIs was reported by Komelj and Krakauer (2015). They investigated the sensitivity of the superconducting critical temperature of the  $\text{H}_3\text{S}$  phase of sulfur hydride to the exchange and correlation functionals. They found that the PBE0 functional enhances the critical temperature by up to 25% as compared to PBE, bringing  $T_c$  from 201–217 K to 253–270 K (the spread in values is related to the choice of the parameter  $\mu^*$ ).

Mandal, Cohen, and Haule (2014) reported work on the superconductor FeSe based on dynamical mean-field theory (DMFT). In this case DMFT yielded a threefold enhancement of the coupling strength for selected modes.

As mentioned in Sec. IX.A.1, Antonius *et al.* (2014) performed  $GW$  calculations of the electron-phonon coupling in diamond using a frozen-phonon approach. They found that quasiparticle corrections lead to a uniform enhancement of the electron-phonon matrix elements. The net effect is an increase of the zero-point renormalization of the band gap by 40% as compared to standard LDA calculations. Monserrat (2016a)

confirmed this result and found a *GW* correction of comparable magnitude in the case of silicon. However, Monserrat also found that the *GW* corrections to the zero-point band gap renormalization of LiF, MgO, and TiO<sub>2</sub> are very small ( $\sim 5\%$  of the PBE value); therefore at present it is not possible to draw general conclusions.

Finally, we mention that Faber *et al.* (2015) examined possible strategies for systematically incorporating *GW* corrections in electron-phonon calculations. By using diamond, graphene, and C<sub>60</sub> as test cases, they showed that a “constant screening” approximation is able to reproduce complete *GW* results with an error below 10% at reduced computational cost. This approximation amounts to evaluating the variation of the Green’s function *G* in a frozen-phonon calculation, while retaining the screened Coulomb interaction *W* of the unperturbed ground state.

All these recent developments point to the need of moving beyond local exchange and correlation functionals in the study of electron-phonon interactions from first principles. In the future, it will be important to devise accurate computational methods for calculating not only the intraband electron-phonon matrix elements (as in the frozen-phonon method) but also matrix elements between all states and for scattering across the entire Brillouin zone.

For completeness we emphasize that the underestimation of the EPI matrix elements by semilocal DFT functionals does not propagate in the same way into different materials properties. This is readily understood by examining two fundamental quantities, the Allen-Heine renormalization of electron bands, Eq. (167), and the adiabatic phonon frequencies, as obtained from Eqs. (126) and (127). In the former case the electronic screening enters as  $\epsilon_{\infty}^{-2}$ ; in the latter case the screening contributes through a term which scales with  $\epsilon_{\infty}^{-1/2}$ . As a result, in the hypothetical case of a semiconductor for which DFT underestimated the electronic permittivity by 20%, we would have an error of  $\sim 40\%$  in the energy renormalization, and of  $\sim 10\%$  in the phonon frequencies. This example is an oversimplification of the problem, but it shows that different properties relating to the EPI could be affected to a very different degree by the inherent limitations of DFT functionals.

### XIII. CONCLUSIONS

The study of electron-phonon interactions has a long and distinguished history, but it is only during the past two decades that quantitative and predictive calculations have become possible. First-principles calculations of electron-phonon couplings are finding an unprecedented variety of applications in many areas of condensed matter and materials physics, from spectroscopy to transport, from metals to semiconductors and superconductors. In this article we discussed the standard DFT formalism for performing calculations of electron-phonon interactions, showed how most equations can be derived from a field-theoretic framework using a few well-defined approximations, and reviewed recent applications of the theory to many materials of current interest.

As calculation methods improve relentlessly and quantitative comparisons between theory and experiment become

increasingly refined, new and more complex questions arise. Much is still left to do, both in the fundamental theory of electron-phonon interactions and in the development of more accurate and more efficient computational methods.

For one, we are still using theories where the coupling matrix elements are calculated using the adiabatic local density approximation to DFT. The need for moving beyond standard DFT and beyond the adiabatic approximation can hardly be overemphasized. Progress is being made on the incorporation of nonlocal corrections into electron-phonon matrix elements, for example, using hybrid functionals or *GW* techniques, but very little is known about retardation effects. It is expected that such effects may be important in the study of heavily doped oxides and semiconductors, in both their normal and superconducting states (Mahan, 1993, Sec. 6.3.A), but *ab initio* investigations are currently missing. This is truly uncharted territory and a systematic effort in this direction is warranted.

In this article we emphasized that it is possible to formulate a compact, unified theory of electron-phonon interactions starting from a fully *ab initio* field-theoretic approach. The only assumption which is absolutely crucial to the theory is the *harmonic* approximation. Abandoning the harmonic approximation leads to the appearance of several new terms in the equations, and the resulting formalism becomes considerably more complex than in Table I. Despite these difficulties, given the importance of anharmonic effects in many systems of current interest, extending the theory to the case of anharmonic phonons and multiphonon interactions constitutes a pressing challenge. *Ab initio* investigations of anharmonic effects on the temperature dependence of band gaps have recently been reported (Monserrat, Drummond, and Needs, 2013; Antonius *et al.*, 2015). Since these studies rely on nonperturbative adiabatic calculations in supercells, it is highly desirable to establish a clear formal connection of these methods with the rigorous field-theoretic approach of Sec. IV. Along the same line, it is important to clarify the relation between many-body approaches, adiabatic supercells calculations, and more traditional classical or path-integral molecular dynamics simulations.

The study of electron-phonon interactions has long been dominated by Fröhlich-like Hamiltonians, whereby the electron-phonon coupling is retained only to linear order in the atomic displacements. This is the case for all the model Hamiltonians mentioned in Sec. V.B.6. It is now clear that quadratic couplings, leading to the Debye-Waller contributions in the optical spectra of semiconductors, are by no means negligible and should be investigated more systematically. For example, in the current literature it is invariably assumed that Debye-Waller contributions are negligible in metals near the Fermi surface; while this is probably the case for the simplest elemental metals, what happens in the case of multiple Fermi-surface sheets is far from clear and should be tested by direct calculations.

The identification of the correct matrix elements to be calculated is not always a trivial task as discussed for the case of the nonadiabatic phonon self-energy. In the future it will be important to pay attention to these aspects, especially in view of detailed comparison with experiment. For now, the issue of whether the phonon self-energy arising from EPIs should be

calculated using bare or screened EPI matrix elements (Sec. VII) is to be considered an open question and calls for further investigation.

The theory and applications reviewed in this article focused on nonmagnetic systems. The rationale for this choice is that a complete many-body theory of electron-phonon interactions for magnetic systems is not available yet. Recent investigations of spin-phonon couplings were conducted by assuming that the spin and the vibrational degrees of freedom can be decoupled, as in the Born-Oppenheimer approximation. Under this assumption it is possible to investigate how the spin configuration responds to a frozen phonon or, alternatively, how the vibrational frequencies depend on the spin configuration [see, for example, Chan *et al.* (2007), Łażewski *et al.* (2010), Lee and Rabe (2011), and Cao, Giustino, and Radaelli (2015)]. In all these cases it is desirable to employ a more rigorous many-body theory of spin-phonon interactions. The Hedin-Baym equations discussed in Sec. IV maintain their validity in the case of spin-polarized systems, provided collinear spins are assumed. In more general situations, where it is important to consider noncollinear spins, external magnetic perturbations, or spin-dependent interactions such as spin-orbit and Rashba-Dresselhaus couplings, it becomes necessary to generalize the equations in Table I. Although such a generalization has not been reported yet, the work of Aryasetiawan and Biermann (2008) constitutes a promising starting point. In that work the Schwinger functional derivative technique (see Sec. IV.B.1) was used to extend Hedin's equations at clamped nuclei to systems containing spin-dependent interactions. Generalizing Aryasetiawan and Biermann's work to incorporate nuclear vibrations will be important for the study of electron-phonon interactions in many systems of current interest, from multifunctional materials to topological quantum matter.

At this time it is not possible to predict how this fast-moving field will evolve over the years to come. However, the impressive progress made during the past decade gives us confidence that this interesting research area will continue to thrive and will keep surprising us with fascinating challenges and exciting new opportunities.

## ACKNOWLEDGMENTS

I am indebted to Samuel Poncé, Carla Verdi, and Marios Zacharias for a critical reading of the manuscript, and to many colleagues who have kindly provided feedback on the arXiv preprint version of this article. This article is the result of many stimulating discussions with friends and colleagues over the last few years. I thank, in particular, Philip Allen, Stefano Baroni, Lilia Boeri, Nicola Bonini, Fabio Caruso, Hyounghoon Choi, Marvin Cohen, Michel Côté, Andrea Dal Corso, Claudia Draxl, Asier Eguren, Paolo Giannozzi, Stefano de Gironcoli, Giulia Galli, Xavier Gonze, Göran Grimvall, Eberhard Gross, Branko Gumhalter, Sohrab Ismail-Beigi, Emmanuil Kioupakis, Georg Kresse, Steven Louie, Roxana Margine, Andrea Marini, Miguel Marques, Nicola Marzari, Takashi Miyake, Bartomeu Monserrat, Jeffrey Neaton, Richard Needs, Jesse Noffsinger, Cheol-Hwan Park, Christopher Patrick, Warren Pickett, Samuel Poncé, Paolo Radaelli, Lucia Reining, John Rehr, Patrick Rinke, Angel

Rubio, Matthias Scheffler, Young-Woo Son, Alexander Tkatchenko, Chris Van de Walle, Carla Verdi, Matthieu Verstraete, and Marios Zacharias. I also thank Marco Bernardi, Asier Eguren, Xavier Gonze, Eberhard Gross, Wu Li, Steven Louie, and Francesco Mauri for kindly allowing me to reproduce their figures. This work was supported by the Leverhulme Trust (Grant No. RL-2012-001), the Graphene Flagship (Horizon 2020 Grant No. 696656-GrapheneCore), and the UK Engineering and Physical Sciences Research Council (Grants No. EP/J009857/1 and No. EP/M020517/1).

## APPENDIX A: BORN-VON KÁRMÁN BOUNDARY CONDITIONS

In this Appendix we provide more details on the notation related to the Born-von Kármán boundary conditions used throughout this article. The crystalline unit cell is defined by the primitive lattice vectors  $\mathbf{a}_i$  with  $i = 1, 2, 3$ , and the  $p$ th unit cell is identified by the vector  $\mathbf{R}_p = \sum_i n_i \mathbf{a}_i$ , with  $n_i$  integers between 0 and  $N_i - 1$ . The BvK supercell contains  $N_p = N_1 \times N_2 \times N_3$  unit cells. The primitive vectors of the reciprocal lattice are denoted by  $\mathbf{b}_j$  and fulfill the duality condition  $\mathbf{a}_i \cdot \mathbf{b}_j = 2\pi\delta_{ij}$ . We consider Bloch wave vectors  $\mathbf{q}$  belonging to a uniform grid in one unit cell of the reciprocal lattice  $\mathbf{q} = \sum_j (m_j/N_j)\mathbf{b}_j$ , with  $m_j$  being integers between 0 and  $N_j - 1$ . This grid contains the same number of  $\mathbf{q}$  vectors as the number of unit cells in the BvK supercell. From these definitions the standard sum rules follow:

$$\begin{aligned} \sum_{\mathbf{q}} \exp(i\mathbf{q} \cdot \mathbf{R}_p) &= N_p \delta_{p0}, \\ \sum_p \exp(i\mathbf{q} \cdot \mathbf{R}_p) &= N_p \delta_{\mathbf{q}0}. \end{aligned} \quad (\text{A1})$$

If  $\mathbf{G}$  is a reciprocal lattice vector, the replacement of any of the  $\mathbf{q}$  vectors by  $\mathbf{q} + \mathbf{G}$  in these expressions and in all expressions presented in this article is inconsequential, since  $\exp(i\mathbf{G} \cdot \mathbf{R}_p) = 1$ . Similarly, any replacement of  $\mathbf{R}_p$  by  $\mathbf{R}_p + \mathbf{T}$ , where  $\mathbf{T}$  is a lattice vector of the BvK supercell is inconsequential. Owing to these properties we are at liberty to replace the  $\mathbf{q}$  grid with a Wigner-Seitz grid, i.e., the first Brillouin zone, and the supercell with a Wigner-Seitz supercell. These choices are useful for practical calculations in order to exploit the symmetry operations of the crystal and to truncate the interatomic force constants, given by Eq. (13), outside a Wigner-Seitz supercell.

## APPENDIX B: LADDER OPERATORS IN EXTENDED SYSTEMS

In this Appendix we describe the construction of the phonon ladder operators  $\hat{a}_{\mathbf{q}\nu}/\hat{a}_{\mathbf{q}\nu}^\dagger$  and derive the phonon Hamiltonian given by Eq. (22). We show how the definition of the ladder operators depends on the behavior of the wave vector  $\mathbf{q}$  under inversion.

The normal modes introduced in Eq. (15) can be used to define a linear coordinate transformation of the ionic displacements as follows:

$$z_{\mathbf{q}\nu} = N_p^{-1/2} \sum_{\kappa\alpha p} e^{-i\mathbf{q}\cdot\mathbf{R}_p} (M_\kappa/M_0)^{1/2} e_{\kappa\alpha,\nu}^*(\mathbf{q}) \Delta\tau_{\kappa\alpha p}. \quad (\text{B1})$$

Here  $z_{\mathbf{q}\nu}$  is referred to as a ‘‘complex normal coordinate’’ (Brüesch, 1982). The exponential and the masses in Eq. (B1) are chosen so as to obtain Eq. (22) starting from Eq. (12). Since there are  $3MN_p$  degrees of freedom, and since the complex normal coordinates correspond to  $2 \times 3MN_p$  real variables, this coordinate transformation carries some redundancy. Indeed by combining Eqs. (18) and (B1) it is seen that

$$z_{-\mathbf{q}\nu} = z_{\mathbf{q}\nu}^*. \quad (\text{B2})$$

The inverse relation of Eq. (B1) is

$$\Delta\tau_{\kappa\alpha p} = N_p^{-1/2} (M_0/M_\kappa)^{1/2} \sum_{\mathbf{q}} e^{i\mathbf{q}\cdot\mathbf{R}_p} e_{\kappa\alpha,\nu}(\mathbf{q}) z_{\mathbf{q}\nu}. \quad (\text{B3})$$

The right-hand side is real valued after Eqs. (18) and (B2). In preparation for the transition to a quantum description of lattice vibrations, it is useful to identify  $3MN_p$  independent normal coordinates. This can be done by partitioning the grid of  $\mathbf{q}$  vectors into three sets. We call  $\mathcal{A}$  the set of vectors which are invariant under inversion, that is,  $-\mathbf{q} + \mathbf{G} = \mathbf{q}$  for some reciprocal lattice vector  $\mathbf{G}$  (including  $|\mathbf{G}| = 0$ ). The center of the Brillouin zone and the centers of its faces belong to this set. The remaining vectors can be separated further into  $\mathcal{B}$  and  $\mathcal{C}$ , in such a way that all the vectors in  $\mathcal{C}$  are obtained from those in  $\mathcal{B}$  by inversion (modulo a reciprocal lattice vector). After defining  $z_{\mathbf{q}\nu} = x_{\mathbf{q}\nu} + iy_{\mathbf{q}\nu}$ , Eq. (B3) can be rewritten as

$$\Delta\tau_{\kappa\alpha p} = N_p^{-1/2} (M_0/M_\kappa)^{1/2} \left[ \sum_{\mathbf{q} \in \mathcal{A}, \nu} e_{\kappa\alpha,\nu}(\mathbf{q}) x_{\mathbf{q}\nu} + 2\text{Re} \sum_{\mathbf{q} \in \mathcal{B}, \nu} e^{i\mathbf{q}\cdot\mathbf{R}_p} e_{\kappa\alpha,\nu}(\mathbf{q}) (x_{\mathbf{q}\nu} + iy_{\mathbf{q}\nu}) \right]. \quad (\text{B4})$$

The  $\mathbf{q}$  vectors of the set  $\mathcal{C}$  have been grouped together with those in  $\mathcal{B}$  by taking the real part in the second line. It can be verified that in this expression there are exactly  $3MN_p$  real coordinates; therefore we can choose the  $x_{\mathbf{q}\nu}$  for  $\mathbf{q}$  in  $\mathcal{A}$  and the pairs  $x_{\mathbf{q}\nu}, y_{\mathbf{q}\nu}$  for  $\mathbf{q}$  in  $\mathcal{B}$  as the independent variables. These variables are referred to as ‘‘real normal coordinates’’ (Brüesch, 1982).

Using Eqs. (12)–(18), (A1), and (B4) the nuclear Hamiltonian can be written in terms of  $3MN_p$  independent harmonic oscillators in the real normal coordinates:

$$\hat{H}_p = \frac{1}{2} \sum_{\mathbf{q} \in \mathcal{B}, \nu} \hbar\omega_{\mathbf{q}\nu} (-\partial^2/\partial\tilde{x}_{\mathbf{q}\nu}^2 - \partial^2/\partial\tilde{y}_{\mathbf{q}\nu}^2 + \tilde{x}_{\mathbf{q}\nu}^2 + \tilde{y}_{\mathbf{q}\nu}^2) + \frac{1}{2} \sum_{\mathbf{q} \in \mathcal{A}, \nu} \hbar\omega_{\mathbf{q}\nu} (-\partial^2/\partial\tilde{x}_{\mathbf{q}\nu}^2 + \tilde{x}_{\mathbf{q}\nu}^2), \quad (\text{B5})$$

where for ease of notation we performed the scaling

$$\tilde{x}_{\mathbf{q}\nu} = x_{\mathbf{q}\nu}/2l_{\mathbf{q}\nu} \quad \text{for } \mathbf{q} \text{ in } \mathcal{A}, \quad (\text{B6})$$

$$\tilde{x}_{\mathbf{q}\nu} = x_{\mathbf{q}\nu}/l_{\mathbf{q}\nu}, \quad \tilde{y}_{\mathbf{q}\nu} = y_{\mathbf{q}\nu}/l_{\mathbf{q}\nu} \quad \text{for } \mathbf{q} \text{ in } \mathcal{B}, \quad (\text{B7})$$

with  $l_{\mathbf{q}\nu}$  the zero-point displacement amplitude of Eq. (21). In the case of  $|\mathbf{q}| = 0$  there are three normal modes for which  $\omega_{\mathbf{q}\nu} = 0$ , and the corresponding potential terms  $\tilde{x}_{\mathbf{q}\nu}^2$  must be removed from Eq. (B5).

The eigenstates of Eq. (B5) are found by introducing the real ladder operators for each normal coordinate (Cohen-Tannoudji, Diu, and Laloe, 1977):

$$\hat{a}_{\mathbf{q}\nu,x} = (\tilde{x}_{\mathbf{q}\nu} + \partial/\partial\tilde{x}_{\mathbf{q}\nu})/\sqrt{2}, \quad (\text{B8})$$

and similarly for  $\hat{a}_{\mathbf{q}\nu,y}$ . With these definitions Eq. (B5) becomes

$$\hat{H}_p = \sum_{\mathbf{q} \in \mathcal{B}, \nu} \hbar\omega_{\mathbf{q}\nu} (\hat{a}_{\mathbf{q}\nu,x}^\dagger \hat{a}_{\mathbf{q}\nu,x} + \hat{a}_{\mathbf{q}\nu,y}^\dagger \hat{a}_{\mathbf{q}\nu,y} + 1) + \sum_{\mathbf{q} \in \mathcal{A}, \nu} \hbar\omega_{\mathbf{q}\nu} (\hat{a}_{\mathbf{q}\nu,x}^\dagger \hat{a}_{\mathbf{q}\nu,x} + 1/2). \quad (\text{B9})$$

The eigenstates of this Hamiltonian are products of simple harmonic oscillators (Merzbacher, 1998), and the ground state is

$$\chi_0(\{\tau_{\kappa p}\}) = A e^{-1/2(\sum_{\mathbf{q} \in \mathcal{A}, \nu} \tilde{x}_{\mathbf{q}\nu}^2 + \sum_{\mathbf{q} \in \mathcal{B}, \nu} \tilde{x}_{\mathbf{q}\nu}^2 + \tilde{y}_{\mathbf{q}\nu}^2)}, \quad (\text{B10})$$

with  $A$  a normalization constant. The relations between the positions  $\tau_{\kappa p}$  and the normal coordinates  $\tilde{x}_{\mathbf{q}\nu}, \tilde{y}_{\mathbf{q}\nu}$  are given by Eqs. (B1), (B6), (B7), and (21).

The eigenstates of  $\hat{H}_p$  can be generated by applying  $\hat{a}_{\mathbf{q}\nu,x}^\dagger$  and  $\hat{a}_{\mathbf{q}\nu,y}^\dagger$  to the ground state  $\chi_0$ . However this approach is not entirely satisfactory, since we cannot assign separate quantum numbers to modes with wave vectors  $\mathbf{q}$  or  $-\mathbf{q}$ . In order to avoid this inconvenience we observe that, for each normal mode, the first line of Eq. (B5) defines an effective isotropic two-dimensional harmonic oscillator. The degenerate eigenstates of this oscillator can be combined to form eigenstates of the angular momentum. This leads to right and left circular quanta with the same energy and definite angular momentum (Cohen-Tannoudji, Diu, and Laloe, 1977). This analogy motivates the consideration of the following linear combinations, for  $\mathbf{q}$  in  $\mathcal{B}$ :

$$\hat{a}_{\mathbf{q}\nu}^+ = (\hat{a}_{\mathbf{q}\nu,x} + i\hat{a}_{\mathbf{q}\nu,y})/\sqrt{2}, \quad (\text{B11})$$

$$\hat{a}_{\mathbf{q}\nu}^- = (\hat{a}_{\mathbf{q}\nu,x} - i\hat{a}_{\mathbf{q}\nu,y})/\sqrt{2}. \quad (\text{B12})$$

Since both  $\hat{a}_{\mathbf{q}\nu,x}$  and  $\hat{a}_{\mathbf{q}\nu,y}$  lower the energy of an eigenstate by the same quantum of energy  $\hbar\omega_{\mathbf{q}\nu}$ , the resulting states are degenerate and their linear combinations are also eigenstates for the same eigenvalue. As a consequence we can generate all the eigenstates of the Hamiltonian  $\hat{H}_p$  by acting on the ground state  $\chi_0$  with the creation operators  $\hat{a}_{\mathbf{q}\nu}^{+\dagger}$  and  $\hat{a}_{\mathbf{q}\nu}^{-\dagger}$ . In this reasoning the wave vectors  $\mathbf{q}$  belong to  $\mathcal{B}$ ; if we now consider Eqs. (B2), (B11), and (B12), we see that formally we also have  $\hat{a}_{\mathbf{q}\nu}^- = \hat{a}_{-\mathbf{q}\nu}^+$ . Therefore it is natural to associate  $\hat{a}_{\mathbf{q}\nu}^-$  to phonons propagating along the direction  $-\mathbf{q}$ .

These observations suggest replacing the real ladder operators of Eq. (B8) by the complex ladder operators  $\hat{a}_{\mathbf{q}\nu}^+$  and  $\hat{a}_{-\mathbf{q}\nu}^-$  for  $\mathbf{q}$  in  $\mathcal{B}$  and  $\mathcal{C}$ , respectively. In the case of  $\mathbf{q}$  in  $\mathcal{A}$  we keep the real operators  $\hat{a}_{\mathbf{q}\nu,x}$ . These definitions can be turned into the following compact expressions:

$$\hat{a}_{\mathbf{q}\nu} = \hat{a}_{\mathbf{q}\nu,x} \quad \text{for } \mathbf{q} \text{ in } \mathcal{A}, \quad (\text{B13})$$

$$\hat{a}_{\mathbf{q}\nu} = (\hat{a}_{\mathbf{q}\nu,x} + i\hat{a}_{\mathbf{q}\nu,y})/\sqrt{2} \quad \text{for } \mathbf{q} \text{ in } \mathcal{B}, \mathcal{C}. \quad (\text{B14})$$

Using these operators the nuclear Hamiltonian of Eq. (B5) takes the well-known form given by Eq. (22). Any eigenstate of  $\hat{H}_p$  can now be generated as  $\prod_{\mathbf{q}\nu} (n_{\mathbf{q}\nu}!)^{-1/2} (\hat{a}_{\mathbf{q}\nu}^\dagger)^{n_{\mathbf{q}\nu}} \chi_0$ . In this form we see that it is possible to independently assign a number of phonons  $n_{\mathbf{q}\nu}$  to each wave vector  $\mathbf{q}$  and each mode  $\nu$ . Using Eqs. (B6)–(B8), (B13), and (B14) we also have the basic identity

$$z_{\mathbf{q}\nu} = l_{\mathbf{q}\nu} (\hat{a}_{\mathbf{q}\nu} + \hat{a}_{-\mathbf{q}\nu}^\dagger). \quad (\text{B15})$$

By combining this last expression with Eq. (B3) we obtain Eq. (20).

## REFERENCES

- Abrikosov, A. A., L. P. Gor'kov, and I. E. Dzyaloshinski, 1975, *Methods of Quantum Field Theory in Statistical Physics* (Dover, New York).
- Adler, S. L., 1962, "Quantum theory of the dielectric constant in real solids," *Phys. Rev.* **126**, 413–420.
- Agapito, L. A., A. Ferretti, A. Calzolari, S. Curtarolo, and M. Buongiorno Nardelli, 2013, "Effective and accurate representation of extended Bloch states on finite Hilbert spaces," *Phys. Rev. B* **88**, 165127.
- Akashi, R., and R. Arita, 2013a, "Development of density-functional theory for a plasmon-assisted superconducting state: Application to lithium under high pressures," *Phys. Rev. Lett.* **111**, 057006.
- Akashi, R., and R. Arita, 2013b, "Nonempirical study of superconductivity in alkali-doped fullerides based on density functional theory for superconductors," *Phys. Rev. B* **88**, 054510.
- Akashi, R., M. Kawamura, S. Tsuneyuki, Y. Nomura, and R. Arita, 2015, "First-principles study of the pressure and crystal-structure dependences of the superconducting transition temperature in compressed sulfur hydrides," *Phys. Rev. B* **91**, 224513.
- Akashi, R., K. Nakamura, R. Arita, and M. Imada, 2012, "High-temperature superconductivity in layered nitrides  $\beta$ -Li<sub>x</sub>MNCl ( $M = \text{Ti, Zr, Hf}$ ): Insights from density functional theory for superconductors," *Phys. Rev. B* **86**, 054513.
- Albers, R. C., L. Bohlin, M. Roy, and J. W. Wilkins, 1976, "Normal and umklapp phonon decay rates due to phonon-phonon and electron-phonon scattering in potassium at low temperatures," *Phys. Rev. B* **13**, 768–786.
- Alexandrov, A. S., 2008, *Polarons in advanced materials* (Springer, New York).
- Alexandrov, A. S., and J. T. Devreese, 2010, *Advances in polaron physics* (Springer, New York).
- Allen, P. B., 1972a, "Electron-phonon coupling and pseudopotentials: The mass enhancements of the noble metals," *Phys. Rev. B* **5**, 3857–3862.
- Allen, P. B., 1972b, "Neutron spectroscopy of superconductors," *Phys. Rev. B* **6**, 2577–2579.
- Allen, P. B., 1976, "Fermi-surface harmonics: A general method for nonspherical problems. application to Boltzmann and Eliashberg equations," *Phys. Rev. B* **13**, 1416–1427.
- Allen, P. B., 1978, "Solids with thermal or static disorder. I. one-electron properties," *Phys. Rev. B* **18**, 5217–5224.
- Allen, P. B., and M. Cardona, 1981, "Theory of the temperature dependence of the direct gap of germanium," *Phys. Rev. B* **23**, 1495–1505.
- Allen, P. B., and M. Cardona, 1983, "Temperature dependence of the direct gap of Si and Ge," *Phys. Rev. B* **27**, 4760–4769.
- Allen, P. B., and M. L. Cohen, 1969, "Pseudopotential calculation of the mass enhancement and superconducting transition temperature of simple metals," *Phys. Rev.* **187**, 525–538.
- Allen, P. B., and M. L. Cohen, 1970, "Calculation of the temperature dependence of the electron-phonon mass enhancement," *Phys. Rev. B* **1**, 1329–1336.
- Allen, P. B., M. L. Cohen, L. M. Falicov, and R. V. Kasowski, 1968, "Superconductivity and band structure from a single pseudopotential: Zinc and cadmium," *Phys. Rev. Lett.* **21**, 1794–1796.
- Allen, P. B., and R. C. Dynes, 1975, "Transition temperature of strongly coupled superconductors reanalyzed," *Phys. Rev. B* **12**, 905–922.
- Allen, P. B., and V. Heine, 1976, "Theory of the temperature dependence of electronic band structures," *J. Phys. C* **9**, 2305.
- Allen, P. B., and J. C. K. Hui, 1980, "Thermodynamics of solids: Corrections from electron-phonon interactions," *Z. Phys. B* **37**, 33–38.
- Allen, P. B., and M. J. G. Lee, 1972, "Phase-shift pseudopotentials and the electron-phonon interaction: Theory and results for alkali metals," *Phys. Rev. B* **5**, 3848–3857.
- Allen, P. B., and B. Mitrovic, 1982, *Theory of Superconducting  $T_c$* , edited by F. Seitz, D. Turnbull, and H. Ehrenreich, Solid State Physics, Vol. 37 (Academic, New York).
- Allen, P. B., and J. P. Nery, 2017, "Low temperature semiconductor band-gap thermal shifts:  $T^4$  shifts from ordinary acoustic and  $T^2$  from piezoacoustic coupling," *Phys. Rev. B* **95**, 035211.
- Animalu, A. O. E., and V. Heine, 1965, "The screened model potential for 25 elements," *Philos. Mag.* **12**, 1249–1270.
- Antončík, E., 1955, "On the theory of temperature shift of the absorption curve in non-polar crystals," *Czech. J. Phys.* **5**, 449–461.
- Antonius, G., S. Poncé, P. Boulanger, M. Côté, and X. Gonze, 2014, "Many-body effects on the zero-point renormalization of the band structure," *Phys. Rev. Lett.* **112**, 215501.
- Antonius, G., S. Poncé, E. Lantagne-Hurtubise, G. Auclair, X. Gonze, and M. Côté, 2015, "Dynamical and anharmonic effects on the electron-phonon coupling and the zero-point renormalization of the electronic structure," *Phys. Rev. B* **92**, 085137.
- Aperis, A., P. Maldonado, and P. M. Oppeneer, 2015, "Ab initio theory of magnetic-field-induced odd-frequency two-band superconductivity in  $\text{MgB}_2$ ," *Phys. Rev. B* **92**, 054516.
- Aryasetiawan, F., and S. Biermann, 2008, "Generalized Hedin's equations for quantum many-body systems with spin-dependent interactions," *Phys. Rev. Lett.* **100**, 116402.
- Aryasetiawan, F., and O. Gunnarsson, 1998, "The GW method," *Rep. Prog. Phys.* **61**, 237.
- Aryasetiawan, F., L. Hedin, and K. Karlsson, 1996, "Multiple plasmon satellites in Na and Al spectral functions from ab initio cumulant expansion," *Phys. Rev. Lett.* **77**, 2268–2271.
- Ashcroft, N. W., and N. D. Mermin, 1976, *Solid State Physics* (Harcourt, New York).
- Ashcroft, N. W., and J. W. Wilkins, 1965, "Low temperature electronic specific heat of simple metals," *Phys. Lett.* **14**, 285–287.

- Audouze, C., F. Jollet, M. Torrent, and X. Gonze, 2006, "Projector augmented-wave approach to density-functional perturbation theory," *Phys. Rev. B* **73**, 235101.
- Bachelet, G. B., D. R. Hamann, and M. Schlüter, 1982, "Pseudopotentials that work: From H to Pu," *Phys. Rev. B* **26**, 4199–4228.
- Bardeen, J., 1937, "Conductivity of monovalent metals," *Phys. Rev.* **52**, 688–697.
- Bardeen, J., and D. Pines, 1955, "Electron-phonon interaction in metals," *Phys. Rev.* **99**, 1140–1150.
- Bardeen, J., and W. Shockley, 1950, "Deformation potentials and mobilities in non-polar crystals," *Phys. Rev.* **80**, 72–80.
- Baroni, S., S. de Gironcoli, A. Dal Corso, and P. Giannozzi, 2001, "Phonons and related crystal properties from density-functional perturbation theory," *Rev. Mod. Phys.* **73**, 515–562.
- Baroni, S., P. Giannozzi, and A. Testa, 1987, "Green's-function approach to linear response in solids," *Phys. Rev. Lett.* **58**, 1861–1864.
- Bassani, G. F., and G. P. Parravicini, 1975, *Electronic states and optical transitions in solids* (Pergamon, New York).
- Bauer, R., A. Schmid, P. Pavone, and D. Strauch, 1998, "Electron-phonon coupling in the metallic elements Al, Au, Na, and Nb: A first-principles study," *Phys. Rev. B* **57**, 11276–11282.
- Baym, G., 1961, "Field-theoretic approach to the properties of the solid state," *Ann. Phys. (N.Y.)* **14**, 1–42.
- Becker, M., and H. Y. Fan, 1949, "Optical properties of semiconductors. III. Infra-red transmission of silicon," *Phys. Rev.* **76**, 1531–1532.
- Berciu, M., 2006, "Green's function of a dressed particle," *Phys. Rev. Lett.* **97**, 036402.
- Berger, E., P. Valášek, and W. von der Linden, 1995, "Two-dimensional Hubbard-Holstein model," *Phys. Rev. B* **52**, 4806–4814.
- Bernardi, M., D. Vigil-Fowler, J. Lischner, J. B. Neaton, and S. G. Louie, 2014, "Ab initio study of hot carriers in the first picosecond after sunlight absorption in silicon," *Phys. Rev. Lett.* **112**, 257402.
- Bernardi, M., D. Vigil-Fowler, C. S. Ong, J. B. Neaton, and S. G. Louie, 2015, "Ab initio study of hot electrons in GaAs," *Proc. Natl. Acad. Sci. U.S.A.* **112**, 5291–5296.
- Bersier, C., A. Floris, A. Sanna, G. Profeta, A. Continenza, E. K. U. Gross, and S. Massidda, 2009, "Electronic, vibrational, and superconducting properties of CaBeSi: First-principles calculations," *Phys. Rev. B* **79**, 104503.
- Bhosale, J., A. K. Ramdas, A. Burger, A. Muñoz, A. H. Romero, M. Cardona, R. Lauck, and R. K. Kremer, 2012, "Temperature dependence of band gaps in semiconductors: Electron-phonon interaction," *Phys. Rev. B* **86**, 195208.
- Bianchi, M., E. D. L. Rienks, S. Lizzit, A. Baraldi, R. Balog, L. Hørnækær, and Ph. Hofmann, 2010, "Electron-phonon coupling in potassium-doped graphene: Angle-resolved photoemission spectroscopy," *Phys. Rev. B* **81**, 041403.
- Bloch, F., 1929, "Über die Quantenmechanik der Elektronen in Kristallgittern," *Z. Phys.* **52**, 555–600.
- Blöchl, P. E., 1994, "Projector augmented-wave method," *Phys. Rev. B* **50**, 17953–17979.
- Bogoliubov, N. N., 1958, "A new method in the theory of superconductivity. I," *Sov. Phys. JETP* **7**, 41–46 [<http://www.jetp.ac.ru/cgi-bin/e/index/e/7/1/p41?a=list>].
- Bonini, N., M. Lazzeri, N. Marzari, and F. Mauri, 2007, "Phonon anharmonicities in graphite and graphene," *Phys. Rev. Lett.* **99**, 176802.
- Born, M., and K. Huang, 1954, *Dynamical Theory of Crystal Lattices* (Clarendon, Oxford).
- Born, M., and R. Oppenheimer, 1927, "Zur Quantentheorie der Molekeln," *Ann. Phys. (Berlin)* **389**, 457–484.
- Borysenko, K. M., J. T. Mullen, E. A. Barry, S. Paul, Y. G. Semenov, J. M. Zavada, M. Buongiorno Nardelli, and K. W. Kim, 2010, "First-principles analysis of electron-phonon interactions in graphene," *Phys. Rev. B* **81**, 121412.
- Bostwick, A., T. Ohta, T. Seyller, K. Horn, and E. Rotenberg, 2007, "Quasiparticle dynamics in graphene," *Nat. Phys.* **3**, 36–40.
- Botti, S., and M. A. L. Marques, 2013, "Strong renormalization of the electronic band gap due to lattice polarization in the GW formalism," *Phys. Rev. Lett.* **110**, 226404.
- Broido, D. A., M. Malorny, G. Birner, N. Mingo, and D. A. Stewart, 2007, "Intrinsic lattice thermal conductivity of semiconductors from first principles," *Appl. Phys. Lett.* **91**, 231922.
- Brouder, C., G. Panati, M. Calandra, C. Mourougane, and N. Marzari, 2007, "Exponential localization of Wannier functions in insulators," *Phys. Rev. Lett.* **98**, 046402.
- Briesch, P., 1982, *Phonons, Theory and Experiments*, Vol. 1 (Springer-Verlag, Berlin).
- Bungaro, C., K. Rapcewicz, and J. Bernholc, 2000, "Ab initio phonon dispersions of wurtzite AlN, GaN, and InN," *Phys. Rev. B* **61**, 6720–6725.
- Butler, W. H., F. J. Pinski, and P. B. Allen, 1979, "Phonon linewidths and electron-phonon interaction in Nb," *Phys. Rev. B* **19**, 3708–3721.
- Calandra, M., and F. Mauri, 2007, "Electron-phonon coupling and electron self-energy in electron-doped graphene: Calculation of angular-resolved photoemission spectra," *Phys. Rev. B* **76**, 205411.
- Calandra, M., G. Profeta, and F. Mauri, 2010, "Adiabatic and nonadiabatic phonon dispersion in a Wannier function approach," *Phys. Rev. B* **82**, 165111.
- Campbell, D. K., T. A. DeGrand, and S. Mazumdar, 1984, "Soliton energetics in Peierls-Hubbard models," *Phys. Rev. Lett.* **52**, 1717–1720.
- Cancellieri, C., A. S. Mishchenko, U. Aschauer, A. Filippetti, C. Faber, O. S. Barišić, V. A. Rogalev, T. Schmitt, N. Nagaosa, and V. N. Strocov, 2016, "Polaronic metal state at the LaAlO<sub>3</sub>/SrTiO<sub>3</sub> interface," *Nat. Commun.* **7**, 10386.
- Cannuccia, E., and A. Marini, 2011, "Effect of the quantum zero-point atomic motion on the optical and electronic properties of diamond and trans-polyacetylene," *Phys. Rev. Lett.* **107**, 255501.
- Cannuccia, E., and A. Marini, 2012, "Zero point motion effect on the electronic properties of diamond, trans-polyacetylene and polyethylene," *Eur. Phys. J. B* **85**, 320.
- Cannuccia, E., and A. Marini, 2013, "Ab initio study of the effects induced by the electron-phonon scattering in carbon based nanostructures," [arXiv:1304.0072](https://arxiv.org/abs/1304.0072).
- Cao, K., F. Giustino, and P. G. Radaelli, 2015, "Theory of electromagnons in CuO," *Phys. Rev. Lett.* **114**, 197201.
- Capaz, R. B., C. D. Spataru, P. Tangney, M. L. Cohen, and S. G. Louie, 2005, "Temperature dependence of the band gap of semiconducting carbon nanotubes," *Phys. Rev. Lett.* **94**, 036801.
- Carbotte, J. P., and R. C. Dynes, 1967, "Resistivity of solid aluminum," *Phys. Lett. A* **25**, 532–533.
- Cardona, M., 2001, "Renormalization of the optical response of semiconductors by electron-phonon interaction," *Phys. Status Solidi A* **188**, 1209–1232.
- Cardona, M., 2005, "Electron-phonon interaction in tetrahedral semiconductors," *Solid State Commun.* **133**, 3–18.
- Cardona, M., and M. L. W. Thewalt, 2005, "Isotope effects on the optical spectra of semiconductors," *Rev. Mod. Phys.* **77**, 1173–1224.
- Caruso, F., and F. Giustino, 2015, "Spectral fingerprints of electron-plasmon coupling," *Phys. Rev. B* **92**, 045123.
- Caruso, F., H. Lambert, and F. Giustino, 2015, "Band structures of plasmonic polarons," *Phys. Rev. Lett.* **114**, 146404.

- Caudal, N., A. M. Saitta, M. Lazzeri, and F. Mauri, 2007, “Kohn anomalies and nonadiabaticity in doped carbon nanotubes,” *Phys. Rev. B* **75**, 115423.
- Ceperley, D. M., and B. J. Alder, 1980, “Ground state of the electron gas by a stochastic method,” *Phys. Rev. Lett.* **45**, 566–569.
- Chakraborty, B., and P. B. Allen, 1978, “Theory of temperature dependence of optical properties of solids,” *J. Phys. C* **11**, L9.
- Chan, K. T., J. D. Sau, P. Zhang, and M. L. Cohen, 2007, “*Ab initio* calculations of phonon splitting in antiferromagnetic  $\text{ZnCr}_2\text{O}_4$ ,” *Phys. Rev. B* **75**, 054304.
- Chang, K. J., and M. L. Cohen, 1986, “Electron-phonon interactions and superconductivity in Si, Ge, and Sn,” *Phys. Rev. B* **34**, 4552–4557.
- Chang, K. J., M. M. Dacorogna, M. L. Cohen, J. M. Mignot, G. Chouteau, and G. Martinez, 1985, “Superconductivity in high-pressure metallic phases of Si,” *Phys. Rev. Lett.* **54**, 2375–2378.
- Chen, C., J. Avila, E. Frantzeskakis, A. Levy, and M. C. Asensio, 2015, “Observation of a two-dimensional liquid of Fröhlich polarons at the bare  $\text{SrTiO}_3$  surface,” *Nat. Commun.* **6**, 8585.
- Choi, H. J., M. L. Cohen, and S. G. Louie, 2003, “Anisotropic Eliashberg theory of  $\text{MgB}_2$ :  $T_c$ , isotope effects, superconducting energy gaps, quasiparticles, and specific heat,” *Physica C (Amsterdam)* **385**, 66–74.
- Choi, H. J., S. G. Louie, and M. L. Cohen, 2009a, “Anisotropic Eliashberg theory for superconductivity in compressed and doped  $\text{MgB}_2$ ,” *Phys. Rev. B* **79**, 094518.
- Choi, H. J., S. G. Louie, and M. L. Cohen, 2009b, “Prediction of superconducting properties of  $\text{CaB}_2$  using anisotropic Eliashberg theory,” *Phys. Rev. B* **80**, 064503.
- Choi, H. J., D. Roundy, H. Sun, M. L. Cohen, and S. G. Louie, 2002a, “First-principles calculation of the superconducting transition in  $\text{MgB}_2$  within the anisotropic Eliashberg formalism,” *Phys. Rev. B* **66**, 020513.
- Choi, H. J., D. Roundy, H. Sun, M. L. Cohen, and S. G. Louie, 2002b, “The origin of the anomalous superconducting properties of  $\text{MgB}_2$ ,” *Nature (London)* **418**, 758–760.
- Cohen, M. H., and V. Heine, 1961, “Cancellation of kinetic and potential energy in atoms, molecules, and solids,” *Phys. Rev.* **122**, 1821–1826.
- Cohen, M. L., and T. K. Bergstresser, 1966, “Band structures and pseudopotential form factors for fourteen semiconductors of the diamond and zinc-blende structures,” *Phys. Rev.* **141**, 789–796.
- Cohen-Tannoudji, C., B. Diu, and F. Laloe, 1977, *Quantum Mechanics*, Vol. 1 (Wiley, New York).
- Cowley, R. A., 1963, “The lattice dynamics of an anharmonic crystal,” *Adv. Phys.* **12**, 421–480.
- Cudazzo, P., G. Profeta, A. Sanna, A. Floris, A. Continenza, S. Massidda, and E. K. U. Gross, 2008, “*Ab initio* description of high-temperature superconductivity in dense molecular hydrogen,” *Phys. Rev. Lett.* **100**, 257001.
- Cudazzo, P., G. Profeta, A. Sanna, A. Floris, A. Continenza, S. Massidda, and E. K. U. Gross, 2010, “Electron-phonon interaction and superconductivity in metallic molecular hydrogen. II. superconductivity under pressure,” *Phys. Rev. B* **81**, 134506.
- Dacorogna, M. M., K. J. Chang, and M. L. Cohen, 1985, “Pressure increase of the electron-phonon interaction in superconducting hexagonal silicon,” *Phys. Rev. B* **32**, 1853–1855.
- Dacorogna, M. M., M. L. Cohen, and P. K. Lam, 1985, “Self-consistent calculation of the  $q$  dependence of the electron-phonon coupling in aluminum,” *Phys. Rev. Lett.* **55**, 837–840.
- Dal Corso, A., 2007, “Density functional perturbation theory for lattice dynamics with fully relativistic ultrasoft pseudopotentials: Application to fcc-Pt and fcc-Au,” *Phys. Rev. B* **76**, 054308.
- Dal Corso, A., 2008, “*Ab initio* phonon dispersions of face centered cubic Pb: effects of spin-orbit coupling,” *J. Phys. Condens. Matter* **20**, 445202.
- Dal Corso, A., 2013, “*Ab initio* phonon dispersions of transition and noble metals: effects of the exchange and correlation functional,” *J. Phys. Condens. Matter* **25**, 145401.
- Dal Corso, A., S. Baroni, R. Resta, and S. de Gironcoli, 1993, “*Ab initio* calculation of phonon dispersions in II-VI semiconductors,” *Phys. Rev. B* **47**, 3588–3592.
- Dal Corso, A., and S. de Gironcoli, 2000, “*Ab initio* phonon dispersions of Fe and Ni,” *Phys. Rev. B* **62**, 273–277.
- Dal Corso, A., A. Pasquarello, and A. Baldereschi, 1997, “Density-functional perturbation theory for lattice dynamics with ultrasoft pseudopotentials,” *Phys. Rev. B* **56**, R11369–R11372.
- Damascelli, A., Z. Hussain, and Z.-X. Shen, 2003, “Angle-resolved photoemission studies of the cuprate superconductors,” *Rev. Mod. Phys.* **75**, 473–541.
- Debernardi, A., S. Baroni, and E. Molinari, 1995, “Anharmonic phonon lifetimes in semiconductors from density-functional perturbation theory,” *Phys. Rev. Lett.* **75**, 1819–1822.
- de Gironcoli, S., 1995, “Lattice dynamics of metals from density-functional perturbation theory,” *Phys. Rev. B* **51**, 6773–6776.
- Deinzer, G., G. Birner, and D. Strauch, 2003, “*Ab initio* calculation of the linewidth of various phonon modes in germanium and silicon,” *Phys. Rev. B* **67**, 144304.
- Della Sala, F., R. Rousseau, A. Görling, and D. Marx, 2004, “Quantum and thermal fluctuation effects on the photoabsorption spectra of clusters,” *Phys. Rev. Lett.* **92**, 183401.
- Díaz-Sánchez, L. E., A. H. Romero, and X. Gonze, 2007, “Phonon band structure and interatomic force constants for bismuth: Crucial role of spin-orbit interaction,” *Phys. Rev. B* **76**, 104302.
- D’Innocenzo, V., G. Grancini, M. J. P. Alcocer, A. R. S. Kandada, S. D. Stranks, M. M. Lee, G. Lanzani, H. J. Snaith, and A. Petrozza, 2014, “Excitons versus free charges in organo-lead tri-halide perovskites,” *Nat. Commun.* **5**, 3586.
- Dumke, W. P., 1956, “Deformation potential theory for  $n$ -type Ge,” *Phys. Rev.* **101**, 531–536.
- Dynes, R. C., and J. P. Carbotte, 1968, “Phonon-limited resistivity of some simple metals,” *Phys. Rev.* **175**, 913–919.
- Eiguren, A., and C. Ambrosch-Draxl, 2008a, “Complex quasiparticle band structure induced by electron-phonon interaction: Band splitting in the  $1 \times 1\text{H}/\text{W}(110)$  surface,” *Phys. Rev. Lett.* **101**, 036402.
- Eiguren, A., and C. Ambrosch-Draxl, 2008b, “Wannier interpolation scheme for phonon-induced potentials: Application to bulk  $\text{MgB}_2$ , W, and the  $1 \times 1$  H-covered  $\text{W}(110)$  surface,” *Phys. Rev. B* **78**, 045124.
- Eiguren, A., C. Ambrosch-Draxl, and P. M. Echenique, 2009, “Self-consistently renormalized quasiparticles under the electron-phonon interaction,” *Phys. Rev. B* **79**, 245103.
- Eiguren, A., S. de Gironcoli, E. V. Chulkov, P. M. Echenique, and E. Tosatti, 2003, “Electron-phonon interaction at the  $\text{Be}(0001)$  surface,” *Phys. Rev. Lett.* **91**, 166803.
- Eiguren, A., and I. G. Gurtubay, 2014, “Helmholtz Fermi surface harmonics: an efficient approach for treating anisotropic problems involving Fermi surface integrals,” *New J. Phys.* **16**, 063014.
- Eiguren, A., B. Hellsing, E. V. Chulkov, and P. M. Echenique, 2003, “Phonon-mediated decay of metal surface states,” *Phys. Rev. B* **67**, 235423.
- Eiguren, A., B. Hellsing, F. Reinert, G. Nicolay, E. V. Chulkov, V. M. Silkin, S. Hüfner, and P. M. Echenique, 2002, “Role of bulk and surface phonons in the decay of metal surface states,” *Phys. Rev. Lett.* **88**, 066805.



- Eliashberg, G. M., 1960, "Interactions between electrons and lattice vibrations in a superconductor," *Sov. Phys. JETP* **11**, 696–702 [<http://www.jetp.ac.ru/cgi-bin/e/index/e/11/3/p696?a=list>].
- Eliashberg, G. M., 1963, "The low temperature specific heat of metals," *Sov. Phys. JETP* **16**, 780–781.
- Emin, D., 2013, *Polarons* (Cambridge University Press, Cambridge, England).
- Engel, E. A., B. Monserrat, and R. J. Needs, 2015, "Vibrational renormalisation of the electronic band gap in hexagonal and cubic ice," *J. Chem. Phys.* **143**, 244708.
- Engelsberg, S., and J. R. Schrieffer, 1963, "Coupled electron-phonon system," *Phys. Rev.* **131**, 993–1008.
- Errea, I., M. Calandra, and F. Mauri, 2014, "Anharmonic free energies and phonon dispersions from the stochastic self-consistent harmonic approximation: Application to platinum and palladium hydrides," *Phys. Rev. B* **89**, 064302.
- Errea, I., B. Rousseau, and A. Bergara, 2011, "Anharmonic stabilization of the high-pressure simple cubic phase of calcium," *Phys. Rev. Lett.* **106**, 165501.
- Faber, C., P. Boulanger, C. Attaccalite, E. Cannuccia, I. Duchemin, T. Deutsch, and X. Blase, 2015, "Exploring approximations to the *GW* self-energy ionic gradients," *Phys. Rev. B* **91**, 155109.
- Faber, C., J. Laflamme Janssen, M. Côté, E. Runge, and X. Blase, 2011, "Electron-phonon coupling in the  $C_{60}$  fullerene within the many-body *GW* approach," *Phys. Rev. B* **84**, 155104.
- Fan, H. Y., 1951, "Temperature dependence of the energy gap in semiconductors," *Phys. Rev.* **82**, 900–905.
- Fetter, A. L., and J. D. Walecka, 2003, *Quantum theory of many-particle systems* (Dover Publications, New York).
- Florentini, M., and N. Bonini, 2016, "Thermoelectric coefficients of *n*-doped silicon from first principles via the solution of the Boltzmann transport equation," *Phys. Rev. B* **94**, 085204.
- Flores-Livas, J. A., and A. Sanna, 2015, "Superconductivity in intercalated group-IV honeycomb structures," *Phys. Rev. B* **91**, 054508.
- Floris, A., S. de Gironcoli, E. K. U. Gross, and M. Cococcioni, 2011, "Vibrational properties of MnO and NiO from DFT+*U*-based density functional perturbation theory," *Phys. Rev. B* **84**, 161102.
- Floris, A., G. Profeta, N. N. Lathiotakis, M. Lüders, M. A. L. Marques, C. Franchini, E. K. U. Gross, A. Continenza, and S. Massidda, 2005, "Superconducting properties of  $MgB_2$  from first principles," *Phys. Rev. Lett.* **94**, 037004.
- Floris, A., A. Sanna, S. Massidda, and E. K. U. Gross, 2007, "Two-band superconductivity in Pb from *ab initio* calculations," *Phys. Rev. B* **75**, 054508.
- Franceschetti, A., 2007, "First-principles calculations of the temperature dependence of the band gap of Si nanocrystals," *Phys. Rev. B* **76**, 161301.
- Frenkel, J., 1932, *Wave Mechanics: Elementary Theory* (Oxford University Press, New York).
- Friedrich, M., A. Riefer, S. Sanna, W. G. Schmidt, and A. Schindlmayr, 2015, "Phonon dispersion and zero-point renormalization of  $LiNbO_3$  from density-functional perturbation theory," *J. Phys. Condens. Matter* **27**, 385402.
- Frohlich, H., 1937, "Theory of electrical breakdown in ionic crystals," *Proc. R. Soc. A* **160**, 230–241.
- Frohlich, H., 1954, "Electrons in lattice fields," *Adv. Phys.* **3**, 325–361.
- Frohlich, H., and N. F. Mott, 1939, "The mean free path of electrons in polar crystals," *Proc. R. Soc. A* **171**, 496–504.
- Frohlich, H., H. Pelzer, and S. Zienau, 1950, "Properties of slow electrons in polar materials," *Philos. Mag. Ser. 7* **41**, 221–242.
- Giannozzi, P., S. de Gironcoli, P. Pavone, and S. Baroni, 1991, "*Ab initio* calculation of phonon dispersions in semiconductors," *Phys. Rev. B* **43**, 7231–7242.
- Giannozzi, P., *et al.*, 2009, "QUANTUM ESPRESSO: a modular and open-source software project for quantum simulations of materials," *J. Phys. Condens. Matter* **21**, 395502.
- Gillis, N. S., 1970, "Self-consistent phonons and the coupled electron-phonon system," *Phys. Rev. B* **1**, 1872–1876.
- Giuliani, G., and G. Vignale, 2005, *Quantum Theory of the Electron Liquid* (Cambridge University Press, Cambridge, England).
- Giustino, F., 2014, *Materials Modelling using Density Functional Theory* (Oxford University Press, New York).
- Giustino, F., M. L. Cohen, and S. G. Louie, 2007, "Electron-phonon interaction using Wannier functions," *Phys. Rev. B* **76**, 165108.
- Giustino, F., M. L. Cohen, and S. G. Louie, 2008, "Small phonon contribution to the photoemission kink in the copper oxide superconductors," *Nature (London)* **452**, 975–978.
- Giustino, F., S. G. Louie, and M. L. Cohen, 2010, "Electron-phonon renormalization of the direct band gap of diamond," *Phys. Rev. Lett.* **105**, 265501.
- Giustino, F., J. R. Yates, I. Souza, M. L. Cohen, and S. G. Louie, 2007, "Electron-phonon interaction via electronic and lattice wannier functions: Superconductivity in boron-doped diamond reexamined," *Phys. Rev. Lett.* **98**, 047005.
- Göbel, A., T. Ruf, M. Cardona, C. T. Lin, J. Wrzesinski, M. Steube, K. Reimann, J.-C. Merle, and M. Joucla, 1998, "Effects of the isotopic composition on the fundamental gap of CuCl," *Phys. Rev. B* **57**, 15183–15190.
- Gonze, X., 1995a, "Perturbation expansion of variational principles at arbitrary order," *Phys. Rev. A* **52**, 1086–1095.
- Gonze, X., 1995b, "Adiabatic density-functional perturbation theory," *Phys. Rev. A* **52**, 1096–1114.
- Gonze, X., 1997, "First-principles responses of solids to atomic displacements and homogeneous electric fields: Implementation of a conjugate-gradient algorithm," *Phys. Rev. B* **55**, 10337–10354.
- Gonze, X., D. C. Allan, and M. P. Teter, 1992, "Dielectric tensor, effective charges, and phonons in  $\alpha$ -quartz by variational density-functional perturbation theory," *Phys. Rev. Lett.* **68**, 3603–3606.
- Gonze, X., P. Boulanger, and M. Côté, 2011, "Theoretical approaches to the temperature and zero-point motion effects on the electronic band structure," *Ann. Phys. (Berlin)* **523**, 168–178.
- Gonze, X., and C. Lee, 1997, "Dynamical matrices, Born effective charges, dielectric permittivity tensors, and interatomic force constants from density-functional perturbation theory," *Phys. Rev. B* **55**, 10355–10368.
- Gonze, X., *et al.*, 2009, "ABINIT: First-principles approach to material and nanosystem properties," *Comput. Phys. Commun.* **180**, 2582–2615.
- Gopalan, S., P. Lautenschlager, and M. Cardona, 1987, "Temperature dependence of the shifts and broadenings of the critical points in GaAs," *Phys. Rev. B* **35**, 5577–5584.
- Gor'kov, L. P., 1958, "On the energy spectrum of superconductors," *Sov. Phys. JETP* **7**, 505–508 [<http://www.jetp.ac.ru/cgi-bin/e/index/e/7/3/p505?a=list>].
- Grimvall, G., 1969, "New aspects on the electron-phonon system at finite temperatures with an application on lead and mercury," *Phys. Kondens. Mater.* **9**, 283–299.
- Grimvall, G., 1975, "Electron-electron renormalization of the electronic heat capacity in simple metals," *Phys. Scr.* **12**, 337.
- Grimvall, G., 1981, *The electron-phonon interaction in metals* (North-Holland, Amsterdam).
- Grüneis, A., C. Attaccalite, A. Rubio, D. V. Vyalikh, S. L. Molodtsov, J. Fink, R. Follath, W. Eberhardt, B. Büchner, and T. Pichler, 2009a,

- “Electronic structure and electron-phonon coupling of doped graphene layers in  $KC_8$ ,” *Phys. Rev. B* **79**, 205106.
- Grüneis, A., *et al.*, 2009b, “Phonon surface mapping of graphite: Disentangling quasi-degenerate phonon dispersions,” *Phys. Rev. B* **80**, 085423.
- Gumhalter, B., V. Kovač, F. Caruso, H. Lambert, and F. Giustino, 2016, “On the combined use of GW approximation and cumulant expansion in the calculations of quasiparticle spectra: The paradigm of Si valence bands,” *Phys. Rev. B* **94**, 035103.
- Gunnarsson, O., B. I. Lundqvist, and J. W. Wilkins, 1974, “Contribution to the cohesive energy of simple metals: Spin-dependent effect,” *Phys. Rev. B* **10**, 1319–1327.
- Gunst, T., T. Markussen, K. Stokbro, and M. Brandbyge, 2016, “First-principles method for electron-phonon coupling and electron mobility: Applications to two-dimensional materials,” *Phys. Rev. B* **93**, 035414.
- Guzzo, M., J. J. Kas, F. Sottile, M. G. Silly, F. Sirotti, J. J. Rehr, and L. Reining, 2012, “Plasmon satellites in valence-band photoemission spectroscopy,” *Eur. Phys. J. B* **85**, 324.
- Guzzo, M., J. J. Kas, L. Sponza, C. Giorgetti, F. Sottile, D. Pierucci, M. G. Silly, F. Sirotti, J. J. Rehr, and L. Reining, 2014, “Multiple satellites in materials with complex plasmon spectra: From graphite to graphene,” *Phys. Rev. B* **89**, 085425.
- Guzzo, M., G. Lani, F. Sottile, P. Romaniello, M. Gatti, J. J. Kas, J. J. Rehr, M. G. Silly, F. Sirotti, and L. Reining, 2011, “Valence electron photoemission spectrum of semiconductors: *Ab initio* description of multiple satellites,” *Phys. Rev. Lett.* **107**, 166401.
- Hall, L. H., J. Bardeen, and F. J. Blatt, 1954, “Infrared absorption spectrum of germanium,” *Phys. Rev.* **95**, 559–560.
- Hamann, D. R., M. Schlüter, and C. Chiang, 1979, “Norm-conserving pseudopotentials,” *Phys. Rev. Lett.* **43**, 1494–1497.
- Han, P., and G. Bester, 2013, “Large nuclear zero-point motion effect in semiconductor nanoclusters,” *Phys. Rev. B* **88**, 165311.
- Hannewald, K., and P. A. Bobbert, 2004a, “*Ab initio* theory of charge-carrier conduction in ultrapure organic crystals,” *Appl. Phys. Lett.* **85**, 1535–1537.
- Hannewald, K., and P. A. Bobbert, 2004b, “Anisotropy effects in phonon-assisted charge-carrier transport in organic molecular crystals,” *Phys. Rev. B* **69**, 075212.
- Hayman, B., and J. P. Carbotte, 1971, “Resistivity and electron-phonon interaction in Li,” *J. Phys. F* **1**, 828.
- He, L., F. Liu, G. Hautier, M. J. T. Oliveira, M. A. L. Marques, F. D. Vila, J. J. Rehr, G.-M. Rignanese, and A. Zhou, 2014, “Accuracy of generalized gradient approximation functionals for density-functional perturbation theory calculations,” *Phys. Rev. B* **89**, 064305.
- He, L., and D. Vanderbilt, 2001, “Exponential decay properties of Wannier functions and related quantities,” *Phys. Rev. Lett.* **86**, 5341–5344.
- Hedin, L., 1965, “New method for calculating the one-particle Green’s function with application to the electron-gas problem,” *Phys. Rev.* **139**, A796–A823.
- Hedin, L., and B. I. Lundqvist, 1971, “Explicit local exchange-correlation potentials,” *J. Phys. C* **4**, 2064.
- Hedin, L., and S. Lundqvist, 1969, *Effects of electron-electron and electron-phonon interactions on the one-electron states of solids*, edited by F. Seitz, D. Turnbull, and H. Ehrenreich, Solid State Physics, Vol. 23 (Academic, New York).
- Heid, R., K.-P. Bohnen, I. Yu. Sklyadneva, and E. V. Chulkov, 2010, “Effect of spin-orbit coupling on the electron-phonon interaction of the superconductors Pb and Tl,” *Phys. Rev. B* **81**, 174527.
- Heid, R., K.-P. Bohnen, R. Zeyher, and D. Manske, 2008, “Momentum dependence of the electron-phonon coupling and self-energy effects in superconducting  $YBa_2Cu_3O_7$  within the local density approximation,” *Phys. Rev. Lett.* **100**, 137001.
- Heine, V., and I. Abarenkov, 1964, “A new method for the electronic structure of metals,” *Philos. Mag.* **9**, 451–465.
- Hellman, O., I. A. Abrikosov, and S. I. Simak, 2011, “Lattice dynamics of anharmonic solids from first principles,” *Phys. Rev. B* **84**, 180301.
- Hellman, O., P. Steneteg, I. A. Abrikosov, and S. I. Simak, 2013, “Temperature dependent effective potential method for accurate free energy calculations of solids,” *Phys. Rev. B* **87**, 104111.
- Hengsberger, M., D. Purdie, P. Segovia, M. Garnier, and Y. Baer, 1999, “Photoemission study of a strongly coupled electron-phonon system,” *Phys. Rev. Lett.* **83**, 592–595.
- Herring, C., 1940, “A new method for calculating wave functions in crystals,” *Phys. Rev.* **57**, 1169–1177.
- Heyd, J., G. E. Scuseria, and M. Ernzerhof, 2003, “Hybrid functionals based on a screened Coulomb potential,” *J. Chem. Phys.* **118**, 8207.
- Himmetoglu, B., and A. Janotti, 2016, “Transport properties of  $KTaO_3$  from first principles,” *J. Phys. Condens. Matter* **28**, 065502.
- Himmetoglu, B., A. Janotti, H. Peelaers, A. Alkauskas, and C. G. Van de Walle, 2014, “First-principles study of the mobility of  $SrTiO_3$ ,” *Phys. Rev. B* **90**, 241204.
- Hoddeson, L. H., and G. Baym, 1980, “The development of the quantum mechanical electron theory of metals: 1900–28,” *Proc. R. Soc. A* **371**, 8–23.
- Hohenberg, P., and W. Kohn, 1964, “Inhomogeneous electron gas,” *Phys. Rev.* **136**, B864–B871.
- Holstein, T., 1959, “Studies of polaron motion: Part I. The molecular-crystal model,” *Ann. Phys. (N.Y.)* **8**, 325–342.
- Hone, D., 1960, “Exchange and correlation corrections to the electron-phonon interaction,” *Phys. Rev.* **120**, 1600–1606.
- Hong, J., A. Stroppa, J. Íñiguez, S. Picozzi, and D. Vanderbilt, 2012, “Spin-phonon coupling effects in transition-metal perovskites: A DFT + *U* and hybrid-functional study,” *Phys. Rev. B* **85**, 054417.
- Hooton, D. J., 1955, “LI. a new treatment of anharmonicity in lattice thermodynamics: I,” *Philos. Mag.* **46**, 422–432.
- Hybertsen, M. S., and S. G. Louie, 1986, “Electron correlation in semiconductors and insulators: Band gaps and quasiparticle energies,” *Phys. Rev. B* **34**, 5390–5413.
- Ichinokura, S., K. Sugawara, A. Takayama, T. Takahashi, and S. Hasegawa, 2016, “Superconducting calcium-intercalated bilayer graphene,” *ACS Nano* **10**, 2761–2765.
- Johnson, P. D., *et al.*, 2001, “Doping and temperature dependence of the mass enhancement observed in the cuprate  $Bi_2Sr_2CaCu_2O_{8+\delta}$ ,” *Phys. Rev. Lett.* **87**, 177007.
- Kaasbjerg, K., K. S. Thygesen, and K. W. Jacobsen, 2012, “Phonon-limited mobility in *n*-type single-layer  $MoS_2$  from first principles,” *Phys. Rev. B* **85**, 115317.
- Kadanoff, L. P., and G. Baym, 1962, *Quantum statistical mechanics: Green’s function methods in equilibrium and nonequilibrium problems*, Frontiers in Physics (Benjamin, New York).
- Karki, B. B., R. M. Wentzcovitch, S. de Gironcoli, and S. Baroni, 2000, “*Ab initio* lattice dynamics of  $MgSiO_3$  perovskite at high pressure,” *Phys. Rev. B* **62**, 14750–14756.
- Kas, J. J., J. J. Rehr, and L. Reining, 2014, “Cumulant expansion of the retarded one-electron Green function,” *Phys. Rev. B* **90**, 085112.
- Kato, T., T. Kobayashi, and M. Namiki, 1960, “Formal theory of Green functions,” *Prog. Theor. Phys.* **15**, 3–60.
- Kaveh, M., and N. Wiser, 1972, “Temperature and volume dependence of the electrical resistivity of sodium and potassium,” *Phys. Rev. B* **6**, 3648–3658.
- Kawai, H., K. Yamashita, E. Cannuccia, and A. Marini, 2014, “Electron-electron and electron-phonon correlation effects on the

- finite-temperature electronic and optical properties of zinc-blende GaN," *Phys. Rev. B* **89**, 085202.
- Keating, P. N., 1968, "Dielectric screening and the phonon spectra of metallic and nonmetallic crystals," *Phys. Rev.* **175**, 1171–1180.
- Kheifets, A. S., V. A. Sashin, M. Vos, E. Weigold, and F. Aryasetiawan, 2003, "Spectral properties of quasiparticles in silicon: A test of many-body theory," *Phys. Rev. B* **68**, 233205.
- Kim, T. Y., C.-H. Park, and N. Marzari, 2016, "The electronic thermal conductivity of graphene," *Nano Lett.* **16**, 2439–2443.
- King-Smith, R. D., R. J. Needs, V. Heine, and M. J. Hodgson, 1989, "A first-principle calculation of the temperature dependence of the indirect band gap of silicon," *Europhys. Lett.* **10**, 569.
- King-Smith, R. D., and D. Vanderbilt, 1993, "Theory of polarization of crystalline solids," *Phys. Rev. B* **47**, 1651–1654.
- Kioupakis, E., P. Rinke, A. Schleife, F. Bechstedt, and C. G. Van de Walle, 2010, "Free-carrier absorption in nitrides from first principles," *Phys. Rev. B* **81**, 241201.
- Kioupakis, E., D. Steiauf, P. Rinke, K. T. Delaney, and C. G. Van de Walle, 2015, "First-principles calculations of indirect auger recombination in nitride semiconductors," *Phys. Rev. B* **92**, 035207.
- Kittel, C., 1963, *Quantum Theory of Solids* (Wiley, New York).
- Kittel, C., 1976, *Introduction to Solid State Physics* (Wiley, New York), 5th ed.
- Koehler, T. R., 1966, "Theory of the self-consistent harmonic approximation with application to solid neon," *Phys. Rev. Lett.* **17**, 89–91.
- Kohn, W., 1959a, "Analytic properties of Bloch waves and Wannier functions," *Phys. Rev.* **115**, 809–821.
- Kohn, W., 1959b, "Image of the fermi surface in the vibration spectrum of a metal," *Phys. Rev. Lett.* **2**, 393–394.
- Kohn, W., and L. J. Sham, 1965, "Self-consistent equations including exchange and correlation effects," *Phys. Rev.* **140**, A1133–A1138.
- Komelj, M., and H. Krakauer, 2015, "Electron-phonon coupling and exchange-correlation effects in superconducting H<sub>3</sub>S under high pressure," *Phys. Rev. B* **92**, 205125.
- Kreibich, T., and E. K. U. Gross, 2001, "Multicomponent density-functional theory for electrons and nuclei," *Phys. Rev. Lett.* **86**, 2984–2987.
- Kreibich, T., R. van Leeuwen, and E. K. U. Gross, 2008, "Multicomponent density-functional theory for electrons and nuclei," *Phys. Rev. A* **78**, 022501.
- Kresse, G., J. Furthmüller, and J. Hafner, 1995, "Ab initio force constant approach to phonon dispersion relations of diamond and graphite," *Europhys. Lett.* **32**, 729.
- Laflamme Janssen, J., M. Côté, S. G. Louie, and M. L. Cohen, 2010, "Electron-phonon coupling in C<sub>60</sub> using hybrid functionals," *Phys. Rev. B* **81**, 073106.
- Lam, P. K., M. M. Dacorogna, and M. L. Cohen, 1986, "Self-consistent calculation of electron-phonon couplings," *Phys. Rev. B* **34**, 5065–5069.
- Langreth, D. C., 1970, "Singularities in the X-ray spectra of metals," *Phys. Rev. B* **1**, 471–477.
- Lanzara, A., *et al.*, 2001, "Evidence for ubiquitous strong electron-phonon coupling in high-temperature superconductors," *Nature (London)* **412**, 510–514.
- LaShell, S., E. Jensen, and T. Balasubramanian, 2000, "Nonquasi-particle structure in the photoemission spectra from the Be(0001) surface and determination of the electron self energy," *Phys. Rev. B* **61**, 2371–2374.
- Lautenschlager, P., P. B. Allen, and M. Cardona, 1985, "Temperature dependence of band gaps in Si and Ge," *Phys. Rev. B* **31**, 2163–2171.
- Lautenschlager, P., P. B. Allen, and M. Cardona, 1986, "Phonon-induced lifetime broadenings of electronic states and critical points in Si and Ge," *Phys. Rev. B* **33**, 5501–5511.
- Lautenschlager, P., M. Garriga, S. Logothetidis, and M. Cardona, 1987, "Interband critical points of GaAs and their temperature dependence," *Phys. Rev. B* **35**, 9174–9189.
- Lautenschlager, P., M. Garriga, L. Vina, and M. Cardona, 1987, "Temperature dependence of the dielectric function and interband critical points in silicon," *Phys. Rev. B* **36**, 4821–4830.
- Lax, M., 1952, "The Franck-Condon principle and its application to crystals," *J. Chem. Phys.* **20**, 1752–1760.
- Łażewski, J., P. Piekarczyk, J. Toboła, B. Wiendlocha, P. T. Jochym, M. Sternik, and K. Parlinski, 2010, "Phonon mechanism of the magnetostructural phase transition in MnAs," *Phys. Rev. Lett.* **104**, 147205.
- Lazzeri, M., C. Attaccalite, L. Wirtz, and F. Mauri, 2008, "Impact of the electron-electron correlation on phonon dispersion: Failure of LDA and GGA DFT functionals in graphene and graphite," *Phys. Rev. B* **78**, 081406.
- Lazzeri, M., M. Calandra, and F. Mauri, 2003, "Anharmonic phonon frequency shift in MgB<sub>2</sub>," *Phys. Rev. B* **68**, 220509.
- Lazzeri, M., and F. Mauri, 2006, "Nonadiabatic Kohn anomaly in a doped graphene monolayer," *Phys. Rev. Lett.* **97**, 266407.
- Lazzeri, M., S. Piscanec, F. Mauri, A. C. Ferrari, and J. Robertson, 2006, "Phonon linewidths and electron-phonon coupling in graphite and nanotubes," *Phys. Rev. B* **73**, 155426.
- Lee, C., P. Ghosez, and X. Gonze, 1994, "Lattice dynamics and dielectric properties of incipient ferroelectric TiO<sub>2</sub> rutile," *Phys. Rev. B* **50**, 13379–13387.
- Lee, J. H., and K. M. Rabe, 2011, "Large spin-phonon coupling and magnetically induced phonon anisotropy in SrMO<sub>3</sub> perovskites (M = V, Cr, Mn, Fe, Co)," *Phys. Rev. B* **84**, 104440.
- Lee, K.-H., and K. J. Chang, 1996, "Linear-response calculation of the Coulomb pseudopotential  $\mu^*$  for Nb," *Phys. Rev. B* **54**, 1419–1422.
- Lee, K.-H., K. J. Chang, and M. L. Cohen, 1995, "First-principles calculations of the Coulomb pseudopotential  $\mu^*$ : Application to Al," *Phys. Rev. B* **52**, 1425–1428.
- Lee, T. D., F. E. Low, and D. Pines, 1953, "The motion of slow electrons in a polar crystal," *Phys. Rev.* **90**, 297–302.
- Leonardo, A., I. Yu. Sklyadneva, V. M. Silkin, P. M. Echenique, and E. V. Chulkov, 2007, "Ab initio calculation of the phonon-induced contribution to the electron-state linewidth on the Mg(0001) surface versus bulk Mg," *Phys. Rev. B* **76**, 035404.
- Li, W., 2015, "Electrical transport limited by electron-phonon coupling from boltzmann transport equation: An ab initio study of Si, Al, and MoS<sub>2</sub>," *Phys. Rev. B* **92**, 075405.
- Li, X., J. T. Mullen, Z. Jin, K. M. Borysenko, M. Buongiorno Nardelli, and K. W. Kim, 2013, "Intrinsic electrical transport properties of monolayer silicene and MoS<sub>2</sub> from first principles," *Phys. Rev. B* **87**, 115418.
- Liao, B., B. Qiu, J. Zhou, S. Huberman, K. Esfarjani, and G. Chen, 2015, "Significant reduction of lattice thermal conductivity by the electron-phonon interaction in silicon with high carrier concentrations: A first-principles study," *Phys. Rev. Lett.* **114**, 115901.
- Linscheid, A., A. Sanna, F. Essenberg, and E. K. U. Gross, 2015, "Ab initio theory of superconductivity in a magnetic field. I. spin density functional theory for superconductors and eliashberg equations," *Phys. Rev. B* **92**, 024505.
- Linscheid, A., A. Sanna, A. Floris, and E. K. U. Gross, 2015, "First-principles calculation of the real-space order parameter and

- condensation energy density in phonon-mediated superconductors,” *Phys. Rev. Lett.* **115**, 097002.
- Linscheid, A., A. Sanna, and E. K. U. Gross, 2015, “*Ab initio* theory of superconductivity in a magnetic field. II. numerical solution,” *Phys. Rev. B* **92**, 024506.
- Lischner, J., D. Vigil-Fowler, and S. G. Louie, 2013, “Physical origin of satellites in photoemission of doped graphene: An *ab initio* GW plus cumulant study,” *Phys. Rev. Lett.* **110**, 146801.
- Liu, A. Y., and A. A. Quong, 1996, “Linear-response calculation of electron-phonon coupling parameters,” *Phys. Rev. B* **53**, R7575–R7579.
- Lloyd-Williams, J. H., and B. Monserrat, 2015, “Lattice dynamics and electron-phonon coupling calculations using nondiagonal supercells,” *Phys. Rev. B* **92**, 184301.
- Lüders, M., M. A. L. Marques, N. N. Lathiotakis, A. Floris, G. Profeta, L. Fast, A. Continenza, S. Massidda, and E. K. U. Gross, 2005, “*Ab initio* theory of superconductivity. I. Density functional formalism and approximate functionals,” *Phys. Rev. B* **72**, 024545.
- Luttinger, J. M., 1960, “Fermi surface and some simple equilibrium properties of a system of interacting fermions,” *Phys. Rev.* **119**, 1153–1163.
- Luttinger, J. M., and J. C. Ward, 1960, “Ground-state energy of a many-fermion system. II,” *Phys. Rev.* **118**, 1417–1427.
- Mahan, G. D., 1993, *Many-Particle Physics* (Plenum, New York), 2nd ed.
- Maksimov, E. G., 1976, “A self-consistent description of the electron-phonon system in metals and the problem of lattice stability,” *Sov. Phys. JETP* **42**, 1138–1143 [<http://www.jetp.ac.ru/cgi-bin/e/index/e/42/6/p1138?a=list>].
- Maksimov, E. G., and S. V. Shulga, 1996, “Nonadiabatic effects in optical phonon self-energy,” *Solid State Commun.* **97**, 553–560.
- Mandal, S., R. E. Cohen, and K. Haule, 2014, “Strong pressure-dependent electron-phonon coupling in FeSe,” *Phys. Rev. B* **89**, 220502.
- Maradudin, A. A., and S. H. Vosko, 1968, “Symmetry properties of the normal vibrations of a crystal,” *Rev. Mod. Phys.* **40**, 1–37.
- Margine, E. R., and F. Giustino, 2013, “Anisotropic Migdal-Eliashberg theory using Wannier functions,” *Phys. Rev. B* **87**, 024505.
- Margine, E. R., and F. Giustino, 2014, “Two-gap superconductivity in heavily *n*-doped graphene: *Ab initio* Migdal-Eliashberg theory,” *Phys. Rev. B* **90**, 014518.
- Margine, E. R., H. Lambert, and F. Giustino, 2016, “Electron-phonon interaction and pairing mechanism in superconducting Ca-intercalated bilayer graphene,” *Sci. Rep.* **6**, 21414.
- Marini, A., 2008, “*Ab initio* finite-temperature excitons,” *Phys. Rev. Lett.* **101**, 106405.
- Marini, A., S. Poncé, and X. Gonze, 2015, “Many-body perturbation theory approach to the electron-phonon interaction with density-functional theory as a starting point,” *Phys. Rev. B* **91**, 224310.
- Marques, M. A. L., M. Lüders, N. N. Lathiotakis, G. Profeta, A. Floris, L. Fast, A. Continenza, E. K. U. Gross, and S. Massidda, 2005, “*Ab initio* theory of superconductivity. II. Application to elemental metals,” *Phys. Rev. B* **72**, 024546.
- Marsiglio, F., M. Schossmann, and J. P. Carbotte, 1988, “Iterative analytic continuation of the electron self-energy to the real axis,” *Phys. Rev. B* **37**, 4965–4969.
- Martin, R. M., 2004, *Electronic structure: Basic theory and practical methods* (Cambridge University Press, Cambridge, England).
- Marzari, N., A. A. Mostofi, J. R. Yates, I. Souza, and D. Vanderbilt, 2012, “Maximally localized Wannier functions: Theory and applications,” *Rev. Mod. Phys.* **84**, 1419–1475.
- Marzari, N., and D. Vanderbilt, 1997, “Maximally localized generalized Wannier functions for composite energy bands,” *Phys. Rev. B* **56**, 12847–12865.
- Mauri, F., O. Zakharov, S. de Gironcoli, S. G. Louie, and M. L. Cohen, 1996, “Phonon softening and superconductivity in tellurium under pressure,” *Phys. Rev. Lett.* **77**, 1151–1154.
- McMillan, W. L., 1968, “Transition temperature of strong-coupled superconductors,” *Phys. Rev.* **167**, 331–344.
- Mermin, N. D., 1966, “A short simple evaluation of expressions of the Debye-Waller form,” *J. Math. Phys. (N.Y.)* **7**, 1038–1038.
- Merzbacher, E., 1998, *Quantum Mechanics* (Wiley, New York), 3rd ed.
- Migdal, A. B., 1958, “Interaction between electrons and lattice vibrations in a normal metal,” *Sov. Phys. JETP* **7**, 996–1001 [<http://www.jetp.ac.ru/cgi-bin/e/index/e/7/6/p996?a=list>].
- Monserrat, B., 2016a, “Correlation effects on electron-phonon coupling in semiconductors: Many-body theory along thermal lines,” *Phys. Rev. B* **93**, 100301.
- Monserrat, B., 2016b, “Vibrational averages along thermal lines,” *Phys. Rev. B* **93**, 014302.
- Monserrat, B., N. D. Drummond, and R. J. Needs, 2013, “Anharmonic vibrational properties in periodic systems: energy, electron-phonon coupling, and stress,” *Phys. Rev. B* **87**, 144302.
- Monserrat, B., N. D. Drummond, C. J. Pickard, and R. J. Needs, 2014, “Electron-phonon coupling and the metallization of solid helium at terapascal pressures,” *Phys. Rev. Lett.* **112**, 055504.
- Monserrat, B., E. A. Engel, and R. J. Needs, 2015, “Giant electron-phonon interactions in molecular crystals and the importance of nonquadratic coupling,” *Phys. Rev. B* **92**, 140302.
- Monserrat, B., and R. J. Needs, 2014, “Comparing electron-phonon coupling strength in diamond, silicon, and silicon carbide: First-principles study,” *Phys. Rev. B* **89**, 214304.
- Mooradian, A., and G. B. Wright, 1966, “Observation of the interaction of plasmons with longitudinal optical phonons in GaAs,” *Phys. Rev. Lett.* **16**, 999–1001.
- Morel, P., and P. W. Anderson, 1962, “Calculation of the superconducting state parameters with retarded electron-phonon interaction,” *Phys. Rev.* **125**, 1263–1271.
- Moser, S., *et al.*, 2013, “Tunable polaronic conduction in anatase TiO<sub>2</sub>,” *Phys. Rev. Lett.* **110**, 196403.
- Mostofi, A. A., J. R. Yates, Y.-S. Lee, I. Souza, D. Vanderbilt, and N. Marzari, 2008, “Wannier90: A tool for obtaining maximally-localised Wannier functions,” *Comput. Phys. Commun.* **178**, 685–699.
- Mott, N. F., and H. Jones, 1936, *The Theory of the Properties of Metals and Alloys* (Oxford University Press, New York).
- Nagamatsu, J., N. Nakagawa, T. Muranaka, Y. Zenitani, and J. Akimitsu, 2001, “Superconductivity at 39 K in magnesium diboride,” *Nature (London)* **410**, 63–64.
- Nakajima, S., 1954, *Proceedings of the International Conference on Theoretical Physics at Kyoto-Tokyo* (Science Council of Japan).
- Nambu, Y., 1960, “Quasi-particles and gauge invariance in the theory of superconductivity,” *Phys. Rev.* **117**, 648–663.
- Nery, J. P., and P. B. Allen, 2016, “Influence of Fröhlich polaron coupling on renormalized electron bands in polar semiconductors: Results for zinc-blende GaN,” *Phys. Rev. B* **94**, 115135.
- Noffsinger, J., F. Giustino, S. G. Louie, and M. L. Cohen, 2009, “Origin of superconductivity in boron-doped silicon carbide from first principles,” *Phys. Rev. B* **79**, 104511.
- Noffsinger, J., F. Giustino, B. D. Malone, C.-H. Park, S. G. Louie, and M. L. Cohen, 2010, “EPW: A program for calculating the electron-phonon coupling using maximally localized wannier functions,” *Comput. Phys. Commun.* **181**, 2140–2148.

- Noffsinger, J., E. Kioupakis, C. G. Van de Walle, S. G. Louie, and M. L. Cohen, 2012, “Phonon-assisted optical absorption in silicon from first principles,” *Phys. Rev. Lett.* **108**, 167402.
- Nordheim, L., 1931, “Zur Elektronentheorie der Metalle. I,” *Ann. Phys. (Berlin)* **401**, 607–640.
- Olguin, D., M. Cardona, and A. Cantarero, 2002, “Electron-phonon effects on the direct band gap in semiconductors: LCAO calculations,” *Solid State Commun.* **122**, 575–589.
- Onida, G., L. Reining, and A. Rubio, 2002, “Electronic excitations: density-functional versus many-body Green’s-function approaches,” *Rev. Mod. Phys.* **74**, 601–659.
- Ortmann, F., F. Bechstedt, and K. Hannewald, 2009, “Theory of charge transport in organic crystals: Beyond Holstein’s small-polaron model,” *Phys. Rev. B* **79**, 235206.
- Park, C.-H., N. Bonini, T. Sohier, G. Samsonidze, B. Kozinsky, M. Calandra, F. Mauri, and N. Marzari, 2014, “Electron-phonon interactions and the intrinsic electrical resistivity of graphene,” *Nano Lett.* **14**, 1113–1119.
- Park, C.-H., F. Giustino, M. L. Cohen, and S. G. Louie, 2007, “Velocity renormalization and carrier lifetime in graphene from the electron-phonon interaction,” *Phys. Rev. Lett.* **99**, 086804.
- Park, C.-H., F. Giustino, M. L. Cohen, and S. G. Louie, 2008, “Electron-phonon interactions in graphene, bilayer graphene, and graphite,” *Nano Lett.* **8**, 4229–4233.
- Park, C.-H., F. Giustino, J. L. McChesney, A. Bostwick, T. Ohta, E. Rotenberg, M. L. Cohen, and S. G. Louie, 2008, “Van Hove singularity and apparent anisotropy in the electron-phonon interaction in graphene,” *Phys. Rev. B* **77**, 113410.
- Park, C.-H., F. Giustino, C. D. Spataru, M. L. Cohen, and S. G. Louie, 2009, “Angle-resolved photoemission spectra of graphene from first-principles calculations,” *Nano Lett.* **9**, 4234–4239.
- Parr, R. G., and W. Yang, 1994, *Density-functional theory of atoms and molecules* (Oxford University Press, New York).
- Patrick, C. E., and F. Giustino, 2013, “Quantum nuclear dynamics in the photophysics of diamondoids,” *Nat. Commun.* **4**, 2006.
- Patrick, C. E., and F. Giustino, 2014, “Unified theory of electron-phonon renormalization and phonon-assisted optical absorption,” *J. Phys. Condens. Matter* **26**, 365503.
- Patrick, C. E., K. W. Jacobsen, and K. S. Thygesen, 2015, “Anharmonic stabilization and band gap renormalization in the perovskite  $\text{CsSn}_3$ ,” *Phys. Rev. B* **92**, 201205.
- Peelaers, H., E. Kioupakis, and C. G. Van de Walle, 2015, “Free-carrier absorption in transparent conducting oxides: Phonon and impurity scattering in  $\text{SnO}_2$ ,” *Phys. Rev. B* **92**, 235201.
- Pekar, S. I., 1946, “Autolocalization of an electron in a dielectric inertially polarizing medium,” *Zh. Eksp. Teor. Fiz.* **16**, 335–340.
- Perdew, J. P., K. Burke, and M. Ernzerhof, 1996, “Generalized gradient approximation made simple,” *Phys. Rev. Lett.* **77**, 3865–3868.
- Perdew, J. P., M. Ernzerhof, and K. Burke, 1996, “Rationale for mixing exact exchange with density functional approximations,” *J. Chem. Phys.* **105**, 9982–9985.
- Perdew, J. P., and Alex Zunger, 1981, “Self-interaction correction to density-functional approximations for many-electron systems,” *Phys. Rev. B* **23**, 5048–5079.
- Perroni, C. A., E. Piegari, M. Capone, and V. Cataudella, 2004, “Polaron formation for nonlocal electron-phonon coupling: A variational wave-function study,” *Phys. Rev. B* **69**, 174301.
- Phillips, J. C., 1958, “Energy-band interpolation scheme based on a pseudopotential,” *Phys. Rev.* **112**, 685–695.
- Phillips, J. C., and L. Kleinman, 1959, “New method for calculating wave functions in crystals and molecules,” *Phys. Rev.* **116**, 287–294.
- Pick, R. M., M. H. Cohen, and R. M. Martin, 1970, “Microscopic theory of force constants in the adiabatic approximation,” *Phys. Rev. B* **1**, 910–920.
- Pines, D., and D. Bohm, 1952, “A collective description of electron interactions: II. collective vs individual particle aspects of the interactions,” *Phys. Rev.* **85**, 338–353.
- Pisana, S., M. Lazzeri, C. Casiraghi, K. S. Novoselov, A. K. Geim, A. C. Ferrari, and F. Mauri, 2007, “Breakdown of the adiabatic Born-Oppenheimer approximation in graphene,” *Nat. Mater.* **6**, 198–201.
- Piscanec, S., M. Lazzeri, J. Robertson, A. C. Ferrari, and F. Mauri, 2007, “Optical phonons in carbon nanotubes: Kohn anomalies, peierls distortions, and dynamic effects,” *Phys. Rev. B* **75**, 035427.
- Pizzi, G., D. Volja, B. Kozinsky, M. Fornari, and N. Marzari, 2014, “BoltzWann: A code for the evaluation of thermoelectric and electronic transport properties with a maximally-localized Wannier functions basis,” *Comput. Phys. Commun.* **185**, 422–429.
- Poncé, S., G. Antonius, P. Boulanger, E. Cannuccia, A. Marini, M. Côté, and X. Gonze, 2014, “Verification of first-principles codes: Comparison of total energies, phonon frequencies, electron-phonon coupling and zero-point motion correction to the gap between ABINIT and QE/Yambo,” *Comput. Mater. Sci.* **83**, 341–348.
- Poncé, S., G. Antonius, Y. Gillet, P. Boulanger, J. Laflamme Janssen, A. Marini, M. Côté, and X. Gonze, 2014, “Temperature dependence of electronic eigenenergies in the adiabatic harmonic approximation,” *Phys. Rev. B* **90**, 214304.
- Poncé, S., Y. Gillet, J. Laflamme Janssen, A. Marini, M. Verstraete, and X. Gonze, 2015, “Temperature dependence of the electronic structure of semiconductors and insulators,” *J. Chem. Phys.* **143**, 102813.
- Poncé, S., E. R. Margine, C. Verdi, and F. Giustino, 2016, “EPW: Electron-phonon coupling, superconducting and transport properties using maximally localized wannier functions,” *Comput. Phys. Commun.* **209**, 116–133.
- Prange, R. E., and L. P. Kadanoff, 1964, “Transport theory for electron-phonon interactions in metals,” *Phys. Rev.* **134**, A566–A580.
- Prendergast, D., and S. G. Louie, 2009, “Bloch-state-based interpolation: An efficient generalization of the Shirley approach to interpolating electronic structure,” *Phys. Rev. B* **80**, 235126.
- Profeta, G., C. Franchini, N. N. Lathiotakis, A. Floris, A. Sanna, M. A. L. Marques, M. Lüders, S. Massidda, E. K. U. Gross, and A. Continenza, 2006, “Superconductivity in lithium, potassium, and aluminum under extreme pressure: A first-principles study,” *Phys. Rev. Lett.* **96**, 047003.
- Quong, A. A., and B. M. Klein, 1992, “Self-consistent-screening calculation of interatomic force constants and phonon dispersion curves from first principles,” *Phys. Rev. B* **46**, 10734–10737.
- Ralph, H. I., 1970, “Theoretical phonon limited electron mobility in silicon,” *Solid State Commun.* **8**, 1095–1098.
- Ramírez, R., C. P. Herrero, and E. R. Hernández, 2006, “Path-integral molecular dynamics simulation of diamond,” *Phys. Rev. B* **73**, 245202.
- Ramírez, R., C. P. Herrero, E. R. Hernández, and M. Cardona, 2008, “Path-integral molecular dynamics simulation of  $3C - \text{SiC}$ ,” *Phys. Rev. B* **77**, 045210.
- Resta, R., 1994, “Macroscopic polarization in crystalline dielectrics: the geometric phase approach,” *Rev. Mod. Phys.* **66**, 899–915.
- Restrepo, O. D., K. E. Krymowski, J. Goldberger, and W. Windl, 2014, “A first principles method to simulate electron mobilities in 2D materials,” *New J. Phys.* **16**, 105009.

- Restrepo, O.D., K. Varga, and S.T. Pantelides, 2009, "First-principles calculations of electron mobilities in silicon: Phonon and coulomb scattering," *Appl. Phys. Lett.* **94**, 212103.
- Richter, W., 1984, "Coupled plasmon-polar phonon modes in semiconductors: Spatial dispersion and other properties," in *Polarons and Excitons in Polar Semiconductors and Ionic Crystals*, Vol. 127, edited by J.T. Devreese and F. Peeters (Springer, New York), pp. 209–243.
- Rickayzen, G., 1980, *Green's Functions and Condensed Matter, Techniques of Physics* (Academic, New York).
- Ridley, B.K., 1993, *Quantum Processes in Semiconductors* (Clarendon, Oxford).
- Rösch, O., and O. Gunnarsson, 2004, "Electron-phonon interaction in the  $t$ - $J$  model," *Phys. Rev. Lett.* **92**, 146403.
- Saito, M., 2002, "Electron-phonon coupling of electron- or hole-injected  $C_{60}$ ," *Phys. Rev. B* **65**, 220508.
- Saitta, A. M., M. Lazzeri, M. Calandra, and F. Mauri, 2008, "Giant nonadiabatic effects in layer metals: Raman spectra of intercalated graphite explained," *Phys. Rev. Lett.* **100**, 226401.
- Sangalli, D., and A. Marini, 2015a, "Complete collisions approximation to the Kadanoff-Baym equation: a first-principles implementation," *J. Phys. Conf. Ser.* **609**, 012006.
- Sangalli, D., and A. Marini, 2015b, "Ultra-fast carriers relaxation in bulk silicon following photo-excitation with a short and polarized laser pulse," *Europhys. Lett.* **110**, 47004.
- Sanna, A., C. Franchini, A. Floris, G. Profeta, N.N. Lathiotakis, M. Lüders, M. A. L. Marques, E. K. U. Gross, A. Continenza, and S. Massidda, 2006, "*Ab initio* prediction of pressure-induced superconductivity in potassium," *Phys. Rev. B* **73**, 144512.
- Sanna, A., S. Pittalis, J.K. Dewhurst, M. Monni, S. Sharma, G. Umrigar, S. Massidda, and E.K.U. Gross, 2012, "Phononic self-energy effects and superconductivity in  $CaC_6$ ," *Phys. Rev. B* **85**, 184514.
- Sanna, A., G. Profeta, A. Floris, A. Marini, E. K. U. Gross, and S. Massidda, 2007, "Anisotropic gap of superconducting  $CaC_6$ : A first-principles density functional calculation," *Phys. Rev. B* **75**, 020511.
- Sano, W., T. Koretsune, T. Tadano, R. Akashi, and R. Arita, 2016, "Effect of Van Hove singularities on high- $T_c$  superconductivity in  $H_3S$ ," *Phys. Rev. B* **93**, 094525.
- Savini, G., A. C. Ferrari, and F. Giustino, 2010, "First-principles prediction of doped graphane as a high-temperature electron-phonon superconductor," *Phys. Rev. Lett.* **105**, 037002.
- Savrasov, S. Y., 1992, "Linear response calculations of lattice dynamics using muffin-tin basis sets," *Phys. Rev. Lett.* **69**, 2819–2822.
- Savrasov, S. Y., and G. Kotliar, 2003, "Linear response calculations of lattice dynamics in strongly correlated systems," *Phys. Rev. Lett.* **90**, 056401.
- Savrasov, S. Y., D. Y. Savrasov, and O. K. Andersen, 1994, "Linear-response calculations of electron-phonon interactions," *Phys. Rev. Lett.* **72**, 372–375.
- Scalapino, D. J., 1969, *The Electron-Phonon Interaction and Strong-Coupling Superconductors*, edited by R. D. Parks, Superconductivity, Vol. 1 (Dekker).
- Schäfer, W., and M. Wegener, 2002, *Semiconductor Optics and Transport Phenomena*, Advanced Texts in Physics (Springer, New York).
- Schrieffer, J. R., 1983, *Theory of Superconductivity*, Advanced Book Program Series (Perseus).
- Schwinger, J., 1951, "On the Green's functions of quantized fields. I," *Proc. Natl. Acad. Sci. U.S.A.* **37**, 452–455.
- Sham, L. J., 1961, "Electron-phonon interaction by the method of pseudo-potentials," *Proc. Phys. Soc. London* **78**, 895.
- Sham, L. J., and J. M. Ziman, 1963, *Effects of electron-electron and electron-phonon interactions on the one-electron states of solids*, edited by F. Seitz and D. Turnbull, Solid State Physics, Vol. 15 (Academic, New York).
- Shockley, W., and J. Bardeen, 1950, "Energy bands and mobilities in monatomic semiconductors," *Phys. Rev.* **77**, 407–408.
- Shuey, R. T., 1965, "Electron-phonon interaction for indirect interband transitions in germanium," *Phys. Rev.* **139**, A1675–A1684.
- Shukla, A., *et al.*, 2003, "Phonon dispersion and lifetimes in  $MgB_2$ ," *Phys. Rev. Lett.* **90**, 095506.
- Sjakste, J., N. Vast, M. Calandra, and F. Mauri, 2015, "Wannier interpolation of the electron-phonon matrix elements in polar semiconductors: Polar-optical coupling in GaAs," *Phys. Rev. B* **92**, 054307.
- Sjakste, J., N. Vast, and V. Tyuterev, 2007, "*Ab initio* method for calculating electron-phonon scattering times in semiconductors: Application to GaAs and GaP," *Phys. Rev. Lett.* **99**, 236405.
- Sjölander, A., and R. Johnson, 1965, *Proceedings of the International Conference on Inelastic Scattering of Neutrons, Bombay, 1964* (IAEA Proceedings Series) Vol. 1, pp. 61–76.
- Sklyadneva, I. Yu., E. V. Chulkov, W.-D. Schöne, V. M. Silkin, R. Keyling, and P. M. Echenique, 2005, "Role of electron-phonon interactions versus electron-electron interactions in the broadening mechanism of the electron and hole linewidths in bulk Be," *Phys. Rev. B* **71**, 174302.
- Sklyadneva, I. Yu., A. Leonardo, P. M. Echenique, S. V. Eremeev, and E. V. Chulkov, 2006, "Electron-phonon contribution to the phonon and excited electron (hole) linewidths in bulk Pd," *J. Phys. Condens. Matter* **18**, 7923.
- Sohier, T., M. Calandra, C.-H. Park, N. Bonini, N. Marzari, and F. Mauri, 2014, "Phonon-limited resistivity of graphene by first-principles calculations: Electron-phonon interactions, strain-induced gauge field, and boltzmann equation," *Phys. Rev. B* **90**, 125414.
- Souza, I., N. Marzari, and D. Vanderbilt, 2001, "Maximally localized Wannier functions for entangled energy bands," *Phys. Rev. B* **65**, 035109.
- Sternheimer, R. M., 1954, "Electronic polarizabilities of ions from the Hartree-Fock wave functions," *Phys. Rev.* **96**, 951–968.
- Story, S. M., J. J. Kas, F. D. Vila, M. J. Verstraete, and J. J. Rehr, 2014, "Cumulant expansion for phonon contributions to the electron spectral function," *Phys. Rev. B* **90**, 195135.
- Su, W. P., J. R. Schrieffer, and A. J. Heeger, 1979, "Solitons in polyacetylene," *Phys. Rev. Lett.* **42**, 1698–1701.
- Troullier, N., and J. L. Martins, 1991, "Efficient pseudopotentials for plane-wave calculations," *Phys. Rev. B* **43**, 1993–2006.
- Tyuterev, V. G., S. V. Obukhov, N. Vast, and J. Sjakste, 2011, "*Ab initio* calculation of electron-phonon scattering time in germanium," *Phys. Rev. B* **84**, 035201.
- Valla, T., A. V. Fedorov, P. D. Johnson, and S. L. Hulbert, 1999, "Many-body effects in angle-resolved photoemission: Quasiparticle energy and lifetime of a Mo(110) surface state," *Phys. Rev. Lett.* **83**, 2085–2088.
- Valla, T., A. V. Fedorov, P. D. Johnson, B. O. Wells, S. L. Hulbert, Q. Li, G. D. Gu, and N. Koshizuka, 1999, "Evidence for quantum critical behavior in the optimally doped cuprate  $Bi_2Sr_2CaCu_2O_{8+\delta}$ ," *Science* **285**, 2110–2113.
- Vanderbilt, D., 1990, "Soft self-consistent pseudopotentials in a generalized eigenvalue formalism," *Phys. Rev. B* **41**, 7892–7895.
- van Leeuwen, R., 2004, "First-principles approach to the electron-phonon interaction," *Phys. Rev. B* **69**, 115110.
- Varga, B. B., 1965, "Coupling of plasmons to polar phonons in degenerate semiconductors," *Phys. Rev.* **137**, A1896–A1902.

- Varshni, Y. P., 1967, “Temperature dependence of the energy gap in semiconductors,” *Physica (Utrecht)* **34**, 149–154.
- Verdi, C., F. Caruso, and F. Giustino, 2017, “Origin of the crossover from polarons to Fermi liquids in transition metal oxides” (unpublished).
- Verdi, C., and F. Giustino, 2015, “Fröhlich electron-phonon vertex from first principles,” *Phys. Rev. Lett.* **115**, 176401.
- Verstraete, M. J., M. Torrent, F. Jollet, Z. Gilles, and X. Gonze, 2008, “Density functional perturbation theory with spin-orbit coupling: Phonon band structure of lead,” *Phys. Rev. B* **78**, 045119.
- Villegas, C. E. P., A. R. Rocha, and A. Marini, 2016, “Anomalous temperature dependence of the band gap in black phosphorus,” *Nano Lett.* **16**, 5095–5101.
- Vogl, P., 1976, “Microscopic theory of electron-phonon interaction in insulators or semiconductors,” *Phys. Rev. B* **13**, 694–704.
- von Barth, U., and L. Hedin, 1972, “A local exchange-correlation potential for the spin polarized case. I,” *J. Phys. C* **5**, 1629.
- Vukmirović, N., C. Bruder, and V. M. Stojanović, 2012, “Electron-phonon coupling in crystalline organic semiconductors: Microscopic evidence for nonpolaronic charge carriers,” *Phys. Rev. Lett.* **109**, 126407.
- Walker, C. T., and G. A. Slack, 1970, “Who named the ON’s?,” *Am. J. Phys.* **38**, 1380–1389.
- Walter, J. P., R. R. L. Zucca, M. L. Cohen, and Y. R. Shen, 1970, “Temperature dependence of the wavelength-modulation spectra of GaAs,” *Phys. Rev. Lett.* **24**, 102–104.
- Wang, X., D. Vanderbilt, J. R. Yates, and I. Souza, 2007, “Fermi-surface calculation of the anomalous hall conductivity,” *Phys. Rev. B* **76**, 195109.
- Wang, X., J. R. Yates, I. Souza, and D. Vanderbilt, 2006, “*Ab initio* calculation of the anomalous Hall conductivity by Wannier interpolation,” *Phys. Rev. B* **74**, 195118.
- Wang, Z., S. Wang, S. Obukhov, N. Vast, J. Sjakste, V. Tyuterev, and N. Mingo, 2011, “Thermoelectric transport properties of silicon: Toward an *ab initio* approach,” *Phys. Rev. B* **83**, 205208.
- Wang, Z., *et al.*, 2016, “Tailoring the nature and strength of electron-phonon interactions in the SrTiO<sub>3</sub>(001) 2D electron liquid,” *Nat. Mater.* **15**, 835–839.
- Wannier, G. H., 1937, “The structure of electronic excitation levels in insulating crystals,” *Phys. Rev.* **52**, 191–197.
- Williams, F. E., 1951, “Theoretical low temperature spectra of the thallium activated potassium chloride phosphor,” *Phys. Rev.* **82**, 281–282.
- Wiser, N., 1963, “Dielectric constant with local field effects included,” *Phys. Rev.* **129**, 62–69.
- Wright, A. D., C. Verdi, R. L. Milot, G. E. Eperon, M. A. Pérez-Osorio, H. J. Snaith, F. Giustino, M. B. Johnston, and L. M. Herz, 2016, “Electron-phonon coupling in hybrid lead halide perovskites,” *Nat. Commun.* **7**, 11755.
- Xu, B., and M. J. Verstraete, 2014, “First principles explanation of the positive Seebeck coefficient of lithium,” *Phys. Rev. Lett.* **112**, 196603.
- Yates, J. R., X. Wang, D. Vanderbilt, and I. Souza, 2007, “Spectral and Fermi surface properties from Wannier interpolation,” *Phys. Rev. B* **75**, 195121.
- Yin, M. T., and M. L. Cohen, 1982, “*Ab initio* calculation of the phonon dispersion relation: Application to Si,” *Phys. Rev. B* **25**, 4317–4320.
- Yin, Z. P., A. Kutepov, and G. Kotliar, 2013, “Correlation-enhanced electron-phonon coupling: Applications of *GW* and screened hybrid functional to bismuthates, chloronitrides, and other high-*T<sub>c</sub>* superconductors,” *Phys. Rev. X* **3**, 021011.
- Yu, P. Y., and M. Cardona, 2010, *Fundamentals of Semiconductors* (Springer, New York), 4th ed.
- Zacharias, M., and F. Giustino, 2016, “One-shot calculation of temperature-dependent optical spectra and phonon-induced band-gap renormalization,” *Phys. Rev. B* **94**, 075125.
- Zacharias, M., C. E. Patrick, and F. Giustino, 2015, “Stochastic approach to phonon-assisted optical absorption,” *Phys. Rev. Lett.* **115**, 177401.
- Zhang, P., S. G. Louie, and M. L. Cohen, 2007, “Electron-phonon renormalization in cuprate superconductors,” *Phys. Rev. Lett.* **98**, 067005.
- Zheng, J.-J., and E. R. Margine, 2016, “First-principles calculations of the superconducting properties in Li-decorated monolayer graphene within the anisotropic Migdal-Eliashberg formalism,” *Phys. Rev. B* **94**, 064509.
- Zhou, J. S., J. J. Kas, L. Sponza, I. Reshetnyak, M. Guzzo, C. Giorgetti, M. Gatti, F. Sottile, J. J. Rehr, and L. Reining, 2015, “Dynamical effects in electron spectroscopy,” *J. Chem. Phys.* **143**, 184109.
- Zhou, S. Y., G.-H. Gweon, A. V. Fedorov, P. N. First, W. A. De Heer, D.-H. Lee, F. Guinea, A. H. Castro Neto, and A. Lanzara, 2007, “Substrate-induced bandgap opening in epitaxial graphene,” *Nat. Mater.* **6**, 770–775.
- Ziman, J. M., 1960, *Electrons and phonons: the theory of transport phenomena in solids* (Clarendon, Oxford).
- Zollner, S., M. Cardona, and S. Gopalan, 1992, “Isotope and temperature shifts of direct and indirect band gaps in diamond-type semiconductors,” *Phys. Rev. B* **45**, 3376–3385.