

Colloquium: Superheavy elements: Oganesson and beyond

S. A. Giuliani, Z. Matheson, W. Nazarewicz,* and E. Olsen

Department of Physics and Astronomy and FRIB Laboratory, Michigan State University, East Lansing, Michigan 48824, USA

P.-G. Reinhard

Institut für Theoretische Physik II, Universität Erlangen-Nürnberg, 91058 Erlangen, Germany

J. Sadhukhan

*Variable Energy Cyclotron Centre, Kolkata 700064, India
and Homi Bhabha National Institute, Mumbai 400094, India*

B. Schuetrumpf

*GSI Helmholtzzentrum für Schwerionenforschung, 64291 Darmstadt, Germany
and Institut für Kernphysik, Technische Universität Darmstadt, 64289 Darmstadt, Germany*

N. Schunck

Nuclear and Chemical Science Division, Lawrence Livermore National Laboratory, Livermore, California 94551, USA

P. Schwerdtfeger

*Centre for Theoretical Chemistry and Physics, The New Zealand Institute for Advanced Study, Massey University Auckland, 0632 Auckland, New Zealand
and Centre for Advanced Study at the Norwegian Academy of Science and Letters, 0271 Oslo, Norway* (published 22 January 2019)

During the last decade, six new superheavy elements were added into the seventh period of the periodic table, with the approval of their names and symbols. This milestone was followed by proclaiming 2019 the International Year of the Periodic Table of Chemical Elements by the United Nations General Assembly. According to theory, due to their large atomic numbers, the new arrivals are expected to be qualitatively and quantitatively different from lighter species. The questions pertaining to superheavy atoms and nuclei are in the forefront of research in nuclear and atomic physics and chemistry. This Colloquium offers a broad perspective on the field and outlines future challenges.

DOI: [10.1103/RevModPhys.91.011001](https://doi.org/10.1103/RevModPhys.91.011001)**CONTENTS**

I. Introduction	1	X. Perspectives and Expectations	15
II. The Superheavy Territory	2	Acknowledgments	17
III. Proton and Neutron Densities and Shell Structure	3	References	17
IV. Collectivity	5		
V. α Decay	7		
VI. Fission	8		
VII. Chemistry and Relativistic Effects	10		
A. Atomic and chemical properties of superheavy elements: Experiment and theory	11		
B. Bulk properties of superheavy elements	12		
VIII. Beyond $Z = 120$: Extended Periodic Table	12		
A. At and beyond the critical nuclear charge	13		
B. Nuclear physics and the limits of the periodic table	14		
IX. Cosmic Origin of Superheavy Nuclei	14		

I. INTRODUCTION

In 2012 and 2016, six new synthetic elements with atomic numbers $Z = 113$ (Nh, nihonium), $Z = 114$ (Fl, flerovium), $Z = 115$ (Mc, moscovium), $Z = 116$ (Lv, livermorium), $Z = 117$ (Ts, tennessine), and $Z = 118$ (Og, oganesson) joined the periodic table (Barber *et al.*, 2011; Karol *et al.*, 2016a, 2016b). They were all produced by scientists in the period 1998–2010 using heavy-ion fusion reactions. The road to their final confirmations took a dedicated experimental effort worldwide. But how significant were these recent additions? And how relevant is this to fundamental knowledge of nuclear and electronic shell structure? As we argue in this Colloquium, there are strong indications that the atoms and nuclei at the

*witek@frib.msu.edu

limits of mass and charge have the potential to transform and challenge our understanding of atomic and nuclear physics and chemistry.

The science drivers of superheavy element research are aligned with the overarching questions that motivate much of nuclear and atomic science (Freedman, 2012): How did visible matter come into being and how does it evolve? How does subatomic matter organize itself and what phenomena emerge? Are the fundamental interactions that are basic to the structure of matter fully understood? The specific questions asked in the context of superheavy elements are as follows:

- Where do nuclei and elements come from? Can superheavy nuclei be produced in the cosmos?
- How are superheavy nuclei and atoms made and organized? Do very long-lived superheavy nuclei exist in nature?
- What are the heaviest nuclei that can exist? Where does the periodic table of elements end?
- How can superheavy nuclei and atoms be exploited to reveal the fundamental symmetries of nature? What is the interplay between strong and electromagnetic interactions as the product of the fine-structure constant and atomic number approaches 1 ($\alpha Z \rightarrow 1$)?
- What are the chemical and physical properties of superheavy atoms?

The term “superheavy” usually refers to transactinides—chemical elements with $Z \geq 104$ (Seaborg and Loveland, 1990; Hoffman, Ghiorso, and Seaborg, 2000). All currently known superheavy nuclei are radioactive; they have been synthesized in nuclear reactions by scientists. The current superheavy territory consists of two disjoint areas. The *lower superheavy region* includes the lighter isotopes of elements $Z = 110$ – 113 , which were discovered during 1994–2004 in the so-called “cold-fusion” reactions involving lead and bismuth targets irradiated by medium-mass projectiles such as nickel and zinc (Hofmann and Münzenberg, 2000; Münzenberg and Morita, 2015). The rapid drop in production cross section with Z for cold-fusion reactions (it drops down to 0.02 pb for nihonium) suggested that a different approach to reach even higher atomic numbers was needed. The new strategy has relied on “hot-fusion” reactions with the rare ^{48}Ca beam and actinide targets. Relatively large cross sections (in the range of picobarns) associated with those reactions have enabled the discovery of over 50 isotopes of new elements with $Z = 114$ – 118 during 1998–2010 (Oganessian and Utyonkov, 2015a; Oganessian, Sobiczewski, and Ter-Akopian, 2017). These new isotopes form the *upper superheavy region*, which is currently not connected to the lower superheavy region via known nuclear decay chains.

Theoretically, it had been concluded by the end of the 1960s (Myers and Swiatecki, 1966; Sobiczewski, Gareev, and Kalinkin, 1966; Nilsson *et al.*, 1969; Nix, 1972a) that the existence of the heaviest nuclei with $Z > 104$ was primarily determined by shell effects due to the quantum-mechanical motion of protons and neutrons inside the nucleus. These early calculations predicted the nucleus with $Z = 114$, $N = 184$ to be the center of an island of long-lived superheavy nuclei with lifetimes ranging from minutes to millions of years, and this is how the term “island of stability” was coined (Myers and Swiatecki, 1966; Viola and Seaborg, 1966). This suggestion stayed practically unchallenged until the late 1990s when more

refined models, based on realistic effective interactions, were applied to superheavy nuclei and revised early predictions.

The heaviest element studied chemically to date is flerovium. Its relatively long half-life, 1–2 s, enables chemical studies with several atoms/day, which marks the limit of chemistry today (Türler, Eichler, and Yakushev, 2015; Düllmann, 2017b). The nucleus ^{294}Og , produced in Dubna (Oganessian *et al.*, 2006) with a ~ 0.5 pb cross section, marks the current limit of nuclear charge and mass. It decays to ^{290}Lv by α decay with a half-life of 0.58 ms (Brewer *et al.*, 2018), which is currently too short for “atom-at-a-time” chemical studies. As a result, computing its electronic and nuclear structure is the only option currently available to study such elements.

There have been a number of reviews on superheavy element research (Möller and Nix, 1994; Hofmann and Münzenberg, 2000; Ćwiok, Heenen, and Nazarewicz, 2005; Hofmann, 2015; Oganessian and Utyonkov, 2015a; Schädel, 2015; Oganessian, Sobiczewski, and Ter-Akopian, 2017) as well as more popular overviews (Oganessian and Rykaczewski, 2015; Düllmann and Block, 2018; Nazarewicz, 2018). The purpose of this Colloquium is to offer a broad perspective on the field, focused on theoretical expectations.

II. THE SUPERHEAVY TERRITORY

The superheavy nuclei occupy the upper region of the table of nuclides. As shown in Fig. 1, the known superheavy nuclei lie on the left flank of the valley of beta stability. Since those are all proton-rich systems, they can in principle decay by means of electron capture or β^+ decay; see, e.g., Heßberger *et al.* (2016). So far, however, no weak decay modes have been observed in the upper superheavy region. The main observed decay modes are alpha decay and spontaneous fission (SF) (Oganessian and Utyonkov, 2015a; Oganessian, Sobiczewski, and Ter-Akopian, 2017), and this is consistent with theoretical estimates (Ćwiok *et al.*, 1996; Möller, Nix, and Kratz, 1997; Sheng *et al.*, 2014; Heenen *et al.*, 2015).

As seen in Fig. 1, the territory of superheavy isotopes is vast and primarily unexplored, except for a small proton-rich corner. The isotopic chain of Og is a case in point. According to nuclear density functional theory (DFT) (Bender, Heenen, and Reinhard, 2003) based on realistic energy density functionals (EDFs) (Erler, Birge *et al.*, 2012; Agbemava *et al.*, 2014), about 100 isotopes of Og are expected to exist between $N \approx 170$ (proton drip line) and $N \approx 270$ (neutron drip line). The center of the β -stability valley for Og is predicted at $N \approx 192$ (Ćwiok *et al.*, 1996; Möller, Nix, and Kratz, 1997; Sheng *et al.*, 2014; Heenen *et al.*, 2015). Since, as discussed later, there are currently no obvious ways to synthesize neutron-rich superheavy systems, all information about those nuclei must come from theoretical modeling that involves large extrapolations.

As far as atomic territory is concerned, the currently known superheavy elements belong to the seventh period of the periodic table, with Og completing it (Reedijk, 2018). Og belongs to group 18, but, as we shall discuss in Sec. VII, strong relativistic effects make it a very unusual addition to the periodic table. Can the table of nuclides and periodic table of elements be further extended in charge and mass? This question will be addressed in Sec. VIII through various theoretical scenarios.




FIG. 1. Landscape of nucleon-bound nuclei in the (Z, N) plane. The stable isotopes are shown as black squares and those known experimentally are marked in green. Particle drip lines predicted by theory are shown together with their uncertainties (red). The upper inset shows the details of the superheavy region ($Z > 104$ and $N > 160$). The isotopes synthesized in heavy-ion fusion reactions are indicated (Oganessian and Utyonkov, 2015a; Oganessian, Sobczewski, and Ter-Akopian, 2017) together with the anticipated valley of β stability (Ćwiok *et al.*, 1996; Möller, Nix, and Kratz, 1997; Sheng *et al.*, 2014; Heenen *et al.*, 2015). The lower inset marks the placement of the recently named six superheavy elements in the periodic table. The element oganesson completes the seventh period of the periodic table. It is the only element of group 18 of the periodic table (noble gases), which does not naturally occur. Adapted from Nazarewicz, 2018.

III. PROTON AND NEUTRON DENSITIES AND SHELL STRUCTURE

Nucleonic density distributions and their moments contain key information on bulk properties of nuclei. However, direct experimental information on proton and neutron distributions and their moments in superheavy nuclei is nonexistent because of their very short lifetimes. In this respect, experimental laser spectroscopy studies of nuclear charge radii and quadrupole moments, recently extended to $^{252-254}\text{No}$




FIG. 2. Differential root mean square charge radii for Th, U, Pu, Cm, and No isotopes. The measurements (filled symbols) are compared to nuclear DFT calculations with two EDF models: UNEDF1 (dashed line) and SV-min (solid line). Adapted from Raeder *et al.*, 2018.

($Z = 102$), carry great promise (Düllmann and Block, 2018; Raeder *et al.*, 2018); see also Backe *et al.* (2015), Laatiaoui *et al.* (2016), and Ferrer *et al.* (2017). However, we still have a long way to go to reach higher atomic numbers in such studies. For the time being, the new data can be used to test the predictive power of current nuclear models when it comes to charge radii in the actinide nuclei. This is done for two different DFT models in Fig. 2. The results are encouraging as they show that DFT predictions in the region of very heavy nuclei are reliable.

Figure 3 shows density profiles for three Og isotopes. An intriguing feature is the depression of proton density in the nuclear interior. This effect, referred to as semibubble distribution, is driven in superheavy nuclei by strong Coulomb repulsion and manifests itself foremost in heavy nuclei above Pb ($Z = 82$). Speculations on such exotic nuclear structures (bubbles, tori, rings) came up rather early in connection with superheavy nuclei; see, e.g., Davies, Wong, and Krieger (1972), Wong (1973), Dietrich and Pomorski (1998), Dechargé *et al.* (1999), Schuetrumpf, Nazarewicz, and Reinhard (2017), and Staszczak, Wong, and Kosior (2017). The origin of such exotic geometries of nuclear densities is the mismatch of ranges of two opposing forces: the short-range attractive nuclear interaction and the long-range Coulomb repulsion. This competition, known as Coulomb frustration, is also known to produce a great variety of involved phases in bulk nuclear matter, coined nuclear “pasta” (Bonche and Vautherin, 1981; Williams and Koonin, 1985; Pethick and Ravenhall, 1995; Sonoda *et al.*, 2008; Fattoyev, Horowitz, and Schuetrumpf, 2017). The situation in finite superheavy




FIG. 3. Proton (left) and neutron (right) densities of ^{294}Og (top), ^{302}Og (middle), and ^{326}Og (bottom) calculated with SV-min in the (x, z) plane at $y = 0$. The densities are normalized to the maximum density. The central depression of the proton density (semibubble structures) is clearly seen in all three cases. From Schuetrumpf, Nazarewicz, and Reinhard, 2017.

systems is more subtle. The leading mechanism to release Coulomb pressure is nuclear fission; see Sec. VI. Superheavy nuclei can be sufficiently long lived only by virtue of extra binding through shell effects, and this puts a stringent limit on the existence of exotic geometries. The example considered in Fig. 3 is on the safe side in this respect: semibubbles in $^{294,326}\text{Og}$ represent stable configurations in spite of significant quadrupole deformations. [It is interesting to note that the charge distributions of $^{252,254}\text{No}$ are expected to have appreciable central depressions (Raeder *et al.*, 2018). The computed charge radii shown in Fig. 2 take this effect into account.] For discussion of shell effects related to semibubble configurations, see Afanasjev and Frauendorf (2005).

As mentioned in the Introduction, quantum shell stabilization is the basis for the existence of superheavy systems. Early studies of shell structure in superheavy nuclei used empirical shell model potentials (Myers and Swiatecki, 1966; Sobiczewski, Gareev, and Kalinkin, 1966; Mosel and Greiner, 1969; Nilsson *et al.*, 1969; Sobiczewski *et al.*, 1971; Nix, 1972b); see Ćwiok *et al.* (1996) for an overview. These studies were focused on defining new “magic shell closures” beyond ^{208}Pb . However, as was realized later through more microscopic studies, in which the Coulomb interaction was treated self-consistently, the shell structure of superheavy nuclei differs significantly from that of lighter species.

As discussed by Ćwiok *et al.* (1996), Bender *et al.* (1999), Bender, Nazarewicz, and Reinhard (2001), Berger, Hirata, and Girod (2003), Afanasjev and Frauendorf (2005), Shi *et al.* (2014), and Agbemava *et al.* (2015), the general pattern of single-particle energies undergoes significant changes in superheavy systems. Figure 4 shows the single-particle energies for the doubly magic nucleus ^{132}Sn and superheavy nuclei ^{302}Og and $^{472}164$ predicted with the Skyrme-DFT models SV-min and UNEDF1. (It should be emphasized that while the details of shell structure in superheavy nuclei show strong model dependence, general features are robust.) While for ^{132}Sn the magic gaps are sizable, around 4 MeV, the neutron gap $N = 184$ for ^{302}Og is about half that size. This is even more pronounced for the proton spectra where well-isolated shell gaps can hardly be found. First, the level density of single-particle states is large; in fact it grows faster than expected from oscillator scaling $A^{1/3}$ (Agbemava *et al.*, 2015). Consequently, small changes in theoretical modeling can impact shell structure substantially. Second, the shell structure of superheavy nuclei is influenced by Coulomb frustration and the concept of magic shell gaps fades away. This phenomenon of shell diffusion has also been analyzed by Jerabek *et al.* (2018) where it was demonstrated that the nucleon localization functions of superheavy nuclei approach the Fermi-gas limit in the valence region.

As seen in Fig. 4, high- j shells tend to cluster, which results in regions of quantum stabilization. Actually, the presence of a central depression strongly affects high- j orbits due to their large single-particle radii (Ćwiok *et al.*, 1996; Dechargé *et al.*, 1999, 2003; Afanasjev and Frauendorf, 2005; Pei, Xu, and Stevenson, 2005). The proton level scheme for ^{302}Og in Fig. 4 shows a few low- j states near the Fermi level, which inhibits strict shell closure but still maintains a low level density (due to low multiplicities), hence large shell stabilization. On the other hand, the low- j intruders reduce the impact of the $i_{13/2}$ state to drive bubble shapes. The situation looks extremely favorable for the hypothetical superheavy nucleus $^{472}164$ where high- j states cluster around the Fermi level. Its proton density shows a well-developed semibubble with a central density coming down to 66% of the maximum density (Schuetrumpf, Nazarewicz, and Reinhard, 2017).

In spite of the fact that current self-consistent nuclear models do not have spectroscopic quality when it comes to the description of known levels in heavy nuclei, most of those models predict shell-stabilized regions at $N \approx 172\text{--}184$ and $Z \approx 112\text{--}126$ (Ćwiok *et al.*, 1996; Bender *et al.*, 1999; Kruppa *et al.*, 2000; Bender, Nazarewicz, and Reinhard, 2001; Agbemava *et al.*, 2015) as well as around $N = 258$, $N = 308$, and $N = 410$ (Bender, Nazarewicz, and Reinhard, 2001; Zhang *et al.*, 2005; Afanasjev, Agbemava, and Gyawali, 2018) [see also Denisov (2005) and Ismail *et al.* (2016) for estimates based on empirical average potentials].

The importance of Coulomb pressure increases with increasing system size. This favors a reduction of nuclear density around the center and so may give way to exotic nuclear topologies such as bubbles or toroids. Figure 5 illustrates the competition between various configurations in the superheavy nucleus $^{780}254_{526}$ (Nazarewicz *et al.*, 2002) involving normal profiles (similar to densities of stable nuclei), bubbles which have a void at the center, and bandlike




FIG. 4. Proton (top) and neutron (bottom) Hartree-Fock single-particle energies relative to the Fermi level ϵ_F predicted with the Skyrme-DFT models SV-min and UNEDF1 for ^{132}Sn (left), ^{302}Og (middle), and $^{472}164$ (right). The line thickness is proportional to the orbital's degeneracy $2j + 1$ and the numbers indicate shell model occupations. While the details of the shell structure in superheavy nuclei show strong model dependence, general features are robust.

toroids. In many cases, such exotic profiles are predicted to be energetically competitive. However, their stability in superheavy nuclei remains an open question. Some recent calculations suggest that many such forms are unstable against triaxial distortions and fission (Brodziński and Skalski, 2013; Afanasjev, Agbemava, and Gyawali, 2018).

Although questions related to exotic configurations in $Z \gg 120$ systems may seem hardly relevant from the perspective of laboratory experiments, they can gain relevance in the exotic (neutron-rich) atmosphere of stellar explosions or in neutron stars, where exotic forms of nuclear matter are expected (Schuetrumpf *et al.*, 2015). In any case, such massive extrapolations require models which are well tested against information from the heaviest nuclei which are experimentally accessible.

IV. COLLECTIVITY

There are numerous indications that many superheavy nuclei are deformed in their ground states. Theoretical predictions of quadrupole deformations for the transfermium nuclei summarized by Afanasjev and Abdurazakov (2013) and Heenen *et al.* (2015) are consistent with quadrupole moment measurements recently extended to $^{252-254}\text{No}$ (Raeder *et al.*, 2018) and observed large moments of inertia (Herzberg and Greenlees, 2008; Greenlees *et al.*, 2012). As discussed in Sec. V, energies of α particles emitted by superheavy nuclei are impacted by deformation effects. For instance, the measured α -decay energies have provided confirmation of the prolate-deformed subshell closure around




FIG. 5. Coexisting configurations associated with different density distributions calculated in the SLy6 Skyrme-DFT model for the hypothetical superheavy nucleus $^{780}_{254}_{526}$. Three topologies are considered: normal nuclear densities similar to those found in stable nuclei, bubble nuclei distinguished by a substantial dip at the center, and band configurations out of a thin band of nuclear matter wound up to a torus. The contour plots of the total densities are given in the boxes, the arrows connect the profile with the corresponding point on the energy surface, and the colors help to further relate the topologies. All contours are axially symmetric (around the z axis) and the equidensity lines in the contour plots lie at 0.01, 0.03, 0.06, 0.09, 0.12, and 0.15 nucleons/ fm^3 . Adapted from Nazarewicz *et al.*, 2002.

$N = 162$ – 164 predicted by theory (Ćwiok *et al.*, 1983; Möller and Nix, 1994). The transition toward spherical shapes happens around $N = 184$. Located in between is a transitional region of deformation-soft and triaxial nuclei, exhibiting rich shape isomerism (Bürvenich *et al.*, 1998; Ren, 2002; Ćwiok, Heenen, and Nazarewicz, 2005; Prassa *et al.*, 2012; Agbemava *et al.*, 2015; Heenen *et al.*, 2015; Jachimowicz, Kowal, and Skalski, 2017b). The softness of the potential energy surface (PES) generally grows with the level density at the Fermi level, due to reduced shell effects. Consequently, as the level density increases with increasing system size, deformation softness and shape coexistence become important issues in superheavy nuclei. In addition to that rich deformation scenario, there is another type of coexistence in deformed superheavy nuclei, related to K isomerism (Xu *et al.*, 2004; Liu, Walker, and Xu, 2014; David *et al.*, 2015; Jachimowicz, Kowal, and Skalski, 2015).

As a representative example, the PESs of Og isotopes are shown in Fig. 6. The upper panel of the figure shows a two-dimensional PES in the triaxial quadrupole deformation plane. There are two axial competing minima, prolate ($\gamma = 0^\circ$) and oblate ($\gamma = 60^\circ$) separated by a triaxial barrier. The sequence of the axial PESs in the lower panel illustrates the transition from a well-deformed $^{290}\text{Og}_{172}$, over the transitional $^{294}\text{Og}_{176}$, to the spherical $^{302}\text{Og}_{184}$. Increasing the neutron number further, the PESs again become extremely soft and the barriers become so low that these isotopes are expected to have extremely short lifetimes. While topological features of PESs




FIG. 6. Top: Triaxial PES in the Q_{20} - Q_{22} plane for ^{290}Og obtained with the SLy4 Skyrme-DFT model. Adapted from Ćwiok, Heenen, and Nazarewicz, 2005. Bottom: Axial PESs for selected Og isotopes obtained with the Skyrme functionals SLy4, SV-min, and UNEDF1. All models predict very similar PES topology.

are robust, the barrier heights depend sensitively on both model and parametrization. As a consequence, the predictions for fission lifetimes can differ by many orders of magnitude; see Brodziński and Skalski (2013) and Agbemava *et al.* (2017) and Sec. VI. The span of predictions can be reduced by establishing connections with other observables better accessible to experiment; see, e.g., Reinhard (2018).

Figure 7 shows the predicted energies of the low-lying 2^+ collective states. The trend confirms again what we have seen in Fig. 6. Namely, the spherical isotopes around $N \approx 184$ show larger energies while the rather soft PESs farther from $N = 184$ yield significantly lower 2^+ excitations. Note that these excitation energies are rather low as compared to the 2^+ states in lighter nuclei. For more discussion of collective excited states in superheavy nuclei, see Mişicu *et al.* (2002), Sharipov




FIG. 7. Excitation energies of the 2^+ collective state computed for the three parametrizations as in Fig. 6 with the method from Klüpfel *et al.* (2008).




FIG. 8. Nuclei with ground-state octupole deformations predicted with different models. Adapted from Agbemava and Afanasjev, 2017.

and Ermamatov (2005), Sun *et al.* (2008), and Baran and Staszczak (2013).

Ground-state reflection-asymmetric shapes characterized by octupole deformations are well known in light actinides around $Z = 90$ and $N = 132$ (Butler and Nazarewicz, 1996). They appear as a result of the $\Delta\ell = \Delta j = 3$ coupling between the proton $0i_{13/2}$ and $1f_{7/2}$ and the neutron $0j_{15/2}$ and $1g_{9/2}$ shells. As shown in Fig. 8, the next regions of strong octupole coupling are expected around $N = 196$ ($0k_{17/2}$ and $1h_{11/2}$ neutron shells), $Z = 96$ ($0i_{13/2}$ and $1f_{7/2}$ proton shells), and $Z = 120$ ($0j_{15/2}$ and $1g_{9/2}$ proton shells) (Möller *et al.*, 1995, 2008; Erler, Langanke *et al.*, 2012; Warda and Egido, 2012; Agbemava and Afanasjev, 2017; Afanasjev, Abusara, and Agbemava, 2018). However, since the energies of the octupole-driving shells show considerable model dependence, boundaries and extensions of the predicted octupole deformed regions of heavy actinides and superheavy nuclei differ in detail. As discussed in Sec. III, such a situation is typical to superheavy nuclei due to the diffused shell structure (Bender, Nazarewicz, and Reinhard, 2001). In any case, reflection-asymmetric ground states are expected to appear in certain regions of superheavy nuclei and can potentially impact their lifetimes (Erler, Langanke *et al.*, 2012; Warda and Egido, 2012; Jachimowicz, Kowal, and Skalski, 2017b).

Nuclear structure observables strongly depend on the underlying shell structure. Unfortunately, current empirical shell model potentials as well as self-consistent DFT models do not produce robust, consistent explanations of the steadily growing body of spectroscopic information on transactinides (Rudolph *et al.*, 2013, 2015; Ackermann, 2015; Gates *et al.*, 2015; Theisen *et al.*, 2015; Herzberg, 2016; Heßberger, 2016; Ackermann and Theisen, 2017). This strong model dependence can be attributed to large single-particle level density and the fact that theoretical input still lacks spectroscopic quality (Shi, Dobaczewski, and Greenlees, 2014; Shi *et al.*, 2014; Dobaczewski *et al.*, 2015).

V. α DECAY

As discussed in Sec. II, the dominant decay modes of superheavy nuclei are α decay and spontaneous fission. As a result, superheavy nuclei are often identified through α -decay chain analysis (Hofmann and Münzenberg, 2000; Oganessian *et al.*, 2006). Information on the daughter decays comes from both experimental data and theoretical calculations of the energy of the emitted alpha particle, or Q_α values. In order to properly tune future experiments for the identification of new superheavy nuclei, theoretical predictions of Q_α values are essential.

Numerous calculations of Q_α energies for superheavy nuclei have been performed with microscopic-macroscopic and DFT methods (Ćwiok, Nazarewicz, and Heenen, 1999; Bender, 2000; Muntian *et al.*, 2003; Typel and Brown, 2003; Ćwiok, Heenen, and Nazarewicz, 2005; Gambhir, Bhagwat, and Gupta, 2005; Erler, Langanke *et al.*, 2012; Warda and Egido, 2012; Jachimowicz, Kowal, and Skalski, 2014; Agbemava *et al.*, 2015; Wang *et al.*, 2015; Tolokonnikov *et al.*, 2017; Adamian, Antonenko, and Lenske, 2018); comparisons between the results of these methods have also been carried out (Sobiczewski and Pomorski, 2007; Heenen *et al.*, 2015).




FIG. 9. Q_α values along the α -decay chain of ^{294}Og computed in nuclear DFT with different energy density functionals. Experimental and empirical extrapolated values (Wang *et al.*, 2017) are marked by stars. In ^{286}Fl , α decay competes with fission and the latter takes over in ^{282}Cn (Brewer *et al.*, 2018); hence, direct experimental information on Q_α values is not available below ^{282}Cn .

Microscopic calculations of α -decay half-lives are demanding since the α -spectroscopic (or preformation) factor is difficult to compute (Id Betan and Nazarewicz, 2012). The commonly used strategy is to estimate the barrier penetrability by adjusting a core + α potential to reproduce experimental Q_α values and estimate the reduced widths from measured lifetimes in known nuclei. This approach has resulted in various empirical formulas (Viola and Seaborg, 1966; Brown, 1992; Muntian *et al.*, 2003; Parkhomenko and Sobczewski, 2005; Chowdhury, Samanta, and Basu, 2008; Royer and Zhang, 2008; Dong, Zuo, and Scheid, 2011; Koura, 2012; Budaca, Budaca, and Silisteanu, 2016; Santhosh and Nithya, 2018b) and models (Ward, Carlsson, and Åberg, 2015; Clark and Rudolph, 2018) for the α -decay half-life.

In general, predicted Q_α values are fairly robust in the regions away from potential shell closures (Sobczewski and Pomorski, 2007; Heenen *et al.*, 2015). This is illustrated in Fig. 9 for the α -decay chain of ^{294}Og using several Skyrme-DFT models. For Fl and Lv, the results are within the bounds of the EDFs, while the results for Og are all underestimated. Overall, we observe a reasonable consistency between theoretical predictions. The increase in Q_α up to Cn followed by a sudden drop at Fl is explained by the abrupt shape transition from prolate to oblate deformations (see Fig. 10) through the region of triaxial shapes around $N = 174$ (Ćwiok, Heenen, and Nazarewicz, 2005); see Sec. IV for more discussion.

Figure 10 shows the Q_α values and regions of quadrupole deformations (prolate, oblate, spherical) of the even-even isotopes from Fm to Og predicted with the UNEDF2 model. The shape transition from oblate to spherical shapes is due to the predicted region of spherical shapes and enhanced shell-stability near $N = 184$ (Ćwiok *et al.*, 1996; Bender *et al.*, 1999; Kruppa *et al.*, 2000); see also Secs. III and IV. The large gap in values between Hs and Ds is due to the deformed neutron subshell closure around $N = 164$ (Ćwiok *et al.*, 1983;




FIG. 10. Q_α values of the even-even isotopic chains from Fm to Og for the UNEDF2 Skyrme-DFT model. Regions of different shapes (prolate, oblate, spherical) are marked. Experimental values (Oganessian and Utyonkov, 2015b; Wang *et al.*, 2017) are represented by circles and match the color of the corresponding theoretical predictions (lines).

Möller and Nix, 1994). Given the large extrapolations in mass and charge involved, the agreement between experiment and theory is reasonable. In the future, machine learning techniques relying on the statistical modeling of deviations between experiment and predictions (Neufcourt *et al.*, 2018) have the potential to improve the quality of theoretical extrapolations of Q_α values to unknown superheavy nuclei.

For α decays involving odd A , odd-odd nuclei, and isomeric states, half-lives can be significantly increased due to hindrance associated with configuration changes between parent and daughter nuclei (Ćwiok, Nazarewicz, and Heenen, 1999; Xu *et al.*, 2004). Estimates of this hindrance can be found in Delion, Liotta, and Wyss (2007), Ward, Carlsson, and Åberg (2015), Clark and Rudolph (2018), and Jachimowicz, Kowal, and Skalski (2018). In this context, experimental studies of high- K isomers in deformed superheavy nuclei (Hofmann *et al.*, 2001; Robinson *et al.*, 2011; Sulignano *et al.*, 2012; Ackermann, 2015; Asai, Heßberger, and Lopez-Martens, 2015; David *et al.*, 2015) are of great interest.

VI. FISSION

Although α decay is the dominant decay mode for all superheavy nuclei which have been observed through their associated α -decay chains, SF often competes with it and terminates the chain. For example, the observed isotopes for Fl and Ts undergo SF after several α emissions (Khuyagbaatar *et al.*, 2014; Utyonkov *et al.*, 2018). Other, more neutron-rich superheavy nuclei are predicted to decay directly via SF (Staszczak, Baran, and Nazarewicz, 2013; Afanasjev, Aghemava, and Gyawali, 2018; Giuliani, Martínez-Pinedo, and Robledo, 2018) (see Fig. 11).

Accurate modeling of nuclear fission—a quantum-mechanical process involving large-amplitude nuclear collective motion—poses enormous challenges to nuclear theory




FIG. 11. Dominant decay modes for even-even superheavy elements in the range $108 \leq Z \leq 126$, $148 \leq N \leq 188$ predicted with the Skyrme-DFT model SkM*. Spontaneous fission half-lives were computed using the semiclassical approximation while α -decay half-lives were estimated using the Viola-Seaborg formula (Viola and Seaborg, 1966). The contours show the predicted half-lives in logarithmic scale. Adapted from Staszczak, Baran, and Nazarewicz, 2013.

(Schunck and Robledo, 2016). To give just two examples: spontaneous fission half-lives in heavy and superheavy elements span a range of over 35 orders of magnitude (Krappe and Pomorski, 2012), while at the same time, small variations of a few percent of specific fission product yields can have a dramatic impact in modeling critical assemblies in nuclear reactors (Gooden *et al.*, 2016). While the fission of actinide nuclei is, at least semiquantitatively, relatively well understood, superheavy nuclei offer specific challenges. Because of their large atomic numbers, the fissility parameter $x = E_{\text{Coul}}/2E_{\text{surf}}$ takes values greater than 1 in the superheavy region, inner fission barriers are generally smaller than in actinides, and outer fission barriers are expected to be low or nonexistent due to increasing Coulomb pressure. Consequently, the familiar picture of penetration through a double-humped fission barrier (Bjørnholm and Lynn, 1980) undergoes serious revisions in the superheavy region. These revisions have an impact on fission observables such as the SF lifetimes, the distribution of fission fragments, and the characteristics of the fission spectrum (neutrons, γ , β , etc.).

Theoretical estimates of SF lifetimes for superheavy nuclei depend strongly on the computed PESs (see also Sec. IV). A considerable effort has been devoted to systematic predictions of PESs and their saddle points (fission barriers) in the superheavy region (Bürvenich *et al.*, 1998, 2004; Möller *et al.*, 2009; Karatzikos *et al.*, 2010; Kowal, Jachimowicz, and Sobieczewski, 2010; Abusara, Afanasjev, and Ring, 2012; Erler, Langanke *et al.*, 2012; Prassa *et al.*, 2012; Staszczak, Baran, and Nazarewicz, 2013; Jachimowicz, Kowal, and Skalski, 2015; Agbemava *et al.*, 2017; Pomorski *et al.*, 2018), including structures of excited configurations in compound nuclei (Pei *et al.*, 2009; Sheikh, Nazarewicz, and Pei, 2009; Zhu and Pei, 2017); see also Baran *et al.* (2015), Heßberger (2017), and Reinhard (2018) for recent overviews.

Spontaneous fission lifetimes of superheavy nuclei are typically calculated by estimating the tunneling probability through the multidimensional PES (Brack *et al.*, 1972). While there have been attempts to do this within the macroscopic-microscopic approach (Brack *et al.*, 1972; Möller, Nix, and Swiatecki, 1989; Smolańczuk, Skalski, and Sobieczewski, 1995;

Gherghescu *et al.*, 1999), a more microscopic description based, e.g., on nuclear DFT, provides more consistency since all of the ingredients for the calculations (PES and collective inertia tensor) are derived from a single energy functional (Baran *et al.*, 2011; Giuliani and Robledo, 2013). Such microscopic calculations of SF lifetimes have become increasingly predictive in recent years (Schindzielorz *et al.*, 2009; Erler, Langanke *et al.*, 2012; Warda and Egido, 2012; Staszczak, Baran, and Nazarewicz, 2013; Sadhukhan, Nazarewicz, and Schunck, 2016). In this context, the prediction of SF lifetimes depends upon several factors, such as the nuclear zero-point energy, the number of collective coordinates considered, and nuclear pairing (Giuliani, Robledo, and Rodríguez-Guzmán, 2014; Sadhukhan *et al.*, 2014; Zhao *et al.*, 2016; Rodríguez-Guzmán and Robledo, 2017; Reinhard, 2018).

Figure 11 shows a comprehensive theoretical survey of the dominant decay modes of superheavy nuclei calculated within the Skyrme-DFT framework (Staszczak, Baran, and Nazarewicz, 2013) [see also Erler, Langanke *et al.* (2012) and Warda and Egido (2012) for other results]. A region of superheavy isotopes with very short SF lifetimes is predicted in a corridor separating the upper and lower regions of superheavy nuclei. This is consistent with systematics of experimental SF lifetimes (Oganessian and Utyonkov, 2015a), which show the minimum of SF lifetimes at ^{282}Cn .

Similar to the case of α decay discussed in Sec. V, SF lifetimes are expected to be substantially increased in many-quasiparticle states of superheavy nuclei, such as levels of odd- A nuclei or low-lying K isomers (Xu *et al.*, 2004; Liu, Walker, and Xu, 2014; David *et al.*, 2015; Jachimowicz, Kowal, and Skalski, 2015; Rodríguez-Guzmán and Robledo, 2017). This opens an intriguing possibility of the existence of very long-lived isomers of superheavy isotopes.

An interesting, albeit yet experimentally unexplored, decay mode of superheavy nuclei is cluster radioactivity. This phenomenon, first observed in 1984 (Rose and Jones, 1984) as the emission of ^{14}C from ^{223}Ra , has since been confirmed in a number of heavy nuclei with $Z > 86$, which decay by emitting light clusters between ^{14}C and ^{34}Si , as well as in superheavy nuclei with $Z \leq 116$ (Kozulin *et al.*, 2014; Itkis *et al.*, 2015). For superheavy nuclei with $Z \geq 118$, cluster radioactivity is expected to become competitive with alpha decay and spontaneous fission (Poenaru, Gherghescu, and Greiner, 2011, 2012, 2013; Poenaru *et al.*, 2015; Poenaru and Gherghescu, 2018; Santhosh and Nithya, 2018a; Zhang and Wang, 2018). Microscopically, cluster emission can be considered as an extremely asymmetric fission, with the heavy fragment corresponding to a nucleus in the neighborhood of the doubly magic ^{208}Pb (Warda and Robledo, 2011). Nuclear DFT calculations (Warda and Robledo, 2011; Warda, Zdeb, and Robledo, 2018) of nuclei with the same N/Z ratio as known cluster emitters clearly show a lowering of the cluster-decay potential energy valley with respect to the fission valley as Z increases.

The nucleus ^{294}Og is expected to be an excellent candidate for cluster radioactivity as the emitted cluster is the magic nucleus ^{86}Kr with $N = 50$, and consequently the asymmetric channel is expected to be particularly favored (see Fig. 12) (Warda and Egido, 2012; Baran *et al.*, 2015; Matheson *et al.*, 2018). The corresponding SF half-life is predicted to be ~ 13 min (Staszczak, Baran, and Nazarewicz, 2013). This should be




FIG. 12. Potential energy surface of ^{294}Og in the (Q_{20}, Q_{30}) coordinates calculated in nuclear DFT using the functionals UNEDF1_{HFB} and D1S. Two SF pathways are indicated: nearly symmetric (symm) and strongly asymmetric corresponding to cluster decay (cluster). From Matheson *et al.*, 2018.

compared with the measured α -decay half-life of $580 \mu\text{s}$ (Brewer *et al.*, 2018).

VII. CHEMISTRY AND RELATIVISTIC EFFECTS

It was long assumed that electrons in valence shells responsible for chemical properties are not affected much by relativistic effects because valence electrons move rather slowly as compared to the velocity of light (Dirac, 1929; Glashow, 1988); i.e., $v_{\text{val}} \ll c$ should hold for all elements in the periodic table including the latest superheavy additions. While in a classical sense this is correct, quantum theory tells us that this simple picture is insufficient and that even “slow moving” valence electrons have a substantial electron density close to the nucleus where relativistic perturbation operators act. Low angular momentum wave functions, such as s orbitals, are therefore directly affected by these operators and are drawn inward by their tails (the direct relativistic effect) resulting in an orbital contraction and stabilization. In fact, it has been known since the 1970s that relativistic perturbation operators act almost exclusively in the K -shell region (Sobel’Man, 1972; Dzuba, Flambaum, and Sushkov, 1983; Schwarz *et al.*, 1989). This indicates that relativistic effects are intertwined throughout the whole electron cloud and cannot be argued away only by energetic arguments. In the following, we will address three important relativistic effects taking place in superheavy atoms.

The first relativistic effect is related to the $1s$ state and overall stability of the atom. Relativistic quantum theory for fermions is fundamentally based on the Dirac equation. For hydrogenlike atoms with a pointlike nucleus, the discrete spectrum of the Dirac operator is given by the Sommerfeld fine-structure formula:

$$\frac{E_{nk}}{mc^2} = \left(1 + \frac{(Z\alpha)^2}{[n - k + \sqrt{k^2 - (Z\alpha)^2}]^2} \right)^{-1/2}, \quad (1)$$

where $k = j + 1/2$. The positive- and negative-energy continuum corresponds to $E \geq mc^2$ and $E \leq -mc^2$, respectively.

We immediately see that for $Z \rightarrow 1/\alpha$ the ground-state energy $E_{1,1} \rightarrow 0$. Beyond this limit, the lowest energy eigenvalue becomes imaginary (the so-called $1s$ catastrophe). That is, the $1s$ level should be viewed as a resonance embedded in the negative-energy continuum. While this does not strictly hold for multi-electron systems, the screening of the nuclear charge by all electrons in an atom is not sufficient and will only slightly shift the $Z_c = 1/\alpha \approx 137$ limit. While this caused some concern in the relativistic electronic structure community more than 30 years ago, predicting that the Dirac equation fails in the superheavy element region, it was pointed out early (Pomeranchuk and Smorodinsky, 1945) that the finite size of the nucleus increases the value of critical nuclear charge at which the catastrophe occurs to $Z_c = 169$. In this case, however, for known functional forms of the nuclear charge distribution, analytical eigenvalues are no longer available (Andrae, 2000).

The second relativistic effect concerns contraction of outer shells. In addition to interpretational difficulties with the Dirac equation, it initially came as a surprise that the outer ns shells with higher principal quantum number n can undergo large relativistic contractions when compared to the inner ns shells with low n values (Desclaux and Pyykkö, 1976; Pitzer, 1979; Pyykkö and Desclaux, 1979) thus contradicting the original orthogonality argument that the outer shells contract only because of the inner-shell contractions (Pyykkö, 1988). However, these shell effects can be rather subtle and strongly dependent on the nuclear charge Z , quantum numbers $(n\ell j)$, and shell occupancies (Schwarz *et al.*, 1990; Autschbach *et al.*, 2002).

Figure 13(a) demonstrates this for the ratio $q_{n\ell j} = \langle r \rangle_{n\ell j}^{\text{R}} / \langle r \rangle_{n\ell j}^{\text{NR}}$ of the relativistic (R) to nonrelativistic (NR) expectation values for the element Cn ($Z = 112$; $[\text{Rn}]5f^{14}6d^{10}7s^2$ configuration). The rather large $7s$ valence shell contraction (stabilization) shown in the figure makes Cn chemically more inert compared to the lighter congener Hg (Pitzer, 1975; Gaston *et al.*, 2007; Eichler *et al.*, 2008;




FIG. 13. Ratio of the relativistic (R) to nonrelativistic (NR) expectation values of orbital radii for different Dirac-Hartree-Fock $(n\ell j)$ orbitals in Cn ($Z = 112$, top) and $Z = 120$ (bottom).

Steenbergen *et al.*, 2017), where relativistic effects are known to be large; they are responsible for Hg being the only elemental liquid metal at room temperature (Calvo *et al.*, 2013; Steenbergen, Pahl, and Schwerdtfeger, 2017). Relativistic effects are also responsible for changing the ground-state configuration of Rg ($Z = 111$) from $6d^{10}7s^1$ ($^2S_{1/2}$) to $6d^97s^2$ ($^2D_{5/2}$) and halving its atomic size, making Rg as small as copper in the same periodic group (Eliav *et al.*, 1994).

The rather large ns -shell contractions and stabilizations observed for the group 11 and 12 elements (compared, for example, to the group 1 and 2 elements) originate from the occupation of the lower-lying diffuse $(n-1)d$ shell, which does not efficiently screen the nucleus, thus increasing the effective nuclear charge Z_{eff} experienced by the valence ns electrons. Figure 13 shows the predicted rather large valence $7s$ shell contraction in Cn compared to the $8s$ shell contraction in element $Z = 120$. Moreover, direct shell contractions (mainly the ns and $np_{1/2}$ shells) lead to additional screening of the nucleus, which lowers the effective nuclear charge Z_{eff} and results in an expansion or destabilization of the more diffuse higher angular momentum states such as the f , g and $d_{5/2}$ orbitals (known as the indirect relativistic effect) (Pyykkö, 1988). The $p_{3/2}$ and $d_{3/2}$ orbitals may contract or expand depending on the nuclear charge and shell occupancy. A more detailed discussion about these subtle shell-structure effects can be found in Autschbach *et al.* (2002).

The third principal relativistic effect comes from the fine-structure splitting (mainly *spin-orbit coupling*) of orbitals with angular momentum $\ell > 0$ (Pyykkö, 1988). For Og^+ , the $^2P_{3/2} - ^2P_{1/2}$ splitting is 10.1 eV at the Fock-space coupled-cluster level (Jerabek *et al.*, 2018), i.e., it is larger than most bond dissociation energies. The relativistic $7p_{3/2}$ expansion and destabilization ($q_{7p_{3/2}} = 1.052$ for neutral Og) and the large relativistic $8s$ contraction and stabilization ($q_{8s} = 0.739$ for Og in the $7p_{3/2}^3 8s^1$ configuration) lead to a positive electron affinity of 0.064 eV (with a sizable QED correction of -0.006 eV) (Goedenko *et al.*, 2003), a novelty within group 18 of noble gases. Spin-orbit effects make flerovium a closed-shell atom ($[\text{Rn}]5f^{14}6d^{10}7s^2 p_{1/2}^2$ configuration) with zero electron affinity (Borschevsky *et al.*, 2009), and chemically more inert compared to the lighter group 14 elements (Pitzer, 1975). The large $p_{1/2}/p_{3/2}$ separation for all shells in a superheavy element is also reflected in the q_{np} values shown in Fig. 13.

A. Atomic and chemical properties of superheavy elements: Experiment and theory

The chemistry of superheavy elements and the influence of relativistic effects have been reviewed by, e.g., Schädel (2006, 2015), Türler and Pershina (2013), Pershina (2015), and Schwerdtfeger *et al.* (2015). In the following, we emphasize those chemical and physical properties of the superheavy elements that are strongly influenced by relativistic shell-structure effects leading to anomalies and often unexpected features when comparison is made to their lighter congeners in the periodic table (Schwerdtfeger and Seth, 1998). So far,

chemical experiments at the atom-at-a-time scale have only been carried out for transactinides up to Hs (Türler and Pershina, 2013; Nagame, Kratz, and Schädel, 2016) and for Cn and Fl (Eichler *et al.*, 2008, 2010; Yakushev *et al.*, 2014), which all have reasonably long-lived isotopes in the time range of seconds.

Experimental studies giving direct information about the electronic shell structure are available only up to the heaviest actinides including Fm, No, and Lr (Sewtz *et al.*, 2003; Sato *et al.*, 2015; Laatiaoui *et al.*, 2016; Chhetri *et al.*, 2017, 2018; Raeder *et al.*, 2018). However, optical lines of actinide atoms up to Es have been identified in the visual echelle spectrography of the main-sequence star HD 101065 (Przybylski's star) (Gopka *et al.*, 2008). It may thus be speculated that the spectra of neutron-rich superheavy elements created in neutron star mergers or supernovae can be observed (see Sec. IX). This requires accurate theoretical predictions of atomic spectra for superheavy elements, which currently is a major challenge for open-shell many-electron systems (Sewtz *et al.*, 2003). In this context, large isotope shifts and hyperfine structure in electronic spectra of superheavy elements has been investigated by Dzuba, Flambaum, and Webb (2017), comparing different isotopes measured in the laboratory (which are neutron poor) with those in interstellar space (which are mostly stable and neutron rich). The reach of laser spectroscopy measurements of isotope shifts and nuclear moments in the heaviest elements corresponds to $^{252-254}\text{No}$ (Raeder *et al.*, 2018). The electronic spectrum of neutral Og was computed by Lackenby, Dzuba, and Flambaum (2018).

Precision mass measurements of unstable atoms provide a direct measure of the nuclear and electronic binding energy. Direct mass measurements of transfermium nuclei have been carried out with the Penning trap mass spectrometer for $^{252-255}\text{No}$ and $^{255,256}\text{Lr}$ (Block *et al.*, 2010; Ramirez *et al.*, 2012). Recently mass excesses of several transfermium nuclei were obtained by means of time-of-flight mass spectrometry (Ito *et al.*, 2018), which enabled mass determination of heavier nuclei, up to ^{261}Bh and ^{266}Mt . The mass of an atom M_A can be decomposed into $M_A = Z(m_p + m_e) + Nm_n - (B_{\text{nuc}} + B_{\text{el}})/c^2$, where m_p , m_e , and m_n are, respectively, proton, electron, and neutron masses known to high precision, and B_{nuc} and B_{el} are the nuclear and electronic binding energies. Note that the electronic binding energy is not negligible for superheavy elements. For instance, for Og the Dirac-Coulomb Hartree-Fock calculations predict $B_{\text{el}} = 1.487$ MeV. At the nonrelativistic level the electronic binding energy is significantly smaller, $B_{\text{el}}^{\text{NR}} = 1.260$ MeV. The difference of 0.227 MeV originates mostly from the inner shells close to the nucleus (Jerabek *et al.*, 2018). Interestingly, the electron density gets smeared out due to relativistic effects, making the valence electron cloud of Og a Fermi-like gas as Fig. 14 illustrates.

Gas phase chromatographical setups can be easily coupled to accelerators and are ideal to study the volatility of superheavy elements (Gäggeler, 2011). The most recent adsorption studies of Cn and Fl on gold surfaces suggest that both elements bind weakly to the surface, i.e., $-\Delta H_{\text{ads}}(\text{Au})$ (in kJ/mol) = 98 ± 3 (Hg) (Soverna *et al.*, 2005), 52_{-3}^{+4} (Cn) (Eichler *et al.*, 2008), ≥ 48 (Fl) (Yakushev *et al.*, 2014), and 20 (Rn) (Soverna *et al.*, 2005). This is in good




FIG. 14. Electron localization functions C_e from nonrelativistic (NR, left) and Dirac-Hartree-Fock calculations (R, right) for the heavy rare gas atoms Xe (top), Rn (middle), and Og (bottom). From Jerabek *et al.*, 2018.

agreement with the latest theoretical studies (Perschina *et al.*, 2008a, 2008c; Pershina, Anton, and Jacob, 2009; Rampino, Storch, and Belpassi, 2015; Pershina, 2018). From an empirical relationship between $\Delta H_{\text{ads}}(\text{Au})$ and the cohesive energy of the bulk material E_{coh} , one estimates $E_{\text{coh}} = 38_{-12}^{+10}$ kJ/mol for Cn (Eichler *et al.*, 2008). This agrees with the recent incremental relativistic coupled-cluster theory (Steenbergen *et al.*, 2017), which yields $E_{\text{coh}} = 39.6$ kJ/mol. Consequently, Cn is predicted to be rather volatile (Pershina *et al.*, 2010) with an expected melting point around that of Hg, but perhaps being still a liquid at standard conditions. For Fl only very few atoms were identified on a gold surface, and the predictions are not yet accurate enough to determine $\Delta H_{\text{ads}}(\text{Au})$. Recent solid-state calculations using relativistic density functional theory predict $E_{\text{coh}} = 48.6$ kJ/mol (Hermann *et al.*, 2010), in rather good agreement with the experimental estimated value (Yakushev *et al.*, 2014). Thus one might speculate that Fl is liquid at room temperature. The possibility of future adsorption experiments of Nh on an inert surface has been explored theoretically (Pershina *et al.*, 2008b; Pershina, 2016).

Relativistic effects can have a decisive influence on the stability of oxidation states. For example, the relativistic $7s$ stabilization together with the indirect relativistic $6d_{5/2}$ destabilization supports efficient sd mixing in chemical bonding, thus stabilizing higher oxidation states in compounds of Rg (as high as the oxidation state +5 in RgF_5^-)

(Seth *et al.*, 1998), Cn (oxidation state +4 as in CnF_4) (Seth, Schwerdtfeger, and Dolg, 1997), and possibly also for Ds (Waber and Averill, 1974). In contrast, the strong relativistic $7s$ stabilization diminishes sp mixing in chemical bonding for the $7p$ block elements (known as the relativistic inert pair effect) (Schwerdtfeger *et al.*, 1992), and thus reduces the stability of high oxidation states in Nh and Fl (and possibly in Mc as well) (Seth, Faegri, and Schwerdtfeger, 1998; Seth, Schwerdtfeger, and Faegri, 1999; Schwerdtfeger and Seth, 2002). In contrast to XeF_6 , which has been isolated and identified (Hoyer, Emmeler, and Seppelt, 2006), the large spin-orbit splitting between the $7p_{1/2}$ and $7p_{3/2}$ levels will most likely restrict the oxidation state in Og to +4. Note that spin-orbit effects change the structure of OgF_4 from the expected D_{4h} symmetry to T_d (Nash and Bursten, 1999).

B. Bulk properties of superheavy elements

Thermodynamic properties of the bulk superheavy elements from Nh to Og have been predicted empirically by extrapolation by Bonchev and Kamenska (1981), and for Ts and Og by Takahashi (2002). From these it is expected that Fl and element 119 could be either liquid or solid under normal conditions, and Og is a gas. On the other hand, based on atomic polarizability arguments (Nash and Bursten, 1999) Og is likely to be solid at room temperature. Indeed, Og is calculated to have a rather high dipole polarizability ($\alpha_d = 58.0$ a.u.) as compared to the other rare gas elements, which increases dispersive interactions (Jerabek *et al.*, 2018). Methods are now available for the accurate determination of bulk properties including phase transitions for the rare gas elements (Pahl *et al.*, 2008; Schwerdtfeger and Hermann, 2009; Schwerdtfeger, 2016). For example, density functional theory predicts solid-state properties for the $6d$ metals from Lr to Rg (Gyanchandani and Sikka, 2011). According to these calculations, the next nearest neighbor distance in the solid is $R(5d) < R(6d)$ for the elements within the same group of the periodic table; hence, no evidence for strong relativistic effects is expected for this quantity in contrast to many other chemical properties (Pershina, 2002; Türlér and Pershina, 2013; Wang *et al.*, 2016; Iliáš and Pershina, 2017). Recently, using parallel tempering Monte Carlo simulations within a many-body expansion for the atomic interaction potential (Smits *et al.*, 2018), it was possible to confirm the melting temperature of $T_m = 202$ K of ^{222}Rn originally measured in 1909 (Gray and Ramsay, 1909). Preliminary simulations suggest $T_m \approx 320$ K for Og (Smits *et al.*, 2018). For Cn and Fl, many-body effects in the interaction potential are so important that melting simulations become prohibitively expensive [see recent successful simulations for Hg (Steenbergen, Pahl, and Schwerdtfeger, 2017)].

VIII. BEYOND Z = 120: EXTENDED PERIODIC TABLE

The first attempt to extend the periodic table beyond element 120 goes back to Mann (1969) and Mann and Waber (1970). Soon afterward Fricke, Greiner, and Waber (1971) pointed out that beyond element 120 the $7d_{3/2}$, $8p_{1/2}$, $6f_{5/2}$, and $5g_{7/2}$ shells are expected to be filled more or less simultaneously, making a clear placement in the periodic table difficult. They predicted the filling of the $5g$ shell to be complete at element 144, whereas a newer version of the

periodic table (Pyykkö, 2011, 2016) completes the g -shell filling at element 138. Clearly, accurate multireference configuration-interaction calculations are required to determine the correct ground state of the elements beyond $Z = 120$ (Lu *et al.*, 1971). Accurate Fock-space coupled-cluster calculations for the superheavy elements 121 and 122 (Eliav, Fritzsche, and Kaldor, 2015) show that the ground states are $^2P_{1/2}(8s^28p_{1/2})$ for element 121 and $8s^27d_{3/2}8p_{1/2}(J = 2)$ for element 122 (with the $8s^28p_{1/2}^2$ state being close in energy to $\Delta E = 0.185$ eV). Thus, according to Eliav, Fritzsche, and Kaldor (2015), the g shell is not occupied up to $Z = 122$. In earlier Hartree-Fock-Slater calculations, g -shell occupation starts at element 125 (Mann, 1969; Mann and Waber, 1970; Lu *et al.*, 1971). Note that for element 140 the ground-state configuration could be either $8s^28p^27d^16f^35g^{14}$ or $8s^28p^46f^15g^{15}$ according to Indelicato, Bieroń, and Jönsson (2011). Such calculations are already at the limit of available computer resources due to the large number of possible configurations involved. In spite of the difficulties of correctly placing superheavy elements into the periodic table, a new periodic table was obtained by Pyykkö (2011, 2016) from average level multireference Dirac-Fock calculations with successive occupation of electronic shells as shown in Fig. 15. The anomalies to the Madelung rule (Wong, 1979) shown in the figure are due to the strong spin-orbit stabilization of the $8p_{1/2}$ and $9p_{1/2}$ shells.

Little is known about the chemistry of the elements beyond $Z = 120$ (Fricke, Greiner, and Waber, 1971). The Dirac-Coulomb DFT calculations for the hexafluorides of the elements 125–129 (Dognon and Pyykkö, 2017) suggest g -level occupations for these compounds. However, besides the relativistic expansion of the $5g$ orbitals (for 140^{2+} with a full g shell one predicts $q_{5g_{7/2}} = 1.190$ and $q_{5g_{9/2}} = 1.394$), the g electrons do not participate in the chemical bonding and are therefore corelike, similar to the $4f$ electrons in the lanthanide series.




FIG. 15. Successive shell filling [according to Pyykkö (2011, 2016)] up to the heaviest noble gas element with $Z = 172$ according to the Madelung rule. The anomalies in expected shell filling are highlighted in blue and result from strong relativistic stabilization effects of the $p_{1/2}$ shells at high atomic numbers.




FIG. 16. Orbital energy dependence on nuclear charge Z for H-like atoms. The $1s_{1/2}$ orbit dives into the negative-energy continuum at $-2m_e c^2$ at Z_{crit} (positive-energy continuum set at zero energy). Extrapolations into the negative-energy continuum region are marked by dashed lines. Adapted from Schwerdtfeger *et al.*, 2015.

A. At and beyond the critical nuclear charge

As discussed in Sec. VII, the current electronic shell structure predictions end at elements with critical nuclear charge $Z_{\text{crit}} \sim 172$ – 173 , where the lowest $1s$ level enters the negative energy continuum (supercritical region) as predicted by a mean-field treatment of the Dirac equation (Pomeranchuk and Smorodinsky, 1945; Indelicato, Bieroń, and Jönsson, 2011; Schwerdtfeger *et al.*, 2015); see Fig. 16. In multi-electron systems, since the nucleus becomes screened by all the electrons including those in higher ($n\ell j$) shells, Z_{crit} shifts to slightly larger values. In this supercritical region, the small (lower) component of the Dirac spinor becomes appreciable as shown in Fig. 17, and the terms “small” and “large” for the two Dirac components are no longer valid. Near the critical nuclear charge, the single-particle approximation (e.g., Dirac-Coulomb-Hartree-Fock) breaks down and a proper treatment




FIG. 17. The fraction of the $1s$ electron density due to the small component $\Delta\rho_{1s}^S$ (dashed line) and the fraction of the $1s$ electron density contained within the nuclear radius $\Delta\rho_{1s}^{\text{nuc}}$ (solid line) as a function of nuclear charge Z for H-like atoms. Adapted from Schwerdtfeger *et al.*, 2015.

involves the full QED Lagrangian. Moreover, in strong Coulomb fields, the $Z\alpha$ expansion used to treat QED shifts for one-particle levels also breaks down (Soff *et al.*, 1982). Nonetheless, effective QED operators obtained at nuclear charges $Z \ll 137$ are usually continued into the highly non-perturbative domain (Indelicato, Bieroń, and Jönsson, 2011; Schwerdtfeger *et al.*, 2015). It was noted that QED corrections to the $1s$ level for $Z \sim 172$ are rather small compared to $2m_e c^2$ as self-energy and vacuum-polarization contributions almost cancel out in such supercritical fields (Soff *et al.*, 1982).

The extension of the periodic system into new areas has been investigated by Greiner (2008), who hypothesized that in the supercritical region with $Z \sim 172$ – 173 , where the $1s$ state becomes a resonance embedded in the negative-energy continuum of the Dirac equation, spontaneous e^+e^- pair creation becomes possible because of vacuum fluctuations (Gershtein and Zel'dovich, 1969; Pieper and Greiner, 1969; Reinhardt, Greiner, and Arenhövel, 1971). In a simple picture, an electron from the Dirac sea occupies the $1s$ state leaving a hole in the sea, which escapes as a positron while the electron's charge remains near the source. In this process, energy is conserved because the occupation of the $1s$ level frees up an energy of $> 2m_e c^2$. One can view this as the inverse pair annihilation, which is well described by ordinary Feynman diagrams. From the energy-time uncertainty principle one can estimate that such a process would happen on the time scale of $\sim 10^{-21}$ s. Since the $1s$ orbit is occupied by two electrons, the inverse pair annihilation can happen twice. However, once the $1s$ state becomes doubly occupied, the vacuum becomes stable due to the Pauli principle (Reinhardt, Müller, and Greiner, 1981; Müller-Nehler and Soff, 1994; Greiner, 2008). We note that the e^+e^- pair creation process has never been observed in a collision of atoms with high nuclear charges (Ahmad *et al.*, 1997, 1999) and may require collision experiments with fully stripped ions. The electron-positron pair creation in low-energy collisions of heavy bare nuclei has been investigated only recently (Maltsev *et al.*, 2015).

It is currently unclear how to accurately treat single- or multielectron systems beyond Z_{crit} . However, there are no compelling arguments suggesting that Z_{crit} should determine the end of the electronic shell stability and therefore the end of the periodic table, even though Pyykkö's analysis ends at Z_{crit} (Pyykkö, 2011).

B. Nuclear physics and the limits of the periodic table

The existence of elements with $Z > 118$ also depends on nuclear physics. For instance, as the $1s$ density starts to accumulate around the nucleus in strong Coulomb fields (see Fig. 17), the probability of electron capture increases. One might speculate that the most difficult hurdle to the stabilization of the electronic $1s$ shell as one gets closer to Z_{crit} , and ultimately the cause of the end of the superheavy elements, could be K capture (Schwerdtfeger *et al.*, 2015).

Another important factor that needs to be considered is the large difference between electronic and nuclear time scales. According to the report of the *Transferrmium Working Group* (Wapstra *et al.*, 1991), in order to talk about a new element, the corresponding nuclide with an atomic number Z must exist for at least 10^{-14} s, which is a reasonable estimate of the time it

takes a nucleus to acquire its outer electrons, bearers of the chemical properties. Consequently, if for all isotopes of some superheavy element, including isomeric states (Heenen *et al.*, 2015; Jachimowicz, Kowal, and Skalski, 2017a), nuclear lifetimes are shorter than 10^{-14} s, the corresponding element does not exist. On the other hand, in order to define a nuclide, its lifetime should be longer than the single-particle time scale $T_{\text{s.p.}} \approx 1.3 \times 10^{-22}$ s (Goldanskii, 1966; Thoennessen, 2004) that corresponds to the time scale needed to create the nuclear mean field. Consequently, there is no chemistry for nuclides with lifetimes between 10^{-14} and 10^{-22} s.

Unfortunately, reliable predictions of lifetimes of nuclei with $Z > 118$ are currently not available, primarily due to difficulties related to the assessment of Coulomb frustration effects on fission. Indeed, since it is difficult to say at present whether the exotic topologies of nuclear density illustrated in, e.g., Fig. 5, can occur as long-lived (metastable) states, the limits of nuclear mass and charge are presently unknown (Brodziński and Skalski, 2013; Afanasjev, Agbemava, and Gyawali, 2018). It is thus possible that the periodic table will end well before reaching Z_{crit} for purely nuclear reasons (Gambhir, Bhagwat, and Gupta, 2015).

IX. COSMIC ORIGIN OF SUPERHEAVY NUCLEI

The heavy elements of Es and Fm were discovered in 1952 in the debris from the thermonuclear explosion conducted at Eniwetok Atoll in the Pacific Ocean. The fermium nucleus was made through the capture of 17 neutrons by ^{238}U followed by subsequent beta decays (Seaborg, 1959; Ghiorso, 2003). In nature, a similar process known as the rapid neutron-capture process (or r process) is invoked in order to explain the existence of roughly half of the nuclei heavier than iron, including all of the actinides (Burbidge *et al.*, 1957; Cameron, 1957; Cowan, Thielemann, and Truran, 1991). This nucleosynthesis process occurs in astrophysical scenario(s) where the time scale for neutron capture is much shorter than the time scale for beta decay (Arnould, Goriely, and Takahashi, 2007; Qian and Wasserburg, 2007). The recent observation of the electromagnetic transient associated with the GW170807 neutron star merger (Abbott *et al.*, 2017; Pian *et al.*, 2017) confirmed that material with high opacities can be produced during such events, suggesting neutron star mergers as primary candidate sites for hosting the r process (Lattimer and Schramm, 1974, 1976; Symbalisty and Schramm, 1982; Eichler *et al.*, 1989; Davies *et al.*, 1994). Numerical simulations show that high neutron densities can be achieved in the dynamical ejecta of this astrophysical scenario (Rosswog *et al.*, 1999; Goriely, Bauswein, and Janka, 2011; Korobkin *et al.*, 2012; Bauswein, Goriely, and Janka, 2013; Sekiguchi *et al.*, 2015; Radice *et al.*, 2016; Kasen *et al.*, 2017) and in the accretion disk formed after the merging of two neutron stars (Siegel and Metzger, 2017a, 2017b; Fernández *et al.*, 2018). These high neutron densities allow the seed nuclei that are present in the environment to undergo multiple neutron captures before beta decaying, leading to the synthesis of neutron-rich nuclei [see also Freiburghaus, Rosswog, and Thielemann (1999), Goriely (2015), Lippuner and Roberts (2015), de Jesús Mendoza-Temis *et al.* (2015), and Thielemann *et al.* (2017)]. The termination of

the r -process path is determined by the region of the nuclear chart where fission is the dominant decay mode and where the fragments produced cycle the material to lower mass regions (Panov *et al.*, 2005; Beun *et al.*, 2008). The fundamental question of whether superheavy nuclei can be produced in nature is thus intrinsically related to determining which neutron-rich nuclei are stable against fission.

During the initial phase of the r process, the main competing decays are neutron captures, beta decays, and, for fissioning nuclei, neutron-induced fission (Panov and Thielemann, 2004; Martínez-Pinedo *et al.*, 2007; Panov, 2016, 2018). As long as neutron captures dominate over neutron-induced fission, the r -process path can proceed toward heavier and more neutron-rich regions of the nuclear chart, but as soon as it enters in a region of low fission barriers, the fragments formed in fission are cycled back (Beun *et al.*, 2008). Since the excitation energy of a nucleus after capturing a low-energy neutron is roughly given by the neutron separation energy, the competition between neutron captures and fission barriers is roughly driven by the difference between the height of the fission barrier and the neutron separation energy $B_f - S_n$ (Panov *et al.*, 2010). Several systematic calculations of the fission properties of r -process nuclei showed that close to the neutron drip line, fission barriers tend to increase with increasing neutron number, leading to positive values of $B_f - S_n$ (Petermann *et al.*, 2012; Möller *et al.*, 2015; Giuliani, Martínez-Pinedo, and Robledo, 2018). This suggests that the production of heavier nuclei could be enhanced if the r -process path proceeds closer to the neutron drip line, where neutron-induced fission is hindered. While superheavy nuclei produced in the r process are expected to be short lived, they may impact the observed r -process abundances and electromagnetic transients produced by the radioactive decay of r -process nuclei (kilonova) (Wanajo, 2018; Wu *et al.*, 2018; Zhu *et al.*, 2018).

Because of the competition between neutron capture, neutron-induced fission, and beta decay, the neutron number $N = 184$ may play a major role in determining the production of superheavy nuclei (Martínez-Pinedo *et al.*, 2007). Indeed, as discussed in Sec. IV, most nuclear models predict the region of increased shell stability around $N = 184$. Consequently, the neutron separation energy sharply drops for $N > 184$; this pushes the material toward nuclei with higher proton number ($Z \sim 100$), where several nuclear structure models predict nuclei with rather low fission barriers (Erler, Langanke *et al.*, 2012; Baran *et al.*, 2015; Goriely, 2015; Möller *et al.*, 2015; Giuliani, Martínez-Pinedo, and Robledo, 2018). As shown in Fig. 18, low fission barriers could prevent the production of nuclei beyond $N = 184$ due to neutron-induced fission.

In spite of the general lowering of fission barriers above $N = 184$, some r -process simulations show that for a particular combination of nuclear masses and fission barriers, nuclei with $Z \geq 104$ could be produced during the first seconds of the r process (Petermann *et al.*, 2012). In this case, other fission channels that could prevent the formation of superheavy nuclei are beta-delayed fission and spontaneous fission (Korneev and Panov, 2011; Erler, Langanke *et al.*, 2012; Panov *et al.*, 2013; Mumpower *et al.*, 2018). The beta-delayed fission channel is powered by the large Q_β value of neutron-rich nuclei, which can populate the daughter nuclei at excitation energies close to the fission barrier




FIG. 18. Dominant decay channels as a function of proton and neutron numbers predicted in the BCPM-DFT model for typical conditions ($T = 0.9$ GK, $n_n = 10^{28}$ cm $^{-3}$) of the r process in neutron star mergers: spontaneous fission, neutron capture, neutron-induced α emission, neutron-induced fission, and neutron-induced two-neutron emission. α decay is not competitive at the astrophysical conditions assumed. Adapted from Giuliani, Martínez-Pinedo, and Robledo, 2018.

(Thielemann, Metzinger, and Klapdor, 1983; Mumpower *et al.*, 2018). Moreover, a region of vanishing fission barriers is predicted around $Z/N \approx 104/188$ where spontaneous fission becomes the dominant decay channel (Erler, Langanke *et al.*, 2012; Giuliani, Martínez-Pinedo, and Robledo, 2018). This region seems to cross the beta-decay path of superheavy nuclei produced during the r process, thus blocking their decays toward stability.

The successful synthesis of superheavy nuclei in the laboratory revived the question of whether these nuclei can be found in nature. There have been searches for traces of superheavy elements in astrophysical data, including galactic cosmic rays, meteorites, and terrestrial ores (Donnelly *et al.*, 2012; Ter-Akopian and Dmitriev, 2015). In galactic cosmic ray studies, the most conclusive results were obtained by extracting olivine crystals from pallasite meteorites and analyzing the deposited nuclear tracks (Perelygin, Bondar *et al.*, 2003; Perelygin *et al.*, 2003; Polukhina, 2012). A recent study from the OLIMPIYA experiment claimed that three tracks belonging to superheavy nuclei with charge number 119_{-6}^{+10} have been found in meteorite olivine crystals (Alekseev *et al.*, 2017), but this result has not yet been confirmed by independent studies. In another development, the search for superheavy elements in terrestrial and extra-terrestrial samples was carried out with accelerator mass spectrometry techniques (Dellinger *et al.*, 2011; Korschinek and Kutschera, 2015) but the outcome of those investigations has been negative so far. In short, in spite of numerous efforts, no definitive proof of existence of superheavy nuclei in nature has been found (Ter-Akopian and Dmitriev, 2015).

X. PERSPECTIVES AND EXPECTATIONS

There is an exciting world beyond oganesson; major expansions of the nuclear chart and periodic table are on the horizon. In the short term, the search for the new elements $Z = 119$ and 120 will be carried out in several laboratories (Düllmann, 2016, 2017a; Hofmann *et al.*, 2016; Heßberger and Ackermann, 2017; Roberto and Rykaczewski, 2018). The reactions considered involve beams of ^{44}Ca , ^{50}Ti , ^{51}V , ^{54}Cr , ^{58}Fe , and ^{64}Ni (Nasirov *et al.*, 2011; Wang *et al.*, 2012; Liu and

Bao, 2013; Zhu, Xie, and Zhang, 2014; Hofmann, 2015; Zagrebaev and Greiner, 2015; Düllmann, 2016; Hofmann *et al.*, 2016; Adamian, Antonenko, and Lenske, 2018; Li *et al.*, 2018) and actinide targets (Roberto *et al.*, 2015; Roberto and Rykaczewski, 2018).

Since the lower and upper superheavy regions are still disjointed experimentally, i.e., they are not connected by known decays, the Z and A assignments of superheavy nuclei produced in hot-fusion reactions rely on indirect techniques. Another short-term goal is, therefore, to connect these two superheavy regions to provide a direct mass or charge identification. This can be achieved through, e.g., x-ray spectroscopy (Rudolph *et al.*, 2013; Ackermann, 2015) and direct mass measurements (Gates, 2016; Gregorich, 2016). Recently, the first direct measurements of the mass numbers of ^{288}Mc and ^{284}Nh were carried out at Berkeley (Gates *et al.*, 2018), confirming the previous indirect mass-number assignments.

The excursion toward the $N = 184$ region of longer-lived superheavies is the longer-term goal. Current experimental lifetimes are indicative of increasing stability for $Z > 110$ when adding neutrons. But to get closer to $N = 184$ is not going to be easy. To find the optimal production methods, systematic fusion reaction studies are being carried out (Itkis *et al.*, 2015; Loveland, 2016; Hinde *et al.*, 2017; Khuyagbaatar *et al.*, 2018). One proposed route involves an enriched ^{251}Cf target and ^{58}Fe beam (Rykaczewski *et al.*, 2016). Reactions using multinucleon transfer and radioactive neutron-rich beams are also being considered (Loveland, 2007; Zagrebaev and Greiner, 2008; Schädel, 2016; Karpov, Rachkov, and Saiko, 2018). Here the challenge for theory is to develop predictive models of superheavy nuclei production, capable of guiding future experimental searches (Zagrebaev and Greiner, 2015; Giardina *et al.*, 2018).

Another major challenge will be in atomic structure calculations at the accuracy of a few wave numbers including accurate values for transition dipole moments. Here sophisticated electron correlation frameworks including quantum electrodynamic effects are required to correctly predict the dense valence-electron spectrum of superheavy elements, required for future experimental atom-at-a-time studies (Sewtz *et al.*, 2003) in, e.g., cold traps. While such an accuracy has been already achieved for closed-shell atoms or for few-valence electron systems (Eliav, Fritzsche, and Kaldor, 2015; Schwerdtfeger *et al.*, 2015; Pařteka *et al.*, 2017), and even for small closed-shell molecules with lighter atoms (Owens *et al.*, 2015, 2018), this is not yet the case for complex multielectron, open-shell systems where a very large multireference and configuration-interaction space within a relativistic and QED framework is required to correctly describe both static and dynamic electron correlation effects. Such calculations will give a deep insight into the electronic shell structure with the correct prediction of ground-state symmetry (configuration) for the superheavy elements beyond oganesson, and into the regime of the critical nuclear charge Z_{crit} . The correct description of the elements beyond Z_{crit} involves the treatment of resonance states, which are well established for the positive-energy continuum (Jagau *et al.*, 2015; Fosseze *et al.*, 2016).

On the chemical side, experimental studies of Cn and Fl have illuminated the importance of relativistic effects on the properties of superheavy atoms, thus making the search for

chemistry beyond the standard periodic table the major science driver. Here fully relativistic quantum calculations, including QED effects, will continue providing guidance and stimulation for future atom-at-a-time experiments. The accurate simulation of the chemistry of superheavy elements in solution, gas phase and the bulk, or adsorbed on surfaces, with both density functional and wave-function-based methods, using relativistic pseudopotential theory including QED (Schwerdtfeger, 2011; Hangele *et al.*, 2012; Hangele, Dolg, and Schwerdtfeger, 2013), will remain a major challenge for the chemistry (well) beyond the standard periodic table (Türler and Pershina, 2013; Schwerdtfeger *et al.*, 2015).

Concerning atom-at-a-time chemistry, the future will see new chemical compounds being formed for the transactinide elements up to Og and possibly beyond. For example, there are currently worldwide efforts to study the carbonyls of Sg and beyond (Even *et al.*, 2014, 2015; Malli, 2015; Iliáš and Pershina, 2017) and many other transactinides. Another difficult task on the horizon is to synthesize more neutron-rich, longer-lived isotopes of known superheavy elements with $Z = 110\text{--}118$, with the goals of moving closer to $N = 184$ and enabling chemical studies. The discovery of new elements beyond Og will add the eighth period to the periodic table.

The field of superheavy element research puts nuclear and atomic theory to the test. There are strong theoretical suggestions that superheavy atoms and nuclei differ from lighter species because of their large charges and masses. The presence of large electrostatic forces gives rise to strong Coulomb frustration effects in the nuclear system and large relativistic effects in the atomic system; both present unusual challenges for many-body theory. Will next-generation hyperfine studies help to pin down the presence of exotic topologies? Since theories of superheavy nuclei heavily rely on extrapolations, it is essential to provide uncertainty quantification on predictions. To constrain nuclear models in the superheavy region, new high-quality data on bulk properties and spectroscopy of superheavy systems are required. High-quality experimental data have been accumulated on global properties of superheavy nuclei and their spectroscopy (Rudolph *et al.*, 2013, 2015; Ackermann, 2015; Theisen *et al.*, 2015; Herzberg, 2016; Heßberger, 2016; Ackermann and Theisen, 2017).

Whether superheavy nuclei can be produced during the r -process nucleosynthesis, and whether the observed elemental abundances contain unambiguous information on the role of superheavy elements, can only be addressed through systematic reaction network calculations. On the nuclear physics side, such calculations require a consistent set of nuclear reaction rates, binding energies, beta- and gamma-decay decay rates, fission rates, and fission fragment distributions, as well as a consistent treatment of weak interactions (Horowitz *et al.*, 2018). From an experimental point of view, further constraints will be provided by the data on neutron-rich rare isotopes from next-generation radioactive ion beam facilities. This new experimental information is expected to significantly reduce the uncertainties arising from the prediction of nuclear properties of some of the r -process nuclei (Ahearn and Freedman, 2007; Langanke and Schatz, 2013; Horowitz *et al.*, 2018; Surman and Mumpower, 2018). Unfortunately, very heavy, neutron-rich nuclei that are key for our understanding of fission in the r process are likely to

remain out of experimental reach in the foreseeable future. Consequently, the progress in this area will heavily rely on high-fidelity theoretical simulations.

The long term prospects in the unexplored regions of mass and charge are fascinating. They include the exploration of the region of long-lived superheavy nuclei around $N = 184$, the bold expansion of the chart of the nuclides, pinning down the presence of voids and other exotic topologies of nucleonic densities due to Coulomb frustration, delineating the role of superheavy nuclei in nucleosynthesis, and carrying out atomic and chemistry studies in the regime governed by large relativistic effects. The outstanding discovery potential has greatly motivated worldwide development of new facilities and novel experimental tools. New-generation, high-current stable-beam accelerators will enable new discoveries at the picobarn level. The dedicated facilities in Dubna (Dmitriev, Itkis, and Oganessian, 2016) and RIKEN (Haba, 2016) will substantially increase the production of superheavy species for physics and chemistry. Other major players in different areas of superheavy nuclei research include GSI/FAIR in Germany (Münzenberg, 2015, 2017), ORNL (Roberto *et al.*, 2015; Roberto and Rykaczewski, 2018) and Berkeley (Gates *et al.*, 2015, 2018; Gates, 2016) in the USA, GANIL in France (Ackermann and Theisen, 2017; Theisen, 2017), and several other major laboratories worldwide (Zhang *et al.*, 2012; Yang *et al.*, 2013; Back, 2017; Hinde *et al.*, 2017).

The scientific expedition continues into the uncharted regions of atomic number and nuclear mass. It was launched in the 1960s, fueled by the dream of the island of stability inhabited by very long-lived superheavy species. While this idea has been revised, present-day scientific drivers are, arguably, far more exciting. The prospects for discoveries in the interdisciplinary field of superheavy nuclei and atoms, on the intersection of nuclear physics, atomic physics, chemistry, and astrophysics are outstanding.

ACKNOWLEDGMENTS

Discussions with Yuri Oganessian and Krzysztof Rykaczewski and comments from Anatoli Afanasjev are gratefully appreciated. This work was supported by the U.S. Department of Energy under Awards No. DOE-DE-NA0002847 (NNSA, the Stewardship Science Academic Alliances program), No. DE-SC0013365 (Office of Science), No. DE-SC0018083 (Office of Science, NUCLEI SciDAC-4 Collaboration), and No. DE-AC52-07NA27344 (with the Lawrence Livermore National Laboratory). It was also supported by the BMBF-Verbundforschungsprojekt No. 05P15RDFN1 and the Royal Society of New Zealand in terms of a Marsden Fund (No. 17-MAU-021). Computing support for this work came from the Lawrence Livermore National Laboratory Institutional Computing Grand Challenge program.

REFERENCES

- Abbott, B. P., *et al.*, 2017, “Multi-messenger observations of a binary neutron star merger,” *Astrophys. J. Lett.* **848**, L12.
- Abusara, H., A. V. Afanasjev, and P. Ring, 2012, “Fission barriers in covariant density functional theory: Extrapolation to superheavy nuclei,” *Phys. Rev. C* **85**, 024314.
- Ackermann, D., 2015, “Nuclear spectroscopy in nuclei with $Z > 110$,” *Nucl. Phys. A* **944**, 376–387.
- Ackermann, D., and Ch. Theisen, 2017, “Nuclear structure features of very heavy and superheavy nuclei - tracing quantum mechanics towards the ‘island of stability’,” *Phys. Scr.* **92**, 083002.
- Adamian, G. G., N. V. Antonenko, and H. Lenske, 2018, “Estimates of production and structure of nuclei with $Z = 119$,” *Nucl. Phys. A* **970**, 22–28.
- Afanasjev, A. V., and O. Abdurazakov, 2013, “Pairing and rotational properties of actinides and superheavy nuclei in covariant density functional theory,” *Phys. Rev. C* **88**, 014320.
- Afanasjev, A. V., H. Abusara, and S. E. Agbemava, 2018, “Octupole deformation in neutron-rich actinides and superheavy nuclei and the role of nodal structure of single-particle wavefunctions in extremely deformed structures of light nuclei,” *Phys. Scr.* **93**, 034002.
- Afanasjev, A. V., S. E. Agbemava, and A. Gyawali, 2018, “Hyperheavy nuclei: existence and stability,” *Phys. Lett. B* **782**, 533–540.
- Afanasjev, A. V., and S. Frauendorf, 2005, “Central depression in nuclear density and its consequences for the shell structure of superheavy nuclei,” *Phys. Rev. C* **71**, 024308.
- Agbemava, S. E., and A. V. Afanasjev, 2017, “Octupole deformation in the ground states of even-even $Z \sim 96$, $N \sim 196$ actinides and superheavy nuclei,” *Phys. Rev. C* **96**, 024301.
- Agbemava, S. E., A. V. Afanasjev, D. Ray, and P. Ring, 2014, “Global performance of covariant energy density functionals: Ground state observables of even-even nuclei and the estimate of theoretical uncertainties,” *Phys. Rev. C* **89**, 054320.
- Agbemava, S. E., A. V. Afanasjev, D. Ray, and P. Ring, 2017, “Assessing theoretical uncertainties in fission barriers of superheavy nuclei,” *Phys. Rev. C* **95**, 054324.
- Agbemava, S. E., A. V. Afanasjev, T. Nakatsukasa, and P. Ring, 2015, “Covariant density functional theory: Reexamining the structure of superheavy nuclei,” *Phys. Rev. C* **92**, 054310.
- Ahearne, J. F., and S. J. Freedman, National Research Council, 2007, *Scientific Opportunities with a Rare-Isotope Facility in the United States* (The National Academies Press, Washington, DC).
- Ahmad, I., *et al.*, 1997, “Search for monoenergetic positron emission from heavy-ion collisions at Coulomb-barrier energies,” *Phys. Rev. Lett.* **78**, 618–621.
- Ahmad, I., *et al.*, 1999, “Positron-electron pairs produced in heavy-ion collisions,” *Phys. Rev. C* **60**, 064601.
- Alekseev, V. A., *et al.*, 2017, “Search for the ‘stability island’ of superheavy nuclei using natural track detectors,” *Bulletin of the Lebedev Physics Institute* **44**, 336–339.
- Andrae, D., 2000, “Finite nuclear charge density distributions in electronic structure calculations for atoms and molecules,” *Phys. Rep.* **336**, 413–525.
- Arnould, M., S. Goriely, and K. Takahashi, 2007, “The r-process of stellar nucleosynthesis: Astrophysics and Nucl. Phys. Achievements and mysteries,” *Phys. Rep.* **450**, 97–213.
- Asai, M., F. P. Heßberger, and A. Lopez-Martens, 2015, “Nuclear structure of elements with $100 \leq Z \leq 109$ from alpha spectroscopy,” *Nucl. Phys. A* **944**, 308–332.
- Autschbach, J., S. Siekierski, M. Seth, P. Schwerdtfeger, and W. H. E. Schwarz, 2002, “Dependence of relativistic effects on electronic configuration in the neutral atoms of *d*- and *f*-block elements,” *J. Comput. Chem.* **23**, 804–813.
- Back, B. B., 2017, “The AGFA and AIRIS separators at ATLAS,” *Eur. Phys. J. Web Conf.* **163**, 00003.
- Backe, H., W. Lauth, M. Block, and M. Laatiaoui, 2015, “Prospects for laser spectroscopy, ion chemistry and mobility measurements

- of superheavy elements in buffer-gas traps,” *Nucl. Phys. A* **944**, 492–517.
- Baran, A., M. Kowal, P.-G. Reinhard, L. M. Robledo, A. Staszczak, and M. Warda, 2015, “Fission barriers and probabilities of spontaneous fission for elements with $Z \geq 100$,” *Nucl. Phys. A* **944**, 442–470.
- Baran, A., J. A. Sheikh, J. Dobaczewski, W. Nazarewicz, and A. Staszczak, 2011, “Quadrupole collective inertia in nuclear fission: Cranking approximation,” *Phys. Rev. C* **84**, 054321.
- Baran, A., and A. Staszczak, 2013, “Rotational 2^+ states of superheavy elements in the Skyrme-Hartree-Fock-Bogoliubov model,” *Phys. Scr.* **T154**, 014027.
- Barber, R. C., P. J. Karol, H. Nakahara, E. Vardaci, and E. W. Vogt, 2011, “Discovery of the elements with atomic numbers greater than or equal to 113 (IUPAC technical report),” *Pure Appl. Chem.* **83**, 1485.
- Bauswein, A., S. Goriely, and H.-T. Janka, 2013, “Systematics of dynamical mass ejection, nucleosynthesis, and radioactively powered electromagnetic signals from neutron-star mergers,” *Astrophys. J.* **773**, 78.
- Bender, M., P.-H. Heenen, and P.-G. Reinhard, 2003, “Self-consistent mean-field models for nuclear structure,” *Rev. Mod. Phys.* **75**, 121.
- Bender, M., W. Nazarewicz, and P.-G. Reinhard, 2001, “Shell stabilization of super- and hyperheavy nuclei without magic gaps,” *Phys. Lett. B* **515**, 42–48.
- Bender, M., K. Rutz, P. G. Reinhard, J. A. Maruhn, and W. Greiner, 1999, “Shell structure of superheavy nuclei in self-consistent mean-field models,” *Phys. Rev. C* **60**, 034304.
- Bender, Michael, 2000, “ α -decay chains of $^{289}_{175}114$ and $^{293}_{175}118$ in the relativistic mean-field model,” *Phys. Rev. C* **61**, 031302.
- Berger, J. F., D. Hirata, and M. Girod, 2003, “Microscopic description of superheavy nuclei with the gogny effective interaction,” *Acta Phys. Pol. B* **34**, 1909 [<http://www.actaphys.uj.edu.pl/fulltext?series=Reg&vol=34&page=1909>].
- Beun, J., G. C. McLaughlin, R. Surman, and W. R. Hix, 2008, “Fission cycling in a supernova r process,” *Phys. Rev. C* **77**, 035804.
- Bjørnholm, S., and J. E. Lynn, 1980, “The double-humped fission barrier,” *Rev. Mod. Phys.* **52**, 725–931.
- Block, M., *et al.*, 2010, “Direct mass measurements above uranium bridge the gap to the island of stability,” *Nature (London)* **463**, 785–788.
- Bonche, P., and D. Vautherin, 1981, “A mean-field calculation of the equation of state of supernova matter,” *Nucl. Phys. A* **372**, 496–526.
- Bonchev, D., and V. Kamenska, 1981, “Predicting the properties of the 113-120 transactinide elements,” *J. Phys. Chem.* **85**, 1177–1186.
- Borschevsky, A., V. Pershina, E. Eliav, and U. Kaldor, 2009, “Electron affinity of element 114, with comparison to Sn and Pb,” *Chem. Phys. Lett.* **480**, 49–51.
- Brack, M., J. Damgaard, A. S. Jensen, H. C. Pauli, V. M. Strutinsky, and C. Y. Wong, 1972, “Funny hills: The shell-correction approach to nuclear shell effects and its applications to the fission process,” *Rev. Mod. Phys.* **44**, 320.
- Brewer, N. T., *et al.*, 2018, “Search for the heaviest atomic nuclei among the products from reactions of mixed-Cf with a ^{48}Ca beam,” *Phys. Rev. C* **98**, 024317.
- Brodziński, W., and J. Skalski, 2013, “Predictions for superheavy elements beyond $Z = 126$,” *Phys. Rev. C* **88**, 044307.
- Brown, B. A., 1992, “Simple relation for alpha decay half-lives,” *Phys. Rev. C* **46**, 811–814.
- Budaca, A. I., R. Budaca, and I. Silisteanu, 2016, “Extended systematics of alpha decay half lives for exotic superheavy nuclei,” *Nucl. Phys. A* **951**, 60–74.
- Burbidge, E. M., G. R. Burbidge, W. A. Fowler, and F. Hoyle, 1957, “Synthesis of the elements in stars,” *Rev. Mod. Phys.* **29**, 547–650.
- Bürvenich, T., M. Bender, J. A. Maruhn, and P.-G. Reinhard, 2004, “Systematics of fission barriers in superheavy elements,” *Phys. Rev. C* **69**, 014307.
- Bürvenich, T., K. Rutz, M. Bender, P.-G. Reinhard, J. A. Maruhn, and W. Greiner, 1998, “Superheavy nuclei in deformed mean-field calculations,” *Eur. Phys. J. A* **3**, 139.
- Butler, P. A., and W. Nazarewicz, 1996, “Intrinsic reflection asymmetry in atomic nuclei,” *Rev. Mod. Phys.* **68**, 349–421.
- Calvo, F., E. Pahl, M. Wormit, and P. Schwerdtfeger, 2013, “Evidence for low-temperature melting of mercury owing to relativity,” *Angew. Chem., Int. Ed.* **52**, 7583–7585.
- Cameron, A. G. W., 1957, “Stellar evolution, nuclear astrophysics, and nucleogenesis,” Technical Report [Atomic Energy of Canada Ltd. Chalk River Project, Chalk River, Ontario (Canada)].
- Chhetri, P., *et al.*, 2017, “Impact of buffer gas quenching on the $^1S_0 \rightarrow ^1P_1$ ground-state atomic transition in nobelium,” *Eur. Phys. J. D* **71**, 195.
- Chhetri, P., *et al.*, 2018, “Precision measurement of the first ionization potential of nobelium,” *Phys. Rev. Lett.* **120**, 263003.
- Chowdhury, P. Roy, C. Samanta, and D. N. Basu, 2008, “Search for long lived heaviest nuclei beyond the valley of stability,” *Phys. Rev. C* **77**, 044603.
- Clark, R. M., and D. Rudolph, 2018, “ α decay of high- K isomers in ^{270}Ds and ^{266}Hs in a superfluid tunneling model,” *Phys. Rev. C* **97**, 024333.
- Cowan, J. J., F.-K. Thielemann, and J. W. Truran, 1991, “The r -process and nucleochronology,” *Phys. Rep.* **208**, 267–394.
- Ćwiok, S., V. V. Pashkevich, J. Dudek, and W. Nazarewicz, 1983, “Fission barriers of transfermium elements,” *Nucl. Phys. A* **410**, 254–270.
- Ćwiok, S., J. Dobaczewski, P. H. Heenen, P. Magierski, and W. Nazarewicz, 1996, “Shell structure of the superheavy elements,” *Nucl. Phys. A* **611**, 211–246.
- Ćwiok, S., P. H. Heenen, and W. Nazarewicz, 2005, “Shape coexistence and triaxiality in the superheavy nuclei,” *Nature (London)* **433**, 705–709.
- Ćwiok, S., W. Nazarewicz, and P. H. Heenen, 1999, “Structure of odd- N superheavy elements,” *Phys. Rev. Lett.* **83**, 1108–1111.
- David, H. M., *et al.*, 2015, “Decay and fission hindrance of two- and four-quasiparticle K isomers in ^{254}Rf ,” *Phys. Rev. Lett.* **115**, 132502.
- Davies, K. T. R., C. Y. Wong, and S. J. Krieger, 1972, “Hartree-Fock calculations of bubble nuclei,” *Phys. Lett. B* **41**, 455–457.
- Davies, M. B., W. Benz, T. Piran, and F. K. Thielemann, 1994, “Merging neutron stars. I. Initial results for coalescence of non-rotating systems,” *Astrophys. J.* **431**, 742–753.
- Dechargé, J., J. F. Berger, M. Girod, and K. Dietrich, 2003, “Bubbles and semi-bubbles as a new kind of superheavy nuclei,” *Nucl. Phys. A* **716**, 55–86.
- Dechargé, J., J.-F. Berger, K. Dietrich, and M. S. Weiss, 1999, “Superheavy and hyperheavy nuclei in the form of bubbles or semi-bubbles,” *Phys. Lett. B* **451**, 275–282.
- de Jesús Mendoza-Temis, J., M.-R. Wu, K. Langanke, G. Martínez-Pinedo, A. Bauswein, and H.-T. Janka, 2015, “Nuclear robustness of the r process in neutron-star mergers,” *Phys. Rev. C* **92**, 055805.
- Delion, D. S., R. J. Liotta, and R. Wyss, 2007, “ α decay of high-spin isomers in superheavy nuclei,” *Phys. Rev. C* **76**, 044301.
- Dellinger, F., O. Forstner, R. Golser, A. Priller, P. Steier, A. Wallner, G. Winkler, and W. Kutschera, 2011, “Ultrasensitive search for long-lived superheavy nuclides in the mass range $A = 288$ to $A = 300$ in natural Pt, Pb, and Bi,” *Phys. Rev. C* **83**, 065806.

- Denisov, V. Y., 2005, “Magic numbers of ultraheavy nuclei,” *Phys. At. Nucl.* **68**, 1133.
- Desclaux, J. P., and P. Pyykkö, 1976, “Dirac-Fock one-centre calculations. the molecules CuH, AgH and AuH including p-type symmetry functions,” *Chem. Phys. Lett.* **39**, 300–303.
- Dietrich, K., and K. Pomorski, 1998, “Stability of bubble nuclei through shell effects,” *Phys. Rev. Lett.* **80**, 37–40.
- Dirac, P. A. M., 1929, “Quantum mechanics of many-electron systems,” *Proc. R. Soc. A* **123**, 714–733.
- Dmitriev, S., M. Itkis, and Y. Oganessian, 2016, “Status and perspectives of the Dubna superheavy element factory,” *Eur. Phys. J. Web Conf.* **131**, 08001.
- Dobaczewski, J., A. V. Afanasjev, M. Bender, L. M. Robledo, and Yue Shi, 2015, “Properties of nuclei in the nobelium region studied within the covariant, Skyrme, and Gogny energy density functionals,” *Nucl. Phys. A* **944**, 388–414.
- Dognon, J.-P., and P. Pyykkö, 2017, “Chemistry of the 5g elements: Relativistic calculations on hexafluorides,” *Angew. Chem., Int. Ed.* **56**, 10132–10134.
- Dong, J., W. Zuo, and W. Scheid, 2011, “New approach for alpha decay half-lives of superheavy nuclei and applicability of WKB approximation,” *Nucl. Phys. A* **861**, 1–13.
- Donnelly, J., A. Thompson, D. O’Sullivan, J. Daly, L. Drury, V. Domingo, and K.-P. Wenzel, 2012, “Actinide and ultra-heavy abundances in the local galactic cosmic rays: an analysis of the results from the LDEF ultra-heavy cosmic-ray experiment,” *Astrophys. J.* **747**, 40.
- Düllmann, C. E., 2016, “On the search for elements beyond $Z = 118$. An outlook based on lessons from the heaviest known elements,” *Eur. Phys. J. Web Conf.* **131**, 08004.
- Düllmann, C. E., 2017a, “How elements up to 118 were reached and how to go beyond,” *Eur. Phys. J. Web Conf.* **163**, 00015.
- Düllmann, C. E., 2017b, “Studying chemical properties of the heaviest elements: One atom at a time,” *Nucl. Phys. News* **27**, 14–20.
- Düllmann, C. E., and M. Block, 2018, “The quest for superheavy elements and the island of stability,” *Sci. Am.* **318**, 46.
- Dzuba, V. A., V. V. Flambaum, and O. P. Sushkov, 1983, “Energy levels of francium,” *Phys. Lett. A* **95**, 230–232.
- Dzuba, V. A., V. V. Flambaum, and J. K. Webb, 2017, “Isotope shift and search for metastable superheavy elements in astrophysical data,” *Phys. Rev. A* **95**, 062515.
- Eichler, D., M. Livio, T. Piran, and D. N. Schramm, 1989, “Nucleosynthesis, neutrino bursts and γ -rays from coalescing neutron stars,” *Nature (London)* **340**, 126–128.
- Eichler, R., *et al.*, 2008, “Thermochemical and physical properties of element 112,” *Angew. Chem., Int. Ed.* **47**, 3262–3266.
- Eichler, R., *et al.*, 2010, “Indication for a volatile element 114,” *Radiochim. Acta* **98**, 133–139.
- Eliav, E., S. Fritzsche, and U. Kaldor, 2015, “Electronic structure theory of the superheavy elements,” *Nucl. Phys. A* **944**, 518–550.
- Eliav, E., U. Kaldor, P. Schwerdtfeger, B. A. Hess, and Y. Ishikawa, 1994, “Ground state electron configuration of element 111,” *Phys. Rev. Lett.* **73**, 3203–3206.
- Erler, J., N. Birge, M. Kortelainen, W. Nazarewicz, E. Olsen, A. M. Perhac, and M. Stoitsov, 2012, “The limits of the nuclear landscape,” *Nature (London)* **486**, 509.
- Erler, J., K. Langanke, H. P. Loens, G. Martínez-Pinedo, and P.-G. Reinhard, 2012, “Fission properties for r -process nuclei,” *Phys. Rev. C* **85**, 025802.
- Even, J., *et al.*, 2014, “Synthesis and detection of a seaborgium carbonyl complex,” *Science* **345**, 1491–1493.
- Even, J., *et al.*, 2015, “In situ synthesis of volatile carbonyl complexes with short-lived nuclides,” *J. Radioanal. Nucl. Chem.* **303**, 2457–2466.
- Fattoyev, F. J., C. J. Horowitz, and B. Schuetrumpf, 2017, “Quantum nuclear pasta and nuclear symmetry energy,” *Phys. Rev. C* **95**, 055804.
- Fernández, R., A. Tchekhovskoy, E. Quataert, F. Foucart, and D. Kasen, 2019, “Long-term GRMHD simulations of neutron star merger accretion disks: Implications for electromagnetic counterparts,” *Mon. Not. R. Astron. Soc.* **482**, 3373–3393.
- Ferrer, R., *et al.*, 2017, “Towards high-resolution laser ionization spectroscopy of the heaviest elements in supersonic gas jet expansion,” *Nat. Commun.* **8**, 14520.
- Fossez, K., Xingze Mao, W. Nazarewicz, N. Michel, W. R. Garrett, and M. Płoszajczak, 2016, “Resonant spectra of quadrupolar anions,” *Phys. Rev. A* **94**, 032511.
- Freedman, S. J., NAS Report, 2012, Nuclear Physics: Exploring the Heart of Matter. Report of the Committee on the Assessment of and Outlook for Nuclear Physics (The National Academies Press, Washington, DC).
- Freiburghaus, C., S. Rosswog, and F.-K. Thielemann, 1999, “ r -process in neutron star mergers,” *Astrophys. J.* **525**, L121–L124.
- Fricke, B., W. Greiner, and J. T. Waber, 1971, “The continuation of the periodic table up to $Z = 172$ the chemistry of superheavy elements,” *Theor. Chim. Acta* **21**, 235–260.
- Gäggeler, H. W., 2011, “Gas chemical properties of heaviest elements,” *Radiochim. Acta* **99**, 503–513.
- Gambhir, Y. K., A. Bhagwat, and M. Gupta, 2005, “ α -decay half-lives of the observed superheavy nuclei ($Z = 108$ – 118),” *Phys. Rev. C* **71**, 037301.
- Gambhir, Y. K., A. Bhagwat, and M. Gupta, 2015, “The highest limiting Z in the extended periodic table,” *J. Phys. G* **42**, 125105.
- Gaston, N., I. Opahle, H. W. Gäggeler, and P. Schwerdtfeger, 2007, “Is eka-mercury (element 112) a group 12 metal?,” *Angew. Chem., Int. Ed.* **46**, 1663–1666.
- Gates, J. M., 2016, “Prospects of A and Z identification experiments at LBNL,” *Eur. Phys. J. Web Conf.* **131**, 08003.
- Gates, J. M., *et al.*, 2015, “Decay spectroscopy of element 115 daughters: $^{280}\text{Rg} \rightarrow ^{276}\text{Mt}$ and $^{276}\text{Mt} \rightarrow ^{272}\text{Bh}$,” *Phys. Rev. C* **92**, 021301.
- Gates, J. M., *et al.*, 2018, “First direct measurements of superheavy element mass numbers,” *Phys. Rev. Lett.* **121**, 222501.
- Gershtein, S. S., and Ya. B. Zel’dovich, 1969, “Positron production during the mutual approach of heavy nuclei and the polarization of the vacuum,” *Sov. Phys. JETP* **30**, 358 [<http://adsabs.harvard.edu/abs/1969JETP...30..358G>].
- Gherghescu, R. A., J. Skalski, Z. Patyk, and A. Sobczewski, 1999, “Non-axial shapes in spontaneous fission of superheavy nuclei,” *Nucl. Phys. A* **651**, 237–249.
- Ghiorso, A., 2003, “Einsteinium and fermium,” *Chem. Eng. News* **81**, 174–175.
- Giardina, G., G. Mandaglio, A. K. Nasirov, A. Anastasi, F. Curciarello, and G. Fazio, 2018, “Uncertainties and understanding of experimental and theoretical results regarding reactions forming heavy and superheavy nuclei,” *Nucl. Phys. A* **970**, 169–207.
- Giuliani, S. A., G. Martínez-Pinedo, and L. M. Robledo, 2018, “Fission properties of superheavy nuclei for r -process calculations,” *Phys. Rev. C* **97**, 034323.
- Giuliani, S. A., and L. M. Robledo, 2013, “Fission properties of the Barcelona-Catania-Paris-Madrid energy density functional,” *Phys. Rev. C* **88**, 054325.
- Giuliani, S. A., L. M. Robledo, and R. Rodríguez-Guzmán, 2014, “Dynamic versus static fission paths with realistic interactions,” *Phys. Rev. C* **90**, 054311.

- Glashow, Sheldon, 1988, *Interactions: A Journey Through the Mind of a Particle Physicist and the Matter of This World* (Warner Books, New York).
- Goidenko, I., L. Labzowsky, E. Eliav, U. Kaldor, and P. Pyykkö, 2003, “QED corrections to the binding energy of the eka-radon ($Z = 118$) negative ion,” *Phys. Rev. A* **67**, 020102.
- Goldanskii, V.I., 1966, “Modes of radioactive decay involving proton emission,” *Annu. Rev. Nucl. Sci.* **16**, 1.
- Gooden, M. E., *et al.*, 2016, “Energy dependence of fission product yields from ^{235}U , ^{238}U and ^{239}Pu for incident neutron energies between 0.5 and 14.8 MeV,” *Nucl. Data Sheets* **131**, 319.
- Gopka, V. F., A. V. Yushchenko, V. A. Yushchenko, I. V. Panov, and Ch. Kim, 2008, “Identification of absorption lines of short half-life actinides in the spectrum of Przybylski’s star (HD 101065),” *Kinematics and Physics of Celestial Bodies* **24**, 89–98.
- Goriely, S., 2015, “The fundamental role of fission during r -process nucleosynthesis in neutron star mergers,” *Eur. Phys. J. A* **51**, 22.
- Goriely, S., A. Bauswein, and H.-T. Janka, 2011, “ R -process nucleosynthesis in dynamically ejected matter of neutron star mergers,” *Astrophys. J. Lett.* **738**, L32.
- Gray, R. W., and W. Ramsay, 1909, “CXXII.—some physical properties of radium emanation,” *J. Chem. Soc., Trans.* **95**, 1073–1085.
- Greenlees, P. T., *et al.*, 2012, “Shell-structure and pairing interaction in superheavy nuclei: Rotational properties of the $Z=104$ nucleus ^{256}Rf ,” *Phys. Rev. Lett.* **109**, 012501.
- Gregorich, K., 2016, “How good are superheavy element Z and A assignments?,” *Eur. Phys. J. Web Conf.* **131**, 06002.
- Greiner, W., 2008, “Structure of vacuum and elementary matter: From superheavies via hypermatter to antimatter—the vacuum decay in supercritical fields,” *Adv. Quantum Chem.* **53**, 99–150.
- Gyanchandani, J., and S. K. Sikka, 2011, “Physical properties of the $6d$ -series elements from density functional theory: Close similarity to lighter transition metals,” *Phys. Rev. B* **83**, 172101.
- Haba, H., 2016, “Superheavy element chemistry at GARIS,” *Eur. Phys. J. Web Conf.* **131**, 07006.
- Hangele, T., M. Dolg, M. Hanrath, X. Cao, and P. Schwerdtfeger, 2012, “Accurate relativistic energy-consistent pseudopotentials for the superheavy elements 111 to 118 including quantum electrodynamic effects,” *J. Chem. Phys.* **136**, 214105.
- Hangele, T., M. Dolg, and P. Schwerdtfeger, 2013, “Relativistic energy-consistent pseudopotentials for superheavy elements 119 and 120 including quantum electrodynamic effects,” *J. Chem. Phys.* **138**, 174113.
- Heenen, P.-H., J. Skalski, A. Staszczak, and D. Vretenar, 2015, “Shapes and α - and β -decays of superheavy nuclei,” *Nucl. Phys. A* **944**, 415–441.
- Hermann, A., J. Furthmüller, H. W. Gäggeler, and P. Schwerdtfeger, 2010, “Spin-orbit effects in structural and electronic properties for the solid state of the group-14 elements from carbon to superheavy element 114,” *Phys. Rev. B* **82**, 155116.
- Herzberg, R.-D., 2016, “In-beam spectroscopy of the heaviest elements,” *Eur. Phys. J. Web Conf.* **131**, 02004.
- Herzberg, R.-D., and P. T. Greenlees, 2008, “In-beam and decay spectroscopy of transfermium nuclei,” *Prog. Part. Nucl. Phys.* **61**, 674–720.
- Heßberger, F. P., 2016, “Nuclear structure of the transactinides - investigated by decay spectroscopy,” *Eur. Phys. J. Web Conf.* **131**, 02005.
- Heßberger, F. P., 2017, “Spontaneous fission properties of superheavy elements,” *Eur. Phys. J. A* **53**, 75.
- Heßberger, F. P., and D. Ackermann, 2017, “Some critical remarks on a sequence of events interpreted to possibly originate from a decay chain of an element 120 isotope,” *Eur. Phys. J. A* **53**, 123.
- Heßberger, F. P., *et al.*, 2016, “Investigation of electron capture decay of ^{258}Db and α decay of ^{258}Rf ,” *Eur. Phys. J. A* **52**, 328.
- Hinde, D. J., *et al.*, 2017, “Quasifission dynamics in the formation of superheavy elements,” *Eur. Phys. J. Web Conf.* **163**, 00023.
- Hoffman, D. C., A. Ghiorso, and G. T. Seaborg, 2000, *The Transuranium People* (World Scientific, River Edge, NJ).
- Hofmann, S., 2015, “Super-heavy nuclei,” *J. Phys. G* **42**, 114001.
- Hofmann, S., and G. Münzenberg, 2000, “The discovery of the heaviest elements,” *Rev. Mod. Phys.* **72**, 733–767.
- Hofmann, S., *et al.*, 2001, “The new isotope $^{270}\text{110}$ and its decay products ^{266}Hs and ^{262}Sg ,” *Eur. Phys. J. A* **10**, 5–10.
- Hofmann, S., *et al.*, 2016, “Review of even element super-heavy nuclei and search for element 120,” *Eur. Phys. J. A* **52**, 180.
- Horowitz, C. J., *et al.*, 2018, “ r -Process Nucleosynthesis: Connecting Rare-Isotope Beam Facilities with the Cosmos,” [arXiv:1805.04637](https://arxiv.org/abs/1805.04637).
- Hoyer, S., T. Emmler, and K. Seppelt, 2006, “The structure of xenon hexafluoride in the solid state,” *J. Fluorine Chem.* **127**, 1415–1422.
- Id Betan, R., and W. Nazarewicz, 2012, “ α decay in the complex-energy shell model,” *Phys. Rev. C* **86**, 034338.
- Iliáš, M., and V. Pershina, 2017, “Hexacarbonyls of Mo, W, and Sg: Metal-CO bonding revisited,” *Inorg. Chem.* **56**, 1638–1645.
- Indelicato, P., J. Bieroń, and P. Jönsson, 2011, “Are MCDF calculations 101% correct in the super-heavy elements range?,” *Theor. Chem. Acc.* **129**, 495–505.
- Ismail, M., A. Y. Ellithi, A. Adel, and H. Anwer, 2016, “On magic numbers for super- and ultraheavy systems and hypothetical spherical double-magic nuclei,” *J. Phys. G* **43**, 015101.
- Itkis, M. G., E. Vardaci, I. M. Itkis, G. N. Knyazheva, and E. M. Kozulin, 2015, “Fusion and fission of heavy and superheavy nuclei (experiment),” *Nucl. Phys. A* **944**, 204–237.
- Ito, Y., *et al.*, 2018, “First direct mass measurements of nuclides around $Z = 100$ with a multireflection time-of-flight mass spectrograph,” *Phys. Rev. Lett.* **120**, 152501.
- Jachimowicz, P., M. Kowal, and J. Skalski, 2014, “ Q_α values in superheavy nuclei from the deformed woods-saxon model,” *Phys. Rev. C* **89**, 024304.
- Jachimowicz, P., M. Kowal, and J. Skalski, 2015, “Candidates for long-lived high- K ground states in superheavy nuclei,” *Phys. Rev. C* **92**, 044306.
- Jachimowicz, P., M. Kowal, and J. Skalski, 2017a, “Adiabatic fission barriers in superheavy nuclei,” *Phys. Rev. C* **95**, 014303.
- Jachimowicz, P., M. Kowal, and J. Skalski, 2017b, “Effect of non-axial octupole shapes in heavy and superheavy nuclei,” *Phys. Rev. C* **95**, 034329.
- Jachimowicz, P., M. Kowal, and J. Skalski, 2018, “Hindered α decays of heaviest high- K isomers,” *Phys. Rev. C* **98**, 014320.
- Jagau, T.-C., D. Zuev, K. B. Bravaya, E. Epifanovsky, and A. I. Krylov, 2015, “Correction to “A fresh look at resonances and complex absorbing potentials: Density matrix-based approach”,” *J. Phys. Chem. Lett.* **6**, 3866–3866.
- Jerabek, P., B. Schuetrumpf, P. Schwerdtfeger, and W. Nazarewicz, 2018, “Electron and nucleon localization functions of oganesson: Approaching the Thomas-Fermi limit,” *Phys. Rev. Lett.* **120**, 053001.
- Karatzikos, S., A. V. Afanasjev, G. A. Lalazissis, and P. Ring, 2010, “The fission barriers in actinides and superheavy nuclei in covariant density functional theory,” *Phys. Lett. B* **689**, 72–81.
- Karol, P. J., R. C. Barber, B. M. Sherrill, E. Vardaci, and T. Yamazaki, 2016a, “Discovery of the element with atomic number $Z = 118$ completing the 7th row of the periodic table (IUPAC technical report),” *Pure Appl. Chem.* **88**, 155.
- Karol, P. J., R. C. Barber, B. M. Sherrill, E. Vardaci, and T. Yamazaki, 2016b, “Discovery of the elements with atomic numbers $Z = 113$, 115 and 117 (IUPAC technical report),” *Pure Appl. Chem.* **88**, 139.

- Karpov, A. V., V. A. Rachkov, and V. V. Saiko, 2018, “Formation of neutron-enriched heavy and superheavy nuclei in fusion reactions,” *Phys. Part. Nucl. Lett.* **15**, 247–256.
- Kasen, D., B. Metzger, J. Barnes, E. Quataert, and E. Ramirez-Ruiz, 2017, “Origin of the heavy elements in binary neutron-star mergers from a gravitational-wave event,” *Nature (London)* **551**, 80–84.
- Khuyagbaatar, J., *et al.*, 2014, “ $^{48}\text{Ca} + ^{249}\text{Bk}$ fusion reaction leading to element $Z = 117$: Long-lived α -decaying ^{270}Db and discovery of ^{266}Lr ,” *Phys. Rev. Lett.* **112**, 172501.
- Khuyagbaatar, J., *et al.*, 2018, “Nuclear structure dependence of fusion hindrance in heavy element synthesis,” *Phys. Rev. C* **97**, 064618.
- Klüpfel, P., J. Erler, P.-G. Reinhard, and J. A. Maruhn, 2008, “Systematics of collective correlation energies from self-consistent mean-field calculations,” *Eur. Phys. J. A* **37**, 343.
- Korneev, I. Yu., and I. V. Panov, 2011, “Contribution of fission to heavy-element nucleosynthesis in an astrophysical r -process,” *Astron. Lett.* **37**, 864–873.
- Korobkin, O., S. Rosswog, A. Arcones, and C. Winteler, 2012, “On the astrophysical robustness of the neutron star merger r -process,” *Mon. Not. R. Astron. Soc.* **426**, 1940–1949.
- Korschinek, G., and W. Kutschera, 2015, “Mass spectrometric searches for superheavy elements in terrestrial matter,” *Nucl. Phys. A* **944**, 190–203.
- Koura, H., 2012, “Phenomenological formula for alpha-decay half-lives,” *J. Nucl. Sci. Technol.* **49**, 816–823.
- Kowal, M., P. Jachimowicz, and A. Sobczewski, 2010, “Fission barriers for even-even superheavy nuclei,” *Phys. Rev. C* **82**, 014303.
- Kozulin, E. M., *et al.*, 2014, “Fusion-fission and quasifission of superheavy systems with $Z = 110 - 116$ formed in ^{48}Ca -induced reactions,” *Phys. Rev. C* **90**, 054608.
- Krappe, H. J., and K. Pomorski, 2012, *Theory of Nuclear Fission* (Springer, New York).
- Kruppa, A. T., M. Bender, W. Nazarewicz, P.-G. Reinhard, T. Vertse, and S. Ćwiok, 2000, “Shell corrections of superheavy nuclei: Green’s function Hartree-Fock method,” *Phys. Rev. C* **61**, 034313.
- Laatiaoui, M., *et al.*, 2016, “Atom-at-a-time laser resonance ionization spectroscopy of nobelium,” *Nature (London)* **538**, 495–498.
- Lackenby, B. G. C., V. A. Dzuba, and V. V. Flambaum, 2018, “Atomic structure calculations of super heavy noble element oganesson ($Z = 118$),” *Phys. Rev. A* **98**, 042512.
- Langanke, K., and H. Schatz, 2013, “The role of radioactive ion beams in nuclear astrophysics,” *Phys. Scr.* **T152**, 014011.
- Lattimer, J. M., and D. N. Schramm, 1974, “Black-hole-neutron-star collisions,” *Astrophys. J.* **192**, L145.
- Lattimer, J. M., and D. N. Schramm, 1976, “The tidal disruption of neutron stars by black holes in close binaries,” *Astrophys. J.* **210**, 549.
- Li, F., L. Zhu, Z. H. Wu, X. B. Yu, J. Su, and C. C. Guo, 2018, “Predictions for the synthesis of superheavy elements $Z = 119$ and 120 ,” *Phys. Rev. C* **98**, 014618.
- Lippuner, J., and L. F. Roberts, 2015, “ R -process lanthanide production and heating rates in kilonovae,” *Astrophys. J.* **815**, 82.
- Liu, H. L., P. M. Walker, and F. R. Xu, 2014, “Favored configurations for four-quasiparticle K isomerism in the heaviest nuclei,” *Phys. Rev. C* **89**, 044304.
- Liu, Zu-Hua, and Jing-Dong Bao, 2013, “Possibility to produce element 120 in the $^{54}\text{Cr} + ^{248}\text{Cm}$ hot fusion reaction,” *Phys. Rev. C* **87**, 034616.
- Loveland, W., 2007, “Synthesis of transactinide nuclei using radioactive beams,” *Phys. Rev. C* **76**, 014612.
- Loveland, W., 2016, “Characterizing the mechanism(s) of heavy element synthesis reactions,” *Eur. Phys. J. Web Conf.* **131**, 04003.
- Lu, C. C., T. A. Carlson, F. B. Malik, T. C. Tucker, and C. W. Nestor, 1971, “Relativistic Hartree-Fock-Slater eigenvalues, radial expectation values, and potentials for atoms, $2 \leq Z \leq 126$,” *At. Data Nucl. Data Tables* **3**, 1–31.
- Malli, G. L., 2015, “Relativistic effects for the reaction $\text{Sg} + 6 \text{CO}$ $\text{Sg}(\text{CO})_6$: Prediction of the mean bond energy, atomization energy, and existence of the first organometallic transactinide superheavy hexacarbonyl $\text{Sg}(\text{CO})_6$,” *J. Chem. Phys.* **142**, 064311.
- Maltsev, I. A., V. M. Shabaev, I. I. Tupitsyn, A. I. Bondarev, Y. S. Kozhedub, G. Plunien, and Th. Stöhlker, 2015, “Electron-positron pair creation in low-energy collisions of heavy bare nuclei,” *Phys. Rev. A* **91**, 032708.
- Mann, J. B., 1969, “Stability of $8p$ electrons in superheavy elements,” *J. Chem. Phys.* **51**, 841–842.
- Mann, J. B., and J. T. Waber, 1970, “SCF relativistic Hartree-Fock calculations on the superheavy elements 118–131,” *J. Chem. Phys.* **53**, 2397–2406.
- Martínez-Pinedo, G., D. Mocolj, N. T. Zinner, A. Kelić, K. Langanke, I. Panov, B. Pfeiffer, T. Rauscher, K.-H. Schmidt, and F.-K. Thielemann, 2007, “The role of fission in the r -process,” *Prog. Part. Nucl. Phys.* **59**, 199–205.
- Matheson, Z., S. Giuliani, W. Nazarewicz, J. Sadhukhan, and N. Schunck, 2018, [arXiv:1812.06490](https://arxiv.org/abs/1812.06490).
- Mișicu, Ș., T. Bürvenich, T. Cornelius, and W. Greiner, 2002, “Properties of some collective excitations in spherical nuclei from the superheavy island,” *J. Phys. G* **28**, 1441.
- Möller, P., A. J. Sierk, T. Ichikawa, A. Iwamoto, and M. R. Mumpower, 2015, “Fission barriers at the end of the chart of the nuclides,” *Phys. Rev. C* **91**, 024310.
- Möller, P., R. Bengtsson, B. G. Carlsson, P. Olivius, T. Ichikawa, H. Sagawa, and A. Iwamoto, 2008, “Axial and reflection asymmetry of the nuclear ground state,” *At. Data Nucl. Data Tables* **94**, 758–780.
- Möller, P., and J. R. Nix, 1994, “Stability of heavy and superheavy elements,” *J. Phys. G* **20**, 1681.
- Möller, P., J. R. Nix, and K.-L. Kratz, 1997, “Nuclear properties for astrophysical and radioactive-ion-beam applications,” *At. Data Nucl. Data Tables* **66**, 131–343.
- Möller, P., J. R. Nix, W. D. Myers, and W. J. Swiatecki, 1995, “Nuclear ground-state masses and deformations,” *At. Data Nucl. Data Tables* **59**, 185–381.
- Möller, P., J. R. Nix, and W. J. Swiatecki, 1989, “New developments in the calculation of heavy-element fission barriers,” *Nucl. Phys. A* **492**, 349–387.
- Möller, P., A. J. Sierk, T. Ichikawa, A. Iwamoto, R. Bengtsson, H. Uhrenholt, and S. Åberg, 2009, “Heavy-element fission barriers,” *Phys. Rev. C* **79**, 064304.
- Mosel, U., and W. Greiner, 1969, “On the stability of superheavy nuclei against fission,” *Z. Phys. A* **222**, 261.
- Müller-Nehler, U., and G. Soff, 1994, “Electron excitations in superheavy quasimolecules,” *Phys. Rep.* **246**, 101–250.
- Mumpower, M. R., T. Kawano, T. M. Sprouse, N. Vassh, E. M. Holmbeck, R. Surman, and P. Möller, 2018, “ β -delayed Fission in r -process Nucleosynthesis,” *Astrophys. J.* **869**, 14.
- Muntian, I., S. Hofmann, Z. Patyk, and A. Sobczewski, 2003, “Properties of heaviest nuclei,” *Acta Phys. Pol. B* **34**, 2073 [<http://www.actaphys.uj.edu.pl/fulltext?series=Reg&vol=34&page=2073>].
- Münzenberg, G., 2015, “From bohrium to copernicium and beyond SHE research at SHIP,” *Nucl. Phys. A* **944**, 5–29.
- Münzenberg, G., and K. Morita, 2015, “Synthesis of the heaviest nuclei in cold fusion reactions,” *Nucl. Phys. A* **944**, 3–4.

- Münzenberg, G., *et al.*, 2017, “SHE research with rare-isotope beams, challenges and perspectives, and the new generation of SHE factories,” in *New Horizons in Fundamental Physics*, edited by Stefan Schramm and Mirko Schäfer (Springer International Publishing, Cham), pp. 81–90.
- Myers, W. D., and W. J. Swiatecki, 1966, “Nuclear masses and deformations,” *Nucl. Phys.* **81**, 1–60.
- Nagame, Y., J. V. Kratz, and M. Schädel, 2016, “Chemical properties of rutherfordium (Rf) and dubnium (Db) in the aqueous phase,” *Eur. Phys. J. Web Conf.* **131**, 07007.
- Nash, C. S., and B. E. Bursten, 1999, “Spin-orbit coupling versus the vsepr method: On the possibility of a nonplanar structure for the super-heavy noble gas tetrafluoride (118)F₄,” *Angew. Chem., Int. Ed.* **38**, 151–153.
- Nasirov, A. K., G. Mandaglio, G. Giardina, A. Sobczewski, and A. I. Muminov, 2011, “Effects of the entrance channel and fission barrier in the synthesis of superheavy element $Z = 120$,” *Phys. Rev. C* **84**, 044612.
- Nazarewicz, W., 2018, “The limits of nuclear mass and charge,” *Nat. Phys.* **14**, 537–541.
- Nazarewicz, W., M. Bender, S. Ćwiok, P. H. Heenen, A. T. Kruppa, P.-G. Reinhard, and T. Vertse, 2002, “Theoretical description of superheavy nuclei,” *Nucl. Phys. A* **701**, 165.
- Neufcourt, L., Y. Cao, W. Nazarewicz, and F. Viens, 2018, “Bayesian approach to model-based extrapolation of nuclear observables,” *Phys. Rev. C* **98**, 034318.
- Nilsson, S. G., C.-F. Tsang, A. Sobczewski, Z. Szymański, S. Wycech, C. Gustafson, I.-L. Lamm, P. Möller, and B. Nilsson, 1969, “On the nuclear structure and stability of heavy and super-heavy elements,” *Nucl. Phys. A* **131**, 1–66.
- Nix, J. R., 1972a, “Predictions for superheavy nuclei,” *Phys. Today* **25**, 30.
- Nix, J. R., 1972b, “Calculation of fission barriers for heavy and superheavy nuclei,” *Annu. Rev. Nucl. Sci.* **22**, 65.
- Oganessian, Yu. Ts., A. Sobczewski, and G. M. Ter-Akopian, 2017, “Superheavy nuclei: from predictions to discovery,” *Phys. Scr.* **92**, 023003.
- Oganessian, Yu. Ts., and V. K. Utyonkov, 2015a, “Super-heavy element research,” *Rep. Prog. Phys.* **78**, 036301.
- Oganessian, Yu. Ts., and V. K. Utyonkov, 2015b, “Superheavy nuclei from ⁴⁸Ca-induced reactions,” *Nucl. Phys. A* **944**, 62–98.
- Oganessian, Yu. Ts., *et al.*, 2006, “Synthesis of the isotopes of elements 118 and 116 in the ²⁴⁹Cf and ²⁴⁵Cm + ⁴⁸Ca fusion reactions,” *Phys. Rev. C* **74**, 044602.
- Oganessian, Yuri Ts., and K. Rykaczewski, 2015, “A beachhead on the island of stability,” *Phys. Today* **68**, 32.
- Owens, A., A. Yachmenev, J. Kupper, S. N. Yurchenko, and W. Thiel, 2018, “The rotation-vibration spectrum of methyl fluoride from first principles,” *Phys. Chem. Chem. Phys.*
- Owens, A., S. N. Yurchenko, W. Thiel, and V. Špirko, 2015, “Accurate prediction of the ammonia probes of a variable proton-to-electron mass ratio,” *Mon. Not. R. Astron. Soc.* **450**, 3191–3200.
- Pahl, E., F. Calvo, L. Koči, and P. Schwerdtfeger, 2008, “Accurate Melting Temperatures for Neon and Argon from Ab Initio Monte Carlo Simulations,” *Angew. Chem., Int. Ed.* **47**, 8207.
- Panov, I. V., 2016, “Nucleosynthesis of heavy elements in the r-process,” *Phys. At. Nucl.* **79**, 159–198.
- Panov, I. V., 2018, “Formation of superheavy elements in nature,” *Phys. At. Nucl.* **81**, 68–75.
- Panov, I. V., E. Kolbe, B. Pfeiffer, T. Rauscher, K.-L. Kratz, and F.-K. Thielemann, 2005, “Calculations of fission rates for r-process nucleosynthesis,” *Nucl. Phys. A* **747**, 633–654.
- Panov, I. V., and F.-K. Thielemann, 2004, “Fission and the r-process: Competition between neutron-induced and beta-delayed fission,” *Astron. Lett.* **30**, 647–655.
- Panov, I. V., I. Yu. Korneev, G. Martínez-Pinedo, and F. K. Thielemann, 2013, “Influence of spontaneous fission rates on the yields of superheavy elements in the r-process,” *Astron. Lett.* **39**, 150–160.
- Panov, I. V., I. Yu. Korneev, T. Rauscher, G. Martínez-Pinedo, A. Kelić-Heil, N. T. Zinner, and F.-K. Thielemann, 2010, “Neutron-induced astrophysical reaction rates for translead nuclei,” *Astron. Astrophys.* **513**, A61.
- Parkhomenko, A., and A. Sobczewski, 2005, “Phenomenological formula for α -decay half-lives of heaviest nuclei,” *Acta Phys. Pol. B* **36**, 3095 [<http://www.actaphys.uj.edu.pl/fulltext?series=Reg&vol=36&page=3095>].
- Pašteka, L. F., E. Eliav, A. Borschevsky, U. Kaldor, and P. Schwerdtfeger, 2017, “Relativistic coupled cluster calculations with variational quantum electrodynamics resolve the discrepancy between experiment and theory concerning the electron affinity and ionization potential of gold,” *Phys. Rev. Lett.* **118**, 023002.
- Pei, J. C., W. Nazarewicz, J. A. Sheikh, and A. K. Kerman, 2009, “Fission barriers of compound superheavy nuclei,” *Phys. Rev. Lett.* **102**, 192501.
- Pei, J. C., F. R. Xu, and P. D. Stevenson, 2005, “Density distributions of superheavy nuclei,” *Phys. Rev. C* **71**, 034302.
- Perelygin, V. P., Yu. V. Bondar, W. Ensinger, R. L. Fleischer, P. Vater, and S. G. Stetsenko, 2003, “On search and identification of short-lived super heavy cosmic-ray nuclei ($Z \geq 110$) by fossil track study of the extraterrestrial crystals: Results and perspectives [I],” *Nucl. Phys. A* **718**, 410–412.
- Perelygin, V. P., *et al.*, 2003, “On search and identification of short-lived super heavy cosmic-ray nuclei ($Z \geq 110$) by fossil track study of the extraterrestrial crystals: Results and perspectives [II],” *Nucl. Phys. A* **718**, 422–424.
- Pershina, V., 2002, “Theoretical predictions of properties and chemical behavior of superheavy elements,” *J. Nucl. Radiochem. Sci.* **3**, 137–141.
- Pershina, V., 2015, “Electronic structure and properties of super-heavy elements,” *Nucl. Phys. A* **944**, 578.
- Pershina, V., 2016, “A theoretical study on the adsorption behavior of element 113 and its homologue Tl on a quartz surface: Relativistic periodic DFT calculations,” *J. Phys. Chem. C* **120**, 20232–20238.
- Pershina, V., 2018, “Reactivity of superheavy elements Cn, Nh, and Fl and their lighter homologues Hg, Tl, and Pb, respectively, with a gold surface from periodic DFT calculations,” *Inorg. Chem.* **57**, 3948–3955.
- Pershina, V., J. Anton, and T. Jacob, 2009, “Theoretical predictions of adsorption behavior of elements 112 and 114 and their homologs Hg and Pb,” *J. Chem. Phys.* **131**, 084713.
- Pershina, V., A. Borschevsky, J. Anton, and T. Jacob, 2010, “Theoretical predictions of trends in spectroscopic properties of homonuclear dimers and volatility of the 7p elements,” *J. Chem. Phys.* **132**, 194314.
- Pershina, V., A. Borschevsky, E. Eliav, and U. Kaldor, 2008a, “Adsorption of inert gases including element 118 on noble metal and inert surfaces from ab initio Dirac-Coulomb atomic calculations,” *J. Chem. Phys.* **129**, 144106.
- Pershina, V., A. Borschevsky, E. Eliav, and U. Kaldor, 2008b, “Atomic properties of element 113 and its adsorption on inert surfaces from ab initio Dirac-Coulomb calculations,” *J. Phys. Chem. A* **112**, 13712–13716.
- Pershina, V., A. Borschevsky, E. Eliav, and U. Kaldor, 2008c, “Prediction of the adsorption behavior of elements 112 and 114

- on inert surfaces from ab initio Dirac-Coulomb atomic calculations,” *J. Chem. Phys.* **128**, 024707.
- Petermann, I., K. Langanke, G. Martínez-Pinedo, I. V. Panov, P. G. Reinhard, and F. K. Thielemann, 2012, “Have superheavy elements been produced in nature?,” *Eur. Phys. J. A* **48**, 122.
- Pethick, C. J., and D. G. Ravenhall, 1995, “Matter at large neutron excess and the physics of neutron-star crusts,” *Annu. Rev. Nucl. Part. Sci.* **45**, 429–484.
- Pian, E., *et al.*, 2017, “Spectroscopic identification of r-process nucleosynthesis in a double neutron-star merger,” *Nature (London)* **551**, 67.
- Pieper, W., and W. Greiner, 1969, “Interior electron shells in superheavy nuclei,” *Z. Phys. A* **218**, 327–340.
- Pitzer, K. S., 1975, “Are elements 112, 114, and 118 relatively inert gases?,” *J. Chem. Phys.* **63**, 1032–1033.
- Pitzer, K. S., 1979, “Relativistic effects on chemical properties,” *Acc. Chem. Res.* **12**, 271–276.
- Poenaru, D. N., and R. A. Gherghescu, 2018, “ α decay and cluster radioactivity of nuclei of interest to the synthesis of $Z = 119, 120$ isotopes,” *Phys. Rev. C* **97**, 044621.
- Poenaru, D. N., R. A. Gherghescu, and W. Greiner, 2011, “Heavy-particle radioactivity of superheavy nuclei,” *Phys. Rev. Lett.* **107**, 062503.
- Poenaru, D. N., R. A. Gherghescu, and W. Greiner, 2012, “Cluster decay of superheavy nuclei,” *Phys. Rev. C* **85**, 034615.
- Poenaru, D. N., R. A. Gherghescu, and W. Greiner, 2013, “Heavy-particle radioactivity,” *J. Phys. Conf. Ser.* **436**, 012056.
- Poenaru, D. N., R. A. Gherghescu, W. Greiner, and N. S. Shakib, 2015, “How Rare Is Cluster Decay of Superheavy Nuclei?,” in *Nuclear Physics: Present and Future*, FIAS Interdisciplinary Science Series (Springer, Cham), pp. 131–140.
- Polukhina, N. G., 2012, “Nuclear track detection: advances and potential in astrophysics, particle physics, and applied research,” *Phys. Usp.* **55**, 614.
- Pomeranchuk, I. Y. a., and Ya. A. Smorodinsky, 1945, “On the energy levels of systems with $Z > 137$,” *J. Fiz. USSR* **9**, 97–100 [http://www.physics.princeton.edu/~mcdonald/examples/EP/pomeranchuk_jpusr_9_97_45.pdf].
- Pomorski, K., B. Nerlo-Pomorska, J. Bartel, and C. Schmitt, 2018, “Stability of superheavy nuclei,” *Phys. Rev. C* **97**, 034319.
- Prassa, V., T. Nikšić, G. A. Lalazissis, and D. Vretenar, 2012, “Relativistic energy density functional description of shape transitions in superheavy nuclei,” *Phys. Rev. C* **86**, 024317.
- Pyykkö, P., 1988, “Relativistic effects in structural chemistry,” *Chem. Rev.* **88**, 563–594.
- Pyykkö, P., 2011, “A suggested periodic table up to $Z \leq 172$, based on Dirac-Fock calculations on atoms and ions,” *Phys. Chem. Chem. Phys.* **13**, 161–168.
- Pyykkö, P., 2016, “Is the Periodic Table all right (“PT OK”)?” *Eur. Phys. J. Web Conf.* **131**, 01001.
- Pyykkö, P., and J. P. Desclaux, 1979, “Relativity and the periodic system of elements,” *Acc. Chem. Res.* **12**, 276–281.
- Qian, Y.-Z., and G. Wasserburg, 2007, “Where, oh where has the r-process gone?,” *Phys. Rep.* **442**, 237–268.
- Radice, D., F. Galeazzi, J. Lippuner, L. F. Roberts, C. D. Ott, and L. Rezzolla, 2016, “Dynamical mass ejection from binary neutron star mergers,” *Mon. Not. R. Astron. Soc.* **460**, 3255–3271.
- Raeder, S., *et al.*, 2018, “Probing sizes and shapes of nobelium isotopes by laser spectroscopy,” *Phys. Rev. Lett.* **120**, 232503.
- Ramirez, E. M., *et al.*, 2012, “Direct mapping of nuclear shell effects in the heaviest elements,” *Science* **337**, 1207–1210.
- Rampino, S., L. Storchi, and L. Belpassi, 2015, “Gold-superheavy-element interaction in diatomics and cluster adducts: A combined four-component Dirac-Kohn-Sham/charge-displacement study,” *J. Chem. Phys.* **143**, 024307.
- Reedijk, J., 2018, “Row 7 of the periodic table complete: Can we expect more new elements; and if so, when?,” *Polyhedron* **141**, 1–4.
- Reinhard, P.-G., 2018, “Nuclear density-functional theory and fission of super-heavy elements,” *Eur. Phys. J. A* **54**, 13.
- Reinhard, P.-G., W. Greiner, and H. Arenhövel, 1971, “Electrons in strong external fields,” *Nucl. Phys. A* **166**, 173–197.
- Reinhardt, J., B. Müller, and W. Greiner, 1981, “Theory of positron production in heavy-ion collisions,” *Phys. Rev. A* **24**, 103–128.
- Ren, Z., 2002, “Shape coexistence in even-even superheavy nuclei,” *Phys. Rev. C* **65**, 051304.
- Roberto, J. B., C. W. Alexander, R. A. Boll, J. D. Burns, J. G. Ezold, L. K. Felker, S. L. Hogle, and K. P. Rykaczewski, 2015, “Actinide targets for the synthesis of super-heavy elements,” *Nucl. Phys. A* **944**, 99–116.
- Roberto, J. B., and K. P. Rykaczewski, 2018, “Discovery of element 117: Super-heavy elements and the “island of stability”,” *Sep. Sci. Technol.* **53**, 1813–1819.
- Robinson, A. P., *et al.*, 2011, “Search for a 2-quasiparticle high- K isomer in ^{256}Rf ,” *Phys. Rev. C* **83**, 064311.
- Rodríguez-Guzmán, R., and L. M. Robledo, 2017, “Microscopic description of fission in odd-mass uranium and plutonium nuclei with the Gogny energy density functional,” *Eur. Phys. J. A* **53**, 245.
- Rose, H. J., and G. A. Jones, 1984, “A new kind of natural radioactivity,” *Nature (London)* **307**, 245–247.
- Rosswog, S., M. B. Davies, W. Benz, and T. Piran, 1999, “Mass ejection in neutron star mergers,” *Astron. Astrophys.* **526**, 499–526 [<http://adsabs.harvard.edu/abs/1999A%26A...341..499R>].
- Royer, G., and H. F. Zhang, 2008, “Recent α decay half-lives and analytic expression predictions including superheavy nuclei,” *Phys. Rev. C* **77**, 037602.
- Rudolph, D., *et al.*, 2013, “Spectroscopy of element 115 decay chains,” *Phys. Rev. Lett.* **111**, 112502.
- Rudolph, D., *et al.*, 2015, “Selected spectroscopic results on element 115 decay chains,” *J. Radioanal. Nucl. Chem.* **303**, 1185.
- Rykaczewski, K. P., J. B. Roberto, N. T. Brewer, and V. K. Utyonkov, 2016, “ORNL actinide materials and a new detection system for superheavy nuclei,” *Eur. Phys. J. Web Conf.* **131**, 05005.
- Sadhukhan, J., J. Dobaczewski, W. Nazarewicz, J. A. Sheikh, and A. Baran, 2014, “Pairing-induced speedup of nuclear spontaneous fission,” *Phys. Rev. C* **90**, 061304.
- Sadhukhan, J., W. Nazarewicz, and N. Schunck, 2016, “Microscopic modeling of mass and charge distributions in the spontaneous fission of ^{240}Pu ,” *Phys. Rev. C* **93**, 011304.
- Santhosh, K. P., and C. Nithya, 2018a, “Systematic studies of α and heavy-cluster emissions from superheavy nuclei,” *Phys. Rev. C* **97**, 064616.
- Santhosh, K. P., and C. Nithya, 2018b, “Predictions on the modes of decay of odd Z superheavy isotopes within the range $105 \leq Z \leq 135$,” *At. Data Nucl. Data Tables* **121–122**, 216–255.
- Sato, T. K., *et al.*, 2015, “Measurement of the first ionization potential of lawrencium, element 103,” *Nature (London)* **520**, 209.
- Schädel, M., 2006, “Chemistry of superheavy elements,” *Angew. Chem., Int. Ed.* **45**, 368–401.
- Schädel, M., 2015, “Chemistry of the superheavy elements,” *Phil. Trans. R. Soc. A* **373**, 20140191.
- Schädel, M., 2016, “Prospects of heavy and superheavy element production via inelastic nucleus-nucleus collisions - from $^{238}\text{U} + ^{238}\text{U}$ to $^{18}\text{O} + ^{254}\text{Es}$,” *Eur. Phys. J. Web Conf.* **131**, 04001.

- Schindzielorz, N., J. Erler, P. Klüpfel, P.-G. Reinhard, and G. Hager, 2009, "Fission of super-heavy nuclei explored with Skyrme forces," *Int. J. Mod. Phys. E* **18**, 773–781.
- Schuetrumpf, B., M. A. Klatt, K. Iida, G. E. Schröder-Turk, J. A. Maruhn, K. Mecke, and P.-G. Reinhard, 2015, "Appearance of the single gyroid network phase in nuclear pasta matter," *Phys. Rev. C* **91**, 025801.
- Schuetrumpf, B., W. Nazarewicz, and P.-G. Reinhard, 2017, "Central depression in nucleonic densities: Trend analysis in the nuclear density functional theory approach," *Phys. Rev. C* **96**, 024306.
- Schunck, N., and L. M. Robledo, 2016, "Microscopic theory of nuclear fission: a review," *Rep. Prog. Phys.* **79**, 116301.
- Schwarz, W. H. E., P. Schwerdtfeger, J. G. Snijders, and E. J. Baerends, 1990, "Relativistic atomic orbital contractions and expansions: magnitudes and explanations," *J. Phys. B* **23**, 3225–3240.
- Schwarz, W. H. E., E. M. van Wezenbeek, E. J. Baerends, and J. G. Snijders, 1989, "The origin of relativistic effects of atomic orbitals," *J. Phys. B* **22**, 1515–1530.
- Schwerdtfeger, P., 2011, "The pseudopotential approximation in electronic structure theory," *ChemPhysChem* **12**, 3143–3155.
- Schwerdtfeger, P., 2016, "Toward an accurate description of solid-state properties of superheavy elements - a case study for the element Og ($Z = 118$)," *Eur. Phys. J. Web Conf.* **131**, 07004.
- Schwerdtfeger, P., G. A. Heath, M. Dolg, and M. A. Bennett, 1992, "Low valencies and periodic trends in heavy element chemistry. a theoretical study of relativistic effects and electron correlation effects in group 13 and period 6 hydrides and halides," *J. Am. Chem. Soc.* **114**, 7518–7527.
- Schwerdtfeger, P., and A. Hermann, 2009, "Equation of state for solid neon from quantum theory," *Phys. Rev. B* **80**, 064106.
- Schwerdtfeger, P., L. F. PaŠteka, A. Punnett, and P. O. Bowman, 2015, "Relativistic and quantum electrodynamic effects in super-heavy elements," *Nucl. Phys. A* **944**, 551–577.
- Schwerdtfeger, P., and M. Seth, 1998, "Relativistic effects of the superheavy elements," in *Encyclopedia of Computational Chemistry*, Vol. 4, edited by P. von R. Schleyer, P. R. Schreiner, N. L. Allinger, T. Clark, J. Gasteiger, P. A. Kollman, and H. F. Schaefer, III (John Wiley & Sons, Ltd, New York), pp. 2480–2499.
- Schwerdtfeger, P., and M. Seth, 2002, "Relativistic quantum chemistry of the superheavy elements. closed-shell element 114 as a case study," *J. Nucl. Radiochem. Sci.* **3**, 133–136.
- Seaborg, G. T., 1959, "The transcalifornium elements," *J. Chem. Educ.* **36**, 38.
- Seaborg, G. T., and W. D. Loveland, 1990, *The Elements Beyond Uranium* (Wiley-Interscience, New York).
- Sekiguchi, Y., K. Kiuchi, K. Kyutoku, and M. Shibata, 2015, "Dynamical mass ejection from binary neutron star mergers: Radiation-hydrodynamics study in general relativity," *Phys. Rev. D* **91**, 064059.
- Seth, M., F. Cooke, P. Schwerdtfeger, J.-L. Heully, and M. Pelissier, 1998, "The chemistry of the superheavy elements. ii. the stability of high oxidation states in group 11 elements: Relativistic coupled cluster calculations for the di-, tetra- and hexafluoro metallates of Cu, Ag, Au, and element 111," *J. Chem. Phys.* **109**, 3935–3943.
- Seth, M., K. Faegri, and P. Schwerdtfeger, 1998, "The stability of the oxidation state +4 in group 14 compounds from carbon to element 114," *Angew. Chem., Int. Ed.* **37**, 2493–2496.
- Seth, M., P. Schwerdtfeger, and M. Dolg, 1997, "The chemistry of the superheavy elements. i. pseudopotentials for 111 and 112 and relativistic coupled cluster calculations for (112)H⁺, (112)F₂, and (112)F₄," *J. Chem. Phys.* **106**, 3623–3632.
- Seth, M., P. Schwerdtfeger, and K. Faegri, 1999, "The chemistry of superheavy elements. iii. theoretical studies on element 113 compounds," *J. Chem. Phys.* **111**, 6422–6433.
- Sewtz, M., *et al.*, 2003, "First observation of atomic levels for the element fermium ($Z = 100$)," *Phys. Rev. Lett.* **90**, 163002.
- Sharipov, S. h., and M. J. Ermamatov, 2005, "Ground state and β -vibrational bands of axial-symmetric superheavy nuclei," *Int. J. Mod. Phys. E* **14**, 1235.
- Sheikh, J. A., W. Nazarewicz, and J. C. Pei, 2009, "Systematic study of fission barriers of excited superheavy nuclei," *Phys. Rev. C* **80**, 011302.
- Sheng, Z.-Q., L.-P. Shu, Y. Meng, J.-G. Hu, and J.-F. Qian, 2014, "Competition between α -decay and β -decay for heavy and super-heavy nuclei," *Chin. Phys. C* **38**, 124101.
- Shi, Y., J. Dobaczewski, and P. T. Greenlees, 2014, "Rotational properties of nuclei around ²⁵⁴No investigated using a spectroscopic-quality Skyrme energy density functional," *Phys. Rev. C* **89**, 034309.
- Shi, Y., D. E. Ward, B. G. Carlsson, J. Dobaczewski, W. Nazarewicz, I. Ragnarsson, and D. Rudolph, 2014, "Structure of superheavy nuclei along decay chains of element 115," *Phys. Rev. C* **90**, 014308.
- Siegel, D. M., and B. D. Metzger, 2017a, "Three-dimensional general-relativistic magnetohydrodynamic simulations of remnant accretion disks from neutron star mergers: Outflows and r -process nucleosynthesis," *Phys. Rev. Lett.* **119**, 231102.
- Siegel, D. M., and B. D. Metzger, 2017b, "Three-dimensional grmhd simulations of neutrino-cooled accretion disks from neutron star mergers," *Astrophys. J.* **858**, 52.
- Smits, O. R., P. Jerabek, E. Pahl, and P. Schwerdtfeger, 2018, "A hundred-year-old experiment re-evaluated: Accurate ab initio Monte Carlo simulations of the melting of radon," *Angew. Chem., Int. Ed.* **57**, 9961–9964.
- Smolańczuk, R., J. Skalski, and A. Sobiczewski, 1995, "Spontaneous-fission half-lives of deformed superheavy nuclei," *Phys. Rev. C* **52**, 1871–1880.
- Sobel'Man, I. I., 1972, *Introduction to the Theory of Atomic Spectra: International Series of Monographs in Natural Philosophy*, Vol. 40 (Elsevier, New York).
- Sobiczewski, A., F. A. Gareev, and B. N. Kalinkin, 1966, "Closed shells for $Z > 82$ and $N > 126$ in a diffuse potential well," *Phys. Lett.* **22**, 500–502.
- Sobiczewski, A., T. Krogulski, J. Blocki, and Z. Szymański, 1971, "Closed shells for very heavy nuclei," *Nucl. Phys. A* **168**, 519.
- Sobiczewski, A., and K. Pomorski, 2007, "Description of structure and properties of superheavy nuclei," *Prog. Part. Nucl. Phys.* **58**, 292–349.
- Soff, G., P. Schlüter, B. Müller, and W. Greiner, 1982, "Self-energy of electrons in critical fields," *Phys. Rev. Lett.* **48**, 1465–1468.
- Sonoda, H., G. Watanabe, K. Sato, K. Yasuoka, and T. Ebisuzaki, 2008, "Phase diagram of nuclear "pasta" and its uncertainties in supernova cores," *Phys. Rev. C* **77**, 035806.
- Soverna, S., *et al.*, 2005, "Thermochromatographic studies of mercury and radon on transition metal surfaces," *Radiochim. Acta* **93**, 1–8.
- Staszczak, A., A. Baran, and W. Nazarewicz, 2013, "Spontaneous fission modes and lifetimes of superheavy elements in the nuclear density functional theory," *Phys. Rev. C* **87**, 024320.
- Staszczak, A., C. Y. Wong, and A. Kossior, 2017, "Toroidal high-spin isomers in the nucleus ³⁰⁴120," *Phys. Rev. C* **95**, 054315.
- Steenbergen, K. G., J.-M. Mewes, L. F. Pasteka, H. W. Gäggeler, G. Kresse, E. Pahl, and P. Schwerdtfeger, 2017, "The cohesive energy of superheavy element copernicium determined from accurate

- relativistic coupled-cluster theory,” *Phys. Chem. Chem. Phys.* **19**, 32286–32295.
- Steenbergen, K. G., E. Pahl, and P. Schwerdtfeger, 2017, “Accurate, large-scale density functional melting of Hg: Relativistic effects decrease melting temperature by 160 K,” *J. Phys. Chem. Lett.* **8**, 1407–1412.
- Sulignano, B., *et al.*, 2012, “Investigation of high- K states in ^{252}No ,” *Phys. Rev. C* **86**, 044318.
- Sun, Y., G.-L. Long, F. Al-Khudair, and J. A. Sheikh, 2008, “ γ -vibrational states in superheavy nuclei,” *Phys. Rev. C* **77**, 044307.
- Surman, R., and M. Mumpower, 2018, “Masses and lifetimes for r -process nucleosynthesis: FRIB outlook,” *Eur. Phys. J. Web Conf.* **178**, 04002.
- Symbalisty, E., and D. N. Schramm, 1982, “Neutron star collisions and the r -process,” *Astrophys. Lett.* **22**, 143–145 [<http://adsabs.harvard.edu/abs/1982ApL....22..143S>].
- Takahashi, N., 2002, “Boiling points of the superheavy elements 117 and 118,” *J. Radioanal. Nucl. Chem.* **251**, 299–301.
- Ter-Akopian, G. M., and S. N. Dmitriev, 2015, “Searches for superheavy elements in nature: Cosmic-ray nuclei; spontaneous fission,” *Nucl. Phys. A* **944**, 177–189.
- Theisen, C. h., 2017, “Perspectives of super-heavy nuclei research with the upcoming separator-spectrometer setup S3 at GANIL/SPIRAL2 - The VAMOS Gas-Filled separator and AGATA,” *Eur. Phys. J. Web Conf.* **163**, 00059.
- Theisen, C. h., P. T. Greenlees, T.-L. Khoo, P. Chowdhury, and T. Ishii, 2015, “In-beam spectroscopy of heavy elements,” *Nucl. Phys. A* **944**, 333–375.
- Thielemann, F.-K., M. Eichler, I. V. Panov, and B. Wehmeyer, 2017, “Neutron Star Mergers and Nucleosynthesis of Heavy Elements,” *Annu. Rev. Nucl. Part. Sci.* **67**, 253–274.
- Thielemann, F.-K., J. Metzinger, and H. V. Klapdor, 1983, “Beta-delayed fission and neutron emission: Consequences for the astrophysical r -process and the age of the galaxy,” *Z. Phys. A* **309**, 301–317.
- Thoennessen, M., 2004, “Reaching the limits of nuclear stability,” *Rep. Prog. Phys.* **67**, 1187.
- Tolokonnikov, S. V., I. N. Borzov, M. Kortelainen, Yu. S. Lutostansky, and E. E. Saperstein, 2017, “Alpha-decay energies of superheavy nuclei for the Fayans functional,” *Eur. Phys. J. A* **53**, 33.
- Türler, A., R. Eichler, and A. Yakushev, 2015, “Chemical studies of elements with $Z \geq 104$ in gas phase,” *Nucl. Phys. A* **944**, 640–689.
- Türler, A., and V. Pershina, 2013, “Advances in the production and chemistry of the heaviest elements,” *Chem. Rev.* **113**, 1237–1312.
- Typel, S., and B. A. Brown, 2003, “Skyrme Hartree-Fock calculations for the α -decay Q values of superheavy nuclei,” *Phys. Rev. C* **67**, 034313.
- Utyonkov, V. K., *et al.*, 2018, “Neutron-deficient superheavy nuclei obtained in the $^{240}\text{Pu} + ^{48}\text{Ca}$ reaction,” *Phys. Rev. C* **97**, 014320.
- Viola, V. E., and G. T. Seaborg, 1966, “Nuclear systematics of the heavy elements-II lifetimes for alpha, beta and spontaneous fission decay,” *J. Inorg. Nucl. Chem.* **28**, 741–761.
- Waber, J. T., and F. W. Averill, 1974, “Molecular orbitals of PtF_6 and $\text{E}110_6$ calculated by the self-consistent multiple scattering αx method,” *J. Chem. Phys.* **60**, 4466–4470.
- Wanajo, S., 2018, “Physical Conditions for the r -process I. Radioactive Energy Sources of Kilonovae,” *Astrophys. J.* **868**, 65.
- Wang, M., G. Audi, F. G. Kondev, W. J. Huang, S. Naimi, and X. Xu, 2017, “The AME2016 atomic mass evaluation (ii). tables, graphs and references,” *Chin. Phys. C* **41**, 030003.
- Wang, N., E. G. Zhao, W. Scheid, and S. G. Zhou, 2012, “Theoretical study of the synthesis of superheavy nuclei with $Z = 119$ and 120 in heavy-ion reactions with trans-uranium targets,” *Phys. Rev. C* **85**, 041601(R).
- Wang, Y. L., H. S. Hu, W. L. Li, F. Wei, and J. Li, 2016, “Relativistic effects break periodicity in group 6 diatomic molecules,” *J. Am. Chem. Soc.* **138**, 1126–1129.
- Wang, Y. Z., S. J. Wang, Z. Y. Hou, and J. Z. Gu, 2015, “Systematic study of α -decay energies and half-lives of superheavy nuclei,” *Phys. Rev. C* **92**, 064301.
- Wapstra, A. H., *et al.*, 1991, “Criteria that must be satisfied for the discovery of a new chemical element to be recognized,” *Pure Appl. Chem.* **63**, 879.
- Ward, D. E., B. G. Carlsson, and S. Åberg, 2015, “ α -decay spectra of odd nuclei using the effective Skyrme interaction,” *Phys. Rev. C* **92**, 014314.
- Warda, M., and J. L. Egidio, 2012, “Fission half-lives of superheavy nuclei in a microscopic approach,” *Phys. Rev. C* **86**, 014322.
- Warda, M., and L. M. Robledo, 2011, “Microscopic description of cluster radioactivity in actinide nuclei,” *Phys. Rev. C* **84**, 044608.
- Warda, M., A. Zdeb, and L. M. Robledo, 2018, “Cluster radioactivity in super heavy nuclei,” *Phys. Rev. C* **98**, 041602.
- Williams, R. D., and S. E. Koonin, 1985, “Sub-saturation phases of nuclear matter,” *Nucl. Phys. A* **435**, 844–858.
- Wong, C. Y., 1973, “Toroidal and spherical bubble nuclei,” *Ann. Phys.* **77**, 279–353.
- Wong, D. P., 1979, “Theoretical justification of Madelung’s rule,” *J. Chem. Educ.* **56**, 714.
- Wu, M. R., J. Barnes, G. Martinez-Pinedo, and B. D. Metzger, 2018, “Fingerprints of heavy element nucleosynthesis in the late-time lightcurves of kilonovae,” [arXiv:1808.10459](https://arxiv.org/abs/1808.10459).
- Xu, F. R., E. G. Zhao, R. Wyss, and P. M. Walker, 2004, “Enhanced stability of superheavy nuclei due to high-spin isomerism,” *Phys. Rev. Lett.* **92**, 252501.
- Yakushev, A., *et al.*, 2014, “Superheavy element flerovium (element 114) is a volatile metal,” *Inorg. Chem.* **53**, 1624–1629.
- Yang, J. C., *et al.*, 2013, “High intensity heavy ion accelerator facility (HIAF) in China,” *Nucl. Instrum. Methods Phys. Res., Sect. B* **317**, 263–265.
- Zagrebaev, V., and W. Greiner, 2008, “Synthesis of superheavy nuclei: A search for new production reactions,” *Phys. Rev. C* **78**, 034610.
- Zagrebaev, V. I., and W. Greiner, 2015, “Cross sections for the production of superheavy nuclei,” *Nucl. Phys. A* **944**, 257–307.
- Zhang, W., J. Meng, S. Q. Zhang, L. S. Geng, and H. Toki, 2005, “Magic numbers for superheavy nuclei in relativistic continuum Hartree-Bogoliubov theory,” *Nucl. Phys. A* **753**, 106–135.
- Zhang, Y. L., and Y. Z. Wang, 2018, “Systematic study of cluster radioactivity of superheavy nuclei,” *Phys. Rev. C* **97**, 014318.
- Zhang, Z. Y., *et al.*, 2012, “Observation of superheavy nuclide ^{271}Ds ,” *Chin. Phys. Lett.* **29**, 012502–012503.
- Zhao, J., B. N. Lu, T. Nikšić, D. Vretenar, and S. G. Zhou, 2016, “Multidimensionally-constrained relativistic mean-field study of spontaneous fission: Coupling between shape and pairing degrees of freedom,” *Phys. Rev. C* **93**, 044315.
- Zhu, L., W. J. Xie, and F. S. Zhang, 2014, “Production cross sections of superheavy elements $Z = 119$ and 120 in hot fusion reactions,” *Phys. Rev. C* **89**, 024615.
- Zhu, Y., *et al.*, 2018, “Californium-254 and Kilonova Light Curves,” *Astrophys. J.* **863**, L23.
- Zhu, Y. i., and J. C. Pei, 2017, “Towards microscopic studies of survival probabilities of compound superheavy nuclei,” *Phys. Scr.* **92**, 114001.