

# Massive quantum systems as interfaces of quantum mechanics and gravity

Sougato Bose

*Department of Physics and Astronomy, [University College London](#),  
Gower Street, WC1E 6BT London, United Kingdom*

Ivette Fuentes

*School of Physics and Astronomy, [University of Southampton](#),  
Southampton SO17 1BJ, United Kingdom  
and Keble College, [University of Oxford](#), Oxford OX1 3PG, United Kingdom*

Andrew A. Geraci 

*Center for Fundamental Physics, Department of Physics and Astronomy,  
[Northwestern University](#), Evanston, Illinois 60208, USA  
and Center for Interdisciplinary Exploration and Research in Astrophysics (CIERA),  
[Northwestern University](#), Evanston, Illinois 60208, USA*

Saba Mehsar Khan 

*Department of Physics, [Lancaster University](#), Lancaster LA1 4YB, United Kingdom*

Sofia Qvarfort 

*Nordita, [KTH Royal Institute of Technology and Stockholm University](#),  
Hannes Alfvéns väg 12, SE-106 91 Stockholm, Sweden  
and Department of Physics, [Stockholm University](#), AlbaNova University Center,  
SE-106 91 Stockholm, Sweden*

Markus Rademacher 

*Department of Physics and Astronomy, [University College London](#),  
Gower Street, WC1E 6BT London, United Kingdom*

Muddassar Rashid

*Department of Physics, [King's College London](#), Strand, London WC2R 2LS, United Kingdom*

Marko Toroš 

*Faculty of Mathematics and Physics, [University of Ljubljana](#), Jadranska 19, SI-1000 Ljubljana,  
Slovenia*

Hendrik Ulbricht 

*School of Physics and Astronomy, [University of Southampton](#), Southampton SO17 1BJ,  
United Kingdom*

Clara C. Wanjura

*[Max Planck Institute for the Science of Light](#), Staudtstraße 2, 91058 Erlangen, Germany  
and Cavendish Laboratory, [University of Cambridge](#), Cambridge CB3 0HE, United Kingdom*

 (published 13 February 2025)

The traditional view from particle physics is that quantum-gravity effects should become detectable only at extremely high energies and small length scales. Owing to the significant technological challenges involved, there has been limited progress in identifying experimentally detectable effects that can be accessed in the foreseeable future. However, in recent decades the size and mass of

---

\*Contact author: [sofia.qvarfort@fysik.su.se](mailto:sofia.qvarfort@fysik.su.se)

quantum systems that can be controlled in the laboratory have reached unprecedented scales, enabled by advances in ground-state cooling and quantum-control techniques. Preparations of massive systems in quantum states pave the way for the explorations of a low-energy regime in which gravity can be both sourced and probed by quantum systems. Such approaches constitute an increasingly viable alternative to accelerator-based, laser-interferometric, torsion-balance, and cosmological tests of gravity. In this review an overview of proposals where massive quantum systems act as interfaces between quantum mechanics and gravity is provided. Conceptual difficulties in the theoretical description of quantum systems in the presence of gravity are discussed, tools for modeling massive quantum systems in the laboratory are reviewed, and an overview of the current state-of-the-art experimental landscape is provided. Proposals covered in this review include precision tests of gravity, tests of gravitationally induced wave-function collapse and decoherence, and gravity-mediated entanglement. The review concludes with an outlook and summary of the key questions raised.

DOI: [10.1103/RevModPhys.97.015003](https://doi.org/10.1103/RevModPhys.97.015003)

## CONTENTS

I. Introduction	2	1. Gravitational decoherence	26
II. Consolidating Quantum Mechanics and Gravity	4	2. Nonlinear modifications	27
A. Incorporating gravitational effects into quantum mechanics at low energies	4	3. Nonlinear and stochastic modifications	28
1. Newtonian potential in the Schrödinger equation	4	C. Entanglement mediated by gravity	30
2. Gravity and the quantum harmonic oscillator	5	1. Gravitationally interacting interferometer-based protocol	31
3. Gravity beyond the Schrödinger equation	6	2. Alternative protocols	32
4. Quantum field theory in curved spacetime	7	3. Major challenges	34
5. Harmonic oscillator in the presence of gravity using the Klein-Gordon equation	9	4. Implications	34
6. Entanglement and decoherence in noninertial frames and black holes	9	D. Other tests of gravity	35
7. Perturbative quantum gravity	10	1. Tests of the generalized uncertainty principle	35
B. Summary of challenges	10	2. Tests of the gravitational Aharonov-Bohm effect	36
1. Quantum states and the superposition principle	11	3. Tests of gravity through resonances and phonon excitations	36
2. Quantum state evolution	11	4. Tests of quantum field theory in curved spacetime and analog gravity	37
3. Quantum measurements	11	V. Experimental Pathways toward Tests of Gravity	38
4. Composite quantum systems and entanglement	12	A. State of the art of experimental tests of gravitation with massive systems	38
III. Theoretical Frameworks for Modeling Massive Quantum Systems in the Laboratory	12	1. Tests with atoms and molecules	39
A. Coupling a mechanical mode to a probe	12	2. Tests with neutrons	41
1. Optomechanical interaction	12	3. Tests with torsion balances and clamped mechanical systems	42
2. Coupling to a two-level system	13	4. Tests with levitated mechanical systems	43
B. Open-system dynamics for massive quantum systems	14	5. Approaches with hybrid systems	44
1. Quantum master equations	14	B. Controlling massive mechanical quantum systems in the laboratory	45
2. Langevin equations and input-output formalism	15	1. Squeezing and swapping of mechanics	45
C. Measurement and control of massive quantum systems	17	2. Spatial superpositions of mechanical systems	45
1. Quantum measurements	17	3. Entanglement in mechanical systems	48
2. Feedback and feedforward	18	VI. Outlook	49
D. Quantum metrology with massive quantum systems	19	Acknowledgments	50
1. Langevin description of a quantum sensor	19	References	50
2. Standard quantum limit	20		
3. Classical and quantum Fisher information	20		
IV. Proposed Tests of Gravity with Massive Quantum Systems	21		
A. Precision tests of gravity	22		
1. Weak-force detection with backaction evading measurements	22		
2. Additional weak-force detection schemes	23		
3. Weak-force detection with BECs	24		
4. Deviations from the Newtonian potential	24		
5. Tests of the equivalence principle and dark-matter searches	25		
B. Gravitational decoherence, semiclassical models, self-energy, and gravitationally induced wave-function collapse	26		

## I. INTRODUCTION

Inspired by particle physics, the central paradigm of gravitational physics indicates that quantum-gravity effects should become detectable at extremely high energies and extremely small length scales. Known as the Planck scale, this regime encompasses particle energies of  $10^{19}$  GeV or length scales of  $10^{-35}$  m. However, accessing this parameter regime is extremely challenging, and there are few prospects for achieving the necessary technological progress in the next few decades. Tests of proposals such as string theory (Dienes, 1997; Schwarz and Seiberg, 1999), loop quantum gravity

(Rovelli, 1998; Ashtekar and Bianchi, 2021), and causal dynamical triangulation (Loll, 2020), to mention just a few, therefore remain outstanding.

At lower energies, however, quantum systems with masses several orders of magnitude higher than the atomic mass scale are starting to become accessible in the laboratory. At these scales current theories predict that gravity should start affecting the dynamics of quantum states. A number of proposals and ideas have therefore been put forward. They encompass questions about superpositions of gravitational fields, gravity-induced wave-function collapse via self-gravity or decoherence due to external gravitational fields, as well as the quantum nature of the gravitational field itself. The most encouraging aspect of these proposals is that many of them appear to be experimentally and technologically accessible in the near future.

Historically, the first tests of gravity (beyond drop tests performed by Galileo) were carried out by Cavendish in the 1790s. Here a torsion balance was used to measure the gravitational constant  $G$  (Newton, Bouguer, and Cavendish, 1900). Since then, torsion balance experiments have been a cornerstone of gravitational research, and the most accurate estimates of Newton's constant, namely,  $G = (6.674\,30 \pm 0.000\,15) \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$  is based on a number of torsion balance experiments (Tiesinga *et al.*, 2021). Yet,  $G$  remains one of the least precisely known fundamental constants, and experiments actually disagree on the value of  $G$  more than they should based on the reported uncertainties (Rothleitner and Schlamminger, 2017). The smallest detected gravitational coupling measured to date was observed between 2-mm-radius gold spheres (Westphal *et al.*, 2021), and the smallest separation at which the gravitational potential has been measured is  $52 \text{ }\mu\text{m}$  (Lee *et al.*, 2020).

More focused searches for quantum gravity have been considered in the context of particle accelerators and tests of the standard model. The current energy scale of the Large Hadron Collider is 6.8 TeV per beam and 13.6 TeV during collisions, which is 16 orders of magnitude from the energies of the Planck scale ( $10^{19} \text{ GeV}$ ). Nevertheless, several proposals predict that gravity might interact more strongly at energies below the Planck scale because of the existence of additional dimensions (Arkani-Hamed, Dimopoulos, and Dvali, 1998; Dimopoulos and Landsberg, 2001). Concretely, any signs of gravity should appear mainly as missing energy signatures due to direct graviton production. No such evidence has yet been conclusively found, and the technological challenges involved in reaching higher energy with particle accelerator scales are substantial.

Another way to test gravity is to turn to cosmological observations. Indications of quantum gravity could potentially be found in the signatures of  $\gamma$ -ray bursts (Amelino-Camelia *et al.*, 1998). Another key question is how quantum-gravity effects influenced the early formation of the Universe. While many other effects have likely been washed out during the latter stages of the Universe's expansion, the detection of primordial gravitational waves could shed light on quantum-gravity effects present shortly after the big bang (Kamionkowski and Kovetz, 2016). In addition, non-Gaussian signatures in the cosmic microwave background might provide additional insights into

this period (Komatsu, 2010). Other astrophysical tests of quantum gravity have also been proposed, such as signatures in the light of distant quasars (Lieu and Hillman, 2003; Ragazzoni, Turatto, and Gaessler, 2003). However, no such signals have yet been found.

The detection of gravitational waves by the Laser Interferometer Gravitational-Wave Observatory (LIGO) Collaboration (Abbott *et al.*, 2016a) has opened yet another avenue for tests of quantum gravity. Many theories of quantum gravity [such as loop quantum gravity (Rovelli, 1998; Ashtekar and Bianchi, 2021) and string theory (Dienes, 1997)] require modification to the classical Einstein-Hilbert action, which in turn affects the propagation of gravitational waves (Alexander, Finn, and Yunes, 2008). The detection of primordial gravitational waves could potentially also shed light on which effective theories of gravity are valid at lower scales, which is otherwise known as the so-called swampland problem of string theory (Dias *et al.*, 2019). In addition, the development of Laser Interferometer Space Antenna (LISA) provides prospects for tests of the equivalence principle and Lorentz invariance (Barausse *et al.*, 2020).

In this review we provide an alternative viewpoint to the paradigm of accelerator-based, cosmological, and laser-interferometric tests of gravity. The core question that this review thus addresses is: What aspects of gravity can be tested with massive quantum systems, and what can we learn from the outcome of these tests? Here we define *massive quantum systems* as systems with masses far beyond the single-atom mass scale, such as micromechanical resonators, levitated nanobeads, and Bose-Einstein condensates (BECs). We are primarily concerned with the nonrelativistic regime, where velocities are lower than the speed of light. Our goal is to gather the tools and ideas necessary for testing gravity at low energies with massive quantum systems. Some of these notions are sketched in Fig. 1. We also hope that this review will serve as a useful introduction and overview of the field for those who are setting out to explore these questions. For a summary of tests that can be performed with tabletop experiments, see Carney, Stamp, and Taylor (2019); for a summary




FIG. 1. Intersection of quantum mechanics and gravity. The possible tests of gravity that can be performed with quantum systems are shown. (a) Quantum-enhanced measurements of gravitational effects. (b) Quantum superpositions are unstable owing to gravitational self-energy. (c) An external gravitational field that acts as the environment causes the quantum system to decohere. (d) Two quantum systems in spatial superpositions become entangled through gravity.

of those that can be performed with superconductors, see [Gallerati, Modanese, and Ummarino \(2022\)](#).

The review is structured as follows. In Sec. II we provide an overview of tensions between quantum mechanics and gravity, as well as a conceptual overview of how gravity can be incorporated into quantum mechanics at low energies. In Sec. III we provide an overview of theory tools for modeling massive quantum systems in the laboratory. Many of these tools are directly applicable to the proposed tests that follow in the next section. In Sec. IV we review proposals for tests of gravity with massive quantum systems. These include precision tests of gravity, searches for gravitational decoherence and wave-function collapse, schemes for entangling massive quantum systems through gravity, etc. In Sec. V we provide an overview of state-of-the-art gravity tests and experimental platforms that appear to be promising for tests of gravity. The review concludes with some final remarks in Sec. VI.

Before proceeding, we note that it would be impossible to give a completely balanced overview of such a broad and diverse field. We hope that this review provides a snapshot of the field today and that it ultimately helps focus efforts toward realizing some of the proposals that have been put forward for testing the interplay between quantum mechanics and gravity.

## II. CONSOLIDATING QUANTUM MECHANICS AND GRAVITY

A limitation in developing a theory of quantum gravity has been the inability to resolve the persisting tensions between the fundamental principles of quantum physics and general relativity. Current theories are good approximations in certain regimes. The relations between current theories can be found in the cube of theories in Fig. 2; see [Bronstein \(1933\)](#). A “theory of everything” that combines quantum physics and general relativity is expected to be a theory in which the speed of light  $c$ , Planck’s constant  $\hbar$ , and the gravitational constant  $G$  all play significant roles. Scientists disagree on the need to quantize gravity. In this section we discuss how gravity is different from other forces, and why it has been so difficult to construct a unified theory.

We start by exploring different attempts to incorporate gravity into quantum mechanics at low energies. We cover modifications of quantum dynamics that include gravity as a phase term or driving term. We then move to quantum field theory in curved spacetime and perturbative quantum gravity (Sec. II.A). Then we summarize the challenges that arise when general relativity (GR) is incorporated into the way that we generally perform nonrelativistic quantum mechanics in first quantization (Sec. II.B). Many of these challenges have been discussed throughout the literature. While we provide an outline of the main ideas and research directions here, this is by no means a complete account of the history or challenges that arise in the context of developing a fully fledged theory of quantum gravity.

### A. Incorporating gravitational effects into quantum mechanics at low energies

The goal of this section is to provide a high-level outline of the main theoretical ideas that enable a limited consolidation




FIG. 2. Cube of theories. Current theories are placed in a cube where the axes are small expansion parameters: the speed of light  $1/c$ , Planck’s constant  $\hbar$ , and the gravitational constant  $G$ . The axes are the speed of light  $c$ , Planck’s constant  $\hbar$ , and the gravitational constant  $G$ . While some physicists aim at building a theory of everything that includes quantum physics (depending on  $\hbar$ ) and general relativity (depending on both  $c$  and  $G$ ), others suggest that gravity should not be quantized. This review focuses mainly on the regime of the lower, light-red triangle and the upper diagonal (quantum field theory). The asterisk indicates the interplay of quantum mechanics and gravity at low energies (this could include Schrödinger-Newton-like equations, collapse models, gravitational phase shifts, etc.; see Secs. II.A and IV). Adapted from [Bronstein, 1933](#).

of quantum mechanics and gravity. For each case we detail the underlying assumptions that enable the treatment and discuss the validity and limits of the theory. Every time we introduce a new tool (such as quantum field theory), we discuss the leap beyond the current framework. However, we note that current tools are often not sufficient and that theory must ultimately be guided by experiments.

### 1. Newtonian potential in the Schrödinger equation

The second postulate of quantum mechanics dictates that the evolution of a single or composite quantum wave function  $\Psi(t, \mathbf{x})$  in time is described by the Schrödinger equation. A natural starting point when one attempts to incorporate gravity into the dynamics of a quantum system is to include it as a potential term in the Schrödinger equation. However, time in the Schrödinger equation is absolute, in contrast to general relativity, where time is an observer-dependent quantity. To use the Schrödinger equation, we must therefore make the following assumptions: we consider a single inertial frame where (i) time is well defined, (ii) the gravitational field is weak, and (iii) the quantum particles do not travel at relativistic speeds. With these assumptions it is possible to include the Newtonian potential from a source mass into the Schrödinger equation for a quantum particle with mass  $m$  as follows:

$$i\hbar \frac{\partial}{\partial t} \Psi(t, \mathbf{x}) = \left[ \frac{\mathbf{p}^2}{2m} + V_N(\mathbf{x}) \right] \Psi(\mathbf{x}), \quad (1)$$



where  $\Psi(\mathbf{x})$  is the wave function in the position basis, for which we have denoted the position vector  $\mathbf{x}$ , and the momentum operator  $p_i \equiv -i\hbar\partial/\partial x_i$  for the direction  $x_i$ . The gravitational constant  $G$  explicitly appears in the Newtonian potential  $V_N(\mathbf{x}) = -Gmm_S/|\mathbf{x} - \mathbf{x}_S|$ , where  $m_S$  is the source mass and  $\mathbf{x}_S$  is the position of the source mass. For small radial displacements  $\delta x \gg R$  away from the source located at distance  $R$ , such as for a particle moving in Earth's gravitational field, we may approximate the Newtonian potential as  $Gmm_S/|\mathbf{x} - \mathbf{x}_S| \approx mg\delta x$ , where we have introduced the gravitational acceleration  $g = m_S G/R^2$  (omitting the constant part of the potential and higher-order corrections). For such a linear potential, the solutions to Eq. (1) are given by Airy functions (Griffiths and Schroeter, 2018), and the Feynman path-integral propagator is given by (Sakurai and Commins, 1995)

$$\begin{aligned} & \langle \mathbf{x}_n, t_n | \mathbf{x}_{n-1}, t_{n-1} \rangle \\ &= \sqrt{\frac{m}{2\pi i \hbar \Delta t}} \exp \left[ i \int_{t_{n-1}}^{t_n} dt \frac{m \dot{\mathbf{x}}^2/2 - mg\delta x}{\hbar} \right], \end{aligned} \quad (2)$$

where  $\Delta t = t_n - t_{n-1}$  denotes the time increment.

Indeed, the fact that gravity can be included in the Schrödinger equation as in Eq. (1) has been experimentally verified. In their pioneering experiment Colella, Overhauser, and Werner (COW) demonstrated that neutrons passing through Earth's gravitational field acquire a phase shift that can be observed through interference (Colella, Overhauser, and Werner, 1975), with the initial theoretical proposal outlined by Overhauser and Colella (1974). This effect is sometimes referred to as gravity-induced quantum interference (Rauch and Werner, 2015).

For the description of the COW experiment, we follow Abele and Leeb (2012). The neutrons are placed in a superposition of two spatial locations. Each branch then traverses a path at two different heights above Earth. The two paths enclose a parallelogram; see the original sketch of the experiment in Fig. 3. One branch of the superposition takes the upper path  $A \rightarrow C \rightarrow D$ , and the lower branch takes the path  $A \rightarrow B \rightarrow D$ . The momentum of each neutron is determined through energy conservation, which dictates that the sum of the kinetic and potential energy must remain the same,

$$E_0 = \frac{\hbar^2 k_0^2}{2m_n} = \frac{\hbar^2 k^2}{2m_n} + m_n g H(\phi), \quad (3)$$

where  $m_n$  is the neutron mass,  $g$  is the gravitational acceleration, and  $H(\phi) = H_0 \sin(\phi)$  is the height, which depends on the orientation of the setup  $\phi$  (i.e.,  $\phi = \pi/2$  corresponds to the maximum height difference  $H_0$ , in other words, the height difference between the segments  $AB$  and  $CD$  for vertical orientation). The difference in height between the two paths means that the momentum  $p = \hbar k$  on the higher path  $CD$  must be less than the momentum  $p_0 = \hbar k_0$  on the lower path  $AB$ . The corresponding accumulated phase difference  $\Delta\Phi_{\text{COW}}$  between the two different paths is given by

$$\Delta\Phi_{\text{COW}} = \Delta k S \approx -q_{\text{COW}} \sin(\phi), \quad (4)$$




FIG. 3. Superpositions of neutrons along different paths in a gravitational field. From Colella, Overhauser, and Werner, 1975.

where  $\Delta k = k - k_0$  and  $S$  is the path length of the segments  $AB$  and  $CD$ . It is instructive to write the final phase using the geometric factor  $q_{\text{COW}} = \lambda m_n^2 g A_0 / \hbar^2$ , where  $A_0 = H_0 S$  is the area of the parallelogram (i.e., the interferometric area) and  $\lambda$  is the neutron wavelength. Depending on the value of the phase difference  $\Delta\Phi_{\text{COW}}$ , we then observe either destructive or constructive interference of each individual neutron with itself as it is recombined at the end of the interferometric paths.

Gravitational phase shifts, such as the one in Eq. (4), have sparked numerous investigations, not only with neutrons but also with cold atoms, where experiments span from tests of the equivalence principle to searches of dark matter (Tino, 2021). However, subtleties arise when additional corrections from general relativity are similarly included as phase shifts. We return to this question in Sec. II.A.3.

## 2. Gravity and the quantum harmonic oscillator

Before moving beyond Newtonian gravity, we mention another route for including gravity as a perturbation in the dynamics of a quantum system. One of the few known analytic solutions to the Schrödinger equation apart from the hydrogen atom is the quantum harmonic oscillator (QHO). The potential  $V(\mathbf{x})$  in Eq. (1) is quadratic in  $\mathbf{x}$ , such that in a single spatial dimension  $x$ ,  $V(x) \propto x^2$ . The resulting solutions describe a harmonic oscillator with quantized energy levels. The QHO is important in the context of tests of gravity because gravitational effects can be described as effective interaction terms as modifications to the quadratic trapping term.

In the language of second quantization, the Hamiltonian of the QHO reads

$$\hat{H}_{\text{QHO}} = \frac{1}{2} m \omega_m^2 \hat{x}^2 + \frac{1}{2m} \hat{p}^2, \quad (5)$$

where  $m$  is the mass of the system,  $\omega_m$  is the angular frequency, and  $\hat{x}$  and  $\hat{p}$  are the position and momentum operators. In second quantization  $\hat{x}$  and  $\hat{p}$  are given in terms of the annihilation and creation operators  $\hat{a}$  and  $\hat{a}^\dagger$  as

$$\hat{x} = \sqrt{\frac{\hbar}{2\omega_m m}}(\hat{a}^\dagger + \hat{a}), \quad \hat{p} = i\sqrt{\frac{\hbar\omega_m m}{2}}(\hat{a}^\dagger - \hat{a}), \quad (6)$$

where  $[\hat{a}, \hat{a}^\dagger] = 1$ .

Gravitational effects can be included in the description of the center-of-mass dynamics of these quantum systems. We consider a pointlike gravitational source of mass  $m_S$  situated at a distance  $r_0$  away from the quantum systems (Scala *et al.*, 2013; Qvarfort *et al.*, 2018; Rätzel *et al.*, 2018). The Newtonian potential is given by  $Gmm_S/r_0$ . Assuming that the quantum system that probes the gravitational field is perturbed by a small distance  $\delta x$ , we expand the Newtonian potential in terms of  $\delta x$  as

$$V(r - \delta x) \approx \frac{Gmm_S}{r_0} \left( 1 + \frac{\delta x}{r_0} + \frac{(\delta x)^2}{2r_0^2} + \mathcal{O}[(\delta x)^2] \right). \quad (7)$$

Inserting this potential in the Hamiltonian of the QHO and replacing the perturbation  $\delta x$  with the quantum operator  $\hat{x}$ , we obtained a modified Hamiltonian of the QHO,

$$\hat{H}_{\text{QHO}} = \frac{1}{2}m\omega_m^2\hat{x}^2 + \frac{1}{2m}\hat{p}^2 + \mathcal{G}_1\hat{x} + \mathcal{G}_2\hat{x}^2 + \mathcal{O}(\hat{x}^3), \quad (8)$$

where we have defined

$$\mathcal{G}_1 = \frac{Gmm_S}{r_0^2}, \quad \mathcal{G}_2 = \frac{Gmm_S}{2r_0^3}, \quad (9)$$

and where higher orders of the perturbation can be similarly defined, although the resulting nonlinear equations of motion are generally challenging to solve. We provide an overview of quantum sensing of gravitational fields with quantum optomechanical systems and with Bose-Einstein condensates, where the force enters as described here in Sec. IV.A.

### 3. Gravity beyond the Schrödinger equation

Thus far in our presentation, incorporating gravity into quantum mechanics has been straightforward since both the Newtonian potential and the Schrödinger equation are non-relativistic and share a joint notion of absolute time. However, problems start to arise as we go further and include additional effects from general relativity such as time dilation.

Consider a quantum particle in a spatial superposition where each branch of the superposition follows a different spacetime trajectory, not unlike the COW experiments discussed in Sec. II.A.1. In that case, we assumed that a weak gravitational effect introduces a potential difference. However, if the gravitational effect is strong, such that the background spacetime can no longer be considered flat across the relevant length scale, the two branches of the superposition experience different proper times throughout their trajectories and should therefore evolve at different rates. There is no prescription for how to perform calculations in this scenario in the absence of an external observer. However, some studies have considered using the Schrödinger equation to describe the system only in the reference frame in which the system is measured. That is, any evolution of the quantum states, as seen in the laboratory

frame, can be described as a result of some effective dynamics that arise from gravity. For example, Zych *et al.* (2011) proposed that internal degrees of freedom of particles can act as clocks that record the elapsed proper time. The addition of internal clock states solves the challenge of interpreting a phase shift as either a potential shift or a redshift due to differences in proper time. Similarly, Pikovski *et al.* (2015) showed that the effects of time dilation, as seen by an external observer, result in decoherence in composite particles. That is, by defining a Hamiltonian for the center-of-mass and internal degrees of freedom, general relativistic corrections are incorporated into the full dynamics of the particle. When the superposition branches are brought back together, the effect manifests as decoherence. Several other mechanisms that cause decoherence have been derived using similar semi-classical arguments. We cover these arguments in Sec. IV.B. It has been pointed out in the literature that there are inconsistencies that arise when proper times are introduced in quantum superpositions (Marzlin, 1995; Sonnleitner and Barnett, 2018; Schwartz and Giulini, 2019b). The argument is that classical systems couple to gravity via the *minimum coupling principle*, which is diffeomorphism invariant, and that the coupling is not consistent with Galilean-invariant equations such as the Schrödinger equation (Schwartz and Giulini, 2019a). However, the argument of Pikovski *et al.* (2015) in response is that there is no inconsistency if no nonrelativistic physics is imposed and the correct Schrödinger equation can in fact be derived with relativistic extensions.

Work on describing post-Newtonian phases in atom interferometry considers the free propagation of the atoms along spacetime geodesics. The atom-light interaction is described in a covariant manner to calculate the leading-order general relativistic effects (Dimopoulos *et al.*, 2008; Werner *et al.*, 2024). In addition, these considerations offer a possible reinterpretation of the COW experiment; see Sec. II.A.1. The question becomes: Can the phase shift that is detected by an atom interferometer be interpreted as a gravitational redshift? This interpretation was first suggested by Müller, Peters, and Chu (2010) and was followed by a vigorous debate in the community; see Tino (2021) and references therein. The ambiguity arises because the phase shift in the atom interferometer can be interpreted either as an effective potential shift or as due to redshift that has resulted from the differences in proper times.

Many of the aforementioned works consider interferometry of quantum states in the presence of gravity. Recombining the two states at the end of the interferometry process naturally involves taking a notion of an appropriate inner product. Note that the inner product between quantum states in nonrelativistic quantum mechanics, where there is an absolute time, is not Lorentz invariant. In the position representation, the inner product is given by

$$\int_{\mathbb{R}^3} d\mathbf{x} \psi_j^*(t, x) \psi_\ell(t, x) = \delta_{j,\ell}, \quad (10)$$

where  $\psi_j(t, x)$  are wave functions and  $\delta_{j,\ell}$  is the Kronecker delta function. A consequence is that two quantum states in different inertial frames or different spacetime locations cannot be consistently compared. Since the description of

measurements and averages requires the inner product, the experimental observations cannot be described appropriately with this inner product. This poses a problem that is often overlooked when quantum interferometry is described in curved spacetime. However, as long as a single laboratory frame is considered, the inner product is well defined. In Sec. II.A.4 we introduce the Klein-Gordon inner product, which is appropriate for scalar relativistic quantum fields.

If we want to describe quantum systems in a manner that is consistent with general relativity, we must use a covariant formalism where the equations and the inner products are Lorentz invariant. Quantum field theory in curved spacetime enables such a description in the low-energy regime.

#### 4. Quantum field theory in curved spacetime

Thus far, we have discussed approaches for describing the effects of weak gravitational effects using the Schrödinger equation. Such schemes consider a single inertial frame or study the differences between two inertial frames as an effective Hamiltonian. That is, relativistic corrections are treated as dynamical perturbations in the Schrödinger equation, where time remains absolute. However, in relativity measurements of well-defined quantities must coincide in different frames, and a consistent description in noninertial frames is also required. This is possible only through a covariant formalism. The question becomes: Can quantum systems be described using equations that are Lorentz invariant?

The answer is affirmative within some restrictions. It is possible to describe some aspects of the interplay of quantum physics and general relativity using quantum field theory (QFT) in curved spacetime (CS). QFT in CS is a semiclassical approach that considers the behavior of quantum fields on a classical spacetime background. Spacetime is not quantum, rather, it is a solution of Einstein field equations. The formalism describes multiparticle effects, and it turns out that considering single particles such as single atoms is nontrivial. Quantum field theory in curved spacetime has enabled the study of some effects in quantum physics and quantum information in relativistic settings, including entanglement and its applications in noninertial frames and curved spacetime. Section II.A.6 includes a discussion on the degradation of entanglement as seen by observers in uniform acceleration.

Here we provide an overview of QFT in CS. More comprehensive accounts of this research field were given by Birrell and Davies (1982), Fulling (1989), Schweber (2005), and Parker and Toms (2009). We limit our discussion to a scalar field, which is the simplest case. See Hollands and Wald (2015) for a full review of QFT in CS (including other cases such as the Dirac equation) that describes fermionic fields. In the case of a single scalar field, the Schrödinger equation is replaced by the Klein-Gordon equation, which reads (having set  $\hbar = c = 1$ )

$$(\square_g - m^2)\phi = 0, \quad (11)$$

where  $\square_g = g^{\mu\nu}\nabla_\mu\nabla_\nu$  is the d'Alembertian operator associated with the metric  $g$  and  $\phi$  is the scalar field with mass  $m$ .

Under superficial inspection the Klein-Gordon equation in flat spacetime looks similar to the Schrödinger equation. The main difference is that it has a second derivative in both the spatial and temporal coordinates, making it invariant under Lorentz transformations as required by relativity. Historically, the Klein-Gordon equation has been derived from a relativistic (classical) Hamiltonian of a particle and then interpreting the momentum and position as operators (Schweber, 2005)

$$H_r = \sqrt{(\mathbf{p}c)^2 + (mc^2)^2}, \quad (12)$$

with the momentum vector  $\mathbf{p}$ . To avoid the problem of the square root, which would appear if we inserted  $H_r$  into the Schrödinger equation, the formalism instead considers the squared operator equation

$$-\hbar^2 \frac{\partial^2}{\partial t^2} \phi(t, \mathbf{r}) = [\hat{\mathbf{p}}^2 c^2 + (mc^2)^2] \phi(t, \mathbf{r}). \quad (13)$$

In contrast to the Schrödinger equation, the resulting wave equation is Lorentz invariant,

$$\left[ \left( \frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2 \right) + \left( \frac{mc}{\hbar} \right)^2 \right] \phi(t, \mathbf{r}) = 0. \quad (14)$$

However, interpreting  $\phi(t, \mathbf{r})$  as a wave function is generally problematic. This is because the probability density  $\rho \equiv (i\hbar/2mc^2)(\phi^*[\partial_t\phi] - [\partial_t\phi^*]\phi)$  and the probability current  $j_\ell \equiv (\hbar/2mi)(\phi^*[\partial_\ell\phi] - [\partial_\ell\phi^*]\phi)$  defined such that they satisfy the continuity equation  $\nabla \cdot \mathbf{j} + \partial_\mu \rho = 0$  can take negative values due to the second derivative in time in Eq. (14) (Schweber, 2005). Specifically, the Klein-Gordon inner product for  $\phi(t, \mathbf{r})$  is derived from the continuity equation and is given by

$$(\phi(t, \mathbf{r}), \psi(t, \mathbf{r})) = -i \int_\Sigma [\psi^* \partial_\mu \phi - (\partial_\mu \psi^*) \phi] d\Sigma^\mu,$$

where  $\Sigma$  is a spacelike hypersurface. While the inner product is Lorentz invariant by construction, it yields negative probabilities in some cases. Therefore, interpreting  $\phi(t, \mathbf{r})$  as the wave function of a single particle is inconsistent with quantum mechanics.

The problem can be solved in special cases when the spacetime has specific symmetries. In these cases one can construct an operator-valued function by associating creation and annihilation operators  $\hat{a}_k$  and  $\hat{a}_k^\dagger$  for a mode  $k$  to the positive and negative mode solutions of the Klein-Gordon equation  $u_k$  and  $u_k^*$ ,

$$\hat{\phi}(t, \mathbf{r}) = \int dk (u_k \hat{a}_k + u_k^* \hat{a}_k^\dagger). \quad (15)$$

This operator-valued function obeys the Klein-Gordon equation, and particle states with positive norms are defined by the action of creation operators on the vacuum state. The vacuum state is defined by  $a_k|0\rangle = 0$ . The operators act on the Fock space  $\bigoplus_{n=0}^\infty \mathcal{H}^{\otimes n}$ , where  $|0\rangle \in \mathbb{C}$ ,  $\mathcal{H}$  is the single-particle Hilbert space,  $\mathcal{H}^{\otimes n}$  is the  $n$ -particle sector, and  $\bigoplus, \otimes$  are



the direct sum and the tensor product, respectively. This construction is possible only when the solutions of the Klein-Gordon equation can be classified in positive and negative frequency mode solutions. This classification is possible only when the spacetime admits a timelike Killing vector field. A Killing vector field is the tangent vector space of transformations that leave the metric invariant. Spacetimes that admit a global timelike Killing vector field, such as Minkowski or the Schwarzschild spacetime, are stationary. A consistent theory can be constructed for spacetimes with these symmetries having a well-defined probability distribution. Well-known examples are quantum fields in noninertial frames and eternal black holes (Birrell and Davies, 1982; Schweber, 2005).

A key problem in QFT in CS is that particles are not well defined. Only observers flowing along timelike Killing vector fields can describe particle states in a meaningful way. In general, curved spacetimes do not admit timelike Killing vector fields globally. Moreover, in the case in which the spacetime does have a global timelike Killing vector field, the vector field is not necessarily unique. A consequence of this is that the field can be equivalently quantized in several different bases corresponding to different Killing observers. Using the Klein-Gordon inner product, it is possible to find a unitary transformation, called a Bogoliubov transformation, that relates the solutions to the equation in the different basis. This induces a transformation between the creation and annihilation operators in the old basis  $\hat{a}_j$  and  $\hat{a}_j^\dagger$  and new operators  $\hat{b}_k$  and  $\hat{b}_k^\dagger$  associated with the solutions in a different basis. In the new frame the operators are given by

$$\hat{b}_k = \int dk (\alpha_{kj} \hat{a}_j + \beta_{jk} \hat{a}_j^\dagger), \quad (16)$$

where  $\alpha_{kj}$  and  $\beta_{jk}$  are called Bogoliubov coefficients. A consequence is that the vacua are not equivalent, and the particle content of the field is observer dependent. The vacuum that was annihilated by the mode operator  $\hat{a}_j|0\rangle$  in the first frame no longer appears to be empty in the second frame, since

$$\langle 0 | \hat{b}_k^\dagger \hat{b}_k | 0 \rangle = \int dk |\beta_{kj}|^2 \neq 0. \quad (17)$$

This has important consequences for the study of entanglement in relativistic quantum fields since the notion of subsystems is indispensable for storing information; see Sec. II.A.6.

Some spacetimes do not admit global timelike Killing vector fields but do have spacetime regions where particles can be well defined. An example is the metric that describes a toy model for the expansion of the Universe, which is known as the Friedmann-Lemaître-Robertson-Walker metric. The spacetime is not stationary, and timelike Killing vector fields are defined only in the past and future infinity regions; see Birrell and Davies (1982) and references therein. It is possible to show, using Bogoliubov transformations, that the vacuum state at past infinity has entangled particles in the future infinity region (Ball, Fuentes-Schuller, and Schuller, 2006).

Some particularly interesting consequences of QFT in noninertial frames and CS are the Unruh-Davies-Fulling effect (Fulling, 1973; Davies, 1975; Unruh, 1976) and the closely related Hawking radiation effect (Hawking, 1975). The inertial vacuum appears to be populated by particles in a thermal state for uniformly accelerated observers. A region of spacetime becomes inaccessible to noninertial observers due to their acceleration. Tracing over the field modes in the casually discounted region leads to mixed states. In the case of uniformly accelerated observers in flat spacetime, the Minkowski vacuum corresponds to a thermal state with Unruh temperature

$$T_{\text{Unruh}} = \frac{\hbar a}{2\pi c k_B}, \quad (18)$$

where  $k_B$  is the Boltzmann constant. A similar situation occurs in black-hole spacetimes, where observers hovering outside the horizon lose access to the region inside the black hole. The inertial vacuum state corresponds to a thermal state for observers at a fixed distance from an eternal black hole. This spacetime is stationary and the black-hole mass is constant. In the case of a collapsing star, the spacetime is not stationary and there is energy flux known as Hawking radiation, where

$$T_{\text{Hawking}} = \frac{\hbar c^3}{8\pi G M k_B} \quad (19)$$

is the Hawking temperature for a black hole with mass  $M$ .

These results further emphasize the notion that the vacuum in a curved spacetime is not unique, which has implications for the coherence and entanglement of quantum systems. We explore the consequences for entanglement in curved spacetime in Sec. II.A.6.

A main lesson that we learn from the development of QFT in CS is that fields, not particles, are fundamental. Particles are derived notions that do not always have a viable interpretation. QFT is a multiparticle theory, and single particles can be described using this formalism only when energies are not sufficient to create new particles. In this low-energy case, it is possible to restrict the system to the single-particle sector because the energies present are not high enough to create new particles. In Sec. II.A.5 we discuss attempts to construct a covariant description of the quantum harmonic oscillator in the presence of curved spacetime using the Klein-Gordon equation and the restriction to the aforementioned single-particle sector.

A full reconciliation between quantum mechanics and special relativity requires QFT (in flat spacetime), which has been demonstrated numerous times in particle accelerators. However, QFT in curved space still awaits experimental corroboration. In Sec. IV.D.4 we discuss proposals to test its key predictions using Bose-Einstein condensates. Although QFT in curved spacetime enables the study of some effects at the interface of quantum physics and GR, including entanglement and decoherence (see Sec. IV.B), a full reconciliation between the theories must include the effects of quantum matter on the background metric itself. These effects, known as backreaction, are beyond the scope of QFT in CS. That is,



QFT in CS is limited by a semiclassical description where the spacetime is assumed to be a classical background given by Einstein's equations, and only fields are quantized. Ultimately, the difficulty with including backreaction in a covariant theory of quantum fields is the main difficulty with developing a theory of quantum gravity.

### 5. Harmonic oscillator in the presence of gravity using the Klein-Gordon equation

An alternative approach to describe a harmonic oscillator in the presence of a gravitational field beyond the Newtonian approximation is to use a Klein-Gordon equation and the Klein-Gordon inner product, which, as introduced in Sec. II.A.4, are compatible with both general relativity and quantum physics. The Klein-Gordon equation describes a scalar field in curved spacetime. However, a single particle (such as an atom) in the presence of the gravitational field of a spherical mass can be described by restricting the solutions to the single-particle sector and using the Schwarzschild metric (Marzlin, 1995; Sonnleitner and Barnett, 2018; Schwartz and Giulini, 2019b; Huimann, 2020). The problem with this approach is that the Klein-Gordon equation does not have a trapping potential term.

A solution to this was proposed by Huimann (2020), who designed an effective spacetime metric that not only included the external gravitational field but also mimicked the relevant features of an oscillating trapping potential. The effective metric reduces to the Newtonian potential in the nonrelativistic approximation, and the equation reduces to the Schrödinger equation of a harmonic oscillator in the presence of Newtonian gravity. However, solving the equation beyond this approximation is challenging, and only certain solutions are possible in special cases.

An alternative approach that also uses a restriction of the dynamics to the single-particle sector considers a classical system coupling to gravity via minimum coupling and then quantizes the system via canonical quantization (Schwartz and Giulini, 2019a). This approach was inspired by work computing relativistic corrections of an atom interacting with the electromagnetic field (Sonnleitner and Barnett, 2018) and on studies of the dipole coupling between a system of  $N$  particles with a total charge zero and the electromagnetic field in the presence of a weak gravitational field (Marzlin, 1995). More recently, a full first-order post-Newtonian expansion was performed by Schwartz and Giulini (2019b).

### 6. Entanglement and decoherence in noninertial frames and black holes

Entanglement is a key notion in quantum mechanics and is often regarded as the true herald of quantumness. In a single inertial frame, the Schrödinger equation readily describes how two subsystems interacting via a potential gradually become entangled. However, describing entanglement in relativistic settings is more complicated. Entanglement strongly depends on the notion of subsystem and bipartition. In the quantum theory subsystems can always be defined independently of the observer. As a consequence, entanglement is conserved for moving observers.

In the relativistic case entanglement is invariant only in flat spacetime and only if observers move with a constant velocity. It was shown that the Minkowski vacuum contains spatial correlations that can produce entanglement between initially uncorrelated atoms interacting with the vacuum state (Valentini, 1991; Reznik, 2003; Wang and Blencowe, 2021).

Consider two inertial observers in flat spacetime who are performing an experiment to determine the degree of entanglement between two particles, such as two photons or two fermions. We assume that they find that the systems are maximally entangled. If two uniformly accelerated observers try to determine the degree of entanglement in the same system, they find that there are many particles instead of just two. The notion of the system's bipartition is lost due to the Fulling-Davies-Unruh effect, which was introduced in Sec. II.A.4. The inertial vacuum state corresponds to a thermal state for uniformly accelerated observers. In QFT a well-defined notion of the subsystem is possible only when global bosonic or fermionic modes with sharp frequency are considered. This is because the frequency is invariant, although the number of particles in the mode varies with acceleration. For uniformly accelerated observers a region of spacetime becomes inaccessible and global states become more mixed at higher accelerations, decreasing entanglement. Nonuniform motion and thus gravity produce decoherence (Fuentes-Schuller and Mann, 2005). For localized systems, such as moving cavities (Bruschi, Fuentes, and Louko, 2012) and propagating wave packets in curved spacetime (Bruschi, Ralph *et al.*, 2014), motion and gravity can either degrade states or create entanglement (Friis *et al.*, 2013).

In curved spacetime the situation is even more complex because inertial observers disagree with the particle content of the field. As a consequence, there is no well-defined notion of entanglement in curved spacetime. Entanglement between global modes can be studied only in asymptotically flat spacetimes such as black holes (Adesso and Fuentes, 2009; Jing and Jing, 2023; Wu, Liu *et al.*, 2023) and cosmological toy models (Ball, Fuentes-Schuller, and Schuller, 2006).

The study of the observer-dependent nature of entanglement (Alsing and Fuentes, 2012) in relativistic settings has been a topic of interest in the field of *relativistic quantum information*. For an overview of the field, see Mann and Ralph (2012). The field is concerned with studying relativistic effects on quantum technologies, including quantum communications (Bruschi, Ralph *et al.*, 2014), and on addressing fundamental questions in quantum field theory (Lopp and Martín-Martínez, 2018; Barman *et al.*, 2023; Wu, Zeng, and Wu, 2023), black holes (Ng *et al.*, 2022), cosmology (Bubuianu, Vacaru, and Veliev, 2021), and high-energy physics (Bertlmann and Hiesmayr, 2001; Naikoo *et al.*, 2020) with an information-theoretic perspective.

A good example where notions of quantum information are applicable to fundamental questions is the well-known *information loss paradox* in black holes. Information stored in pure states in the spacetime of a black hole is lost due to states becoming completely mixed after the black hole evaporates via Hawking radiation. Here the interplay of quantum field theory and general relativity leads to a paradox, the resolution of which seems to require giving up fundamental principles such as unitarity, locality, and the equivalence principle.

Quantum fields in black-hole spacetimes give rise to one of the starkest indications of the incompatibility of quantum theory and general relativity. A large amount of work has focused on addressing this problem using quantum information; see the recent papers by [Yoshida \(2019\)](#) and [Penington \(2020\)](#). The question becomes: Could entanglement carry the lost information out of the black hole? The distribution of entanglement, via the monogamy of entanglement, between modes inside and outside of the black hole ([Adesso and Fuentes, 2009](#)) could play a role in the potential resolution to the paradox ([Merali, 2013](#)). However, this resolution requires entanglement to be somehow broken at the horizon. It was conjectured that observers falling into a black hole encounter a fire wall made of high-energy quanta at (or near) the event horizon, which breaks the entanglement ([Almheiri \*et al.\*, 2021](#)). However, there is an ongoing discussion in the community on whether or not this resolves the matter.

## 7. Perturbative quantum gravity

In the preceding sections we assumed that gravity is a background gravitational field obtained by solving Einstein's equations with classical sources (for example, the background gravitational field created by Earth). Such analysis does, however, not take into account that the quantum matter (i.e., the quantum system in the laboratory) can also be a source of gravity. This effect is known as gravitational backreaction and is one of the many challenging problems that a fully fledged quantum theory of gravity should address. The backreaction from a quantum system could be naively included in Einstein field equations with both the spacetime metric and the stress-energy tensors promoted to quantum operators. However, when we try to perturbatively quantize gravity, we are faced with the problem of an infinite number of free parameters coming from the high-energy regime that need to be fixed using experimental data; i.e., we get a theory that is nonrenormalizable and thus does not have predictive power. The full quantization of gravity is an open problem ([Weinberg, 1980](#); [Niedermaier, 2007](#); [Shomer, 2007](#); [Reuter and Saueressig, 2010](#)).

Instead, we limit the discussion of the gravitational backreaction here to the perturbative regime of gravity at low energies. In this regime general relativity can be quantized by following analogous steps, as with any other field theory. This was done in the seminal paper by [Gupta \(1952b\)](#), who used the Gupta-Bleuler formalism ([Bleuler, 1950](#); [Gupta, 1950](#)) applied to the approximate linear form of Einstein's gravitational field and later generalized beyond the linear case ([Gupta, 1952a](#)). Quantum general relativity can be treated as an effective field theory (EFT) at low energies using the covariant Feynman path-integral approach, which allows one to make predictions without full knowledge of the theory at high energies ([Donoghue, 1994](#)). Within this framework we first expand the metric  $g_{\mu\nu}$  as

$$\hat{g}_{\mu\nu} = \eta_{\mu\nu} + \hat{h}_{\mu\nu}, \quad (20)$$

where  $\eta_{\mu\nu}$  is the Minkowski spacetime metric (or in general some other background gravitational field  $\bar{g}_{\mu\nu}$ ) and  $\hat{h}_{\mu\nu}$

contains the fluctuations of the metric that we quantize. Specifically, we can then obtain the graviton propagator,

$$\frac{iP_{\mu\nu,\alpha\beta}}{k^2 + i\epsilon}, \quad (21)$$

where  $k^\mu$  is the four-momenta ( $k^2 = k_\mu k^\mu$ ) and the projection operator is given (in the harmonic gauge) by

$$P_{\mu\nu,\alpha\beta} = \frac{1}{2}(\eta_{\mu\alpha}\eta_{\nu\beta} + \eta_{\mu\beta}\eta_{\nu\alpha} - \eta_{\mu\nu}\eta_{\alpha\beta}). \quad (22)$$

The interaction Lagrangian is given by

$$\mathcal{L}_{\text{int}} = \frac{1}{2}\hat{h}^{\mu\nu}\hat{T}_{\mu\nu}, \quad (23)$$

where  $\hat{T}_{\mu\nu}$  is now the stress-energy tensor produced by quantum systems. Starting with Eq. (23), we can obtain matter-graviton vertices. In addition, we have graviton-graviton vertices as the graviton couples to all energetic particles, including to itself. Once the Feynman rules are obtained, we can then perform calculations as in other quantum field theories; see [Scadron \(2006\)](#).

Suppose that we have two nonrelativistic massive quantum systems. We can write the corresponding stress-energy tensor as

$$\hat{T}_{\mu\nu} = \hat{T}_{\mu\nu}^{(m)} + \hat{T}_{\mu\nu}^{(M)}, \quad (24)$$

where  $\hat{T}_{\mu\nu}^{(m)}$  ( $\hat{T}_{\mu\nu}^{(M)}$ ) is the contribution from a system of mass  $m$  ( $M$ ). Using perturbation theory in the EFT context discussed in Eqs. (20)–(23), we can then find the corrections to the Newtonian potential,

$$\hat{V} = -\frac{GMm}{\hat{r}} \left[ 1 + 3\frac{G(M+m)}{\hat{r}c^2} + \frac{41}{10\pi}\frac{G\hbar}{\hat{r}^2c^3} \right], \quad (25)$$

where  $\hat{r}$  denotes the distance between the two systems. The first term in Eq. (25) is the tree-level contribution, while the second and third terms come from one loop Feynman diagrams. The second and third terms were calculated with three techniques: Feynman diagrams ([Kirilin and Khriplovich, 2002](#); [Bjerrum-Bohr, Donoghue, and Holstein, 2003](#)), unitarity-based methods ([Bjerrum-Bohr, Donoghue, and Vanhove, 2014](#); [Holstein, 2017](#)), and dispersion relations ([Bjerrum-Bohr, Donoghue, and Holstein, 2003](#)). In addition, the result in Eq. (25) applies to particles of any spin and thus is universal ([Holstein and Ross, 2008](#)).

To conclude this section we note that there are many ways in which gravitational effects can be incorporated into the dynamics of quantum systems. To establish which ones are accurate, we must ultimately be guided by experiments.

## B. Summary of challenges

We have outlined ways in which gravity can be consolidated with quantum mechanics in a limited way, although many conceptual and mathematical challenges remain. Here we summarize the challenges by examining the postulates of

quantum mechanics one by one. For each challenge, we mention the resolution when one exists (for example, quantum field theory successfully combines quantum mechanics with special relativity). The remaining challenges must ultimately be determined by experiments.

### 1. Quantum states and the superposition principle

The first postulate states that nonrelativistic quantum mechanics (NRQM) in first quantization associates a Hilbert space with every quantum system by representing the states of a system with vectors in a Hilbert space. To preserve probabilities, physical states  $|\psi_j(t)\rangle$  must be normalized with respect to the inner product  $\langle\psi_j(t)|\psi_\ell(t)\rangle = \delta_{j,\ell}$ , which is shown in Eq. (10). Physical quantities are given in terms of expectation values, which are evaluated using this inner product. A quantum superposition corresponds to a state  $|\Psi(t)\rangle$  that is a linear combination of basis states  $|\psi_j(t)\rangle$  and amplitudes  $c_j$ ,

$$|\Psi(t)\rangle = \sum_j c_j |\psi_j(t)\rangle, \quad (26)$$

in which  $\sum_j |c_j|^2 = 1$  ensures that the superposition state is normalized, hence allowing for a probabilistic interpretation of the theory. Any such superposition remains a valid quantum state.

Several conflicts between this postulate and relativity can be identified:

- (i) To satisfy Lorentz invariance, space and time must enter on an equal footing. As mentioned in Sec. II.A.3, the inner product in Eq. (10) is not Lorentz invariant (Birrell and Davies, 1982), which implies that physical quantities in NRQM are not compatible with physical quantities in relativity.
- (ii) The wave functions  $|\psi_j(t)\rangle$ ,  $|\psi_\ell(t)\rangle$  in Eqs. (10) and (26) are evaluated at equal times  $t$ . Time enters as a global parameter, while in special and general relativity it is a relative concept that depends on the given world line. Furthermore, whereas quantum states can be in a superposition of several spatial locations, in curved spacetime time can pass at different rates in different locations.
- (iii) In both special and general relativity, the times at which events occur are observer dependent. For spacelike events the order in which they occur may change. Such explicit notions of causality are not part of the framework of NRQM but must instead be added by hand.
- (iv) It has been argued that the superposition principle is in conflict with the principle of covariance (Penrose, 1986, 1996) and with the equivalence principle (Howl, Penrose, and Fuentes, 2019). An argument challenging this view was recently put forward (Giacomini and Brukner, 2022).

As we saw in Sec. II.A.4, some of these points can be addressed by moving to quantum field theory and considering fields rather than particles.

### 2. Quantum state evolution

The time evolution of quantum states in NRQM is given by the Schrödinger equation in Eq. (1). There are a number of conflicts with general relativity:

- (i) As we saw in Sec. II.A.4, Lorentz invariance requires derivatives with respect to time and space to be of the same order, which is not the case for the Schrödinger equation in Eq. (1). However, relativistically invariant versions of the Schrödinger equation, such as the Klein-Gordon equation [Eq. (14)] and the Dirac equation, in conjunction with relinquishing the notion of single-particle states are needed to overcome this inconsistency, as illustrated by quantum field theory.
- (ii) In quantum mechanics energies are quantized; while they are not in general relativity, they are closely related to mass and the metric through Einstein's field equations. Two possible approaches to this apparent conflict are to either (i) "quantize gravity," i.e., develop a theory of quantum gravity in which the gravitational field is quantized, or (ii) "gravitize quantum mechanics," i.e., preserve the principles of general relativity, such as the equivalence principle, to modify quantum mechanics. The question of how to resolve these issues remains open.
- (iii) As detailed in Sec. II.A.6, the black-hole information paradox [see Raju (2022) for a review] poses another challenge, as it seems to require one to give up unitarity. This question similarly remains open.

### 3. Quantum measurements

The process of performing measurements in general relativity is straightforward and, up to limitations due to the measurement apparatus, we assume that we can measure with arbitrary precision. However, in quantum mechanics projective measurements are performed according to the Born rule: possible measurement outcomes are the eigenvalues  $\lambda_j$  of Hermitian operators  $\hat{A}$  (observables), and the associated probability to observe this measurement result is the projection of the system's state  $|\psi\rangle$  onto the associated eigenstate  $|\lambda_j\rangle$  of the observable,  $|\langle\lambda_j|\psi\rangle|^2$ . The fact that we measure observables that do not necessarily commute imposes limits to the precision with which we can measure different observables at the same time, most notably captured in the Heisenberg uncertainty for position and momentum,

$$\text{var}(\hat{x}(t))\text{var}(\hat{p}(t)) \geq \frac{\hbar^2}{4}. \quad (27)$$

Several conflicts with general relativity arise from this statement:

- (i) In the context of relativity, we do not encounter the same limitations on measurement precision. Classical variables can be measured to arbitrary precision without state update resulting from the measurement.
- (ii) Without the measurement postulate (that is, external observers), there are no events in quantum



mechanics. However, both special and general relativity are fundamentally based on the notion of events. Quantum superpositions are not compatible with the notion of a single event, such as a measurement, in spacetime. There have, however, been proposals for an event-based formulation of quantum mechanics, which fundamentally modifies the Born rule (Giovannetti, Lloyd, and Maccone, 2023).

- (iii) Problems also arise in QFT in CS. On the one hand, the theory inherits the measurement problem from quantum theory and, on the other hand, new problems arise due to causality. Here it has been shown that projective measurements on quantum fields lead to faster-than-light signaling (Sorkin, 1993). Finding ways to give a resolution to this problem is an active research field; see Fewster and Verch (2020).

Finally, the measurement problem in quantum mechanics, which states that there is no consistent dynamic description of the measurement process, also applies in the context of gravity. The issue is partially addressed by collapse theories, which, while they have not yet been experimentally verified, propose a dynamical mechanism; see Sec. IV.B.3.

#### 4. Composite quantum systems and entanglement

In quantum mechanics we use the tensor product to compose a system out of multiple subsystems, for example,  $|\psi\rangle = |\psi\rangle_A \otimes |\psi\rangle_B$ . We saw in Sec. II.A.4 that in quantum field theory in curved spacetime, the definition of subsystems is problematic since the notion of particle number is observer dependent (Fuentes-Schuller and Mann, 2005; Alsing and Fuentes, 2012). We have already identified the following crucial issues:

- (i) A consequence of the tensor product structure for composite systems in quantum theory is that multipartite systems can be entangled, which means that entanglement becomes an observer-dependent quantity. In the well-known Einstein-Podolsky-Rosen (EPR) paradox, this leads to a violation of causality of locality, i.e., we need to allow for faster-than-light effects if the theory is to remain local.
- (ii) The notion of entanglement requires the Hilbert space partition to be well defined. This is commonly done in terms of particles or modes. However, in curved spacetime, the notion of particles is ill defined. Generally, different inertial observers in curved space see different particle content in the field. Particles can only be well defined in rare spacetimes in which the metric is globally invariant under spatial translations or in which spacetime has regions where the metric has this symmetry. In most cases it is not possible to define a Hilbert space partition and study entanglement in composite systems.

The challenges listed here are all difficult to address, and in many cases their resolution should ultimately be determined through experiments. To enable these experiments, we now review the tools and methods used to model such systems in the laboratory.

### III. THEORETICAL FRAMEWORKS FOR MODELING MASSIVE QUANTUM SYSTEMS IN THE LABORATORY

To test the effects of gravity, which are often extremely small, with massive quantum systems, it is crucial to model the proposed experiment accurately. Here we account for common theoretical tools used to describe mechanical resonators in the laboratory. To begin, we discuss ways in which a probe can interact with the massive system (Sec. III.A). We then cover models of open-system dynamics (Sec. III.B), which are needed to model the experiments. We cover measurements and control schemes necessary for readout (Sec. III.C), as well as quantum metrology tools (Sec. III.D). Snapshots of experiments with massive quantum systems are given in Sec. V.A for testing gravity and in Sec. V.B for generating quantum states.

#### A. Coupling a mechanical mode to a probe

A key challenge in controlling massive systems in the laboratory is the fact that they often cannot be measured directly. To manipulate and control these massive systems, we must first couple them to a probe. We review two such models here.

##### 1. Optomechanical interaction

We first consider the case where a mechanical mode couples to a cavity mode (which can be optical, microwave, electrical, or magnetic). This exposition largely follows the review of Aspelmeyer, Kippenberg, and Marquardt (2014). See also Barzanjeh *et al.* (2022) for further reading. In many systems the frequency of the cavity mode depends on the center-of-mass position of the mechanical resonator. When considering the frequency shift to first order in the position of the mechanical oscillator, we acquire a coupling between the occupation number of the cavity mode and the position of the oscillator. The derivation often depends on platform-specific details. For example, in a Fabry-Pérot moving-end mirror cavity, the cavity deforms due to photon pressure (Law, 1995). For levitated nanoparticles the light-matter interaction can instead be derived by assuming the trapped sphere to be smaller than the laser waist of the beam (Romero-Isart *et al.*, 2011). Similarly, in some electromechanical systems, the motion of the resonator couples to capacitance, which in turn induces a frequency shift (Regal and Lehnert, 2011).

In all of the aforementioned cases, we arrive at the following *cavity-optomechanical Hamiltonian* (we denote it in this way even though the cavity field might not be an optical mode):

$$\hat{H} = \hbar\omega_c \hat{a}^\dagger \hat{a} + \hbar\omega_m \hat{b}^\dagger \hat{b} - \hbar g_0 \hat{a}^\dagger \hat{a} (\hat{b}^\dagger + \hat{b}), \quad (28)$$

where  $\hat{a}$  and  $\hat{a}^\dagger$  are the annihilation and creation operators for the radiation mode with free angular frequency  $\omega_c$  and  $\hat{b}$  and  $\hat{b}^\dagger$  are the annihilation and creation operators for the mechanical mode with free angular frequency  $\omega_m$ . The operators obey the canonical commutator relation  $[\hat{a}, \hat{a}^\dagger] = [\hat{b}, \hat{b}^\dagger] = 1$ . The optomechanical coupling  $g_0$  has units of angular frequency and encodes the strength of the interaction between the cavity

and mechanical modes. In most experimental realizations, the coupling is defined as the optical frequency shift per displacement  $g_0 \equiv -x_{\text{ZPF}}\partial\omega_c/\partial x$ , where  $x_{\text{ZPF}} = \sqrt{\hbar/2m\omega_m}$  is the zero-point fluctuation. The probe fields act as the means for both readout and control. The interaction between the bosonic mode and the mechanical resonator enables the detection of extremely small displacements, for example, due to gravitational effects. The nonlinear quantum dynamics generated by the Hamiltonian in Eq. (28) was first solved by Bose, Jacobs, and Knight (1997) and Mancini, Man'ko, and Tombesi (1997), where it was shown that both the cavity mode and the mechanical mode evolve into highly nonclassical superpositions of coherent states. This is a classic way to generate Schrödinger cat states of the macroscopic mechanical mode (Bose, Jacobs, and Knight, 1999; Marshall *et al.*, 2003; Qvarfort *et al.*, 2018). The solutions were later generalized to time-dependent couplings (Qvarfort *et al.*, 2019). As detailed in Sec. IV.B.1, the system dynamics of this Hamiltonian has been used for a number of proposals related to the detection of gravitational decoherence.

The Hamiltonian in Eq. (28) describes an idealized system isolated from its environment. In a realistic setting, both the bosonic and mechanical modes undergo dissipation, thermalization, and decoherence; see Sec. III.B for details. To replenish the lost quanta from the radiation mode, an external source is used to pump the system. Such a pump is modeled with a bosonic pump term  $\hat{H}_d = \alpha(t)\hat{a} + \alpha^*(t)\hat{a}^\dagger$ , where  $\alpha(t)$  is a complex drive amplitude. However, with the inclusion of such a term, the dynamics induced by the Hamiltonian in Eq. (28) can no longer be solved exactly (Qvarfort and Piovski, 2022). A common method to proceed is to solve the system dynamics perturbatively, or by examining the steady state for weak driving; see Nunnenkamp, Børkje, and Girvin (2011) and Rabl (2011).

For a strong enough pump, the system dynamics can be approximated as linear. Here the term *linear* refers to the fact that the resulting Heisenberg equations of motion contain only linear operator terms. The inclusion of a pump term strongly driving mode  $\hat{a}$  lets us separate  $\hat{a}$  into the classical amplitude of the drive  $\alpha$  and the fluctuations  $\delta\hat{a}$  such that  $\hat{a} = \alpha + \delta\hat{a}$ .<sup>1</sup> The interaction term in Eq. (28) becomes

$$\hat{H}_I = -\hbar g_0(\alpha + \delta\hat{a})^\dagger(\alpha + \delta\hat{a})(\hat{b}^\dagger + \hat{b}). \quad (29)$$

When  $|\alpha| \gg \langle\delta\hat{a}\rangle$ , the cubic term  $-\hbar g_0\delta\hat{a}^\dagger\delta\hat{a}$  can be removed because it is smaller by a factor of  $|\alpha|$  than the other terms. The remaining linear Hamiltonian is

$$\hat{H}_I \approx -\hbar g_0(\alpha^*\delta\hat{a} + \alpha\delta\hat{a}^\dagger)(\hat{b}^\dagger + \hat{b}). \quad (30)$$

Equation (30) is a common starting point for a number of investigations and accurately describes a range of experiments; see Aspelmeier, Kippenberg, and Marquardt (2014) and references therein. For example, within the resolved sideband regime where  $\omega_m \gg \kappa$ , it is possible to engineer either a beam-splitter interaction  $\hat{a}^\dagger\hat{b} + \text{H.c.}$  or a two-mode

squeezing term  $\hat{a}^\dagger\hat{b}^\dagger + \text{H.c.}$  by optically pumping on either the red ( $\omega_c - \omega_m$ ) or blue ( $\omega_c + \omega_m$ ) sideband. The red-sideband interaction is necessary for implementing, for example, sideband cooling (Liu *et al.*, 2013). Another way to couple an optical and mechanical mode is through a dissipative coupling rather than a dispersive one, where the displacement of the mechanical resonator directly modulates the decay rate of the cavity (Elste, Girvin, and Clerk, 2009).

The dynamics of the nonlinear Hamiltonian in Eq. (28) cannot be solved exactly in the presence of a pump term and optical dissipation. However, by engineering the system such that the optical mode dissipates from the cavity on a timescale much faster than that of the mechanical element, the interaction between the optical and mechanical modes can instead be described as an instantaneous interaction. This is also known as the unresolved sideband regime, where  $\omega_m \ll \kappa$ . The framework, often referred to as pulsed optomechanics, was developed by Vanner *et al.* (2011). By considering the Langevin equations in the unresolved sideband regime, one can model the light-matter interaction as an instantaneous unitary operator of the form  $\hat{U} = e^{i\mu\hat{n}_l\hat{X}_m}$  (Piovski *et al.*, 2012), where  $\mu$  is a dimensionless coupling which depends on the pulse shape,  $\hat{n}_l$  is the number of photons in the pulse entering or leaving the cavity, and  $\hat{X}_m = (\hat{b}^\dagger + \hat{b})/\sqrt{2}$  is the mechanical quadrature. For an adiabatic cavity with  $\kappa \gg \tau_{\text{in}}^{-1}$ , where  $\tau_{\text{in}}$  is the characteristic timescale of the input pulse, the value of  $\mu$  becomes  $\mu = \sqrt{8}g_0/\kappa$ , where  $g_0$  is the optomechanical coupling and  $\kappa$  is the optical dissipation rate (Clarke *et al.*, 2023). It is also possible to start from the linearized optomechanical Hamiltonian in Eq. (30) and derive a pulsed interaction that couples the position quadratures of the mechanical mode and the probe field (Khosla *et al.*, 2013; Bennett *et al.*, 2016). The resulting unitary operator is  $\hat{U}_{\text{Lin}} = e^{i\chi\hat{X}_c\hat{X}_m}$ , where  $\chi$  again depends on the pulse shape and  $\hat{X}_c$  is the amplitude quadrature of the input pulse (as opposed to the cavity quadrature). In the adiabatic regime and for an input coherent pulse,  $\chi = 4g_0\sqrt{N}/\kappa$ , where  $N$  is the average number of photons in the input pulse (Vanner *et al.*, 2011). A closely related idea to pulsed optomechanics is that of stroboscopic optomechanics (Brunelli *et al.*, 2020), where a train of short pulses of light is injected into the cavity. Proposals using nonlinear pulsed optomechanics include the generation of cat states (Clarke and Vanner, 2018; Ringbauer *et al.*, 2018), entangled states (Clarke *et al.*, 2020; Neveu *et al.*, 2021), and entangled cat states (Kanari-Naish *et al.*, 2022). State preparation using linearized pulsed optomechanics to generate catlike states has also been put forward using the addition and subtraction of phonons from the mechanical state (Milburn, Kim, and Vanner, 2016) and by swapping the mechanical state with a photon-subtracted state of light (Hoff *et al.*, 2016). Pulsed optomechanics has given rise to a number of protocols intended to test fundamental physics, such as the tests of modified commutator relations that are detailed in Sec. IV.D.1.

## 2. Coupling to a two-level system

Instead of a probe field, it is also possible to couple the mechanical resonator to a two-level system. The advantage of

<sup>1</sup>Depending on the application, we may also consider  $\hat{a} = \langle\hat{a}\rangle + \delta\hat{a}$  and  $\langle\delta\hat{a}\rangle = 0$ . This leads to a Hamiltonian of a similar form; see Aspelmeier, Kippenberg, and Marquardt (2014).

such a coupling is that the high level of control that has been achieved for two-level systems can now be indirectly applied to the mechanical resonator. Collectively, these systems are sometimes known as hybrid optomechanical systems since they couple bosonic continuous degrees of freedom to a two-level qubit system. Examples include nitrogen-vacancy (NV) centers embedded in a nanodiamond (Neukirch *et al.*, 2015; Hoang *et al.*, 2016) and superconducting resonator qubits coupled to mechanical modes (O’Connell *et al.*, 2010; Chu *et al.*, 2018; Satzinger *et al.*, 2018; Arrangoiz-Arriola *et al.*, 2019). See Rogers *et al.* (2014) and Chu and Gröblacher (2020) for dedicated reviews.

In the case of a spin coupled to mechanical motion, the same notions apply to the coupling between the spin of an ion and its center-of-mass position (Cirac and Zoller, 1995). Applying a magnetic field gradient to a trapped ion couples the internal and motional states of the system through the Zeeman effect. The spin-mechanical Hamiltonian reads, to first order in the position operator,

$$\hat{H} = \hbar\omega_m \hat{b}^\dagger \hat{b} + \frac{1}{2} \hbar\omega_0 \hat{\sigma}_z + \frac{\hbar\lambda}{2} (\hat{b}^\dagger + \hat{b}) \hat{\sigma}_z, \quad (31)$$

where  $\omega_0$  is the angular frequency of the qubit system,  $\sigma_z$  is the Pauli operator denoting the free energy,  $\hat{b}$  and  $\hat{b}^\dagger$  denote the annihilation and creation operators of the mechanical mode, and  $\lambda$  is a coupling constant that depends on the platform in question.

Superconducting qubits coupled to a mechanical mode are more commonly modeled using the Jaynes-Cummings Hamiltonian,

$$\hat{H}_{\text{JC}} = \hbar\omega_m \hat{b}^\dagger \hat{b} + \hbar\Omega \frac{\sigma_z}{2} + \frac{\hbar\lambda}{2} (\hat{b}\sigma_+ + \hat{b}^\dagger \sigma_-), \quad (32)$$

where  $\Omega$  is the oscillation frequency of the superconducting qubit,  $\sigma_z$  is the Pauli matrix, and  $\sigma_- = \sigma_x - i\sigma_y$  and  $\sigma_+ = \sigma_x + i\sigma_y$  are the raising and lowering operators in terms of the Pauli matrices. In Eq. (32)  $\lambda$  again denotes the strength of the coupling between the mechanical mode and the qubit. For example, a number of theoretical proposals have utilized the two-level system coupling for enhancing the optomechanical coupling strength (Heikkilä *et al.*, 2014; Pirkkalainen, Cho *et al.*, 2015), state preparation (Yin *et al.*, 2013; Kounalakis, Blanter, and Steele, 2020), and cooling (Martin *et al.*, 2004; Hauss *et al.*, 2008; Jaehne, Hammerer, and Wallquist, 2008; Northombam, Sahoo, and Sarma, 2021).

## B. Open-system dynamics for massive quantum systems

To detect the extremely weak effects of gravity, we must be able to distinguish them against any underlying noise floor. Additionally, proposals such as gravitational decoherence and gravity-induced state reduction stipulate that gravity itself manifests as a noise signature; see Sec. IV.B.1. Both of these considerations necessitate the use of accurate noise models for massive quantum systems. Here we review the main tools for modeling decoherence in massive quantum systems. For a dedicated review on noise models for mechanical

resonators in the quantum regime, see Bachtold, Moser, and Dykman (2022).

## 1. Quantum master equations

Quantum master equations describe the quantum state evolution in a situation in which a system, such as a mechanical oscillator or another probe, is coupled to a larger environment that we cannot control or measure, such as a thermal bath. Apart from modeling the interaction of a probe with its environment, this description is relevant for describing gravitational decoherence, gravitational collapse (see Sec. IV.B), and gravitational entanglement (see Sec. IV.C).

The dynamics of a quantum system coupled to the environment, or bath, can be described with the Hamiltonian  $\hat{H}$ , which contains a term for the system dynamics  $\hat{H}_s$  (e.g. for example, a quantum system in the laboratory, such as a cavity, a mechanical oscillator, or atoms), the environment or bath  $\hat{H}_b$  (for example, a thermal bath), and a term describing the coupling between system and bath  $\hat{H}_{sb}$ . The full Hamiltonian is

$$\hat{H} = \hat{H}_s + \hat{H}_b + \hat{H}_{sb}. \quad (33)$$

The fully evolved state of the system given an initial state  $|\Psi_0\rangle$  is  $|\Psi(t)\rangle = \hat{U}(t)|\Psi_0\rangle$ , where  $\hat{U}(t) = e^{-i\hat{H}t/\hbar}$ . In principle, any type of system-bath coupling  $\hat{H}_{sb}$  is possible, but they might not always lead to analytically solvable dynamics (Gardiner and Zoller, 2000).

In the laboratory we often do not have access to the bath degrees of freedom, which are therefore traced out from the quantum state. The result is a mixed state where the degree of mixedness is captured by the purity  $\text{tr}(\hat{\rho}^2) \leq 1$ , saturating the bound  $\text{tr}(\hat{\rho}^2) = 1$  only when the state is pure.

We now consider a dynamical equation for the evolution of the reduced system density matrix  $\hat{\rho}_s$  (Breuer *et al.*, 2002). This equation is known as a master equation. To derive the simplest possible master equation, we assume (i) that the system and bath are initially in a product state  $\hat{\rho}_{sb} = \hat{\rho}_s \otimes \hat{\rho}_b$ , (ii) that the coupling between the system and the bath (also known as the Born approximation) is weak, (iii) that the environment does not retain a memory of the interactions (the Markov approximation), and (iv) that fast-rotating terms (the secular approximation) can be discarded. By tracing out the bath modes  $\hat{b}_\ell$ , we obtain the Gorini-Kossakowski-Sudarshan-Lindblad master equation for the evolution of  $\hat{\rho}_s(t)$  (commonly referred to simply as the Lindblad equation) (Gorini, Kossakowski, and Sudarshan, 1976; Lindblad, 1976),

$$\begin{aligned} \dot{\hat{\rho}}_s(t) = & -\frac{i}{\hbar} [\hat{H}_s, \hat{\rho}_s(t)] \\ & + \sum_{\ell} \left( \hat{L}_{\ell} \hat{\rho}_s(t) \hat{L}_{\ell}^{\dagger} - \frac{1}{2} \{ \hat{\rho}_s(t), \hat{L}_{\ell}^{\dagger} \hat{L}_{\ell} \} \right). \end{aligned} \quad (34)$$

In Eq. (34)  $\hat{H}_s$  is the system Hamiltonian and  $\hat{L}_{\ell}$  denote the Lindblad *jump operators*. The Lindblad equation can also be written in shorthand as  $\dot{\hat{\rho}}_s = -i[\hat{H}_s, \hat{\rho}_s]/\hbar + \sum_{\ell} \hat{\mathcal{D}}[\hat{L}_{\ell}] \hat{\rho}_s(t)$ , where  $\hat{\mathcal{D}}[\hat{L}_{\ell}] \hat{\rho}(t) = \hat{L}_{\ell} \hat{\rho}(t) \hat{L}_{\ell}^{\dagger} - \hat{L}_{\ell}^{\dagger} \hat{L}_{\ell} \hat{\rho}(t)/2 - \hat{\rho}(t) \hat{L}_{\ell}^{\dagger} \hat{L}_{\ell}/2$  is called the standard dissipator.



For mechanical resonators coupled to probe fields or two-level systems, there are commonly two types of noise that affect the system: dissipation and scattering processes in the probe field and thermalization processes in the phononic modes (Aspelmeyer, Kippenberg, and Marquardt, 2014). Mechanical dissipation and thermalization arise due to processes specific to the system. In clamped systems, for example, unwanted thermal excitations are transferred via the physical point of attachment. Both optical and mechanical noise can be modeled with the Langevin equation in the linear optomechanical regime; see Sec. III.B.2. A common way to model Markovian dissipation and thermalization is to assume Lindblad operators of the form  $\hat{L}_1 = \sqrt{(1 + n_{\text{th}})\Gamma}\hat{b}$  and  $\hat{L}_2 = \sqrt{\Gamma n_{\text{th}}}\hat{b}^\dagger$ , where  $\Gamma$  is the mechanical linewidth and  $n_{\text{th}} = [\exp(\hbar\omega/k_B T) - 1]^{-1}$  is the thermal occupation number of the bath, for which  $\omega$  is the bath frequency,  $k_B$  is Boltzmann's constant, and  $T$  is the temperature of the bath.

In levitated systems environmental noise arises in part due to collisions between the system and the surrounding gas particles. A number of these processes correspond to position localization, which can be described with the following master equation (Romero-Isart, 2011):

$$\langle x|\dot{\hat{\rho}}(t)|x'\rangle = \frac{i}{\hbar}\langle x|[\hat{\rho}(t), \hat{H}]|x'\rangle - \Gamma(x - x')\langle x|\hat{\rho}(t)|x'\rangle. \quad (35)$$

In Eq. (35) the form of  $\Gamma(x - x')$  depends on the nature of the noise. In the limit where the decoherence decay depends quadratically on  $|x - x'|$ , Eq. (35) simplifies to

$$\dot{\hat{\rho}}(t) = \frac{i}{\hbar}[\hat{\rho}(t), \hat{H}] - \Lambda[\hat{x}, [\hat{x}, \hat{\rho}(t)]], \quad (36)$$

where  $\Lambda$  is the dissipation rate. The dynamics was solved for a coupling of a probe field by Bassi, Ippoliti, and Adler (2005), who used a stochastic unraveling method; see Adler, Bassi, and Ippoliti (2005) for details.

We now discuss generalizations to the strong coupling and the nonlinear regime and to non-Markovian dynamics. Commonly, dissipation of the probe field and mechanical mode are treated separately. However, such a treatment is not always valid when the probe field and mechanics are strongly coupled. Instead, we must consider a dressed Lindblad equation (Hu *et al.*, 2015) that was solved by Torres, Betzholtz, and Bienert (2019), who took a damping basis approach (Briegel and Englert, 1993). In the nonlinear regime a solution to the Lindblad master equation was found for weak dissipation (Mancini, Man'ko, and Tombesi, 1997) and was later generalized to arbitrary  $\kappa$  (Qvarfort, Vanner *et al.*, 2021), although a closed-form expression for the evolved state cannot be obtained. The Lindblad equation for an optomechanical system in the nonlinear regime was solved for dissipation of the mechanical mode (Bose, Jacobs, and Knight, 1997; Mancini, Man'ko, and Tombesi, 1997), with Lindblad operators  $\hat{L} = \sqrt{\gamma}\hat{b}^\dagger\hat{b}$ , where  $\gamma$  is the mechanical dissipation rate.

Models assuming a Markovian environment are often sufficient to capture the open dynamics of the system accurately. However, there are certain cases where a non-Markovian

description is necessary. Non-Markovianity arises when the bath retains a memory of the interaction with the system and information can flow back into the system (Breuer *et al.*, 2002). Generally, to model such non-Markovian noise, we must either consider the Caldeira-Leggett master equation for Brownian motion (Caldeira and Leggett, 1983a) or solve a general non-Markovian master equation (Hu, Paz, and Zhang, 1992), although care must be taken since the dynamics predicted by these approaches do not automatically guarantee physical states as a solution (Kohen, Marston, and Tannor, 1997). Non-Markovian dynamics can also be modeled using Lindblad-type equations with time-dependent noise rates (Zhang *et al.*, 2012). Some studies of mechanical resonators indicate the requirement for non-Markovian dynamics. For example, it was shown that in clamped systems the resulting noise spectrum was consistent with a non-Markovian spectrum (Gröblacher, Hammerer *et al.*, 2009). In addition, theoretical works indicate that modeling the effects of damped tunneling two-level systems on a nanomechanical flexing beam resonator gives rise to non-Markovian noise (Remus, Blencowe, and Tanaka, 2009). Optomechanical systems in non-Markovian environments have been modeled, although generally without the use of a master equation. A Feynman-Vernon influence functional method was used to study sideband cooling in non-Markovian environments (Triana, Estrada, and Pachón, 2016), and the influence of non-Markovian noise on the optomechanical nonlinearity was considered by Qvarfort (2023).

Beyond analytic solutions master equations are often solved numerically. Useful tools include the QuTiP Python package (Johansson, Nation, and Nori, 2012). See Campaioli, Cole, and Hapuarachchi (2023) for a tutorial.

## 2. Langevin equations and input-output formalism

The quantum Langevin equations model the nonunitary evolution of the quantum modes in the Heisenberg picture. The interaction between an input mode and the system is imprinted on output fields, which are detected in experiments. The Langevin equations are also a useful tool for modeling quantum metrology (see Sec. III.D) and, in particular, experimental tests of gravity such as weak-force detection (see Sec. IV.A).

The Langevin equations for the bosonic field  $\hat{a}_j$  read [see Caldeira and Leggett (1983b) and Gardiner and Zoller (2000) for a derivation]

$$\dot{\hat{a}}_j = -\frac{\gamma_j + \gamma_{j,\text{in}}}{2}\hat{a}_j - \frac{i}{\hbar}[\hat{H}_s, \hat{a}_j] - \sqrt{\gamma_{j,\text{in}}}\hat{a}_{j,\text{in}}. \quad (37)$$

Note that, following standard convention,  $\hat{a}_{j,\text{in}}$  has units of  $s^{-1/2}$ . This input field could be noise from a heat bath or a coherent probe field. In a concrete setting  $\gamma_{j,\text{in}}$  could be the coupling rate between the system and a waveguide used to couple the probe field to the system. If the system also experiences loss into other channels at rate  $\gamma_j$ , this rate is added to the overall decay rate.

The input fields are connected to the output fields via input-output boundary conditions (Caves, 1982; Gardiner and Collett, 1985; Clerk *et al.*, 2010)

$$\hat{a}_{j,\text{out}} = \hat{a}_{j,\text{in}} + \sqrt{\gamma_{j,\text{in}}} \hat{a}_j. \quad (38)$$

By solving the internal system dynamics as a function of the input fields, one can model the output fields purely as a function of the input fields.

The Langevin equations for a cavity-optomechanical system with a Hamiltonian given by Eq. (28) are (Bowen and Milburn, 2015)

$$\begin{aligned} \dot{\hat{a}} &= -\frac{\kappa}{2} \hat{a} + i\Delta \hat{a} + ig_0 \hat{a} (\hat{b}^\dagger + \hat{b}) - \sqrt{\kappa_{\text{in}}} \hat{a}_{\text{in}}, \\ \dot{\hat{b}} &= -\frac{\Gamma}{2} \hat{b} - i\omega_m \hat{b} + ig_0 \hat{a}^\dagger \hat{a} - \sqrt{\Gamma_{\text{in}}} \hat{b}_{\text{in}}, \end{aligned} \quad (39)$$

in which  $\Delta = \omega - \omega_c$  is the detuning of the light frequency  $\omega_l$  from the cavity frequency  $\omega_c$  [Eq. (28)],  $\kappa$  is the optical linewidth,  $\kappa_{\text{in}}$  is the rate by which the probe field dissipates away from the cavity,  $\Gamma$  is the mechanical linewidth, and  $\Gamma_{\text{in}}$  is the coupling rate for thermal heating. Equations (39) are challenging to solve in the nonlinear regime but can be linearized by considering a strong optical pump field; see Sec. III.A.1. Such a treatment lies at the basis of many models of optomechanical systems (Aspelmeyer, Kippenberg, and Marquardt, 2014).

In a typical experimental setting, we are interested in the frequency-dependent response to an input  $\hat{a}_{j,\text{in}}$ , which we obtain by means of the Fourier transform from Eq. (37),  $\hat{a}(\omega) \equiv (1/\sqrt{2\pi}) \int_{-\infty}^{\infty} dt e^{i\omega t} \hat{a}(t)$ . Experimentally, the Fourier transform is calculated over a finite time window  $[-\tau, \tau]$  that converges to the Fourier integral in the limit  $\tau \rightarrow \infty$ . The Fourier transform is also the main analytic method by which the Langevin equations can be solved, provided that they are linear in terms of the operators they contain. In that case we can apply the Fourier transform to Eq. (37) to derive a scattering matrix

$$S(\omega) = \mathbb{1} + \sqrt{\gamma_{\text{in}}} (i\omega \mathbb{1} + M)^{-1} \sqrt{\gamma_{\text{in}}}, \quad (40)$$

where  $\gamma_{\text{in}} = \text{diag}(\gamma_{1,\text{in}}, \dots, \gamma_{N,\text{in}})$  contains the input noise terms and where the elements of  $M$  are defined via the Langevin equations in Eq. (37) as  $\dot{\hat{a}}_j = \sum_{\ell} M_{j,\ell} \hat{a}_{\ell} - \sqrt{\gamma_j} \hat{a}_{j,\text{in}}$ . The scattering matrix allows us to relate the input and output fields as

$$\hat{a}_{j,\text{out}}(\omega) = \sum_{\ell} S_{j,\ell}(\omega) \hat{a}_{\ell,\text{in}}(\omega), \quad (41)$$

where  $S_{j,\ell}$  are the matrix elements of  $S$ . The transmission between the  $j$ th input and the  $\ell$ th output port is given by  $T_{\ell,j}(\omega) = |S_{\ell,j}(\omega)|^2$ . Interactions such as single- or two-mode squeezing can give rise to  $|S_{\ell,j}(\omega)|^2 \geq 1$ , which is referred to as the gain  $\mathcal{G} = |S_{\ell,j}(\omega)|^2$ . These quantities are relevant for characterizing sensors; devices such as isolators, circulators, and directional amplifiers; and other scattering experiments such as optomechanically induced transparency experiments (Xiong and Wu, 2018).

We now turn our attention to quantum noise in the context of the Langevin equations. For a general treatment of noise, see the designated review by Clerk *et al.* (2010); for a

treatment specifically in the context of cavity optomechanics, see Aspelmeyer, Kippenberg, and Marquardt (2014).

Quantum systems are typically hard to isolate and are susceptible to noise and dissipation. For instance, the real-time motion of a mechanical oscillator subjected to fluctuating thermal Langevin force was measured by Hadjar *et al.* (1999). These forces can be straightforwardly included in the Langevin equations via the input fields  $\hat{a}_{j,\text{in}}$  [Eq. (37)]. Instead of recording real-time trajectories, it is typically more convenient to record the noise power spectral density  $\mathcal{S}_{\hat{O}\hat{O}}(\omega)$  defined for some system operator  $\hat{O}$ . The spectral density describes the intensity of the noise at a given frequency. In practice, we obtain  $\mathcal{S}_{\hat{O}\hat{O}}(\omega)$  by averaging over many experimental runs. According to the Wiener-Khinchine theorem (Wiener, 1930; Khintchine, 1934), this is equivalent to calculating the Fourier transform of the autocorrelation,

$$\mathcal{S}_{\hat{O}\hat{O}}(\omega) \equiv \int_{-\infty}^{\infty} dt e^{i\omega t} \langle \hat{O}^\dagger(t) \hat{O}(0) \rangle. \quad (42)$$

We obtain the noise spectral density from the Langevin equations in Eq. (37) by calculating correlators of, for example, the fields  $\langle \hat{a}_j^\dagger(\omega) \hat{a}_j(\omega) \rangle$  or position  $\langle \hat{x}_j(\omega) \hat{x}_j(\omega) \rangle$ , taking into account the input fluctuations  $\hat{a}_{j,\text{in}}(\omega)$ . The assumption that we made of Markovian noise translates to the fact that  $\hat{a}_{\text{in}}$  are uncorrelated in time. This is also known as Gaussian white noise, and the vacuum fluctuations are given by

$$\langle \hat{a}_{\text{in}}(t) \hat{a}_{\text{in}}^\dagger(t') \rangle = (n_{\text{th}} + 1) \delta(t - t'), \quad (43)$$

$$\langle \hat{a}_{\text{in}}^\dagger(t) \hat{a}_{\text{in}}(t') \rangle = n_{\text{th}} \delta(t - t'), \quad (44)$$

with the number of thermal bosonic excitations. Note that, for the case of an optical probe field, the environment corresponds to the vacuum, meaning that  $n_{\text{th}} = 0$ .

As an example, we consider the position noise of a single harmonic oscillator with a frequency  $\omega_m$  and a damping rate  $\Gamma$ . The spectral density is given by (Clerk *et al.*, 2010)

$$\begin{aligned} \mathcal{S}_{xx} &= 2\pi x_{\text{xpf}}^2 \left( n_{\text{th}}(\hbar\omega_m) \frac{\Gamma}{(\omega_m + \omega)^2 + (\Gamma/2)^2} \right. \\ &\quad \left. + [n_{\text{th}}(\hbar\omega_m) + 1] \frac{\Gamma}{(\omega_m - \omega)^2 + (\Gamma/2)^2} \right), \end{aligned} \quad (45)$$

where  $n_{\text{th}}(\hbar\omega_m)$  is the expected number of particles according to the Bose-Einstein statistic and  $x_{\text{xpf}} = \sqrt{\hbar/(2\omega_m m)}$  is the zero-point fluctuation, for which  $m$  is the mass of the oscillator. The area under the spectral density  $\mathcal{S}_{\hat{x}\hat{x}}(\omega)$ , with  $\hat{x}$  the position operator, is proportional to  $\langle \hat{x}^2 \rangle$ . In thermal equilibrium at large temperatures  $k_B T \gg \hbar\Omega$ ,  $\langle \hat{x}^2 \rangle$  is proportional to the temperature according to the fluctuation-dissipation theorem.

In general, we note that the spectral density in Eq. (45) is not symmetric in  $\omega$ , due to spontaneous emission—which, classically, it would be. As important application to optomechanics, the asymmetry of the noise spectral density of the radiation pressure when driving on the red-detuned sideband

allows the mechanical oscillator to be cooled. The optical noise spectral density then enters in the net optical cooling rate of the mechanical oscillator (Marquardt *et al.*, 2007). If, in addition, the light used to drive the mechanical oscillator is squeezed, sideband cooling allows the cooling of the mechanical mode to the ground state (Clark *et al.*, 2017). The spectral density is also relevant to sensing applications as it determines the signal-to-noise ratio (Clerk *et al.*, 2010; Lau and Clerk, 2018); see Sec. III.D.

### C. Measurement and control of massive quantum systems

To control the massive quantum systems in the laboratory, we need to be able to manipulate their motion and perform accurate measurements. This is particularly important when measuring weak gravitational effects. Here we summarize the key ideas behind different measurement and control schemes that are used in the various proposals covered in Sec. IV. For in-depth discussions of control and measurement of the quantum systems, see the designated reviews by Jacobs and Steck (2006) and Clerk *et al.* (2010), as well as the textbooks by Wiseman and Milburn (2009) and Jacobs (2014).

#### 1. Quantum measurements

To extract information about how gravity affects quantum systems, we must perform a measurement. There are a number of different measurement types and schemes. Here we review the most common ones.

Projective measurements, also known as von Neumann measurements, model the measurement apparatus as a macroscopic pointer that can be read out classically. Strong correlations between the system and the pointer let us determine the state of the system unambiguously by measuring the pointer. However, the measurement destroys the coherence of the wave function, subsequently destroying information about the conjugate observable and leading to backaction quantum noise (Jacobs and Steck, 2006; Clerk *et al.*, 2010).

Quantum nondemolition measurements (Braginsky, Vorontsov, and Thorne, 1980; Peres, 1993; Braginsky and Khalili, 1995, 1996), however, present a special case in which the eigenstates of the observable that we are measuring are also eigenstates of the system or, equivalently, the measured observable  $\hat{A}$  commutes with the Hamiltonian  $\hat{H}$ ,  $[\hat{H}, \hat{A}] = 0$ , and thus  $\hat{H}$  and  $\hat{A}$  are simultaneously diagonalizable. Measuring multiple times yields the same result and allows for improved measurement accuracy, which is crucial for certain force-sensing schemes. We discuss these backaction evasion schemes in more detail in Sec. IV.A.1.

Weak measurements extract only partial information about an observable and thus do not fully destroy the information about the conjugate observable. For a detailed discussion see the pedagogical review by Jacobs and Steck (2006). The main idea is to construct operators  $\hat{P}_m$  such that  $\sum_m \hat{P}_m^\dagger \hat{P}_m = \mathbb{1}$ . The state after the measurement expressed in terms of the projectors  $\hat{P}_n$  and the state  $\hat{\rho}$  before the measurement is then given by

$$\hat{\rho}_f = \frac{\hat{P}_n^\dagger \hat{\rho} \hat{P}_n}{\text{tr}(\hat{P}_n^\dagger \hat{\rho} \hat{P}_n)}, \quad (46)$$

with the probability  $\text{tr}(\hat{P}_n^\dagger \hat{\rho} \hat{P}_n)$ , to obtain this outcome.<sup>2</sup> Note that we recover a von Neumann measurement when measuring in the eigenbasis, i.e., setting  $P_n = |n\rangle\langle n|$ . Rather than measuring in the eigenbasis, we define  $P_n$  as a weighted sum over different eigenstates that peaks at a specific eigenstate but has a certain width (Jacobs and Steck, 2006). A small width corresponds to a strong measurement, with the limit of zero width corresponding to a von Neumann measurement. A large width performs a weak measurement. The measurement strength  $k$  is typically defined as the inverse of this width. It was suggested that weak measurements can lead to more accurate measurements of gravitational forces (Kawana and Ueda, 2019).

Rather than measuring a system once, it can be interesting to continuously extract information from the system. Together with feedback, such measurement strategies can, for instance, be employed to squeeze or cool levitated mechanical systems (Genoni *et al.*, 2015) that can be used for gravity tests; see Sec. V.A.4. A theory for such continuous measurements can be constructed from a sequence of time intervals  $\Delta t$  during which weak measurements are performed with a measurement strength proportional to  $\Delta t$ . In the limit of infinitesimally short time intervals, we obtain a stochastic equation of motion due to the random nature of the measurements; see Jacobs and Steck (2006) for a derivation. The measurement current of the continuously watched observable  $A$  with a weak measurement is then given by

$$dI(t) = \sqrt{k}\langle A(t) \rangle dt + dW(t), \quad (47)$$

in which  $k$  is the measurement strength,  $dW(t)$  is the standard Wiener increment describing the white imprecision noise in the measurement current and fulfills<sup>3</sup>  $\langle\langle dW \rangle\rangle = 0$  and  $\langle\langle dW^2 \rangle\rangle = dt$  (Clerk *et al.*, 2010). The density matrix  $\hat{\rho}_c$  determining the expectation value  $\langle A(t) \rangle$  is now conditional on the measurement current and evolves according to the stochastic master equation

$$d\hat{\rho}_c = -\frac{i}{\hbar}[\hat{H}_s, \hat{\rho}_c]dt + \frac{k}{4}\mathcal{D}[\hat{A}]\hat{\rho}_c dt + \frac{\sqrt{k}}{2}[\hat{A}\hat{\rho}_c + \hat{\rho}_c\hat{A} - 2\langle\hat{A}\rangle\hat{\rho}_c]dW, \quad (48)$$

with  $\mathcal{D}[\cdot]\hat{\rho}_c$  as defined following Eq. (34). The evolution of this conditioned quantum state is referred to as quantum trajectory. Equation (48) can be solved analytically only in a special case (Wiseman, 1996; Jacobs and Steck, 2006); in most cases it has to be solved numerically.

<sup>2</sup>An operator of the form  $\hat{M} = \sum_{n=a}^b \hat{P}_n^\dagger \hat{P}_n$  is a positive operator and  $\text{tr}(\hat{M}\hat{\rho})$  gives the probability that  $n \in [a, b]$ . Therefore, this operator defines a positive operator-valued measure (POVM).

<sup>3</sup>The average over all possible measurement outcomes is denoted by  $\langle\langle \cdot \rangle\rangle$ . For a pedagogic introduction to stochastic calculus, see Jacobs and Steck (2006).



When we average over all possible measurement results (i.e., the observer does not retain the measurement current), Eq. (48) simplifies to

$$d\langle\hat{\rho}_c\rangle = -\frac{i}{\hbar}[\hat{H}_s, \langle\hat{\rho}_c\rangle]dt + \frac{k}{4}\mathcal{D}[\hat{A}]\langle\hat{\rho}_c\rangle dt \quad (49)$$

since  $\langle\hat{\rho}_c dW\rangle = 0$ , as  $\hat{\rho}_c$  and  $dW$  are statistically independent (Jacobs and Steck, 2006).

Continuous measurements can also be described with Langevin equations. Here the recorded measurement current is then determined by the output fields. This is particularly convenient for describing homodyne and heterodyne detection. For a homodyne detection the output at frequency  $\omega_0$  is combined with a signal of a local oscillator at the same frequency  $\omega_0$  in an interferometer. The detected current (in a rotating frame with frequency  $\omega_0$ ) is then given by<sup>4</sup> (Barchielli and Vacchini, 2015)

$$\hat{I}(t) \equiv [e^{-i\phi}\hat{b}_{\text{out}}(t) + e^{i\phi}\hat{b}_{\text{out}}^\dagger(t)]/\sqrt{2}. \quad (50)$$

In Eq. (50)  $\phi$  is a phase difference depending on the optical path that determines the observed quadrature. For  $\phi = 0$  we obtain  $\hat{I}(t) = [\hat{b}_{\text{out}}(t) + \hat{b}_{\text{out}}^\dagger(t)]/\sqrt{2} = q_{\text{out}}(t)$ , while  $\phi = \pi/2$  yields  $\hat{I}(t) = -i[\hat{b}_{\text{out}}(t) - \hat{b}_{\text{out}}^\dagger(t)]/\sqrt{2} = p_{\text{out}}(t)$ .

In a heterodyne detection scheme, the output at frequency  $\omega_0$  is combined with a signal of a local oscillator at a different frequency  $\omega_1$  detuned from the output frequency by  $\Delta \equiv \omega_1 - \omega_0$ , thereby giving rise to the heterodyne current

$$\hat{I}(t) \equiv [e^{-i(\phi-\Delta t)}\hat{b}_{\text{out}}(t) + e^{i(\phi-\Delta t)}\hat{b}_{\text{out}}^\dagger(t)]/\sqrt{2}. \quad (51)$$

As a result, the measured quadrature oscillates in time, providing information about both amplitude and phase. However, this comes at the cost of an added half quantum of noise (Bowen and Milburn, 2015). Homodyne and heterodyne noise spectra were discussed in detail by Barchielli and Vacchini (2015) and Bowen and Milburn (2015).

## 2. Feedback and feedforward

Along with continuous measurements, we can continuously apply operations on the system to steer it toward a desired state. Creating or stabilizing certain quantum states can be advantageous in the context of metrology. In particular, superposition states can be used to test theories about quantum mechanics and gravity. Furthermore, it has been shown that certain entangled states can improve the sensitivity and signal-to-noise ratio (Leibfried *et al.*, 2004; Roos *et al.*, 2006). Feedback is also employed for squeezing or cooling in many of the experimental tests of gravity in Sec. V.A and, in general, for the control of quantum systems; see Sec. V.B. The feedback can either explicitly depend on the measurement current  $I(t)$  obtained in Eq. (47) (closed-loop feedback) or not (open-loop feedback). In addition to applying feedback, we

can also use the continuously measured quantum system to control a second quantum system that is then referred to as feedforward.

The optimal control of quantum systems has been investigated for a long time (Peirce, Dahleh, and Rabitz, 1988; Dahleh, Peirce, and Rabitz, 1990; Judson and Rabitz, 1992; Warren, Rabitz, and Dahleh, 1993; Wiseman and Milburn, 1993; Wiseman, 1994; Magrini *et al.*, 2021). Feedback can be applied either semiclassically—here the measurement current that is used to provide feedback is obtained with a classical sensor—or fully quantum—here the detectors and sensors are all quantum systems. The main idea behind classical, continuous feedback is to apply a semiclassical potential that steers the system coherently toward the desired quantum state (Caves and Milburn, 1987; Wiseman, 1994; Doherty *et al.*, 2000; Lloyd, 2000). For instance, we can measure the position of a mechanical oscillator and then perform a displacement operation to shift its position.

In general, the quantum state evolution now explicitly depends on the stochastic measurement current  $I(t)$ , which will be different in each experimental run, resulting in conditional dynamics described by a stochastic master equation. We obtain the stochastic measurement current [Eq. (47)] by measuring an observable  $\hat{A}$ , which is determined according to the stochastic master equation for continuous feedback [Eq. (48)]. This current [Eq. (47)] is then fed back to drive the system via the Hamiltonian

$$\hat{H}_{\text{fb}} = \sqrt{\kappa_{\text{fb}}}I(t - \tau)\hat{B}, \quad (52)$$

with  $\kappa_{\text{fb}}$  the feedback strength and  $\tau$  some time delay and in which  $\hat{B}$  encodes the operation that is chosen to be applied based on the measurement outcome. Note that the feedback operation  $\hat{B}$  may also involve the measured observable  $\hat{A}$ . Furthermore, the choice of  $\hat{B}$  explicitly depends on the measurement current in the case of closed-loop feedback, while it does not for open-loop feedback.

Instead, we can look at the unconditional dynamics by averaging over all measurement outcomes

$$\begin{aligned} \frac{d\hat{\rho}}{dt} = & -\frac{i}{\hbar}[\hat{H}_s, \hat{\rho}] + \frac{k}{4}\mathcal{D}[\hat{A}]\hat{\rho} + \kappa_{\text{fb}}\mathcal{D}[\hat{B}]\hat{\rho} \\ & - i\frac{\sqrt{\kappa\kappa_{\text{fb}}}}{2}[\hat{B}, \hat{A}\hat{\rho} + \hat{\rho}\hat{A}]. \end{aligned} \quad (53)$$

In Eq. (53)  $-i(\sqrt{\kappa\kappa_{\text{fb}}}/2)[\hat{B}, \hat{A}\hat{\rho} + \hat{\rho}\hat{A}]$  encodes a linear restoring term and dissipation, for example, in an optomechanical system, this could be a restoring force. The term  $\kappa_{\text{fb}}\mathcal{D}[\hat{B}]\hat{\rho}$  describes additional fluctuations as a consequence of the feedback.

Analogous to the previously described scenario, the measurement current of system A can force a second system B in a feedforward scheme. The main difference is that  $\hat{B}$  now denotes an operator of the other system B. For instance, reservoir-engineered nonreciprocity can be thought of as an autonomous feedforward scheme (Metelmann and Clerk, 2017) in which the measurement results of one system are used to drive another system, but not vice versa.

<sup>4</sup>Note that we encounter different conventions for the definition of  $I(t)$ , which is sometimes defined as  $I = (e^{-i\phi}\hat{b}_{\text{out}} + e^{i\phi}\hat{b}_{\text{out}}^\dagger)/2$ , or simply  $I = e^{-i\phi}\hat{b}_{\text{out}} + e^{i\phi}\hat{b}_{\text{out}}^\dagger$ .

Rather than controlling the quantum system based on the classical measurement record, it is also possible to replace the sensors and controllers with quantum systems that coherently interact with the system to be controlled (Lloyd, 2000; Nelson *et al.*, 2000; Nurdin, James, and Petersen, 2009). Coherent feedback protocols can outperform measurement-based schemes (Hamerly and Mabuchi, 2012, 2013) because they can exploit a geodesic path in Hilbert space that is forbidden to measurement-based schemes (Jacobs, Wang, and Wiseman, 2014). A convenient way to describe coherent feedback is with a Langevin equation formalism [Eq. (37)] (Gardiner and Zoller, 2000). Here the output field  $\hat{a}_{\text{out}}$  is fed to the input field  $\hat{b}_{\text{in}}$  of the mode that is to be controlled with some time delay  $\tau$ ,

$$\hat{b}_{\text{in}}(t) = \sqrt{\kappa_{\text{fb}}} \hat{a}_{\text{out}}(t - \tau). \quad (54)$$

A number of schemes have been proposed and implemented to control mechanical oscillators via feedback, such as feedback cooling (Chang *et al.*, 2010; Li, Kheifets, and Raizen, 2011; Hamerly and Mabuchi, 2012, 2013; Genoni *et al.*, 2015; Jain *et al.*, 2016; Vovrosh *et al.*, 2017; Setter *et al.*, 2018; Harwood, Brunelli, and Serafini, 2021; Guo and Gröblacher, 2022; Mansouri *et al.*, 2022; Rademacher, Konopik *et al.*, 2022), schemes to control squeezing, entanglement, and state transfer (Harwood, Brunelli, and Serafini, 2021), or to control the motional state of the mechanical oscillator, its resonance frequency, and its damping rate (Ernzer *et al.*, 2023). Feedback cooling allowed the cooling of a 10 kg mass in the LIGO detector close to its ground state (Abbott *et al.*, 2009; Whittle *et al.*, 2021), and a millimeter-sized membrane resonator was cooled to the ground state with measurement-based feedback (Rossi *et al.*, 2018). Measurement-based feedback cooling has also been demonstrated in electro-mechanical systems (Wang *et al.*, 2023), where it has also been demonstrated that feedback can lead to dynamically stability in situations that would be unstable without feedback. In addition, feedback schemes are employed to equalize mechanical loss rates in experiments with multiple mechanical oscillators (Poggio *et al.*, 2007; del Pino, Slim, and Verhagen, 2022; Wanjura *et al.*, 2023).

## D. Quantum metrology with massive quantum systems

Since gravity is extremely weak, we may sometimes want to quantify the sensitivity of a quantum sensor to ensure that it is powerful enough. The main tool used for this is quantum estimation theory, also referred to as quantum metrology. We outline here the key concepts and refer to the reviews by Paris (2009), Clerk *et al.* (2010), and Tóth and Apellaniz (2014) for further details. Detection methods for optomechanical systems were reviewed by Poot and van der Zant (2012). Many of these tools are used for precision tests of gravity; see Sec. IV.A.

### 1. Langevin description of a quantum sensor

This exposition of a sensing scheme follows (Clerk *et al.*, 2010; Lau and Clerk, 2018). In a typical metrology setting, we attempt to infer an infinitesimal change in a small parameter  $\epsilon$

that the Hamiltonian  $\hat{H}(\epsilon)$  depends on. Expanding  $\hat{H}(\epsilon)$  to first order in  $\epsilon$ , we have

$$\hat{H} = \hat{H}_0 + \epsilon \hat{V} + O(\epsilon^2), \quad (55)$$

where  $\hat{H}_0$  is the free Hamiltonian and  $\hat{V}$  is the operator that encodes  $\epsilon$ . To extract changes in  $\epsilon$ , we probe the system governed by  $\hat{H}$  with a probe field  $\hat{a}_{\text{in}}$  and currents with the response  $\hat{a}_{\text{out}}$ , which then depends on  $\epsilon$ . For small  $\epsilon$  we can write  $\hat{a}_{\text{out}} \cong \hat{a}_{\text{out}}^{(0)} + \lambda \epsilon$ , where  $\lambda$  is a linear response coefficient.

To characterize the resolving power of a quantum sensor, we can calculate the signal-to-noise ratio (SNR) by comparing the integrated signal power to the noise power. For instance, the measurement current can be obtained via homodyne detection; see Sec. III.C.1. The power associated with the signal is then given by the expectation value of the time-integrated measurement current  $\hat{m}(t) \equiv \int_0^t d\tau \hat{I}(\tau)$ , where  $\hat{I}(t)$  is defined in Eq. (50). The measurement current should be compared to the power without the perturbation  $\epsilon$ , so we define the power difference  $\mathcal{P}$  of the signal with and without the perturbation  $\epsilon$  as

$$\mathcal{P} = [\langle \hat{m}(t) \rangle - \langle \hat{m}(t) \rangle|_{\epsilon=0}]^2. \quad (56)$$

The associated noise power is then given by

$$\mathcal{N} \equiv \langle \delta \hat{m}(t) \delta \hat{m}(t) \rangle = t S_{II}(0), \quad (57)$$

where  $\delta \hat{m}(t) = \hat{m}(t) - \langle \hat{m}(t) \rangle$ . In Eq. (57)  $S_{II}(0) = (1/2) \int d\epsilon e^{i\omega t} \langle \{ \delta I(t), \delta I(0) \} \rangle$  is the noise spectral density defined in Eq. (42) of the measurement current at  $\omega = 0$ . Note that Eq. (57) is linear at time  $t$  because we consider the integrated measurement current. The SNR is then given by the ratio of  $\mathcal{P}$  and  $\mathcal{N}$ ,  $\rho_{\text{SNR}} \equiv \mathcal{P}/\mathcal{N}$ . For applications such as gravitational-wave detection (Caves, 1979), force sensing (Caves *et al.*, 1980), and force gradient sensing (Rudolph *et al.*, 2022), it is vital to ensure that the signal is stronger than the noise such that  $\rho_{\text{SNR}} \geq 1$ . Since the noise increases with  $t$ , we require the measurement current to also accumulate information about  $\epsilon$  at the same rate. Therefore, it is crucial to retain long coherence times in the system so that a strong signal can be retained throughout.

Quantum mechanics puts a limit (Caves *et al.*, 2012) on the added noise  $\mathcal{A}$  of an amplifier when we are referring to the amplification gain  $\mathcal{G}$ . In particular, we find for the variance  $\langle \Delta \hat{a}_{\text{out}} \rangle \equiv \langle \hat{a}_{\text{out}}^\dagger \hat{a}_{\text{out}} \rangle - |\langle \hat{a}_{\text{out}} \rangle|^2$  the expression  $\langle \Delta \hat{a}_{\text{out}} \rangle = \mathcal{G}(\langle \Delta \hat{a}_{\text{in}} \rangle + \mathcal{A})$  in which quantum mechanics restricts  $\mathcal{A} \geq 1/2$ .

Nonreciprocity and non-Hermitian topology have been proposed as promising resources for quantum sensors; the former allows fundamental constraints on the signal-to-noise ratio of conventional sensors to be overcome (Lau and Clerk, 2018; Kononchuk *et al.*, 2022; Slim *et al.*, 2024); in addition, the latter can lead to an exponentially enhanced sensitivity (McDonald and Clerk, 2020; Koch and Budich, 2022). Both nonreciprocity (Metelmann and Clerk, 2015) and non-Hermitian topological chains (McDonald, Pereg-Barnea, and Clerk, 2018; Wanjura, Brunelli, and Nunnenkamp, 2020) can be engineered in driven-dissipative quantum

systems, for instance, based on optomechanics (de Lépina *et al.*, 2020; Youssefi, Kono, Bancora *et al.*, 2022; del Pino, Slim, and Verhagen, 2022).

## 2. Standard quantum limit

To perform tests of fundamental physics (see Sec. V.A), it is often crucial to measure the oscillator position accurately. However, Heisenberg's uncertainty principle states that it is impossible to simultaneously know the position and momentum of a single quantum system with high accuracy. A measurement of the mass's position necessarily introduces backaction on its momentum. In this context we often speak of a standard quantum limit (SQL) that limits the accuracy of position measurements as the system evolves in time.

The SQL is straightforward to derive for an effective free mass  $m_{\text{eff}}$  that is harmonically trapped with frequency  $\omega_m$ . Its position quadrature  $\hat{X}_m(t)$  evolves in time as

$$\hat{X}_m(t) = \hat{X}_m(0) \cos(\omega_m t) + \frac{\hat{P}_m(0)}{m_{\text{eff}} \omega_m} \sin(\omega_m t), \quad (58)$$

where  $\hat{X}_m(0)$  and  $\hat{P}_m(0)$  are the position and momentum operators at  $t = 0$ , which remain unchanged during intervals smaller than the damping time. Then, considering the Heisenberg uncertainty principle, we find that

$$\Delta X_m(t) \Delta Y_m(t) \geq \frac{1}{2} |\langle [\hat{X}_m(0), \hat{Y}_m(0)] \rangle| = x_{\text{zpf}}^2, \quad (59)$$

where we have introduced the quadrature  $\hat{Y}_m(t) = \hat{P}_m(t)/m_{\text{eff}} \omega_m$  and  $x_{\text{zpf}} = \sqrt{\hbar/2m_{\text{eff}} \omega_m}$  is the zero-point fluctuation [ $\hat{X}_m(t)$  and  $\hat{Y}_m(t)$  are both expressed in units of length for easier comparison]. Thus, any measurement that tries to measure both quadratures with equal precision is limited to  $\Delta X_m(t) = \Delta Y_m(t) = x_{\text{zpf}}$ . See Caves *et al.* (1980) for a derivation of the SQL for a single quantum oscillator.

When the system is coupled to an external probe field, we consider an *optomechanical SQL* where the position of a mechanical resonator is detected through phase measurements (Bowen and Milburn, 2015). There are small fluctuations in the probe field itself that are known as shot noise. Shot noise can be decreased by increasing the number of quanta in the probe field, which improves the signal-to-noise ratio and makes detection easier. The increase in the field quanta does, however, lead to stronger recoil in the mechanical system known as quantum backaction noise or radiation-pressure noise. The result is a fluctuation force on the mechanical resonator. By balancing these two sources of noise, we arrive at the optomechanical SQL, which sets the limit on the achievable accuracy of position measurements. The SQL can be calculated by deriving the output spectrum of the measured probe field and balancing the resulting shot noise and radiation-pressure noise. See Clerk *et al.* (2010) and Bowen and Milburn (2015) for the full derivation. With the Langevin equations in Eq. (39) to model the dynamics, as well as the input-output relations shown in Eq. (38), we find that the measured quadrature of the mechanical mode in Fourier space results in the following symmetrized power spectral density (Bowen and Milburn, 2015):

$$S_{\text{det}}(\omega) = \frac{1}{8\eta\Gamma|C_{\text{eff}}|} + 2\Gamma|\chi(\omega)|^2|C_{\text{eff}}|, \quad (60)$$

where  $C_{\text{eff}}$  is the effective optomechanical cooperativity, defined as  $C_{\text{eff}} = C/(1 - 2i\omega/\kappa)^2$ , where  $C = 4g_0^2/\kappa\Gamma$  is the optomechanical cooperativity, for which  $g_0$  is the optomechanical coupling and  $\kappa$  and  $\Gamma$  are the optical and mechanical linewidths, respectively.  $\eta$  is the detection efficiency. The expression  $\chi(\omega)$  in Eq. (60) is the mechanical susceptibility, which is given by

$$\chi(\omega) = \frac{\omega_m}{\omega_m^2 - \omega^2 - i\omega\Gamma}. \quad (61)$$

To balance the two terms in Eq. (60), we require the optimal effective cooperativity to be

$$|C_{\text{eff}}^{\text{opt}}| = \frac{1}{4\eta^{1/2}\Gamma|\chi(\omega)|}. \quad (62)$$

The symmetrized spectrum at the SQL is then given by

$$S_{\text{det}}^{\text{SQL}}(\omega) = |\chi(\omega)|, \quad (63)$$

where we have assumed an optimal detection efficiency with  $\eta = 1$ . That is, in the optimum case the spectrum is given by the susceptibility of the resonator  $\chi(\omega)$ .

In practice, there are a number of additional noise sources that can be included in the measured noise spectrum, such as measurement imprecision and amplifier noise; see Sec. III.D.1. For example, Magrini *et al.* (2021) provided a detailed analysis of noise budgeting in an optomechanical experiment for the purpose of quantum-limited measurements, and Martynov *et al.* (2016) listed and characterized a number of relevant noise sources, such as thermal noise, laser noise, and electronic noise, in the LIGO detector that are also relevant to many other experiments. See also Danilishin and Khalili (2012) for a review on how quantum noise can be calculated in a gravitational-wave detector.

Note that the SQL is by no means a fundamental limit, as opposed to the Heisenberg limit; see Sec. III.D.3. It can be evaded using squeezed states, which reduce the noise in the measured quadrature (Bowen and Milburn, 2015).

## 3. Classical and quantum Fisher information

Previously, we focused on how well a detector can probe a signal against a noisy environment. However, we can also ask how much information a quantum system can fundamentally accumulate about a specific effect. This notion is captured by the Fisher information, which is a valuable metrology tool relevant for many of the weak-force detection schemes described in Sec. IV.A. See Paris (2009) and Giovannetti, Lloyd, and Maccone (2011) for comprehensive introductions.

Consider a specific measurement with outcomes  $\{x\}$  performed on the quantum state  $\hat{\rho}(\theta)$ , where  $\theta$  is the parameter that we want to estimate. The distribution of the measurement outcomes is given by  $p(x|\theta) = \text{tr}(\hat{\Pi}_x \hat{\rho}_\theta)$ , where  $\hat{\Pi}_x$  is a POVM element that models the measurements. The classical Fisher information (CFI) corresponds to the amount of



information about  $\theta$  gained from this measurement series. It is given by

$$I_F(\theta) = \int dx p(x|\theta) \left( \frac{\partial \ln p(x|\theta)}{\partial \theta} \right)^2. \quad (64)$$

The CFI can be generalized in the quantum case by optimizing over all possible measurements of the quantum state. This is known as the quantum Fisher information (QFI). The QFI can also be viewed as a distance measure that quantifies the change of the state due to the parameter  $\theta$ . That is, given the two quantum states  $\hat{\rho}_\theta$  and  $\hat{\rho}$ , the most general form of the QFI is

$$I_F(\theta) = 4 \left( \frac{\partial d_B(\hat{\rho}_\theta, \hat{\rho})}{\partial \theta} \Big|_{\epsilon=0} \right), \quad (65)$$

where  $d_B$  is the Bures distance (Helstrom, 1967; Bures, 1969),

$$d_B(\hat{\rho}_1, \hat{\rho}_2) = \sqrt{2[1 - \sqrt{\mathcal{F}(\hat{\rho}_1, \hat{\rho}_2)}]}, \quad (66)$$

for which  $\mathcal{F}$  is the fidelity  $\mathcal{F}(\hat{\rho}_1, \hat{\rho}_2) = (\text{tr}[\sqrt{\sqrt{\hat{\rho}_1} \hat{\rho}_2 \sqrt{\hat{\rho}_1}}])^2$ . For pure states  $|\Psi(\theta)\rangle$  that encode the parameter  $\theta$ , the QFI becomes

$$I_F(\theta) = 4(\langle \partial_\theta \Psi(\theta) | \partial_\theta \Psi(\theta) \rangle - |\langle \Psi(\theta) | \partial_\theta \Psi(\theta) \rangle|^2), \quad (67)$$

where  $\partial_\theta$  denotes the partial derivative with respect to  $\theta$ . The QFI can also be computed for initially mixed states  $\hat{\rho}$  that evolve unitarily, such that  $\hat{\rho}(\theta) = \hat{U}_\theta \hat{\rho}(0) \hat{U}_\theta^\dagger$ . When the initial state can be decomposed in terms of an orthonormal basis  $\hat{\rho}(0) = \sum_n \lambda_n |\lambda_n\rangle \langle \lambda_n|$ , the QFI can be written as (Liu *et al.*, 2014; Pang and Brun, 2014)

$$I_F(\theta) = 4 \sum_n \lambda_n (\langle \lambda_n | \hat{\mathcal{H}}_\theta^2 | \lambda_n \rangle - \langle \lambda_n | \hat{\mathcal{H}}_\theta | \lambda_n \rangle^2) - 8 \sum_{n \neq m} \frac{\lambda_n \lambda_m}{\lambda_n + \lambda_m} |\langle \lambda_n | \hat{\mathcal{H}}_\theta | \lambda_m \rangle|^2, \quad (68)$$

where the second sum is over all terms with  $\lambda_n + \lambda_m \neq 0$ ,  $\lambda_n$  is the eigenvalue of the eigenstate  $|\lambda_n\rangle$ , and the Hermitian operator  $\hat{\mathcal{H}}_\theta$  is defined as  $\hat{\mathcal{H}}_\theta = -i \hat{U}_\theta^\dagger \partial_\theta \hat{U}_\theta$ . Equation (68) can also be extended to the multiparameter case (Liu *et al.*, 2020), where it sometimes is possible to extract more information than in the single-parameter case (Paris, 2009).

When the quantum system interacts with an environment, the QFI can be challenging to compute since Eq. (68) no longer holds. In addition to the general expression in Eq. (65), the QFI can be defined in terms of the symmetric logarithmic derivative  $\hat{L}_\theta$  (Helstrom, 1969; Holevo, 2011),  $I_F = \text{tr}(\hat{\rho} \hat{L}_\theta^2)$ , where  $\hat{L}_\theta$  is given by

$$\partial_\theta \hat{\rho} = \frac{1}{2} (\hat{\rho} \hat{L}_\theta + \hat{L}_\theta \hat{\rho}). \quad (69)$$

If an expression for  $\hat{L}_\theta$  is found, the QFI can be immediately calculated. One way to solve Eq. (69) for  $\hat{L}_\theta$  by treating it as a Lyapunov matrix equation, which has a general solution (Paris, 2009)

$$\hat{L}_\theta = 2 \int_0^\infty dt \exp[-\hat{\rho}_\theta t] \partial_\theta \hat{\rho}_\theta \exp[-\hat{\rho}_\theta t]. \quad (70)$$

When the channel that encodes the parameter  $\theta$  can be represented with Kraus operators, a general upper bound to the QFI can be derived (Escher, de Matos Filho, and Davidovich, 2011).

A key feature of the QFI and the CFI is that they fundamentally relate to the variance of the parameter  $\theta$  through the Cramér-Rao bound (Helstrom, 1969; Braunstein and Caves, 1994)

$$\text{var}(\theta) \geq \frac{1}{\mathcal{M} I_F(\theta)}, \quad (71)$$

where  $\mathcal{M}$  is the number of measurements performed and the inequality is saturated for the QFI. Since  $\mathcal{M}$  is always finite, it is generally desirable to maximize the Fisher information to reduce the variance of the estimated parameter. The Cramér-Rao bound is applicable to any quantum system and provides a generalized uncertainty relation (Braunstein, Caves, and Milburn, 1996), even when no Hermitian operator can be associated with the parameter of interest, such as in the case of phase estimation (Helstrom, 1969; Braunstein and Caves, 1994; Braunstein, Caves, and Milburn, 1996). The Cramér-Rao bound also relates to the so-called Heisenberg limit, which is defined as the scaling of the variance  $\text{var}(\theta)$  of a parameter  $\theta$ . For classical systems, the scaling of  $\text{var}(\theta)$  is at most  $1/N$ , where  $N$  is the number of probes used, in accordance with the central limit theorem (Giovannetti, Lloyd, and Maccone, 2011). As opposed to the standard quantum limit (see Sec. III.D.2), the Heisenberg limit is a hard limit that depends on the number of resources in the system (Zwierz, Pérez-Delgado, and Kok, 2010). These resources can refer either to a number of subsystems or to the translational power of the Hamiltonian, which is higher for nonlinear dynamics (that is, Hamiltonian terms with products of more than quadratic operators). For example, a self-Kerr Hamiltonian with a term proportional to  $(\hat{a}^\dagger \hat{a})^2$  has more translational power than that with just  $\hat{a}^\dagger \hat{a}$ . The Heisenberg scaling does, however, go beyond  $1/N$  for certain initial quantum states. For example, the QFI for phase estimation with NOON states scales as  $1/N^2$  (Bollinger *et al.*, 1996; Dowling, 1998). The QFI has also been used to investigate certain relativistic settings (Pinel *et al.*, 2013; Ahmadi, Bruschi, and Fuentes, 2014; Tian *et al.*, 2015; Hao and Wu, 2016) and has been proposed as a probe of spacetime structure (Du and Mann, 2021).

#### IV. PROPOSED TESTS OF GRAVITY WITH MASSIVE QUANTUM SYSTEMS

Equipped with tools to model massive systems in the laboratory, we now ask the questions: What are the possible ways in which gravity influences quantum systems, and how can these effects be detected? A number of diverse and creative proposals have been put forward that probe the properties of gravity, quantum mechanics, and their interfaces. The goal of this section is to outline the main directions of

research and key proposals that allow tests to be performed with massive quantum systems. We begin by considering gravity from a classical source, such as Earth's gravitational fields, and its detection by quantum systems (Sec. IV.A). Specifically, we focus on proposals for how quantum properties can enhance the sensitivity of the probe. We proceed to consider proposals where gravity causes decoherence of a quantum probe (Sec. IV.B), including different types of decoherence proposals as well as nonlinear modifications of quantum theory. We then review recent proposals for detecting gravitationally induced entanglement (Sec. IV.C). Finally, Sec. IV.D outlines additional proposals that do not strictly fit into the other sections but are still relevant to the topic of this review.

### A. Precision tests of gravity

One approach for precision tests of gravity relies on the use of sensitive mechanical resonators. This section provides a review of weak-force sensing with massive systems in the quantum regime. We restrict ourselves here to proposals where the sensor is in the quantum regime, but where gravity originates from a classical source. This should be contrasted with the tests discussed in Sec. IV.C, where the quantum system itself is considered as a source of the gravitational field. Many precision tests of gravity have already been performed with classical mechanical resonators and atom interferometers. See Sec. V.A for an overview of these experiments and Fig. 7 for a summary of force sensitivities that have been achieved to date.

#### 1. Weak-force detection with backaction evading measurements

The large mass of massive quantum systems (compared with the mass scale of single atoms) means that they couple more strongly to gravity. A common goal of precision gravimetry with massive quantum systems is to resolve the force that affects the center of mass of the mechanical resonator. Usually, a probe field or two-level system is used for the control and readout of the sensor. For a probe field backaction noise and the inherent uncertainty of field fluctuations give rise to the SQL, which we reviewed in Sec. III.D.2, beyond which displacements cannot be resolved. The limits for a moving-end mirror were first discussed by Arcizet *et al.* (2006), and the first experimental observation of radiation pressure due to shot noise was performed by Purdy, Peterson, and Regal (2013).

Backaction evading (BAE) schemes constitute a key resource for weak-force sensing since they allow for an increase in measurement precision without adding additional noise during readout. The effect of measurement backaction can be circumvented if, rather than attempting to measure both of the mechanical quadratures, one merely couples the light field to one of the quadratures such that it becomes a conserved quantity (Thorne *et al.*, 1978; Braginsky, Vorontsov, and Thorne, 1980). Concretely, this means that if we couple only the  $\hat{X}_m = (\hat{b}^\dagger + \hat{b})/\sqrt{2}$  quadrature of the mechanical oscillator to the radiation pressure of the photon field  $\propto \hat{n}$ , the interaction Hamiltonian is  $\hat{H}_{\text{int}} \propto \hat{n} \hat{X}_m$ , which implies that  $[\hat{H}_{\text{int}}, \hat{X}_m] = 0$ . An observable that commutes with the

Hamiltonian is also referred to as a quantum nondemolition (QND) variable since it can be measured repeatedly without destroying the quantum state.

Coupling only one quadrature to the light field is experimentally challenging since it would require a time-dependent coupling between the quadratures and the detected field. This can be achieved in a scheme for a cavity-optomechanical system driven on both the red and the blue sideband (Clerk, Marquardt, and Jacobs, 2008) or by modulating the optomechanical coupling strength (Clerk, Marquardt, and Jacobs, 2008). As discussed in Sec. III.A, the coupling between the cavity mode  $\hat{a}$  and the mechanical oscillator  $\hat{b}$  in the frame rotating with the cavity frequency  $\omega_c$  is given by  $\hat{H}_I \approx -\hbar g_0 (\alpha^* \delta \hat{a} + \alpha \delta \hat{a}^\dagger) (\hat{b}^\dagger + \hat{b})$ . A drive can be modeled by  $\hat{H}_d = \alpha(t) \hat{a} + \alpha^*(t) \hat{a}^\dagger$ , with  $\alpha(t)$  the complex drive amplitude. For a drive on the red sideband at  $\omega_c - \omega_m$ , with  $\omega_m$  the mechanical frequency, we have  $\alpha(t) \propto e^{i(\omega_c - \omega_m)t}$  giving rise to the interaction Hamiltonian  $\hat{H}_I \propto \hat{a}^\dagger \hat{b} + \text{H.c.}$  in the frame rotating with  $\omega_c$  for the photons and  $\omega_m$  for the mechanical modes and in which we neglected counterrotating terms. Similarly, a drive on the blue sideband at  $\omega_c + \omega_m$  and  $\alpha(t) \propto e^{i(\omega_c + \omega_m)t}$  gives rise to the interaction Hamiltonian  $\hat{H}_I \propto \hat{a} \hat{b}^\dagger + \text{H.c.}$  Combining the drives on the red and blue sidebands, we obtain

$$H_I \propto \hat{X}_c \hat{X}_m. \quad (72)$$

That is, the interaction Hamiltonian couples the quadrature  $\hat{X}_c = (\hat{a}^\dagger + \hat{a})/\sqrt{2}$  of the cavity to the quadrature  $\hat{X}_m$  of the mechanical mode such that both  $\hat{X}_m$  and  $\hat{X}_c$  are constants of the motion since  $[H_I, \hat{X}_m] = [H_I, \hat{X}_c] = 0$ . Since the mechanical oscillator couples only to the  $\hat{X}_m$  quadrature of the cavity, information on its motion propagates only into the conjugate cavity quadrature  $\hat{P}_c = i(\hat{a}^\dagger - \hat{a})/\sqrt{2}$ . This allows us to repeatedly (or continuously) measure  $\hat{P}_c$  and infer  $\hat{X}_m$  to arbitrary precision. For a more detailed theoretical discussion of this scheme, see Clerk (2020) and the reviews by Braginsky and Khalili (1996) and Clerk *et al.* (2010). The exact conditional dynamics of an optomechanical system driven on both sidebands were analyzed in detail by Brunelli, Malz, and Nunnenkamp (2019).

The previously described scheme was experimentally implemented in a superconducting electromechanical device by Suh *et al.* (2014), which allowed the detection and reduction of backaction and the detection of a single quadrature below the zero-point fluctuations in the optical domain by Shomroni *et al.* (2019). BAE was also realized in hybrid optomechanical systems of a macroscopic mechanical oscillator and a spin oscillator (Møller *et al.*, 2017). A more elaborate BAE scheme relies on constructing an effective oscillator out of two and measuring a collective variable (Woolley and Clerk, 2013), which was experimentally realized in a microwave circuit with two mechanical oscillators (Ockeloen-Korppi *et al.*, 2016), thereby achieving a measurement precision below the zero-point fluctuations. Backaction noise can also be canceled through the addition of an ensemble of cold atoms that act as a negative-mass oscillator and allow for sensing beyond the SQL

(Motazedifard *et al.*, 2016). Furthermore, in a scheme with four drives (de L  pinay *et al.*, 2021), backaction was evaded and the entanglement between the two oscillators was demonstrated.

Apart from improving the measurement precision, BAE schemes result in the squeezing of the mechanical modes. In a generic, parametrically driven mechanical oscillator, the attainable amount of squeezing is limited due to the onset of parametric instability. However, continuous BAE measurements allow us to overcome this 3 dB limit for squeezing (Lei *et al.*, 2016). Beyond the previously discussed schemes that rely on measuring a single quadrature, an alternative approach to cancel quantum noise and overcome the SQL of force sensing using coherent feedforward quantum control was proposed by Tsang and Caves (2010). Another proposal showed that mechanically driving the oscillator within the BAE scheme leads to a monotonic response of the cavity, which can be beneficial for sensing force gradients (Arvidsson *et al.*, 2024).

Yet another direction for resolving the energy levels of a mechanical oscillator in an electromechanical experiment (Dellantonio *et al.*, 2018) is to use QND measurements to bring us a step closer to understanding how quantum jumps between phonon states work, which is challenging since the coupling to an environment makes it difficult to detect mechanical mode occupation. The notion of QND variables was generalized to a quantum-mechanics-free subsystem (Tsang and Caves, 2012), i.e., subsystems in which all observables commute and their expectation values are governed by classical equations of motion.

## 2. Additional weak-force detection schemes

Theoretical proposals for force sensing with massive quantum systems generally take one of two approaches: either they show that the SQL can be circumvented through novel protocols, such as backaction evading measurements (see Sec. IV.A.1) and the addition of quantum resources, or they consider the fundamental sensitivity that the systems can achieve, often quantified by the classical and quantum Fisher information (see Sec. III.D.3). Both approaches limit the precision of the measurement. Generally, the Fisher information quantifies the precision that can be achieved beyond the SQL.

Apart from backaction evasion, quantum resources such as squeezing and entanglement are required for beating the SQL (Zhang and Zhuang, 2021). In optomechanical systems squeezing of both optical and mechanical motion can be implemented in a number of ways; see Sec. V.B.1 for an overview of experiments. For example, in a mirror-in-the-middle optomechanical setup that results in two coupled cavity modes (Xu and Taylor, 2014), the SQL is surpassed due to the resulting squeezing of the light. The inclusion of a single- or two-mode parametric amplifier (PA) (Mollow and Glauber, 1967) with either  $\hat{a}^{\dagger 2} + \hat{a}^2$  (single mode) or  $\hat{a} \hat{b} + \hat{a}^{\dagger} \hat{b}^{\dagger}$  (two modes) results in sensing precision beyond the SQL (Motazedifard *et al.*, 2019; Zhao *et al.*, 2020). Further, in dissipative optomechanical systems, a PA can counteract the negative effects of mechanical damping, which allows us to go beyond the SQL (Huang and Agarwal, 2017).

Advantages through squeezing can also be achieved by adding a nonlinear medium in the cavity (Peano *et al.*, 2015). Squeezing has also been shown to improve the precision of LIGO (Aasi *et al.*, 2013; Buikema *et al.*, 2020).

Beyond squeezing, entanglement plays a crucial role in sensing and is a key ingredient for achieving a sensitivity that scales with the Heisenberg limit (Zhuang, Zhang, and Shapiro, 2018). Most importantly, by performing measurements with  $N$  entangled sensors, we may go beyond the  $1/\sqrt{N}$  scaling achieved with independent probes and possibly obtain a scaling with  $1/N$ . For example, EPR-entangled states have been proposed for use in LIGO (Ma *et al.*, 2017). Brady *et al.* (2022) proposed the use of an array of mechanical sensors connected by entangled light, with applications for dark-matter searches; see Sec. IV.A.5. However, it has been shown that sensors with multicarrier optical modes do not outperform their single-mode counterparts (Branford, Miao, and Datta, 2018). It has been experimentally demonstrated that using two optically entangled mechanical membranes leads to a 40% improvement in the shot-noise-dominant regime (Xia *et al.*, 2023) and allows a scaling better than  $1/\sqrt{N}$ .

Another method for improving the precision of quantum sensors involves noise mitigation and engineering of the surrounding noise bath. A structured non-Markovian environment was found to amplify the susceptibility for weak-force sensing with an optomechanical sensor (Zhang *et al.*, 2017). More broadly, the use of quantum error correction techniques has been proposed for quantum metrology (D  r *et al.*, 2014; Kessler *et al.*, 2014), even to the extent that the Heisenberg limit can be achieved (Zhou *et al.*, 2018).

Yet another proposal for high-precision sensing, albeit challenging, is through the use of macroscopic superpositions in the sense of a large mass being in a quantum superposition of two distinct spatial locations. A method for generating such superpositions that is particularly effective for large masses is to couple a spin with a mass through a magnetic field gradient (a Stern-Gerlach mechanism, as described in Sec. IV.C) (Scala *et al.*, 2013; Bose, 2016; Wan *et al.*, 2016; Bose *et al.*, 2017; Marshman *et al.*, 2020, 2021; Margalit *et al.*, 2021; Zhou *et al.*, 2022, 2023), which followed from general ideas to couple ancillary systems such as a quantized electromagnetic mode in a cavity with a mechanical object (Bose, Jacobs, and Knight, 1997, 1999; Mancini, Man’ko, and Tombesi, 1997; Marshall *et al.*, 2003; Armata *et al.*, 2017; Qvarfort *et al.*, 2018) and other ancillary quantum systems (superconducting qubits, etc.) (Bose, 2006; Bose and Agarwal, 2006; Johnsson, Brennen, and Twamley, 2016). Such quantum superpositions can be used to detect weak forces to a precision linear in time (essentially Heisenberg scaling), as the accumulated relative phase between the superposed components grows linearly in time. Moreover, at the end of such quantum ancilla-induced interferometry, the phase can be sensed by simply measuring the ancilla. Example applications in the gravitational context involve detection of accelerations to high sensitivity (Johnsson, Brennen, and Twamley, 2016; Qvarfort *et al.*, 2018; Marshman *et al.*, 2020), gravity gradient noise (Toro  s *et al.*, 2021), space debris (Wu, Toro  s *et al.*, 2023), and the possibility to detect gravitational waves with a meter-sized compact interferometer for nano-objects (Marshman *et al.*,



2020) [applications also exist outside the gravitational domain, for example, to detect neutrinos (Kilian *et al.*, 2023)].

The QFI (see Sec. III.D.3) allows us to consider sensitivities beyond the SQL. In the linearized optomechanical regime, the QFI has been considered for squeezed state inputs (Lee *et al.*, 2022). In the nonlinear regime of optomechanics (see Sec. III.A), the QFI was computed for detecting a constant (Armata *et al.*, 2017; Qvarfort *et al.*, 2018), as well as time-dependent gravitational potentials including gravitational waves (Qvarfort, Plato *et al.*, 2021). The QFI was also computed for an opto-magnon-mechanical setup, where the optomechanical system senses small changes in the separation between two magnets (Iakovleva, Sarma, and Twamley, 2023).

For further reading on sensing with mechanical resonators, see the following dedicated reviews on sensing, which cover levitated systems (Rademacher, Millen, and Li, 2020), hybrid optomechanical-BEC systems (Motazedifard, Dalafi, and Naderi, 2021), and cavity optomechanics (Li *et al.*, 2021; Liu, Liu, Ren *et al.*, 2021).

### 3. Weak-force detection with BECs

The mass of a BEC is generally lower than that of a composite quantum resonator, which means that it generally couples more weakly to gravity; see Sec. V.A for a comparison of experimental parameters. However, the fact that all atoms in a BEC are identical makes it possible to control it extremely well in the laboratory. As such, BECs have been explored for force sensing. Gravity sensing with BECs can be done using trapped atoms, or atoms in free fall. In the free-fall case, the precision depends on the time of flight. The time of flight in atom interferometry can be increased using Bragg diffraction and Bloch oscillations of a BEC to slow down the particles (Abend *et al.*, 2016). In such schemes interactions are undesirable because they reduce the coherence time of the interferometer (Pereira dos Santos *et al.*, 2017). However, interactions can be used to prepare initial states that have higher sensitivities (Szigeti *et al.*, 2020). Nevertheless, spatial interferometers cannot be reduced in size without losing precision. An alternative that could resolve this limitation is trapped BECs. Interactions in a trapped BEC give rise to phonons. Phonon modes are sharp in frequency, while the atoms are completely delocalized within the trapped potential. Recent work shows that interferometry in the frequency domain using phonon modes can be used to miniaturize detectors while still retaining high precision (Howl and Fuentes, 2023). In frequency interferometry the precision is limited by the lifetime of the states, not by the size of the system. Squeezed states of phonon modes can be used to measure the gravitational field and its gradient with high precision (Fuentes *et al.*, 2019; Bravo, Rätzel, and Fuentes, 2020) since the frequency of the modes is affected by the gravitational field. Phonon modes can also be used to measure oscillating gravitational fields, such as the acceleration and gradient of an oscillating mass close to the BEC (Rätzel *et al.*, 2018). Single phonon measurement precisions have been reached in BEC analog experiments (Steinhauer, 2022). The most relevant limiting factor is particle loss due to three-body recombination. The resonance of phonons modes

to external gravitational fields has been proposed to detect high-frequency gravitational waves and searches for dark matter (Sabín *et al.*, 2014; Howl and Fuentes, 2023). A BEC trapped in a double well has been proposed in searches of dark energy (Hartley *et al.*, 2024b), and a proposal to show that gravity degrades entanglement between two BEC in a space-based experiment was presented by Bruschi, Sabín *et al.* (2014). While there are only theoretical proposals, the center-of-mass oscillations of a BEC was used to measure Casimir-Polder forces in the lab by Harber *et al.* (2005). Their work shows that BEC technology is useful in measuring small forces.

### 4. Deviations from the Newtonian potential

An open problem in modern physics is the discrepancy between the observation of a small cosmological constant and the predicted value from particle physics theory (Padilla, 2015). Modified gravity theories (MGTs) provide a solution to this dilemma in that some of them predict deviations from general relativity while simultaneously addressing the discrepancy with particle physics. For a review of MGTs, see Clifton *et al.* (2012). To address the fact that no deviations from general relativity have been observed thus far, mechanisms are introduced to explain the absence of large deviations in tests that have been performed thus far. One such proposal, known as a chameleon mechanism (Brax *et al.*, 2004; Khoury and Weltman, 2004a, 2004b), resolves the discrepancy via the introduction of a screening mechanism that depends on the local mass density. Note, however, that it does not solve the cosmological constant problem. According to the proposal, deviations in regions with high density, such as the Solar System, are suppressed. Instead, high vacuum and extremely sensitive laboratory tests are needed. For current bounds on chameleon theories, see the figures given by Burrage and Sakstein (2018).

Most MGTs can be parametrized into the following Yukawa-like modification to Newton's potential:

$$V(r) = -\frac{GM_S m_p}{r} (1 + \alpha e^{-r/\lambda}), \quad (73)$$

where  $M_S$  is the source mass,  $m_p$  is the probe mass (not to be confused with the Planck mass  $M_P$ ),  $\alpha$  is a dimensionless modification to the strength of the potential, and  $\lambda$  is a length scale beyond which the modification is exponentially suppressed. Current Solar System tests of Newton's laws have considerably constrained the free parameters of such modified theories; see Fig. 8 of Murata and Tanaka (2015). The parameter regimes that remain to be excluded include small  $|\alpha|$  and  $\lambda$ , which correspond to the detection of extremely weak forces at short range.

The main avenue for searches for MGTs with massive quantum systems is via precision tests of gravity, which we covered in Secs. IV.A.1–IV.A.3. Several experiments have already been performed with mechanical resonators in the classical regime to bound deviations from the Newtonian potential, including with cantilevers (Chiaverini *et al.*, 2003). A key advantage for mechanical resonators such as levitated systems is that they are relatively confined in space and

therefore can be used to test extremely short length scales. See [Moore and Geraci \(2021\)](#) for a review of searches for new physics with optically levitated sensors. Additional proposals have been put forward for tests with levitated optomechanical devices ([Blakemore \*et al.\*, 2021](#); [Chen, Liu, and Zhu, 2022](#)), as well as specific tests of the chameleon mechanism ([Betz \*et al.\*, 2022](#)). However, while the larger mass of mechanical resonators increases the strength of the Newtonian potential and thus the deviations, their larger volume brings with it additional challenges. For example, in the context of the chameleon mechanism, the large radius of a levitated nanomechanical resonator was found to additionally screen the interaction ([Qvarfort, Rätzel, and Stopyra, 2022](#)). Another major challenge is the presence of Casimir forces, which increase at small distances. See [Onofrio \(2006\)](#) for a review of measurements of Casimir forces in the context of searching for deviations from the Newtonian potential.

### 5. Tests of the equivalence principle and dark-matter searches

The equivalence principle (EP) states that all forms of matter couple to gravity in the same way. An additional formulation known as the weak equivalence principle (WEP) states that gravitational mass and inertial mass are the same. Violation of the EP can be indicative of modified theories of gravity ([Hui, Nicolis, and Stubbs, 2009](#)) or physics beyond the standard model ([Damour, 2012](#)).

As pointed out in Sec. II.B, there is a conflict between the EP and quantum mechanics. Nevertheless, several ideas have been put forward for testing the EP with quantum systems. Perhaps the simplest is a classical test of the Eötvös ratio, which defines the correlation between inertial mass and gravitational mass and which serves as a test of the WEP. The Eötvös ratio is defined as

$$\eta_{A-B} = 2 \frac{|a_A - a_B|}{|a_A + a_B|} = 2 \frac{|(m_i/m_g)_A - (m_i/m_g)_B|}{|(m_i/m_g)_A + (m_i/m_g)_B|}, \quad (74)$$

where  $a_A$  and  $a_B$  are the accelerations of bodies  $A$  and  $B$  and  $m_i$  and  $m_g$  are the inertial and gravitational mass, respectively. The advantage of using quantum systems pertains mainly to the increased precision that they offer as sensors. Most of the measurements of the Eötvös ratio with quantum systems have been carried out through atom interferometry ([Schlippert \*et al.\*, 2014](#); [Duan \*et al.\*, 2016](#); [Overstreet \*et al.\*, 2018](#); [Albers \*et al.\*, 2020](#); [Asenbaum \*et al.\*, 2020](#)). The best Eötvös ratio achieved to date is that of the MICROSCOPE mission, at  $[1.5 \pm 2.3(\text{stat}) \pm 1.5(\text{syst})] \times 10^{-15}$  ([Touboul \*et al.\*, 2022](#)). See also Sec. V.A.1 for an overview of state-of-the-art tests with cold atoms.

In general, there appears to be no clear consensus in the community on how the EP should be formulated for quantum systems, since there are often additional aspects that need to be taken into account. One of the earliest works on this topic showed that, for the simple case of a particle in an external gravitational field, the WEP does not apply to a quantum-mechanical description of the problem ([Greenberger, 1968](#)). While the classical equations of motion can be made independent of mass, the same is not true for quantum mechanics, since mass enters into the quantization rules. However, the

opposite point of view has also been argued. Starting with linearized gravity perturbations as a massless, spin-2 gauge field coupled to itself and to matter, the equivalence principle must hold for quantum systems for the theory to be consistent ([Davies and Falkowski, 1982](#)). It has also been argued that, for quantum particles in free fall, their expectation values for position and momentum are consistent with the WEP ([Viola and Onofrio, 1997](#)).

The influence of internal degrees of freedom of quantum systems on the formulation of the EP has been raised in several works. Since the EP stipulates equivalence between mass and energy, it must take into account the internal (potentially superposed) energy states of a quantum system. Based on this, a quantum formulation of the EP has been proposed that requires equivalence between the rest, inertial, and gravitational internal energy operators ([Zych and Brukner, 2018](#)). An experimental test based on this proposal was put forth by [Orlando \*et al.\* \(2016\)](#), who used trapped spin-1/2 atoms, and [Rosi \*et al.\* \(2017\)](#), who later performed the test using a Bragg atom interferometer. The test provided constraints on the off-diagonal elements of the mass operators and additional constraints of the Etvös ratio for the WEP. Similarly, the WEP can be explicitly considered for internal degrees of freedom. Two distinct formulations of the WEP were proposed by [Anastopoulos and Hu \(2018\)](#). The first formulation states that the probability distribution of position for a free-falling particle is the same as the probability distribution of a free particle (up to a mass-independent shift of its mean). The second states that any two particles with the same velocity wave function behave identically in free fall, regardless of their masses. It has also been stipulated that a quantum version of the EP should be linked to a notion of causality ([Hardy, 2018](#)) since it is always possible to transform to a quantum reference frame in which we have a definite causal structure in the local vicinity of any point. Here the notion of a quantum reference frame refers to frameworks developed by [Guérin and Brukner \(2018\)](#) and [Giacomini, Castro-Ruiz, and Brukner \(2019\)](#) and related work by [de la Hamette \*et al.\* \(2022\)](#).

Tests of the EP can also aid in the search for dark matter since some dark-matter models [such as light scalar dark matter ([Hees \*et al.\*, 2018](#))] predict deviations from the EP. See [Carney \*et al.\* \(2021\)](#) and [Kilian \*et al.\* \(2024\)](#) for overviews of quantum sensing with mechanical resonators for the detection of dark matter. It has been proposed that mechanical oscillators with masses below or around 1 kg operating near the standard quantum limit could be used to detect ultralight dark-matter candidates ([Carney, Hook \*et al.\*, 2021](#)). In addition, an optical cavity with mirrors made of different materials could facilitate coupling channels for vector dark matter ([Manley \*et al.\*, 2021](#)). Heavier dark-matter candidates can also be detected by scattering off a mechanical resonator. A large array of femtogram masses could potentially detect dark-matter candidates at around 10 keV, with the advantage that they also provide directional sensing ([Afek, Carney, and Moore, 2022](#)). A recent white paper by [Brodeur \*et al.\* \(2023\)](#) focused on detecting deviations in momentum kicks resulting from exotic decay processes. They carefully monitored the position of levitated spheres using displacement sensing.

## B. Gravitational decoherence, semiclassical models, self-energy, and gravitationally induced wave-function collapse

Gravity can impart a coherent signal on the quantum state, which can be detected using quantum metrology tools; see Sec. IV.A. However, there are a number of theoretical proposals where the quantum state no longer follows a unitary evolution when interacting with gravity. Here we cover proposals ranging from decoherence arising from quantum and stochastic gravity to modifications to the Schrödinger equation. We also refer the interested reader to the comprehensive reviews dedicated to these topics by Bassi, Großardt, and Ulbricht (2017) and Anastopoulos and Hu (2022).

### 1. Gravitational decoherence

Decoherence is the process by which off-diagonal elements in the density matrix of a quantum system are gradually reduced to zero. There are many proposals for how an external gravitational field interacts with the quantum system to cause decoherence, as well as dissipation and thermalization, in the regime of gravity at low energies. Unlike other sources of decoherence, such as those from fluctuating electromagnetic fields, gravitational decoherence is universal, and its influence cannot be shielded.

The common starting point for most gravitational decoherence proposals is the linearized metric

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}, \quad (75)$$

where  $\eta_{\mu\nu}$  is the Minkowski background spacetime and  $h_{\mu\nu}$  denotes the fluctuations. Fluctuations can emerge within the perturbative quantum theory of gravity [see Eq. (20)], can be postulated in a fundamentally classical theory of gravity [see the discussion near Eq. (89)] or could emerge as a consequence of a minimum length scale of the spacetime fabric (Hossenfelder, 2013). Regardless of its physical origin, the fluctuations of the gravitational field are expected to decohere a quantum system as in any other fluctuating field; see Sec. III.B.

In most proposals the structure of the dynamics of a massive quantum system moving along one axis is captured with the following master equation:

$$\frac{\partial}{\partial t} \hat{\rho} = -\frac{C}{2} [\hat{A}, [\hat{A}, \hat{\rho}]], \quad (76)$$

where we have omitted the Hamiltonian terms for brevity and  $C$  ( $\hat{A}$ ) is a constant prefactor (an operator) specific to the model [see Eq. (34) with only the jump operator  $\hat{L}_1 = \sqrt{C}\hat{A}$ ]. It is then easy to obtain the decoherence rate  $\gamma$  in the eigenbasis of the operators  $\hat{A}$ . Suppose that  $a_1$  and  $a_2$  are two real-valued eigenvalues of the operator  $\hat{A}$ . By applying  $\langle a_L |$  ( $|a_R\rangle$ ) from the left-hand side (right-hand side) of Eq. (76) and multiplying by 2, we readily find that

$$\gamma = C\Delta a^2, \quad (77)$$

where we have defined  $\Delta a = a_L - a_R$  (which can be interpreted as the superposition size). When  $\hat{A}$  is not Hermitian, the analysis in Eqs. (76) and (77) requires generalizations; see the references cited later in the review for further details.

Anastopoulos and Hu (2013) and Blencowe (2013) considered gravitational waves (forming an environmental bath) as a source for the fluctuations  $h_{\mu\nu}$ . It was found that a free particle should decohere with the operators in Eq. (76) given by kinetic energy  $\hat{A} = \hat{p}^2/2m$ , where  $\hat{p}$  is the momentum operator and  $m$  is the particle mass. The characteristic decoherence rate is given by

$$\gamma = \frac{9}{32\pi\tau_P} \frac{T_P}{\Theta} \left( \frac{\Delta E}{E_P} \right)^2, \quad (78)$$

where  $\tau_P$  ( $T_P$ ) is the Planck time (Planck temperature),  $\Delta E$  ( $E_P$ ) is the difference in kinetic energy of a zero momentum and a finite momentum state (Planck energy), and  $\Theta$  is a free parameter of the model. In earlier works a general non-Markovian master equation for the interaction between  $N$  gravitating quantum particles was derived by Anastopoulos (1996), a complementary analysis was given by Oniga and Wang (2016), and a generalization for photons was discussed by Lagouvardos and Anastopoulos (2021).

Decoherence due to the emission of gravitational waves was studied by Suzuki and Queisser (2015). A relation between decoherence and the classical limits in terms of the quadrupole radiation formula and backreaction dissipation was discussed by Oniga and Wang (2017). The analysis from Toroš, Mazumdar, and Bose (2024) recovered the classical results for a linear quadrupole and showed that only systems with quadrupoles would decohere, while a free particle would not decohere. For the simplest case of a harmonically trapped particle (which has a linear quadrupole), it was found that the decoherence operator is  $\hat{A} = \hat{b}^2$  (with  $\hat{b}$  the mode operator of the harmonic oscillator). The associated decoherence rate for number states  $|n\rangle$  is given by

$$\gamma = \frac{64\omega}{15} \left( \frac{E}{E_P} \right)^2 (\langle \hat{n}^2 \rangle - \langle \hat{n} \rangle^2), \quad (79)$$

where  $\omega$  is the frequency of the harmonic trap,  $E = \hbar\omega$  ( $E_P$ ) is the difference between the energy levels (Planck energy), and  $\hat{n}$  denotes the number operator. Starting with Einstein's equivalence principle, it was found that the emission of quantized gravitational waves can happen only via the transition  $n \rightarrow n-2$ , which is prohibited for the states  $|1\rangle$  and  $|0\rangle$ . The number states  $|0\rangle$  and  $|1\rangle$ , or any superposition of these states, are thus protected from decoherence via quantized gravitational waves, resulting in a vanishing decoherence rate in Eq. (79). For a discussion about the bremsstrahlung effects, see Weinberg (1965).

The decoherence effect induced by gravitons in the context of gravitational-wave detectors was discussed by Parikh, Wilczek, and Zahariade (2020) and Parikh, Wilczek, and Zahariade (2021a), while Kanno, Soda, and Tokuda (2021) also considered matter-wave interferometry. The approximate decoherence rate obtained for an interferometer consisting of two paths (i.e., left and right paths) is given by

$$\gamma = 10\Omega_m \left( \frac{mv}{m_P c} \right)^{1/2}, \quad (80)$$



where  $m$  ( $m_p$ ) is the mass of the system (the Planck mass),  $2v$  is the relative speed between the states following the left and right paths of the interferometer, and  $\Omega_m$  is a high-frequency cutoff (which is inversely proportional to the superposition size  $\Delta x$ , i.e.,  $\Omega_m \sim c/\Delta x$ ).

A class of stochastic models can be obtained by considering the nonrelativistic limit of classical field equations and assuming stochastic fluctuations of the metric. Starting with the Klein-Gordon equation, it was found that the decoherence operator is  $\hat{A} = \hat{p}^2/2m$ , where  $\hat{p}$  is the momentum operator and  $m$  is the particle mass (Breuer, Göklü, and Lämmerzahl, 2009). The decoherence rate is given by

$$\gamma = \frac{\tau_c}{\hbar^2} \Delta E^2, \quad (81)$$

where  $\tau_c$  is a free parameter characterizing the correlation time of the stochastic bath and  $\Delta E$  is the difference in kinetic energy. A generalized analysis using the Foldy-Wouthuysen method capturing higher-order corrections was performed by Asprea *et al.* (2021).

Another proposal for decoherence is related to composite quantum particles such that individual parts of the systems follow different geodesics. Piovski *et al.* (2015) showed that gravity entangles the internal and center-of-mass degrees of freedom, which in turn decoheres the center-of-mass degrees of freedom of a system, for example, of a crystal. An equation of the form in Eq. (76) was obtained with the decoherence operator  $\hat{A} = \hat{x}$ , where  $\hat{x}$  is the center-of-mass position operator. The decoherence rate was given as

$$\gamma = \frac{\sqrt{N} g k_b T \Delta x}{\sqrt{2} \hbar c^2}, \quad (82)$$

where  $N$  is the number of degrees of freedom in the crystal,  $g$  is Earth's gravitational acceleration,  $k_b$  is the Boltzmann constant,  $T$  is the temperature, and  $\Delta x$  is the spatial superposition size. The proposal has been the source of much discussion in the community; see Bassi, Großardt, and Ulbricht (2017) and Piovski *et al.* (2017) for a summary of these discussions.

Experimental signatures on matter-wave interferometers were analyzed by Lamine *et al.* (2006), Wang, Bingham, and Mendonça (2006), and Asprea *et al.* (2021). While many of these proposals suggest different mechanisms behind the gravitationally induced decoherence, a large number result in similar reductions of the density matrix elements. As a result, there are tests that search for gravitational decoherence, regardless of its origin. Some of the first proposals for testing gravitational collapse and decoherence with optomechanical systems were given by Bose, Jacobs, and Knight (1999) and Marshall *et al.* (2003). In this protocol a single optical mode is passed through a beam splitter and into an interferometer, wherein one of the arms interacts with a mechanical resonator according to the optomechanical Hamiltonian in Eq. (28). The protocol has been analyzed with the addition of mechanical position-damping noise (Adler, Bassi, and Ippoliti, 2005; Bassi, Ippoliti, and Adler, 2005); see Sec. III.B.1. Additionally, the protocol has been examined in the high-temperature limit (Bernád, Diósi, and Geszti, 2006). It was further developed by Kleckner *et al.* (2008), where a number

of practical aspects were taken into account. Such an experiment has not yet been performed. It has been pointed out that the phases picked up through the optomechanical interaction can be partially reproduced through classical dynamics (Armata *et al.*, 2016).

Should gravity cause decoherence, it is possible that the mechanism itself relies on a modification of quantum mechanics. If the experiments are still modeled using standard quantum mechanics, it can become difficult to distinguish the decoherence effects accurately. This scenario was considered by Pfister *et al.* (2016), who proposed a general information-theoretic measure of decoherence. While many approaches consider continuous variable systems, there are also results for qubit states (Kok and Yurtsever, 2003).

## 2. Nonlinear modifications

A key cornerstone of quantum mechanics is that the Schrödinger equation is a linear equation for the state of the system  $|\psi\rangle$  and that the expectation value of an observable  $\hat{O}$  is a bilinear function of the state, for example,  $\langle\psi|\hat{O}|\psi\rangle$ . We can, however, devise a modification of quantum mechanics where the dynamics becomes nonlinear in the state  $|\psi\rangle$  or construct expectation values with a nonbilinear dependency on the state (Weinberg, 1989a, 1989b). Such modifications are motivated by the measurement problem (see Sec. II.B) and emerge from elementary considerations about semiclassical gravity. Semiclassical gravity is viewed by some as an effective theory, i.e., an approximation to a fundamental quantum theory of gravity. Within this approximation one can investigate the backreaction of matter on the gravitational field, generalizing the results of quantum field theory in curved spacetime (see Sec. II.A.4), and can model classical stochastic fluctuations of the gravitational field (Hu and Verdaguer, 2008). An alternative viewpoint is that semiclassical gravity is not a mere approximation but rather the fundamental theory where gravity remains classical while matter is quantized (Kibble, 1981).

A conceptual start for such a nonlinear modification of quantum mechanics is given in the semiclassical Einstein equation (Møller *et al.*, 1962; Rosenfeld, 1963; Ruffini and Bonazzola, 1969)

$$G_{\mu\nu} = \frac{8\pi G}{c^4} \langle \hat{T}_{\mu\nu} \rangle, \quad (83)$$

where on the left-hand side we have the classical Einstein tensor  $G_{\mu\nu}$ , and on the right-hand side we have the expectation value of the quantum stress-energy tensor  $\hat{T}_{\mu\nu}$  taken with respect to the state of the quantum matter. The coupling in Eq. (83) is arguably the simplest way to couple a classical gravitational field to quantized matter, but more importantly it is the expected theory when the matter is in well-localized states. In such a case matter can still be approximately described using a classical stress-energy tensor  $T_{\mu\nu}$  such that  $T_{\mu\nu} \approx \langle \hat{T}_{\mu\nu} \rangle$ , but beyond this regime, such as when we have spatial superpositions, there is no consensus about its validity, as we later discuss.

In the nonrelativistic regime the gravitational field acted on by a particle in the state  $|\psi\rangle$  thus depends on the value of

$\langle \psi | \hat{T}_{\mu\nu} | \psi \rangle$ . Hence, we expect the associated nonlinear Schrödinger equation to have a cubic dependency on the state of the matter system, i.e.,

$$i\hbar \frac{d}{dt} |\psi\rangle \propto G \langle \psi | \hat{T}_{\mu\nu} | \psi \rangle |\psi\rangle. \quad (84)$$

This equation, which was proposed by Diósi (1984) and Penrose (1996, 1998), has become known as the Schrödinger-Newton equation. A derivation from first principles of Eq. (84) is, however, still a subject of debate (Christian, 1997; Adler, 2007; Giulini and Großardt, 2012; Anastopoulos and Hu, 2014a, 2014b).

Such a hybrid quantum matter-classical gravity model has its appeal in the conceptual simplicity of Eq. (83), with testable predictions differing from those arising from the framework of perturbative quantum gravity; see Sec. II.A.7. However, unlike the latter, which is a fully consistent relativistic theory, deterministic nonlinear modifications of the Schrödinger equation, such as the Schrödinger-Newton equation, are at odds with the requirement of no-faster-than-light signaling to make them, at least conceptually, unsatisfactory (Gisin, 1989; Polchinski, 1991).

Nonetheless, to date, no laboratory experiments have been able to rule out the Schrödinger-Newton equation. Furthermore, it has been suggested that it might be possible to resolve the issue of superluminal signaling if one takes into consideration the measurement problem with a suitable prescription of the wave-function collapse (Bahrami *et al.*, 2014; Bera, Mohan, and Singh, 2015), and hence the predictions of the Schrödinger-Newton equation might still remain valid in specific domains; see Sec. IV.B.3 for a discussion of possible modifications. For a discussion within the context of generalized probabilistic theories and its relation to classical state space, see Mielnik (1974, 1980) and Galley, Giacomini, and Selby (2022). We provide a summary of the current experimental endeavors to test the Schrödinger-Newton equation here.

Starting with the semiclassical Einstein equations in Eq. (83), we obtain in the nonrelativistic limit the one-particle Schrödinger-Newton equation (Diósi, 1984)

$$i\hbar \frac{d}{dt} |\psi_t\rangle = \frac{\hat{p}^2}{2m} |\psi\rangle - Gm^2 \int d\mathbf{s} \frac{|\psi(t, \mathbf{s})|^2}{|\hat{\mathbf{r}} - \mathbf{s}|} |\psi\rangle, \quad (85)$$

where  $m$  is the mass of the particle,  $\hat{\mathbf{r}}$  ( $\hat{p}$ ) is the position (momentum) operator, and we have introduced the wave function  $\psi(t, \mathbf{s}) = \langle \mathbf{s} | \psi_t \rangle$ . The generalization to the  $N$ -particle case can be obtained from Eq. (85) by replacing the source of the gravitational field with  $|\psi(t, \mathbf{s}_1, \dots, \mathbf{s}_N)|^2$ ,

$$i\hbar \frac{d}{dt} |\psi_t\rangle = -Gm^2 \sum_{j,k} \int \prod_l d\mathbf{s}_l \frac{|\psi(t, \mathbf{s}_1, \dots, \mathbf{s}_N)|^2}{|\hat{\mathbf{r}}_j - \mathbf{s}_k|} |\psi\rangle, \quad (86)$$

where we have omitted the kinetic terms for brevity and  $j, k = 1, \dots, N$ .

Equations (85) and (86) form the starting point for a number of experimental proposals. While some analytical results can be obtained (Tod and Moroz, 1999) in most situations, one has to resort to numerical simulations to make quantitative

predictions using the nonlinear Schrödinger-Newton equation, as in the case of the formally similar Gross-Pitaevskii equation (Gross, 1961; Pitaevskii, 1961). The key prediction of the Schrödinger-Newton equation is the modification of the free spreading of the wave function as the last term in Eq. (85), with its Newtonian-like  $1/r$  dependency, can be viewed as a self-gravity term that tends to localize the system in space. A number of papers have investigated the free spreading in space with the required parameter regime for experimental tests (Moroz, Penrose, and Tod, 1998; Carlip, 2008; Giulini and Großardt, 2011; Bahrami *et al.*, 2014; Colin, Durt, and Willox, 2016), and there have been proposals to test secondary effects in harmonic traps such as squeezing (Yang *et al.*, 2013) and energy shifts (Großardt *et al.*, 2016).

Additional dependencies on the state of the system  $|\psi\rangle$  can be introduced to other parts of the quantum formalism (Sorkin, 1994). Motivated by considerations about general covariance, some have argued that all quantities in physical theories must be dynamical (Norton, 1993), suggesting corrections to the Born rule (Berglund *et al.*, 2022). Cubic corrections to the Born rule  $\mathcal{O}(|\psi|^3)$ , i.e., triple interference phenomena, have been theoretically discussed in the context of the Talbot interferometer (Berglund *et al.*, 2023). The class of nonlinear modifications introduced by Weinberg (1989a, 1989b) have also recently been analyzed in the context of gravitationally induced entanglement (Spaventa, Lami, and Plenio, 2023).

### 3. Nonlinear and stochastic modifications

The quest of unifying quantum mechanics and gravity into a single theory and the measurement problem from quantum foundations at first appear to be two distinct problems; see Sec. II.B. However, using an elementary analysis, Penrose (1986) showed that there appears to be a deep conflict between the superposition principle in quantum mechanics and the equivalence principle of general relativity. This result could be viewed as another hint of the necessity to modify gravity, i.e., to construct a quantum theory of gravity. However, any theory where the superposition principle remains valid would not resolve the tension between quantum and classical physics, thereby leaving the measurement problem unanswered. Another option is that the conflict instead signifies the need to also modify quantum mechanics to accommodate notions of gravity, i.e., gravitization of quantum mechanics (Penrose, 1986, 1996, 1998, 2014). Such a theoretical program, while still in its tentative state, suggests that it might be possible to consistently couple classical and quantum systems (in this context gravity and matter, respectively) and to solve the measurement problem simultaneously. We remark, however, that other programs for the emergence of classicality, fully compatible with unmodified quantum mechanics, are also considered in the literature (Giulini, 2000).

The measurement problem of quantum mechanics, which is still a subject of controversy, has its roots in the two prescriptions for the evolution of quantum systems: on the one hand, the Schrödinger equation is deterministic and linear, while, on the other hand, the wave-function collapse postulate induces a stochastic and nonlinear evolution of the state. It was shown that it is possible to combine the two types of prescriptions into a single dynamical law, thus avoiding the

dichotomy, with quantum dynamics the limit for microscopic systems and classical dynamics the limit for macroscopic systems (Ghirardi, Rimini, and Weber, 1986).

The structure of such modifications forms the basis for the family of spontaneous wave-function collapse models (Bassi and Ghirardi, 2003; Bassi *et al.*, 2013), with the basic form given by the following stochastic differential equation (Ghirardi, Pearle, and Rimini, 1990):

$$\frac{d}{dt}|\psi_t\rangle = \sqrt{\lambda}(\hat{A} - \langle\hat{A}\rangle)\frac{dW_t}{dt}|\psi_t\rangle - \frac{\lambda}{2}(\hat{A} - \langle\hat{A}\rangle)^2|\psi_t\rangle, \quad (87)$$

where  $|\psi_t\rangle$  is the state vector,  $\hat{A}$  is the operator,  $\langle\psi_t|\hat{A}|\psi_t\rangle$  is the expectation value,  $dW_t$  is the Wiener increment, and  $\lambda$  is the coupling rate. The models based on Eq. (87) make a series of predictions that are expected to be tested with the next generation of experiments, for example, loss of interferometric visibility, anomalous heating of free systems, and x-ray emission (Bassi *et al.*, 2013; Carlesso *et al.*, 2022). In contrast to the case of deterministic nonlinear modifications (see Sec. IV.B.2), the stochastic nature of the evolution in Eq. (87) conspires with the nonlinear terms to avoid the possibility of superluminal signaling, making such models conceptually more appealing, albeit a relativistic extension of such models is an open problem (Bedingham *et al.*, 2014). Collapse models have already been tested experimentally, including via mechanical systems in a noninterferometric way (Vinante *et al.*, 2016, 2020; Helou *et al.*, 2017; Forstner *et al.*, 2020), which provide the strongest experimental bound to date; see Carlesso *et al.* (2022) for a review on collapse model testing.

As such, the connection between dynamical collapse models of the form in Eq. (87) and gravity remains tentative, with a derivation from first principles an open question (Bahrami, Smirne, and Bassi, 2014). Nonetheless, using an elementary analysis considering a spatial superposition, it has been argued that the system should decohere within a time given by  $\tau = \hbar/E_g$ , where  $E_g$  is the gravitational self-energy of the difference between the mass distributions of the two states in superposition (Penrose, 1986, 1996, 1998, 2014). For example, for a spherical mass distribution, we have the following formula (Penrose, 2014):

$$E_g = \begin{cases} \frac{Gm^2}{R}(2\lambda^2 - \frac{3}{2}\lambda^3 + \frac{1}{3}\lambda^5), & \lambda \leq 1, \\ \frac{Gm^2}{R}(\frac{6}{5} - \frac{1}{2\lambda}), & 1 \leq \lambda, \end{cases} \quad (88)$$

where  $m$  is the total mass,  $R$  is the particle radius,  $\lambda = \Delta x/(2R)$ , and  $\Delta x$  is the superposition size.

Although self-gravity has been extensively investigated in general relativity (Lynden-Bell, 1961), writing a fully consistent relativistic spontaneous collapse model is an open problem. In the nonrelativistic limit, one can nonetheless construct a wave-function collapse model inspired by Newtonian gravity (Diósi, 1987, 1989), which recovers the prediction for Penrose's decoherence time  $\tau$  (Diósi, 2005, 2007). A drawback of the proposed model is, however, that it requires a short-distance cutoff to avoid the divergence for pointlike microscopically mass distributions (Ghirardi, Grassi, and Rimini, 1990). We discuss experimental bounds on the

short-distance cutoff, usually labeled as  $R_0$ , at the end of this section.

There have been a series of investigations aiming to derive the collapse of the wave function from an underlying mechanism related to random fluctuations of spacetime,

$$g_{\mu\nu} = \bar{g}_{\mu\nu} + \delta g_{\mu\nu}, \quad (89)$$

where  $g_{\mu\nu}$  is the spacetime metric,  $\bar{g}_{\mu\nu}$  denotes a fixed background, and  $\delta g_{\mu\nu}$  denotes the stochastic fluctuations. One of the earliest such attempts posited that the wave-function collapse could be induced by real-valued fluctuations of the spacetime metric related to the Planck scale (Karolyhazy, 1966). However, the obtained model is still compatible with the superposition principle, and it seems to be at odds with the predicted x-ray emission from charged particles (Diósi and Lukács, 1993). An alternative idea with complex-valued fluctuations of the spacetime metric was also proposed (Adler, 2004), with a possible model of the basic form of Eq. (87) constructed by Gasbarri *et al.* (2017).

A collapse-like dynamics of the form in Eq. (87) also appears in the context of quantum measurement and control (Wiseman and Milburn, 2009) and has been exploited to construct models of semiclassical gravity. Specifically, by continuously measuring the system, one has access to the signal  $I(t)$  given by Eq. (47). The signal  $I(t)$  is used in experiments to gather information about the state of the system as well as to control the evolution of the system at future times, i.e., by creating a feedback loop. However, in this context the system is not measured by an actual experimentalist or physical measurement apparatus, but it is instead postulated that this dynamics, resembling continuous measurements, is a fundamental law of nature (Diósi, 2018). Such an approach was used in the Kafri-Taylor-Milburn model (Kafri, Taylor, and Milburn, 2014) to construct a semiclassical (linearized) Newtonian interaction. We recall that the quantum interaction arising in standard quantum mechanics is given by

$$-\frac{Gm_1m_2}{|(d+\hat{x}_1)-\hat{x}_2|} \approx \frac{2Gm_1m_2}{d^3}\hat{x}_1\hat{x}_2, \quad (90)$$

where  $d$  is the mean distance between the two masses  $m_1$  and  $m_2$  and  $\hat{x}_1$  and  $\hat{x}_2$  are the position operators, respectively; see Eq. (7). Kafri, Taylor, and Milburn (2014) used a formalism reminiscent of the previously outlined quantum measurement and control. In place of Eq. (90), one finds a modified potential

$$\frac{Gm_1m_2}{d^3}(\hat{x}_1\langle\hat{x}_2\rangle + \langle\hat{x}_1\rangle\hat{x}_2). \quad (91)$$

A related approach was considered by Tilloy and Diósi (2016), who continuously monitored the matter density of the system  $\rho$ , producing the signal given by

$$\rho_t = \langle\rho_t\rangle + \delta\rho_t, \quad (92)$$

where  $\langle\rho_t\rangle$  is the expectation value taken with respect to the state of the system. Tilloy and Diósi (2016) showed that when  $\rho_t$  is the source of Newtonian potential  $\phi$  in the Poisson



equation, i.e.,  $\nabla^2\phi = 4\pi G\rho_t$ , one is able to recover the standard quantum Newtonian interaction among particles in Eq. (90), as well as the terms appearing in the previously discussed Diósi model (Diósi, 1987, 1989).

Another approach that is related to the Diósi model is given by hybrid quantum-classical models (Oppenheim, 2023). In such models working in the Arnowitt-Deser-Misner formalism (Arnowitt, Deser, and Misner, 1959) [see Poisson (2004) for an introduction], one describes the gravitational field using a probability density in classical phase space  $\rho(z)$ , where we attach to each point  $z$  in phase space a distinct density matrix  $\hat{\sigma}(z)$ . As a result, the total state of the system (comprising gravity and matter) is described by the classical-quantum state given by

$$\hat{\rho}_{\text{cq}} = \int d\rho(z)|z\rangle\langle z| \otimes \hat{\sigma}(z). \quad (93)$$

We can construct a master equation for such a dynamics: by tracing away the state of gravitational field  $\rho(z)$ , we obtain a master equation for the matter state (while tracing away the matter system, one obtains a Fokker-Planck-like equation for the gravitational field that provides a correction to general relativity). In the former case, we can, by suitably restricting the general form of the initial equation, recover a master equation arising from Eq. (87) such as the one given by the Diósi model (Oppenheim *et al.*, 2022).

Testing of spontaneous collapse models falls into two broad categories: interferometric and noninterferometric tests. In the former the signature is a loss of interferometric visibility (Toroš and Bassi, 2018) where the current record is provided by experiments with macromolecules (Fein *et al.*, 2019), while the latter is a broad class of all other experiments (Carlesso *et al.*, 2022). To test the idea put forward by Penrose, one must resort to direct tests of the superposition principle in interferometric tests (as a mathematical model has not yet been constructed). One way to test this idea is to use Bose-Einstein condensates (Fuentes and Penrose, 2018; Howl, Penrose, and Fuentes, 2019). However, the model put forward by Diósi, precisely formulated as discussed, can also be tested with indirect noninterferometric tests. The model depends on a single free parameter  $R_0$ , which can be interpreted as the localization length scale, which has been constrained using x-ray emission to values  $R_0 < 5 \times 10^{-10}$  m (Donadi *et al.*, 2021; Arnquist *et al.*, 2022).

### C. Entanglement mediated by gravity

It is not yet known whether gravity is fundamentally a quantum force. In this section we present arguments supporting the view that detecting entanglement induced by gravitational interaction between two masses could finally help to settle this issue. We also review proposals of how to generate such entanglement. The realistic possibility of such an experiment has captured attention only recently. One reason for this raised interest was a proposal that showed that masses as small as micron-sized crystals, which can be isolated by levitation, can be fruitfully combined with spin qubit and quantum gate technologies developed for quantum computation to generate and witness gravitational entanglement (Bose, 2016; Bose

*et al.*, 2017). Simultaneously, a rationale based on a known result from quantum information theory, along with some basic assumptions, was also presented and justified why such entanglement would evidence the quantum nature of gravity (Bose, 2016; Bose *et al.*, 2017; Marshman, Mazumdar, and Bose, 2020); see also Marletto and Vedral (2017) for an alternative rationale. An important fact, noted for the first time by the aforementioned researchers, is that one has to prepare and coherently maintain quantum superpositions having large spatial spreads for the entanglement growth to be realistically measurable. This makes the experiments challenging. We review this class of schemes here, as well as the recent rapid growth of literature that has been triggered by the aforementioned works. These schemes dwell on the need to identify all the necessities needed for a practical realization, alternative proposals, and other ways to justify the conclusions about the quantum features of gravity from such experiments.

In a historic verbal debate with other researchers on whether gravity is quantum, Feynman (2011) advocated a thought experiment involving one mass in a spatial quantum superposition displacing a second mass due to its gravity; see Fig. 11 in Sec. V.B. Although he stressed using quantum amplitudes to describe the setup, he did not explain why such an experiment will demonstrate the quantum character of gravity, nor did he clarify what to measure in order to draw such conclusions. In addition, he did not realize that the second mass also has to be in a highly delocalized state (a spatial quantum superposition of delocalization comparable to the first mass) for the experiment to have a realistic duration. Subsequently, a version of Feynman's setup was suggested by supposing that the large mass in a spatial superposition was a Bose-Einstein condensate (Lindner and Peres, 2005). Such a superposition demands all atoms in one well or another; it is not an easy state to generate. It was then suggested that the interference pattern of a mesoscopic mass in a momentum state be measured (again, this is not easy to prepare), scattering gravitationally from this superposition. However, a single-particle measurement does not suffice to reveal entanglement without extra assumptions or procedures. Moreover, it is not justified exactly why such an interference pattern would constitute a test for quantum-natured gravity. After presenting measurements of gravity for the smallest masses (milligram scale) to date, Schmöle *et al.* (2016) remarked that reaching a quantum coherent regime for such masses can generate entanglement gravitationally. However, an estimate of the required regime and states was not the subject of that work. On the side of logic, Kafri and Taylor (2013) defined a classical force as one that cannot generate entanglement and found that this classical force leads to excess noise. They then suggested a design using tethered torsional oscillators to detect this excess noise, which is challenging and requires  $10^3$  s per run; if that noise is not found, then one can conclude a quantum coherent coupling. Thus, the emphasis in the experimental proposal was to rule out a specific classical interaction rather than verify gravitationally generated entanglement. Similarly, Krisnanda *et al.* (2017) showed that if an inaccessible system entangled two quantum probes, then it would display nonclassical correlations with the probes in the form of quantum discord. They then suggested that this could potentially detect the

nonclassicality of gravity without going into the formulations of any scheme. Thus, an explicit scheme with calculated numbers, which was proposed to exploit contemporary developments in levitation and quantum technologies, along with a simple rationale from quantum information theory was necessary (Bose, 2016; Bose *et al.*, 2017) to inspire confidence in the testability of the quantum character of gravity in the laboratory.

### 1. Gravitationally interacting interferometer-based protocol

We outline here the protocol presented by Bose (2016) and Bose *et al.* (2017). Published the same year was a paper by Marletto and Vedral (2017), although, as they dealt only schematically with the same idea without outlining explicit schemes, we present the following in accordance with Bose *et al.* (2017). We consider two masses, labeled as  $j = 1$  and  $2$ , each with a spin embedded in it. A particularly relevant experimental example for these masses would be a diamond nanocrystal hosting an NV center, which is a highly coherent spin-1 system used in the area of quantum computation (Bar-Gill *et al.*, 2013; Hensen *et al.*, 2015; Wood *et al.*, 2022; March *et al.*, 2023). However, any other crystal with an embedded spin with a long coherence time would suffice and, generically, we require only two spin states, which we label as  $|\uparrow\rangle$  and  $|\downarrow\rangle$ . The two masses, labeled as  $j = 1$  and  $2$ , are each created in a quantum superposition of well-separated Gaussian states  $|L\rangle_j$  and  $|R\rangle_j$  by means of the Stern-Gerlach effect. Thus, we imagine that, ideally, a spin embedded in each mass  $j$  would be placed in a quantum superposition of spin states  $(|\uparrow\rangle_j + |\downarrow\rangle_j)/\sqrt{2}$ , where a spin-dependent force (as in the Stern-Gerlach effect) is applied to the masses so that they move from their initial central positions given by Gaussians  $|C\rangle_j$  to evolve to

$$|\psi\rangle_j = \frac{1}{\sqrt{2}}(|L\rangle_j + |R\rangle_j). \quad (94)$$

This is shown in the upper half of Fig. 4 as the point at which the trajectories achieve their maximal splitting  $\Delta x$ . After achieving a certain maximal splitting  $\Delta x$ , the Stern-Gerlach




FIG. 4. Mechanism of gravitationally generated entanglement. The mechanism of entanglement of two masses through the phase evolution due to gravitational interaction is displayed. The phase evolution due only to the most prominent interaction, between  $|R\rangle_1$  and  $|L\rangle_2$ , is explicitly shown. Adapted from Bose *et al.*, 2017.

force (a spin-dependent splitting process) is stopped, for example, by either switching off the magnetic field or mapping the electronic spins to nuclear spins (this is shown as shoulders in the interferometers of Fig. 4), and the masses are allowed to translate in parallel next to each other. We are going to assume (for simplicity of presentation, although this assumption may be difficult to fulfill) that the superposition is created so fast that the phase accumulation due to gravitational interaction during this time is negligible. During the parallel motion after the creation of the superposition, the four configurations  $LL$ ,  $LR$ ,  $RL$ , and  $RR$  (where  $L$  refers to mass 1 and  $R$  refers to mass 2) have different energies due to their Newtonian interaction. Thus, their quantum phase evolutions happen at different frequencies,

$$\begin{aligned} \omega_{RL} &\sim \frac{Gm_1 m_2}{\hbar(d - \Delta x)}, & \omega_{LR} &\sim \frac{Gm_1 m_2}{\hbar(d + \Delta x)}, \\ \omega_{LL} &= \omega_{RR} \sim \frac{Gm_1 m_2}{\hbar d}. \end{aligned} \quad (95)$$

For simplicity as well as for appreciating the maximal efficiency of the process, we consider the situation when the superposition splitting is much larger than the distance of the closest approach of the masses, i.e.,  $\Delta x \gg d - \Delta x$ . In that case we can simplify to a situation where only  $\omega_{RL}$  is prominent, while the other frequencies are negligible (taken to be zero with respect to  $\omega_{RL}$ ). Then the evolution of the state at a time  $\tau$  is

$$\begin{aligned} |\Psi(t = \tau)\rangle_{12} &= |L\rangle_1 \frac{1}{\sqrt{2}}(|L\rangle_2 + |R\rangle_2) \\ &+ |R\rangle_1 \frac{1}{\sqrt{2}}(e^{-i\omega_{RL}\tau}|L\rangle_2 + |R\rangle_2). \end{aligned} \quad (96)$$

By inspection of the state, we conclude that for any value of  $\omega_{RL}\tau \neq 2k\pi$ , where  $k$  is an integer, the state is entangled, as it cannot be factorized into a product state of the two qubits as  $(|L\rangle_2 + |R\rangle_2) \neq (e^{-i\omega_{RL}\tau}|L\rangle_2 + |R\rangle_2)$ . In fact, for  $\omega_{RL}\tau \sim \pi$  the state is a maximally entangled state of two qubits (the qubit states defined by two orthogonal states  $|L\rangle$  and  $|R\rangle$ ).

To compute the highest possible value of the frequency  $\omega_{RL}$ , one needs to identify the minimum value of  $d - \Delta x$ . This in turn depends on the range within which you can bring the two masses without electromagnetic interactions swamping gravity. It is possible in principle to make masses neutral (by shining UV radiation on the masses or the enclosure). We assume that it is possible to make the masses free of internal charge multipoles (how to achieve this has not yet been fully solved for realistic nanocrystals and microcrystals). Suppose that the electrostatic interactions between the masses are fully eliminated; there is still the Casimir interaction. The ratio of the Casimir to the gravitational interaction is given by

$$\frac{U_{\text{Casimir}}}{U_{\text{gravity}}} \sim \frac{23}{4\pi} \left(\frac{3}{4\pi}\right)^2 \left(\frac{\epsilon - 1}{\epsilon + 2}\right)^2 \frac{m_P^2}{\rho^2(d - \Delta x)^6}, \quad (97)$$

where  $m_P$  is the Planck mass and  $\rho$  is the density of the masses. As we have to use some material to get the dielectric constant and density properties, we choose a diamond, which

is a good candidate for the experiment since it can host an embedded spin as an NV center defect, as stated. If we want gravity to dominate by a factor of 10 over the Casimir interaction, we get the minimum  $d - \Delta x \sim 157 \mu\text{m}$  (van de Kamp *et al.*, 2020; Schut, Geraci *et al.*, 2023). Putting this value into Eq. (95), we get for micron-sized objects (radius  $\sim 1 \mu\text{m}$ , mass  $\sim 10^{-14} \text{ kg}$ )

$$\omega_{RL} \sim \frac{Gm_1m_2}{\hbar(d - \Delta x)} \sim 0.4 \text{ Hz.} \quad (98)$$

Now, how are we going to detect the previously generated entanglement? It is here that the rest of the interferometer shown in Fig. 4 is important. The paths are now recombined once again using Stern-Gerlach forces such that  $|L\downarrow\rangle_j \rightarrow |C\downarrow\rangle_j$  and  $|R\uparrow\rangle_j \rightarrow |C\uparrow\rangle_j$ . Then the state of the two embedded spins becomes

$$|\downarrow\rangle_1 \frac{|\downarrow\rangle_2 + |\uparrow\rangle_2}{\sqrt{2}} + |\uparrow\rangle_1 \frac{e^{-i\omega_{RL}\tau}|\downarrow\rangle_2 + |\uparrow\rangle_2}{\sqrt{2}}. \quad (99)$$

The entanglement of these spins can be verified by measuring spin-spin correlation functions and combining them to construct an entanglement witness. A good entanglement witness in this context [which works for smaller time evolution durations compared to the witness discussed by Bose *et al.* (2017)] is (Chevalier, Paige, and Kim, 2020; Guff, Boulle, and Pivovskii, 2022)

$$\mathcal{W} = 1 - \sigma_x^1 \sigma_x^2 - \sigma_y^1 \sigma_y^2 - \sigma_z^1 \sigma_z^2. \quad (100)$$

If after measuring the correlations the expectation value  $\langle \mathcal{W} \rangle < 0$ , then the state of the two spins is entangled. As the only interaction was gravitational, verifying the entanglement of these spins is equivalent to verifying the gravitationally generated entanglement.

In a real experiment it is possible that  $\Delta x$  is achieved slowly so that a significant contribution to gravitational entanglement happens even during the growth of the superposition. Thus, according to the protocol of entanglement generation,  $\tau$  should be an effective time that correctly captures the total entanglement growth rate during the evolution of the size of the superposition. Moreover, it is also possible that the ideal case of  $\Delta x \gg d - \Delta x$  is not easily achievable. In fact, it is perhaps more likely, at least in the earliest experiments, that  $\Delta x = \chi d$ , where fraction  $0 < \chi < 1$ . In this general case the entanglement developed, as well as the entanglement witness, depends only on a total phase, the “entangling phase,” which is defined as  $\phi_{\text{ent}} = (\omega_{LR} - \omega_{LL})\tau + (\omega_{RL} - \omega_{LL})\tau$ . For the configuration of interferometers given in Fig. 4, we have, for small enough values of the entangling phase  $\langle \mathcal{W} \rangle \sim -\phi_{\text{ent}}$  (Chevalier, Paige, and Kim, 2020),

$$\phi_{\text{ent}} = \frac{Gm_1m_2\tau}{\hbar d} \frac{2\chi^2}{1 - \chi^2}, \quad (101)$$

which for  $\chi \ll 1$  becomes

$$\phi_{\text{ent}} = \frac{2Gm_1m_2(\Delta x)^2\tau}{\hbar d^3}. \quad (102)$$

From Eq. (101) it becomes clear that the fraction  $\chi$  should be chosen to be as close as possible to unity for a higher magnitude of the entanglement witness, enabling a lower number of measurements to determine it. However, in the regime of  $\chi \ll 1$ , from Eq. (102) one observes that we can in principle choose either a light mass  $m$  and a large superposition size  $\Delta x$  (the case discussed in Sec. IV.C) or, alternatively, a heavy mass and a small superposition size without affecting the accumulated entangling phase. For example, with a mass of  $m = 1 \text{ kg}$ , a superposition size of  $\Delta x = 10^{-14} \text{ m}$ , an interparticle separation  $d = 7 \text{ cm}$  (commensurate with the dimensions of such an object taking a standard density for a nanocrystal, say, that of diamond, of  $3.5 \times 10^3 \text{ kg/m}^3$ ), and a time  $\tau = 1 \text{ s}$ ,  $\phi_{\text{ent}} \sim 0.2$  is obtained. However, such masses are usually tethered, which offers extra decoherence channels rather than being levitated. How to achieve, by squeezing and free expansion, a spinless measurement of a two-qubit entanglement witness (in terms of spatial qubits) has also been shown (Yi *et al.*, 2021, 2022).

## 2. Alternative protocols

Instead of an interferometric scheme, we can consider nearby harmonic oscillators with mechanical frequency  $\omega$  that are interacting gravitationally (Krisnanda *et al.*, 2020; Qvarfort, Bose, and Serafini, 2020). We can obtain a figure of merit for the generated entanglement from Eq. (102) by setting the delocalization to be the zero-point motion  $\Delta x = \sqrt{\hbar/(2m\omega)}$  [see Eq. (6)] and the interaction time to be  $t = 1/\omega$ ,

$$\eta = \frac{2Gm}{\omega^2 d^3}, \quad (103)$$

where we have defined  $\eta \equiv \phi_{\text{ent}}$  following the notation from Krisnanda *et al.* (2020). Choosing the separation  $d$  of the masses to be about 1.5 times their radius, Eq. (103) becomes solely dependent on their densities and  $\omega$  (Krisnanda *et al.*, 2020). To achieve considerable entanglement, we again require  $\eta \sim 1$ . For that, for example, even with the densest material (osmium), one has to accomplish the entire protocol with each mass in a  $\omega \sim \text{mHz}$  trap over an interaction time of  $10^3 \text{ s}$ , over which it will be difficult to retain quantum coherence. Thus, to achieve considerable entanglement, one has to use quantum states that are far more spatially spread (essentially similar in spread to the superpositions mentioned earlier) (Krisnanda *et al.*, 2020; Cosco, Pedernales, and Plenio, 2021; Weiss *et al.*, 2021) or non-Gaussian resources (i.e., we need to prepare non-Gaussian initial states) or with non-negligible nonlinear couplings [i.e., cubic or higher-order terms in the position operators beyond the expansion in Eq. (90)], as discussed by Qvarfort, Bose, and Serafini (2020). The generated entanglement can be read out using two optomechanical setups separately monitoring each of the two masses. That is, each mass is a mirror that is coupled to an optical field that can then be measured; see Secs. III.A.1 and III.C. Specific optomechanical configurations to measure the gravitationally induced entanglement have been considered using the single-photon nonlinear regime in a quantum Cavendish experiment (Al Balushi, Cong, and Mann, 2018;



Matsumura and Yamamoto, 2020) and in the linear regime with the cavity driven by a coherent laser field (Miao *et al.*, 2020). Furthermore, Datta and Miao (2021) gave an optomechanical scheme for measuring the differential motion of the two mirrors. They argued that detecting gravitationally induced squeezing of the differential motion should be experimentally more accessible than detecting quantum entanglement between the two masses and, consequently, the conclusions that can be drawn would be different.

We can also consider the experimental situation with unequal masses (Bose *et al.*, 2017). In place of  $(\Delta x)^2$  in Eq. (102), we have  $\Delta x_1 \Delta x_2$ , where  $\Delta x_j$  is the superposition size of the mass  $m_j$  ( $j = 1$  and  $2$ ). To generate substantial entanglement, we again require  $\phi_{\text{ent}} \sim 1$ . One scheme in this direction uses extremely different masses, an atom (in an atom interferometer) and a massive oscillator coupled to it gravitationally (Carney, Müller, and Taylor, 2021). Here treating the atom as a spatial qubit gives a cyclic decoherence-recoherence dynamics of the two-mass system, which is robust to the thermal state of the oscillator, an interesting property that also underpins some previous optomechanical (Bose, Jacobs, and Knight, 1997, 1999; Mancini, Man'ko, and Tombesi, 1997; Marshall *et al.*, 2003; Armata *et al.*, 2017; Qvarfort *et al.*, 2018) and qubit-oscillator schemes (Bose, 2006; Scala *et al.*, 2013). The decoherence recoherence of the spatial qubit is then suggested as evidence of entanglement, although this is reliant on the assumption of the high purity of the joint qubit-oscillator state. Another variant of this is a case with two qubits (Pedernales, Streltsov, and Plenio, 2022): a nanoscale mass as a spatial qubit that is gravitationally coupled to a massive mediating oscillator that is in turn coupled electromagnetically to another qubit. By measuring entanglement between the two qubits, one can test the quantumness of the gravitational interaction. While in a similar manner robustness to the thermal state of the intermediary is present [also reminiscent of geometric phase gates in ion traps (Mølmer and Sørensen, 1999; Solano, de Matos Filho, and Zagury, 1999; Milburn, Schneider, and James, 2000)], one still has the challenge of preparing a sufficiently spread spatial quantum superposition state of a nanoscale mass. Other works have also considered experimental configurations with modified geometries and multidimensional systems (Tilly *et al.*, 2021). Specific schemes with three or four particles (Schut *et al.*, 2022), as well as an array (Miki, Matsumura, and Yamamoto, 2021; Ghosal, Ghosal, and Bandyopadhyay, 2023), have been analyzed.

It has also been noted that the relativistic regime is required to probe the spin nature of the gravitational interaction (Scadron, 2006; Bjerrum-Bohr *et al.*, 2015; Carney, 2022; Biswas *et al.*, 2023). Bose *et al.* (2022) considered the leading-order post-Newtonian terms in an experimental situation with harmonic oscillators, and Aimet, Chevalier, and Kim (2022) let photons from two separate interferometers entangle.

Notably, a hybrid optomechanical scheme testing the quantum counterpart of the light bending by gravity was given by Biswas *et al.* (2023). The scheme consists of a harmonic oscillator of mass  $m$  placed at the origin and of a circular path of radius  $r$  for an optical field confined to

a half-ring cavity (i.e., the mechanical oscillator and the cavity photons are conceptually replacing the two massive interferometers shown in Fig. 4). The interaction between the trapped mass and the photons is purely gravitational, and it reduces to the form of the cavity-optomechanical interaction in Eq. (28) with the coupling  $g_0$ , in this case arising from the quantum light-bending interaction.

A figure of merit similar to the ones in Eqs. (102) and (103) can also be constructed for the latter cases, albeit the details can depend on the specific experimental configuration and on the interaction. To get a tentative idea about the order of magnitude of the entanglement phase with photons, we can use the relation  $m = \hbar\omega/c^2$ . For example, replacing one of the masses in Eq. (102) with  $m = \hbar n\omega/c^2$  (where  $\omega$  is the frequency of the optical field), we find a figure of merit for entanglement between a photon and a massive particle,

$$\phi_{\text{ent}} \sim \frac{2Gm\omega\Delta x^2\tau}{c^2 d^3}, \quad (104)$$

which can be used to gauge the order of magnitude of the entanglement phase. Taking the ratio of Eqs. (104) and (102), we thus see that the effects are suppressed by  $\hbar\omega/(mc^2)$  and, as such, achieving an entanglement phase of order unity requires a large photon number to enhance the effect.

There are also proposals for testing gravitationally induced self-interactions of matter (Anastopoulos and Hu, 2013, 2014a, 2020). While in the previous scheme one was interested in the interaction between two distinct systems  $\propto T_{\mu\nu}^{(A)} T_{\alpha\beta}^{(B)}$  one can also consider self-interaction terms  $\propto T_{\mu\nu}^{(A)} T_{\alpha\beta}^{(A)}$  and  $\propto T_{\mu\nu}^{(B)} T_{\alpha\beta}^{(B)}$ , where  $T_{\mu\nu}^{(A)}$  and  $T_{\mu\nu}^{(B)}$  denote the stress-energy tensors corresponding to systems  $A$  and  $B$ . We can readily see how such terms emerge in the Newtonian limit. The interaction with quantum matter from Eq. (23) reduces to

$$\hat{H}_{\text{int}} = \frac{1}{2} \int d\mathbf{r} \hat{\rho}(\mathbf{r}) \hat{\phi}(\mathbf{r}), \quad (105)$$

where  $\hat{H}_{\text{int}} = - \int \hat{\mathcal{L}}_{\text{int}}(\mathbf{r}) d\mathbf{r}$  is the interaction Hamiltonian,  $\hat{\rho}$  is the matter density, and  $\hat{\phi}$  is the Newtonian potential. The matter density  $\hat{\rho}$  is also a source for  $\hat{\phi}$ , with the solution given by the familiar Newtonian potential. From Eq. (105) we thus find that

$$\mathcal{L}_{\text{int}} = -\frac{G}{2} \int d\mathbf{r} \int d\mathbf{r}' \frac{\hat{\rho}(\mathbf{r}) \hat{\rho}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}. \quad (106)$$

Equation (106) is suggestive for a figure of merit based on the entanglement phase from Eq. (102). Since we have only one system, we set  $\Delta x \sim d$  to find that

$$\phi_{\text{ent}} \sim \frac{Gm^2\tau}{\hbar d}, \quad (107)$$

where  $d$  is interpreted as a characteristic length scale of the problem (for example, the wave-function spread). A scheme with BECs was investigated, where  $\hat{\rho} \propto \hat{a}^\dagger \hat{a}$  (with  $\hat{a}$  the mode of a BEC). We thus find from Eq. (105) a Kerr nonlinearity

that induces non-Gaussianity (Haine, 2021; Howl *et al.*, 2021). Recently, a scheme also considered using the self-interaction of photons (Mehdi, Hope, and Haine, 2023). In place of Eq. (107) we can use  $m = n\hbar\omega/c^2$  (where  $\omega$  is the frequency of the optical field) to find the figure of merit  $\phi_{\text{ent}} \sim 2G\hbar\omega^2\tau/(c^2d)$ , which can again be enhanced by considering a large number of photons. Note that gravitationally mediated entanglement experiments will also be able to test various variants of gravitational theories (Marshman, Mazumdar, and Bose, 2020; Beckering Vinckers, de la Cruz-Dombriz, and Mazumdar, 2023; Chakraborty, Mazumdar, and Pradhan, 2023; Elahi and Mazumdar, 2023). Several foundational questions involving the nonclassical behavior of gravity can be probed with similar setups (Kent and Pitalúa-García, 2021; Etezad-Razavi and Hardy, 2023), including the non-classical behavior of gravity under a measurement (Hanif *et al.*, 2023).

### 3. Major challenges

Proposals for the testing of gravitationally induced quantum phenomena discussed in this section face a series of experimental challenges specific to the experimental implementation. One major difficulty is the achievement of a large superposition for the interferometry-based schemes [for schemes using Gaussian wave packets, this translates to obtaining a large delocalization of the wave packet (Cosco, Pedernales, and Plenio, 2021; Weiss *et al.*, 2021), which is a problem of a similar nature]. A large mass requires a strong force to create a quantum superposition of components separated by  $\Delta x$ . Some of the early protocols of the Stern-Gerlach-based creation of superpositions (Scala *et al.*, 2013; Wan *et al.*, 2016) have been found to have limitations in the achievable growth rate of  $\Delta x$  (Pedernales, Morley, and Plenio, 2020; Marshman *et al.*, 2021). Some solutions have been investigated (Zhou *et al.*, 2022, 2023), and this splitting rate is a work in progress.

If the electromagnetic interactions between the masses can be screened (Schmölle *et al.*, 2016; Schut, Geraci *et al.*, 2023; Schut, Grinin *et al.*, 2023), then the masses can be brought closer (with  $d$  decreased); consequently, the requirement of  $\Delta x$  can be alleviated. For example, the most optimistic results known to us in this context resulted from using both screening and trapping (Schut, Geraci *et al.*, 2023). For a screening material of 1  $\mu\text{m}$  thickness,  $d \sim 11 \mu\text{m}$ , masses  $m_1 \sim m_2 \sim 10^{-14} \text{ kg}$ , then, for  $\Delta x \sim 0.65 \mu\text{m}$  and  $\tau \sim 1 \text{ s}$ ,  $\phi_{\text{ent}} \sim 0.01$  is obtained, which requires  $\sim 10^4$  repetitions of the experiment.

The other important obstacle is maintaining coherence. In short, in the presence of decoherence, at a rate  $\Gamma$  the witness becomes  $\langle \mathcal{W} \rangle \sim \Gamma\tau - \phi_{\text{ent}}$  (Chevalier, Paige, and Kim, 2020; Guff, Boulle, and Pikovski, 2022; Schut, Geraci *et al.*, 2023). Thus, to have a negative expectation value of the witness, one has to keep the growth rate of the entangling phase above the decoherence rate. Here we provide for concreteness some general considerations about noise and decoherence for the figure of merit given in Eq. (102). The requirements on the force noise spectra  $S_{FF}$  can be estimated from the decoherence rate  $\Gamma$  as

$$\Gamma = \frac{S_{FF}(\omega_{\text{exp}})\Delta x^2}{\hbar^2}, \quad (108)$$

where  $\omega_{\text{exp}} = 1/\tau$  is the characteristic frequency of the experiment. We require  $\tau < \Gamma^{-1}$  to have sufficiently long coherence times (Bose *et al.*, 2017). Specific noise and decoherence sources have been considered, as have methods for its mitigation (Pedernales, Morley, and Plenio, 2020; van de Kamp *et al.*, 2020; Rijavec *et al.*, 2021; Toroš *et al.*, 2021; Weiss *et al.*, 2021; Gunnink *et al.*, 2022; Yi *et al.*, 2022; Fragolino *et al.*, 2023; Wu, Zeng, and Wu, 2023). It is perhaps important to emphasize that primary sources are the collisions with background gas, and blackbody radiation emission (Romero-Isart, 2011; van de Kamp *et al.*, 2020). There could also be electromagnetic noise of various forms (Fragolino *et al.*, 2023). Gravitational and inertial noise also plays an important role in the decoherence of large quantum superpositions, and some types of mitigation have been worked out (Toroš *et al.*, 2021). Various systematic noises affect Stern-Gerlach interferometry when it is the mechanism of creating large spatial superpositions, such as those due to phonons (Henkel and Folman, 2022, 2023) and rotations (Japha and Folman, 2022). In this context the coherence of spins is also important, and achieving dynamical decoupling has also been considered (Wood, Bose, and Morley, 2022).

### 4. Implications

Considering that one observes the entanglement between two masses due to their gravitational interaction, what can we conclude from that? Essentially, it verifies one prediction of a fully quantum counterpart of Einstein's equations,

$$\hat{G}_{\mu\nu} = \frac{8\pi G}{c^4} \hat{T}_{\mu\nu}. \quad (109)$$

In this sense it verifies a prediction common to all approaches in which gravity is treated as a quantum field (Donoghue, 1994; Dienes, 1997; Rovelli, 1998). Alternatively, it falsifies all hybrid theories of quantum sources in a state  $|\psi\rangle_{\text{source}}$  leading to classical gravity  $G_{\mu\nu}$  [while Eq. (83) is a special case of this, there could be much more general stochastic theories of the aforementioned hybrid nature (Diósi and Halliwell, 1998; Kafri, Taylor, and Milburn, 2014; Galley, Giacomini, and Selby, 2023; Oppenheim, 2023)]. We present the following arguments in favor of the view that observation of entanglement is inconsistent with gravity being a classical field or curvature even when defined in the aforementioned, highly general sense. We should at once state that such a conclusion is possible if one makes (i) an appropriate (standard) definition of a classical field and (ii) a minimal assumption.

We first define what a classical field is. It is an entity with a probability distribution over fixed values (numbers) at every point in spacetime. Thus, retaining the symbol usually used for the gravitational metric, we would define a classical gravitational field as an entity defined by probabilities  $P^{(i)}$  and corresponding metrics  $g_{\mu\nu}^{(i)}$ ,

$$\{P^{(j)}, g_{\mu\nu}^{(j)}(\mathbf{r}, t)\}, \quad (110)$$

where  $\mu\nu = 0, \dots, 4$  and  $\mathbf{r}, t$  are spacetime points. This definition is broader than just having a unique metric  $g_{\mu\nu}(\mathbf{r}, t)$  defined everywhere in spacetime, as we are allowing for probabilities. The allowance for probabilities makes it possible for gravity to be a statistical field while still being classical. Quantum is more stringent, as it necessitates quantum superpositions of different configurations  $g_{\mu\nu}(\mathbf{r}, t)$ . As long as we disallow superpositions, then even with fluctuations (probabilities) a field is classical. Now comes the assumption. This is the assumption that two masses outside each other's support (by *support*, we mean their positions or, if quantum, their wave functions or, if a second quantized matter field, then the localized mode that they occupy) can interact only with their local field and not directly with each other. This makes the field a *mediator*. Within the domain of nonrelativistic experiments that will be feasible in the foreseeable future, we cannot prove the necessity of the mediator, and we appeal to what is known from the rest of physics, namely, that there is no action at a distance in our known domain of physics.

Under assumptions (i) and (ii), the operations that can happen between the masses due to their interactions with their local gravitational field are local operations and classical communications (LOCC), which cannot create entanglement. Thus, it follows simply that if entanglement is observed between the masses due to their gravitational interaction, then either gravity is not a classical field as per definition (i) (i.e., it is nonclassical) or the assumption of a mediator (ii) is violated. This was the justification presented by [Bose \*et al.\* \(2017\)](#). Within this setting of exchange of a mediator between the masses, only a highly quantum mediator, namely, a virtual off-shell particle (a quantum superposition of all energies) is necessary for the continuously coherent generation of entanglement, as was shown through a fully relativistic treatment by [Marshman, Mazumdar, and Bose \(2020\)](#); for a treatment that also shows the retardation in the growth of entanglement, see [Christodoulou \*et al.\* \(2023\)](#). Alternatively, it has been shown that the presence of entanglement also necessitates an operator-valued interaction between masses, which is not possible with a classical mediator ([Bose \*et al.\*, \(2022\)](#)).

Another way to interpret the results of the experiment is that it evidences a quantum superposition of geometries that one of the masses produces on which the other mass evolves ([Christodoulou and Rovelli, 2019](#)). If the quantum superposition of geometries (i.e., quantum-natured gravity) is disallowed, then no superposition develops. The conditions for justification of a nonclassical nature of gravity within the framework of generalized probability theories have been presented ([Galley, Giacomini, and Selby, 2022](#)). Within the effective quantum field theory description of gravity, it has been argued that once the Newtonian interaction enables entanglement, the other degrees of freedom have to be quantized for consistency ([Belenchia \*et al.\*, 2018](#); [Carney, 2022](#); [Danielson, Satishchandran, and Wald, 2022](#)), along with an expression for a quantum state of gravity associated with a mass in a quantum superposition ([Chen, Giacomini, and Rovelli, 2023](#)). Indeed, if the condition of mediator,

providing the local part of the LOCC, is not imposed, one can still draw conclusions from the generation of entanglement, as discussed by [Fragkos, Kopp, and Pikovski \(2022\)](#), while the modifications needed to draw conclusions about quantum-natured gravity even without assumptions were stated, after a relativistic treatment, by [Martín-Martínez and Rick Perche \(2023\)](#). We note here that LOCC is more restrictive than just disallowing entanglement generation ([Lami, Pedernales, and Plenio, 2023](#)).

#### D. Other tests of gravity

There are a number of ways in which gravity can affect quantum systems beyond the areas of precision gravimetry, decoherence, and entanglement. Here we detail additional tests of gravity and related effects.

##### 1. Tests of the generalized uncertainty principle

Many quantum-gravity theories predict the existence of a finite and minimum length scale at least as small as the Planck length  $l_P = \sqrt{\hbar G/c^3} \approx 1.6 \times 10^{-35}$  m ([Garay, 1995](#); [Hossenfelder, 2013](#)). The emergence of a finite length implies a generalized uncertainty principle (GUP) because the fundamental position uncertainty can no longer be reduced to zero. The widely considered GUP reads

$$\Delta x \Delta p = i\hbar \left[ 1 + \beta_0 \left( \frac{l_P \Delta p}{\hbar} \right)^2 \right], \quad (111)$$

where  $\Delta x$  and  $\Delta p$  denote the uncertainties in the operators and  $\beta$  is a dimensionless constant that indicates the strength of the modification. The minimal length arises because the uncertainty in  $\Delta x$  can no longer be made infinitesimally small. Also associated with the GUP is the modified commutator relation

$$[\hat{x}, \hat{p}] = i\hbar \left[ 1 + \beta_0 \left( \frac{l_P \hat{p}}{\hbar} \right)^2 \right]. \quad (112)$$

Bounding the parameter  $\beta$  in Eq. (111) through experiments also bounds new physics below the length scale  $\sqrt{\beta} l_P$  ([Das and Vagenas, 2008](#)).

The existence of a finite length scale and GUP was first put forward in string theory ([Veneziano, 1986](#); [Amati, Ciafaloni, and Veneziano, 1989](#)) but were later also derived using general mode-independent properties of quantum-gravity theories. For example, a generalized gedanken experiment for the measurement of the area of the apparent horizon of a black hole in quantum gravity leads to the emergence of a GUP ([Maggiore, 1993a](#)). There is also an algebra that gives rise to the modified commutator relation in Eq. (111), just as the operators  $\hat{x}$  and  $\hat{p}$  satisfy  $[\hat{x}, \hat{p}] = i\hbar$  ([Maggiore, 1993b](#)). Model-independent arguments for the measurement of micro-black holes allow us to arrive at a GUP ([Scardigli, 1999](#)). The influence of minimal length scales on quantum states has been widely considered. There are quantum-mechanical implications of a GUP and finite length, which were analyzed by [Kempf, Mangano, and Mann \(1995\)](#), including the localization of wave functions in space and the effects on harmonic oscillators. Harmonic oscillators with



minimal length scales were considered by Chang *et al.* (2002), who also considered the effects on electrons trapped in magnetic fields, as well as by Lewis and Takeuchi (2011). In addition, an equivalent formulation of the GUP but with a maximum observable uncertainty in the momentum, rather than a minimum uncertainty in the position, has been formulated (Petruzziello, 2021).

A number of proposals for laboratory experiments to test GUPs with massive quantum systems have been put forward. Using pulsed optomechanics (see Sec. III.A.1), Pikovski *et al.* (2012) showed that the effects of a GUP should create changes in the trajectories in phase space traced out by a massive system. The scheme was later extended by Kumar and Plenio (2018). Along similar lines, mechanical oscillators near the Planck mass ( $m_P \approx 22$  ng) were analyzed where the modified dynamics was directly compared to the unmodified dynamics (Bawaj *et al.*, 2015). A further proposal considered a pendulum where continuous rf measurements of the frequency of an electromechanical oscillator can help to further bound  $\beta_0$  (Bushev *et al.*, 2019). The radiation-pressure noise can also contain information about the GUP, as proposed by Girdhar and Doherty (2020). The sensitivity to the modified commutator relation was shown to improve in the vicinity of exceptional points (Cui *et al.*, 2021), and quadratic corrections were shown to affect the noise spectrum of an optomechanical system (Sen, Bhattacharyya, and Gangopadhyay, 2022). A recent result is the evaluation of improved constraints on minimum length models using a low-loss phonon cavity (Campbell *et al.*, 2023).

Note that additional considerations suggest that the observed effects from a GUP scale with  $N^{-a}$ , where  $N$  is the number of particles of the composite system and  $a$  is a parameter to be determined (Kumar and Plenio, 2020). The strength of this scaling is unclear but should be taken into account in experiments. It has therefore been proposed that bounds on GUPs can also be obtained through the use of atoms (Chatterjee, Gangopadhyay, and Majumdar, 2021). In another work it was pointed out that different modifications of the canonical commutator yield the same commutator relation in Eq. (112) (Bishop, Lee, and Singleton, 2020), which necessitates the need for caution when interpreting experimental results.

The current leading bound for  $\beta_0$  appearing in Eq. (111) is  $\beta_0 < 5.2 \times 10^6$ , which was calculated using pendulum measurements (Bushev *et al.*, 2019). Including the dependency on the number of constituent particles as discussed would, however, change the bound for  $\beta_0$  as well as its physical interpretation within the considered GUP model (Kumar and Plenio, 2018). Other bounds on  $\beta_0$  have been derived from astronomy (Scardigli and Casadio, 2015) as well as from gravitational waves (Das *et al.*, 2021). See Scardigli and Casadio (2015) for a comparison between bounds obtained from different experiments available at the time.

## 2. Tests of the gravitational Aharonov-Bohm effect

The Aharonov-Bohm effect was originally introduced for electrons in a constant magnetic field, which picks up a phase depending on the spatially dependent vector potential that can be measured in an interferometer. Fundamentally, the phase

difference stems from an action difference between the interferometer arms that would not be accessible classically. Similarly, a gravitational field can induce such action differences (even in the absence of forces), giving rise to a scalar gravitational Aharonov-Bohm effect (Audretsch and Lammerzahl, 1983). Ultracold atoms were proposed as an experimental platform to detect this effect (Hohensee *et al.*, 2012) due to their long coherence times. The gravitational Aharonov-Bohm effect was successfully detected with a light-pulse  $^{87}\text{Rb}$  atom interferometer and a kilogram-scale source mass (Overstreet *et al.*, 2022) that allowed spacetime curvature to be directly probed (Roura, 2022). Further proposals include tests using quantum systems in free fall (Chiao *et al.*, 2023), as well as tests of the vector gravitational Aharonov-Bohm effect (Chiao *et al.*, 2014).

## 3. Tests of gravity through resonances and phonon excitations

The first attempts to detect gravitational waves involved resonant mass antennas (Misner, Thorne, and Wheeler, 1973). Weber searched for gravitational waves using resonances with phononic modes of aluminum bars now known as Weber bars (Weber, 1969; Ferrari *et al.*, 1982). In the experiments Weber's bars reached temperatures of a few kelvins, which is not cold enough to suppress noise sufficiently or to reach the quantum regime. Proposals to use a kilogram-scale superfluid  $^4\text{He}$  resonator promise better sensitivities by reaching millikelvin temperatures (De Lorenzo and Schwab, 2014; Singh *et al.*, 2017).

A quantum version of a resonant antenna has been proposed using resonances of phonon modes in BECs that can reach nanokelvin or picokelvin temperatures (Sabín *et al.*, 2014; Kohlrus *et al.*, 2017). At these temperatures it is possible to prepare highly sensitive quantum states that could be used to detect long-lived high-frequency gravitational waves (between  $10^3$  and  $10^7$  Hz). The quantum thermodynamical properties of resonances in relativistic quantum fields were studied by Bruschi, Morris, and Fuentes (2020), showing that the BEC phonon resonance antenna is a quantum thermal machine capable of extracting energy from the gravitational wave. Resonant effects with BEC phonons can be used to detect gravitational accelerations and gradients in the Newtonian approximation (Rätzel *et al.*, 2018), including the search for modifications of Newtonian dynamics (MOND) (Fernandez-Melendez *et al.*, 2024) and also to measure relativistic corrections (Ahmadi *et al.*, 2014; Lock and Fuentes, 2017; Howl and Fuentes, 2023).

Recent proposals consider searching for signatures of quantum gravity using multiatomic states of cold atoms (see also Sec. IV.C) (Haine, 2021), BEC phonons (Howl *et al.*, 2021), and massive quantum acoustic resonators (Tobar *et al.*, 2023; Tobar, Pikovski, and Tobar, 2024). These proposals aim at detecting single gravitons through their direct interaction with matter, rather than through decoherence. A proposal to detect quantum gravity using Weber bars cooled down to sub-mK temperatures (Aguilar, 2011) suggested searching for quantum-gravity signatures by testing modifications of the energy-momentum uncertainty principle (Bhattacharyya, Gangopadhyay, and Saha, 2020). Signals detected using these new detector concepts could be

correlated with independent classical detections of gravitational waves by laser interferometry to ascertain their origin. There are also new proposals to use laser interferometers to search for signatures of the quantum nature of gravitational waves (Parikh, Wilczek, and Zahariade, 2020, 2021a, 2021b) and looking for geontropic vacuum fluctuations from quantum gravity (Verlinde and Zurek, 2021; Bub *et al.*, 2023). Detecting gravitational waves and gravitons using tabletop experiments that make use of the high sensitivity of quantum technologies promise to open a new area in the study of fundamental physics.

#### 4. Tests of quantum field theory in curved spacetime and analog gravity

Quantum field theory in curved spacetime predicts that spacetime dynamics produces entangled excitations in quantum fields (Ball, Fuentes-Schuller, and Schuller, 2006; Fuentes *et al.*, 2010) and that the presence of horizons gives rise to decoherence (Fuentes-Schuller and Mann, 2005; Adesso and Fuentes, 2009; Alsing and Fuentes, 2012); see Sec. IV.D.4. Underpinning these effects is parametric amplification, where particles are created out of the quantum vacuum by moving boundary conditions or horizons. In this sense there is a deep connection between the dynamical Casimir effect where, by changing the length of a cavity, the vacuum state of the electromagnetic field changes, thereby producing entangled particles (Moore, 1970; Fulling and Davies, 1976; Brusch, Fuentes, and Louko, 2012); parametric down-conversion where a medium change produces entangled photons (Kwiat *et al.*, 1995); and effects of quantum field theory such as Hawking radiation (Hawking, 1974) and the creation of particles by the expansion of the Universe (Birrell and Davies, 1982; Polarski and Starobinsky, 1996), among other effects. This connection is made evident through the mathematical formalism of both quantum optics and quantum field theory in curved spacetime, where Bogoliubov transformations produce mode mixing and two- and single-mode squeezing of modes. Friis *et al.* (2013) applied the formalism of continuous variable quantum information to quantum field theory in curved spacetime to compute entanglement in relativistic settings. Quantum field theory has been demonstrated numerous times in the flat case; however, key predictions of the theory in the presence of gravity are currently out of experimental reach. Systems such as black holes are not accessible to experimentation, and most predicted effects are too small. Consider the dynamical Casimir effect, where producing excitations via oscillating mirrors requires velocities close to the speed of light. Oscillating a microwave mirror at a frequency of 2 GHz with a displacement of 1 nm produces velocities of only  $v \approx 10^{-7}c$ . At these velocities, approximately one photon is produced per day. However, moving the mirror at these speeds requires an input of mechanical power of 100 MW, and at the same time a temperature of  $\approx 20$  mK is needed to ensure that the field is in the vacuum state. For this reason it has become fashionable to simulate effects in the lab. Photon creation by a moving boundary condition was demonstrated using a superconducting circuit where the electromagnetic flux going through a superconducting quantum interference device produced a

boundary condition moving at a third of the speed of light (Wilson *et al.*, 2011).

Analog experiments help test consistencies within a given mathematical model. An alternative that involves using a massive system in a small-scale lab is to test the key predictions of the theory using BECs. Atomic interactions in a BEC produce phonons, which are a massless quantum field that obey a Klein-Gordon equation in an effective curved metric (Fagnocchi *et al.*, 2010; Visser and Molina-París, 2010; Sabín *et al.*, 2014; Hartley *et al.*, 2019a),

$$\square\psi = \frac{1}{\sqrt{-G}}\partial_\mu(\sqrt{-G}G^{\mu\nu}\partial_\nu\psi), \quad (113)$$

where  $G := \det(G)$  is the determinant of the effective metric given by

$$G^{\mu\nu} = \rho \frac{c}{c_s} \left[ g_{\mu\nu} + \left(1 - \frac{c_s^2}{c^2}\right) \frac{u_\mu u_\nu}{|u_\mu||u_\nu|} \right]. \quad (114)$$

In Eq. (114)  $g_{\mu\nu}$  is the spacetime metric,  $\rho$  is the density of the condensate,  $c$  is the speed of light, and  $c_s$  is defined by

$$c_s^2 := \frac{c^2 c_0^2}{|u_\alpha u^\alpha| + c_0^2}, \quad (115)$$

where  $c_0^2 := \lambda \rho \hbar^2 / 2m^2$  and the four-vector  $u_\mu$  is the flow associated with the phase of the wave function of the BEC bulk. The speed of sound in the BEC is  $c_0$ , and the density  $\rho$  of the BEC may depend on space and time coordinates. By choosing or changing the density, the speed of sound, or the velocity field, it is possible to simulate certain spacetime metrics.

One metric that can be simulated this way is that of a black hole, which makes it possible to test Hawking's prediction for black-hole radiation (Hawking, 1974, 1975). By starting with the expression for an irrotational fluid  $\nabla \times \mathbf{v} = 0$ , where  $\mathbf{v}$  is the velocity of the fluid and the analog to the Schwarzschild metric is (Unruh, 1981)

$$ds^2 = \rho \left( (c^2 - v^2) dr^2 - \frac{1}{1 - v^2/c^2} dt^2 - r^2 d\Omega^2 \right), \quad (116)$$

where  $\rho$  is the density of the fluid,  $v$  (as before) is the velocity of the fluid, and  $c$  is the speed of the particles, also referred to as the speed of sound. The Hawking temperature is then given by

$$T_H = \frac{\hbar}{4\pi k_B c} \left[ \frac{d}{dx} (c^2 - v^2) \right]. \quad (117)$$

The  $T_H$  of a BEC should be about 1 nK (Unruh, 1981). A BEC black hole emitting Hawking radiation has been both theoretically simulated (Carusotto *et al.*, 2008) and experimentally realized (Lahav *et al.*, 2010; Steinhauer, 2014, 2022; Steinhauer *et al.*, 2022). Beyond BECs, an analog of Hawking radiation can also be observed in quantum optics (Philbin *et al.*, 2008) using the nonlinear Kerr effect that arises in certain dielectric media.

We note that computer and analog simulations on their own can neither falsify nor verify theories. As a result, current

analog gravity experiments cannot be said to test general relativity directly. However, experimental settings are reaching scales where the key predictions of quantum fields in curved spacetime are becoming testable in the laboratory. Theoretical studies have shown that actual changes of the spacetime metric  $g_{\mu\nu}$  can in principle produce observable effects on phonon states (Sabín *et al.*, 2014). This effect is at the heart of the proposals to detect persistent high-frequency gravitational waves (Sabín *et al.*, 2014) and search for dark matter and dark energy (Howl and Fuentes, 2023), opening the possibility of testing the predictions of quantum field theory in curved spacetime in the lab. Some aspects of the interplay of quantum fields and general relativity could be tested in space-based experiments where photon entanglement has been distributed across thousands of kilometers; for a review see Sidhu *et al.* (2021). At these scales relativity commences since the proper time on Earth is different from the proper time on a satellite. Theoretical studies have shown that spacetime curvature affects the propagation of light wave packets on Earth, in the process affecting quantum communications (Bruschi, Ralph *et al.*, 2014). This effect goes beyond the gravitational phase shift predicted by special relativity. The curvature of the spacetime around Earth can flatten traveling wave packets and decohere quantum states, and these effects can be used to measure spacetime parameters (Bruschi, Datta *et al.*, 2014; Kohlrus *et al.*, 2017; Kohlrus, Bruschi, and Fuentes, 2019).

## V. EXPERIMENTAL PATHWAYS TOWARD TESTS OF GRAVITY

In Sec. IV we reviewed proposals for testing the overlap between quantum mechanics and gravity with massive

quantum systems. Here we review experimental advances toward the regime where the dynamics of quantum systems are affected by gravity.

To showcase the diversity of systems available, we provide a graphical overview of different platforms in Fig. 5. To further demonstrate advances that have been made in terms of controlling massive systems in the laboratory, we plot the masses versus the phonon numbers achieved for a mechanical oscillator in Fig. 6. The symbols there represent the type of system, and the colors indicate the date of publication. For comparison, the largest Bose-Einstein condensates (which are not included in the plot) that have been created thus far contain around  $10^{10}$  atoms (van der Stam *et al.*, 2007), which have a total mass of  $4 \times 10^{-16}$  kg in the case of sodium. We start by reviewing the state-of-the-art experimental tests of gravity today (Sec. V.A). Then we provide an overview of key methods for controlling massive quantum systems in the laboratory, including the preparation of squeezed states, spatial superpositions, and entangled states (Sec. V.B).

### A. State of the art of experimental tests of gravitation with massive systems

Here we review experiments that have made headway toward testing aspects of gravity, such as precision force sensing. In Fig. 7 we plot the force sensitivities that have been achieved to date against the masses of the probe systems. We note that it is not always clear whether the values reported can be compared directly, as we do here. In some cases, such as that of Hofer *et al.* (2023), the values plotted are based on predictions for the ideal experiment. Consult the works in




FIG. 5. Illustration of state-of-the-art mechanical systems. Encapsulated is an array of resonators utilized in experimental efforts aimed at detecting the interplay between quantum mechanics and gravity. The leftmost column showcases mechanical resonators: a 2D clamped resonator (graphene), a 1D clamped resonator (a suspended carbon nanotube), and a 1D singly clamped beam resonator, which together illustrate the diversity in mechanical systems. The central column depicts optical levitation systems: a stand-alone optical levitation and a hybrid Paul-optical levitation system that demonstrate the integration of optical techniques. The rightmost column, which presents an electrical resonator and a magnetic levitation system, represents the incorporation of electromagnetic methodologies. Together these systems exemplify the wide range of experimental apparatuses employed in the quest to uncover the interplay between quantum mechanics and gravity.






FIG. 6. Masses and phonons of massive quantum systems. The masses of systems controlled in the laboratory are plotted against the mechanical occupation number (phonons). Colors indicate the year of publication, while symbols indicate the type of system. Where phonon occupancy numbers were not explicitly stated in the original publication, we have estimated them using  $N = k_B T / (\hbar \omega_m)$ , where  $N$  is the number of thermal phonons,  $k_B$  is Boltzmann's constant,  $T$  is the temperature, and  $\omega_m$  is the reported mechanical frequency. We classify systems as hybrid whenever a qubit or a similar system is used to interface with the mechanical element. The data points correspond to the following parenthetical references: JILA '08 (Teufel *et al.*, 2008), Vienna '09 (Gröblacher, Hertzberg *et al.*, 2009), MPQ '09 (Schliesser *et al.*, 2009), UCSB '10 (O'Connell *et al.*, 2010), EPFL '11 (Riviere *et al.*, 2011), NIST '11 (Teufel *et al.*, 2011), Caltech '11 (Chan *et al.*, 2011), NBI '18 (Rossi *et al.*, 2018), Delft '19 (Guo, Norte, and Gröblacher, 2019), NéeL '21 (Cattiaux *et al.*, 2021), NBI '22 (Seis *et al.*, 2022), EPFL '23 (Youssefi *et al.*, 2023), ICFO '18 (De Bonis *et al.*, 2018), ICFO '14 (Moser *et al.*, 2014), MIT '11 (Schleier-Smith *et al.*, 2011), Soton '17 (Vovrosh *et al.*, 2017), ETH '19 (Windey *et al.*, 2019), ETH '20 (Tebbenjohanns *et al.*, 2020), Vienna '20 (Delić *et al.*, 2020), Tokyo '21 (Kamba *et al.*, 2021), ETH '21 (Tebbenjohanns *et al.*, 2021), Vienna '21 (Magrini *et al.*, 2021), Tokyo '22 (Kamba, Shimizu, and Aikawa, 2022), Florence '22 (Ranfagni *et al.*, 2022), ETH '23 (Piotrowski *et al.*, 2023), UCL '23 (Pontin *et al.*, 2023), and LIGO '21 (Whittle *et al.*, 2021).

question, which are cited in the caption of Fig. 7, for further information.

### 1. Tests with atoms and molecules

Since its early demonstrations and pioneering work of Kasevich and Chu (1991), atom interferometry employing laser-cooled cold atoms (Cronin, Schmiedmayer, and Pritchard, 2009) has been established as a precision technique for sensing (Peters, Chung, and Chu, 1999, 2001; Tino, 2021), with the realization of sensitive gravimeters, gravity gradiometers, and gyroscopes. In addition to measuring the gravitational acceleration due to Earth  $g$  with parts-per-billion precision, atom interferometers have been used to measure the Newton constant  $G_N$  (Lamporesi *et al.*, 2007; Sorrentino *et al.*, 2010; Rosi *et al.*, 2014) at the 150 parts-per-million level and are promising for improving tests of the gravitational inverse square law at laboratory scales (Tino, 2021). Several theories, as described in Sec. IV.A.4, predict modifications of the Newtonian inverse square law with a Yukawa-type deviation below the millimeter length scale, such as that described in Eq. (73). As one particularly well-suited class of modified gravity theories, interferometry with cold atoms is ideal to study Chameleon forces (Jaffe *et al.*, 2017; Sabulsky

*et al.*, 2019) due to the screening effect present in larger-scale test masses. Atom interferometry has also been used to test the Einstein equivalence principle (Schlippert *et al.*, 2014) at the parts-per-trillion level (Asenbaum *et al.*, 2020) and has been proposed as a technique for precision tests of general relativity (Dimopoulos *et al.*, 2007). In addition, atom interferometry has been identified as a promising method to search for gravitational waves in the midband ( $\sim 1$  Hz) (Canuel *et al.*, 2018; Badurina *et al.*, 2020; Abe *et al.*, 2021) between the sensitivity bands of the ground-based interferometer detectors (Abbott *et al.*, 2016a) and LISA (Seoane *et al.*, 2013; Bellei *et al.*, 2017). Recent work has enabled the detection of the gravitational analog of the Aharonov-Bohm effect in precision atom interferometry (Overstreet *et al.*, 2022, 2023).

The sensitivity of atom interferometers as gravimeters are limited by the interrogation time  $T$ , which for free-fall interferometers scales as  $\delta_\phi = k_{\text{eff}} g T^2$ , limiting Earth-based experiments to times of the order of 1 s for a 10 m drop path. Here  $g$  is Earth's gravitational acceleration and  $k_{\text{eff}}$  is the effective wave vector of the momentum transfer in the beam-splitter pulse of a light-pulse atom interferometer. Large momentum transfer beam splitters (Rudolph *et al.*, 2020; Kirsten-Siemß *et al.*, 2023) are a pathway for improved




FIG. 7. Force sensitivities achieved with massive probe systems. BECs, blue; hybrid systems, pink; torsion balances, green; levitated systems, purple; and clamped systems, orange. Plus signs indicate that the system contains less than one thermal phonon. We classify systems as hybrid whenever a qubit or similar system is used to interface with the mechanical element. The data points correspond to the following parenthetical references: Canberra '16 (Hardman *et al.*, 2016), Boulder '03 (Long *et al.*, 2003), Stanford '03 (Chiaverini *et al.*, 2003), Stanford '08 (Geraci *et al.*, 2008), ICL '18 (Pike *et al.*, 2018), ICFO '18 (De Bonis *et al.*, 2018), ICFO '14 (Moser *et al.*, 2014), ICFO '18 (Tavernarakis *et al.*, 2018), UC '99 (Goodkind, 1999), Maryland '02 (Moody, Paik, and Canavan, 2002), Reno '16 (Ranjit *et al.*, 2016), Soton '17 (Hempston *et al.*, 2017), Yale '17 (Monteiro *et al.*, 2017), Yale '20 (Monteiro *et al.*, 2020), UCL '20 (Pontin *et al.*, 2020), Montana '21 (Lewandowski *et al.*, 2021), Vienna '21 (Magrini *et al.*, 2021), Stanford '21 (Blakemore *et al.*, 2021), Tokyo '22 (Kamba, Shimizu, and Aikawa, 2022), Leiden '23 (Fuchs *et al.*, 2024), Vienna '23 (Hofer *et al.*, 2023), ESA '16 (Armano *et al.*, 2016), MICROSCOPE '17 (Touboul *et al.*, 2017), Seattle '12 (Wagner *et al.*, 2012), Seattle '15 (Terrano *et al.*, 2015), Vienna '21 (Westphal *et al.*, 2021), Seattle '22 (Shaw *et al.*, 2022), and ISTA '24 (Agafonova *et al.*, 2024).

sensitivity when they are limited by interrogation time constraints. Recent work has demonstrated momentum transfers of  $102\hbar k$  in  $^{87}\text{Rb}$  (Chiu *et al.*, 2011),  $112\hbar k$  in  $^{174}\text{Yb}$  (Plotkin-Swing *et al.*, 2018), and  $141\hbar k$  in  $^{88}\text{Sr}$  (Rudolph *et al.*, 2020). Space-based approaches may permit significantly longer interrogation times and, alternatively, atom interferometry with atoms trapped in a lattice can extend interrogation times up to 20 s (Xu *et al.*, 2019), with recent work surpassing a minute (Panda *et al.*, 2023).

Atom interferometry has also been performed with ultracold atoms cooled to quantum degeneracy, with both bosonic (van Zoest *et al.*, 2010; Müntinga *et al.*, 2013; Kovachy *et al.*, 2015) and fermionic atomic species (Roati *et al.*, 2004). BECs

of ultracold atoms are a versatile platform that can be used for a variety of precision quantum sensing applications, where their slow wave packet expansion and coherence play an important role. Interferometry with Bose-Einstein condensed atoms has been proposed as a method to search for short-range deviations from Newtonian gravity (Dimopoulos and Geraci, 2003). Bloch oscillations of bosonic Sr atoms have been considered as a method to test short-range gravitational forces (Ferrari *et al.*, 2006). The largest condensates of ultracold atoms have realized atom numbers as large as  $(2-12) \times 10^7$  (Streed *et al.*, 2006; van der Stam *et al.*, 2007), with reports of atom-interferometry with  $5 \times 10^6$  atoms (Hardman *et al.*, 2016). Atom chip-based atomic interferometry experiments with Bose-Einstein condensates have been performed using as many as  $4 \times 10^5$  atoms (Jo *et al.*, 2007). Key advantages when compared with mechanical oscillators are the environmental decoupling and quantum coherence as well as a mature toolbox for quantum state preparation and measurement. BECs and atom interferometers can in principle achieve atom shot-noise-limited sensitivity, with a phase resolution scaling as  $\delta\phi \sim 1/\sqrt{N}$ . Interferometry with phase resolution at the Heisenberg limit  $\delta\phi \sim 1/N$  is also possible with the aid of squeezed and highly entangled states (Szigeti, Hosten, and Haine, 2021). A challenge has to do with the fragility of such highly entangled states due to environmental perturbations such as background gas collisions.

Another modality of sensing with BECs involves observing their center-of-mass oscillations or collective modes. For example, the center-of-mass oscillation of BECs has been used to measure Casimir-Polder surfaces between the condensate and a nearby surface (Harber *et al.*, 2005). The dynamical response of the phonons of BECs has been predicted to be highly sensitive to acceleration due to the gravitational attraction of nearby masses, with sensitivities to oscillating masses at the 100 mg scale at millimeter separations (Rätzel *et al.*, 2018). Experiments employing a BEC in a double well are useful for a variety of fundamental physics tests and could have some advantages when compared to methods using solids (Howl, Penrose, and Fuentes, 2019). BECs can be cooled down to the picokelvin regime, lowering some sources of noise. Another advantage is that atoms are free to tunnel between wells, and states such as two-mode squeezed states involving atom superpositions between the two wells can be prepared (Esteve *et al.*, 2008). However, particularly challenging is to prepare Schrödinger cat or NOON states. These states are highly sensitive to decoherence. The most limiting source of noise is three-body recombination (Tolra *et al.*, 2004).

In addition, atom interferometry with atoms trapped in an optical lattice has been suggested as a possible route toward observing quantum entanglement induced from the gravitational interaction with a mechanical oscillator (Carney, Müller, and Taylor, 2021), albeit with additional assumptions (Hosten, 2022; Ma *et al.*, 2022; Streltsov, Pedernales, and Plenio, 2022).

Finally, matter-wave interferometers for complex molecules formed of many atoms are also promising for tests of gravity, as they benefit from the increase in mass in superposition (Fein *et al.*, 2019). One example is shown in Figs. 8(a) and 8(b), where interferometry measurements assess




FIG. 8. Experimental setup and interferometry measurement. (a) Setup for measuring cesium atoms' acceleration toward a 0.19 kg tungsten cylinder in ultrahigh vacuum. The cylinder is 2.54 cm in height and diameter, with a 0.5 cm axial through-hole and slot width. Differential measurements are used to study the interactions with the mass. (b) A Mach-Zehnder interferometer using Raman transitions in an optical cavity. Three laser pulses manipulate cesium atoms, splitting, reflecting, and recombining them to measure phase differences, providing an ensemble-averaged acceleration over 110 ms across approximately 100 000 atoms. Adapted from Jaffe *et al.*, 2017. (c) Schematic of a Cavendish torsion balance with dumbbell masses ( $M$ ), external forces ( $F$ ), and resulting rotation angle ( $\theta$ ). (d) Torsion pendulum as a gravitational acceleration transducer, containing two gold spheres (1 mm radius, 40 mm apart) on a glass capillary. One sphere serves as a test mass (90.7 mg) and the other serves as a counterbalance (91.5 mg). A silica fiber with a diameter of 4  $\mu\text{m}$  supports the pendulum with a 3.6 mHz torsional resonance. The torsion angle is detected optically. The source mass (92.1 mg) is harmonically moved 3 mm at 12.7 mHz to enhance the gravitational signal. Electrostatic interference is reduced using Faraday shield and discharging techniques. Adapted from Westphal *et al.*, 2021.

the acceleration of cesium atoms toward a tungsten cylindrical mass in an ultrahigh vacuum environment. This setup, which was adapted from Jaffe *et al.* (2017), employs a Mach-Zehnder interferometer with Raman transitions to measure phase differences and determine acceleration. Additionally, the duration of free fall in Earth's gravity ultimately limits the mass of molecules in interferometry, while it is more difficult to pick up a COW-like phase in the typically near-field regime of operation of these large-mass interferometers; see Sec. II.A.1 for a discussion of COW phases on matter waves. Figure 9 illustrates the optical COW experiment conducted in space and demonstrates how a single-photon split by an unbalanced Mach-Zehnder interferometer and transmitted to a satellite reveals gravity-induced phase shifts through observed interference (Mohageg *et al.*, 2022). More details about molecule interferometry experiments are given in Sec. V.B.2.

## 2. Tests with neutrons

Neutrons have proven to be a powerful probe for testing our understanding of gravity. The equivalence principle, which states that all objects fall at the same rate in a gravitational field regardless of their composition or structure, can be scrutinized using neutron matter-wave interferometry (Colella, Overhauser, and Werner, 1975; Rauch *et al.*, 1975; Greenberger, 1983); see also Sec. II.A.1. In these experiments a neutron beam is split into two paths. These two paths then interfere upon recombination, a process that allows for precise determination of the relative gravitational potential experienced by the neutrons in the two paths. The equivalence principle is tested by measuring the phase shift in the interference pattern. This approach has made it possible to verify the equivalence principle to a high degree of precision (Lämmerzahl, 1996, 1998). One of the main advantages of




FIG. 9. Simplified diagram of the optical COW experiment in space. A single photon is split into two paths using an unbalanced Mach-Zehnder interferometer (MZI). This photon is then transmitted to a satellite with an identical MZI. The interference observed at the satellite shows the phase shift caused by gravity. Adapted from Mohageg *et al.*, 2022.



neutrons in these experiments over atoms is that neutrons are uncharged, meaning that they are free from the electromagnetic forces that can influence the movement of atoms. This allows the gravitational interactions to be studied with less interference from other forces, leading to a higher degree of precision in the results.

With the advancement in the field, ultracold neutrons (UCNs) later emerged as a robust tool for testing gravitational theories. The UCNs originated from the insights of Enrico Fermi (Fermi *et al.*, 1936), who recognized the potential of slow neutrons to interact coherently while scattering, thereby creating an effective interaction potential for neutrons passing through matter. This led to the concept of storing neutrons with low kinetic energies, which was initially predicted by Zeldovich (1959) and first realized experimentally by groups in Dubna, Russia (Lushikov *et al.*, 1969), and Munich, Germany (Steyerl, 1969). A significant breakthrough was made by Nesvizhevsky *et al.* (2002), who observed quantized states of matter under the influence of gravity using UCNs. Their work has further opened up possibilities for probing fundamental physics, such as the equivalence principle (Nesvizhevsky *et al.*, 2002). More generally, advancements in this field have led to UCNs becoming a robust tool for testing gravitational theories (Steyerl *et al.*, 2012; Ivanov, Wellenzohn, and Abele, 2021). UCNs cooled to nearly absolute zero can be stored for extended periods, enabling precise measurements of the gravitational behavior of neutrons. In recent years there have been advances in the production of UCNs, with Zimmer, Piegsa, and Ivanov (2011) having reported a world-best UCN density available for users, achieved with a new source based on the conversion of cold neutrons in superfluid helium. Experiments with UCNs aim to measure the gravitational free fall of neutrons with high accuracy, offering a platform to test general relativity and other theories like MOND (Famaey and McGaugh, 2012). UCN spectroscopy has been instrumental in constraining various theories and phenomena, including dark energy, chameleon fields (Jenke *et al.*, 2014), and new short-range forces (Kamiya *et al.*, 2015). In an experiment Haddock *et al.* (2018) deployed a pulsed neutron beam to probe Newton's law of universal gravitation on subnanometer scales. The results set a stringent upper bound on the magnitude of potential unaccounted-for forces, enhancing the foundation upon which we apprehend gravity. Moreover, promising theoretical outlooks are unveiling new paths of exploration, including measurements of the gravitational redshift of neutrons. This involves observing the change in energy of a neutron due to a change in gravitational potential, which establishes a promising technique for testing general relativity (Roura, 2022). Advancements are also foreseen in the precision of measuring the electric dipole moment of neutrons, with potential assistance from quantum sensors based on weak-value amplification (Altarev *et al.*, 1992; Knee *et al.*, 2013; Pendlebury *et al.*, 2015). Other projects, such as those of the *qBOUNCE* and *GRANIT* Collaborations, aim to expand the understanding of gravity at short distances by examining gravitationally bound quantum states of ultracold neutrons (Jenke *et al.*, 2009, 2019; Kreuz *et al.*, 2009). Studies such as these could impose stringent constraints on hypothetical fields and forces, further refining our

understanding of gravity and providing insights that might push beyond the boundaries of currently accepted theories. For a more detailed review of gravity measurements by neutrons, see Pokotilovski (2018).

### 3. Tests with torsion balances and clamped mechanical systems

Sensitive torsion balances are a powerful and proven method for studying exotic short-range gravity (Kapner *et al.*, 2007; Lee *et al.*, 2020), equivalence-principle violation involving ordinary and dark matter (Wagner *et al.*, 2012; Shaw *et al.*, 2022), and novel spin-dependent interactions (Terrano *et al.*, 2015), as well as measuring the Newton constant (Gundlach and Merkowitz, 2000). Torsion balances remain one of the most promising paths forward for these studies as their sensitivity continues to increase and the understanding of background noise and systematic errors from patch charges and other surface forces improves.

Current tests are often limited by environmental vibrations that can “kick” the pendulum, exciting its fundamental and spurious (swing, bounce, and wobble) modes (Wagner *et al.*, 2012). This is particularly true in short-range tests where patch charges couple to the spurious modes producing noise that dominates at small separations and limits the minimum attainable separation (Lee *et al.*, 2020). Time-varying environmental gravity gradients limit equivalence-principle tests. Both of these technical limiting factors could be addressed by the development of a suitable low-vibration underground facility.

Torsion-balance experiments [shown in Figs. 8(c) and 8(d)] have typically employed relatively large source masses, well beyond the scale envisioned for achieving a quantum superposition. Work toward employing sub-mm-scale source masses and similarly miniaturized torsion pendula is underway (Westphal *et al.*, 2021). In this work thus far the smallest source mass that has been used for a gravitational measurement is approximately on the millimeter scale. While far from the scale where macroscopic quantum superpositions have been imagined in interference experiments, this work represents a step in this direction to bridge the gap. These experiments also tend to operate at low frequencies and are limited by the same environmental perturbations and thus could benefit from similar future low-noise facilities.

At even smaller length scales, microcantilevers (Chiaverini *et al.*, 2003; Geraci *et al.*, 2008) and microfabricated torsion oscillators (Long *et al.*, 2003) have been used to obtain bounds on Yukawa-type deviations of the Newton inverse square law at distances ranging from a few microns to tens of microns. Cutting-edge nanofabrication technology is making it possible to routinely design advanced 2D and 1D clamped resonators with massive quality factors, for instance, suspended silicon nitride membranes and carbon nanotube resonators.

Clamped mechanical systems interfaced with superconducting qubits have emerged as a fertile ground for probing the interplay between quantum mechanics and macroscopic objects. Early groundbreaking experiments demonstrated the feasibility of reaching the quantum ground state of mechanical resonators using superconducting circuits (O'Connell *et al.*, 2010; Teufel *et al.*, 2011). Furthermore, laser cooling techniques have been adapted to cool nanomechanical oscillators




FIG. 10. LIGO laser interferometer setup. The interference of light in the two arms leads to the detection of extremely small displacements. Adapted from [Scharpf, 2017](#).

into their quantum ground state ([Chan \*et al.\*, 2011](#)). [Qiu \*et al.\* \(2020\)](#) recently reported laser cooling a clamped oscillator to an average occupation number as low as  $\langle n \rangle = 0.09$  phonons.

Building upon this lineage of research, [Youssefi, Kono, Chegnizadeh, and Kippenberg \(2022\)](#) introduced a hybrid quantum system consisting of a superconducting circuit seamlessly integrated with a micromechanical oscillator. Achieving a thermal decoherence rate of 20.5 Hz and a dephasing rate of 0.09 Hz, they enabled the free evolution of a squeezed mechanical state over milliseconds. We anticipate that such advances will enable exploration of elusive phenomena that arise from the interplay between quantum mechanics and general relativity.

Furthermore, LIGO-style experiments have also contributed significantly to the field. LIGO ([Abbott \*et al.\*, 2016b](#)), as shown in Fig. 10, has provided direct evidence for the existence of black holes and opened up a new avenue for exploring the nature of gravity.

In conclusion, tests with clamped systems, including torsion balances, have proven effective in studying gravity and fundamental physics. Challenges such as environmental vibrations and technical limitations have led to the development of low-vibration facilities and miniaturization efforts. Nanofabrication techniques have enabled advanced resonators, while hybrid quantum systems offer new avenues for investigating quantum physics and dark matter. Additionally, LIGO-style experiments have made groundbreaking contributions to our understanding of gravitational waves. Overall, these advances hold substantial potential to advance our understanding of gravitational phenomena.

#### 4. Tests with levitated mechanical systems

Levitated mechanical systems offer a platform to investigate the interplay between quantum mechanics and gravity in the low-energy nonrelativistic regime. Since levitated systems are highly isolated from their environment the center-of-mass motion can be close to an ideal harmonic oscillator persisting

at large  $Q$  factors ([Geraci, Papp, and Kitching, 2010](#)). This isolation, together with the (in principle) quantum-limited detection of the position and therefore motion of the mechanical system by light (or direct electrical or magnetic interactions), makes them exceptional for testing quantum effects in gravity ([Caves \*et al.\*, 1980](#); [Aspelmeyer, Kippenberg, and Marquardt, 2014](#)).

Recently, the study of levitated mechanical systems has gone beyond the usual limits set by the gravitational law that governs how objects attract each other ([Arndt and Hornberger, 2014](#)). [Moore and Geraci \(2021\)](#) and [Priel \*et al.\* \(2022\)](#) explored the finer points of gravity-related phenomena by probing two-particle interactions with levitated particles beyond established force laws based on precise force and acceleration measurements.

In further detail, levitated systems provide a platform to generate and coherently control quantum effects in their motion via ground-state cooling, measurement-based schemes, etc. ([Aspelmeyer, Kippenberg, and Marquardt, 2014](#)), and at the same time come with sufficient mass for directly testing gravity effects on experimentally accessible time and magnitude scales. Quantum experiments with gravitating particles of the Planck mass ( $m_P = \sqrt{\hbar c/G}$ ) become feasible ([Ulbricht, 2021](#); [Aspelmeyer, 2022](#)). In addition, the levitated system has a full set of only six mechanical modes of translation ( $x$ ,  $y$ , and  $z$ ) and rotation ( $\alpha$ ,  $\beta$ , and  $\gamma$ ) that are developed to be used as quantum probes of gravity in the linear and nonlinear regimes ([Bateman \*et al.\*, 2014](#); [Gosling \*et al.\*, 2024](#); [Kilian \*et al.\*, 2024](#)). Recent experiments achieved the simultaneous cooling of all those modes ([Kamba, Shimizu, and Aikawa, 2023](#); [Pontin \*et al.\*, 2023](#)), which opens the door for quantum state preparation ([Delić \*et al.\*, 2020](#)). Rotational states provide a unique setting with their intrinsic nonlinearities for quantum experiments ([Stickler \*et al.\*, 2018](#); [Schrinski, Stickler, and Hornberger, 2022](#)).

Besides the prospect of using levitated mechanical systems for force ([Winstone \*et al.\*, 2018](#)) and inertial sensing ([Teufel](#)

*etal.*, 2011), direct gravity probes are emerging. Gravity affects levitation directly but is negligible in small-mass particle optical levitation (Ashkin and Dziedzic, 1971; Rademacher, Gosling *et al.*, 2022), while it is relevant in larger-mass Paul ions (Paul, 1990). In Meissner-superconducting and diamagnetic traps,  $g$  influences the trapping position and has a clear effect (Cano *et al.*, 2008). Static gravity was measured with levitated optomechanics by turning off the trap (Frimmer *et al.*, 2017). In addition, a Meissner levitated magnet has been used in a two-mass gravity detection experiment and has measured gravity at the level of attonewton gravity (Fuchs *et al.*, 2024).

In optical levitation nanoparticles and microparticles trapped in an ultrahigh vacuum can be cooled down to their ground state of center-of-mass motion through radiation-pressure forces exerted by optical cavities (Libbrecht and Black, 2004; Barker and Shneider, 2010; Chang *et al.*, 2010; Romero-Isart *et al.*, 2010). This technique has been explored for over a decade to test short-range gravity forces (Geraci, Papp, and Kitching, 2010). By employing optically levitated systems where microspheres are trapped and cooled in a vacuum, it has been possible to probe and measure gravitational effects with unprecedented precision at the micrometer scale (Ranjit *et al.*, 2016). Nanoparticles with a cooled center-of-mass temperature can also serve as a source for matter-wave interferometry experiments (Bateman *et al.*, 2014), which could be used for measuring gravitational acceleration and probing gravity at the micron length scale (Geraci and Goldman, 2015).

Recent experiments (Timberlake *et al.*, 2021) employed levitation via the Meissner effect, where two magnets suspended in a levitated state perturb each other's motion to measure the gravitational attraction between them. These experiments demonstrated the practicality of measuring gravitational acceleration for small masses, showcasing the potential for future improvements in experimental setups. Additionally, the gravitational constant ( $G$ ) can be estimated from such measurements. Coupling to superconducting inductor-capacitor ( $LC$ ) resonators in cryogenic environments. Paul ion trapping provides a stable trap for tunable  $e/m$  ratios (Paul, 1990). The close technological heritage from atomic Paul trapping makes available a set of the center-of-mass motion state-preparation protocols and tools for the manipulation, cooling, and control of charged nanoparticles and microparticles motion via electrodynamical ion levitation (Leibfried *et al.*, 2003; Schneider *et al.*, 2010).

The LISA Pathfinder (LPF) mission, which was designed to detect gravitational waves in space, utilized electrostatic detection of freely falling masses on the level of kilograms (Armano *et al.*, 2016). LPF data, as well as those from Earth-bound gravitational-wave detectors, were employed to establish strong upper bounds on continuous spontaneous localization and Diósi-Penrose models and are space-based derivatives of ground-based optomechanical precision experiments, including gravitational-wave detectors such as LIGO, VIRGO, and GEO600, but also AURIGA (Carlesso *et al.*, 2016). LPF is similar to other space missions such as Gravity Probe B to test Lense-Thirring GR frame-dragging effects (Everitt *et al.*, 2011), and the satellite gradiometry missions GRACE and GOCE (Drinkwater *et al.*, 2003; Tapley *et al.*,

2004). The MICROSCOPE mission (Touboul *et al.*, 2017) aimed to test the weak version of the equivalence principle, the basic principle of Einstein's theory of general relativity. Test masses made of different materials but with equal inertial masses were used in this experiment. By monitoring the motion of these masses over an extended period, MICROSCOPE sought to detect any deviations from the principle. The results of MICROSCOPE provided strong evidence in support of the principle, thereby consolidating the predictions of general relativity.

In conclusion, levitated systems have emerged as a promising platform for investigating the interface between quantum mechanics and gravity. Recent advancements, such as Meissner-effect-based levitation and space-based experiments like LISA Pathfinder and MICROSCOPE, have expanded our capabilities to study fundamental physics principles. The precise measurements achievable in levitated systems, coupled with the microgravity environment of space, contribute to our understanding of gravity, general relativity, and the fundamental laws of physics. To this end the recent MAQRO proposal aims to explore levitated particle dynamics in space, which would open a pathway for matter-wave interference experiments with long interaction times by not falling under Earth's gravity (Belenchia *et al.*, 2022; Kaltenbaek *et al.*, 2022). Future advancements in levitated systems and their applications hold interesting prospects for furthering our knowledge of the quantum-gravity interface (Rademacher, Millen, and Li, 2020).

## 5. Approaches with hybrid systems

Last, there are hybrid mechanical systems (Kolkowitz *et al.*, 2012; Rogers *et al.*, 2014; Treutlein *et al.*, 2014) where systems other than optical fields or photons are coupled to mechanical motion. Hybrid mechanical systems are indeed powerful experiments building on the well-studied Jayens-Cummings Hamiltonian physics of a two-level system coupled to a continuous variable-quantum harmonic oscillator system, as discussed in Sec. III.A.2. Hybrid systems have been at the forefront of demonstration of quantum states of massive systems, while for relevance for probing gravity effects, many of the hybrid systems involve non-center-of-mass motional modes, such as vibration modes of membrane states (O'Connell *et al.*, 2010) or acoustic modes in superconducting qubits (Chu *et al.*, 2017). Hybrid systems are arguably the most established mechanical quantum systems, and the quantum aspects of hybrid systems are discussed more in Sec. V.B.2 as they are used for the generation of massive superpositions as well as for demonstration of quantum entanglement between two large-mass systems. Typically, in hybrid systems the mechanical modes are at high frequency (100 MHz to 10 GHz) which allows for cooling to the quantum ground state by cryogenics, usually in dilution-type refrigerators. Such ground-state cooling gives direct access to the quantum regime and for quantum state preparation and coherent control schemes. Siddiqi (2021) provided an example where this was done using feedback techniques based upon continuous weak measurement (for theory on measurement-based control, see Sec. III.C). These techniques were built on a record-high detection efficiency of a measurement of more



than 80%. A more detailed account of the physics of hybrid systems was given by Aspelmeyer, Kippenberg, and Marquardt (2014).

However, there are approaches for testing aspects of gravity by hybrid systems, including for gravitational-wave detection encompassing coupling atoms to an optomechanical cavity that influences the atom-cavity interaction (Camerer *et al.*, 2011), cooling the system by linking a superconducting qubit to a mechanical resonator for improved detection sensitivity (O’Connell *et al.*, 2010), and integrating quantum dots with mechanical resonators or optical cavities for enhanced detection (Bennett *et al.*, 2010; Yeo *et al.*, 2014). In addition, the use of solid-state systems with mechanical resonators, coupled with optical cavities, proves promising for gravitational-wave detection (Arcizet *et al.*, 2011; Kolkowitz *et al.*, 2012). These approaches leverage the unique properties of different components for higher sensitivity and precision in gravitational-wave detection. An example is the use of optomechanical systems that offer enhanced cooling by constructive quantum interference and suppressed heating by destructive interference, which is essential for precision control and quantum information processing (Chen *et al.*, 2015). Furthermore, modern hybrid systems allow for the exploration of the quantum-classical mechanics interface and demonstrate the potential for a paradigm shift from cryogenic to room temperature quantum experiments using hybrid nanoelectromechanical system resonators (Tavernarakis *et al.*, 2018). Such advancements reflect the rapidly evolving potential of hybrid systems in examining quantum physics at a macroscopic scale and as an avenue for quantum state generation in massive mechanical systems (Akram, Ghafoor, and Saif, 2015; Liu, Liu, Wang *et al.*, 2021).

In conclusion, hybrid systems are expected to facilitate new ways to inject quantum features into large-mass mechanical systems by coupling to qubit systems, and the first concrete steps have already been taken. Hybrid mechanical systems will play a key role when probing into gravity effects within the domain discussed in this review because of the maturity of quantum-controlling large-mass mechanical states.

## B. Controlling massive mechanical quantum systems in the laboratory

Here we review experimental achievements and theoretical proposals for large-mass mechanical systems in the quantum domain by sectioning them into three classics of exemplary quantum states: squeezing, superposition, and entanglement.

### 1. Squeezing and swapping of mechanics

As we are interested in creating quantum states of mechanics, we now discuss the squeezing of mechanical degrees of freedom. We do not, however, discuss the application of squeezed light to mechanical oscillators as it was, for instance, used to advance the gravitational-wave detectors VIRGO (Schnabel, 2017) and, eventually, LIGO (LIGO Scientific Collaboration, 2011; Aasi *et al.*, 2013).

*Squeezing in clamped optomechanics.* An early demonstration of directly squeezing the mechanical mode was given by

Rugar and Grütter (1991). In the demonstration control over the spring constant enabled parametric drive and thus amplified the mechanical motion of the oscillator. This approach allowed for noise suppression of  $-4.9$  dB in one quadrature; see Sec. III.D.2. Improvement in noise suppression has been theoretically proposed using both detuned parametric driving and continuous weak measurement of the mechanical oscillator (Szorkovszky *et al.*, 2011). Experimentally, noise suppression was demonstrated by weak measurement and achieved  $-6.2$  dB in one quadrature (Szorkovszky *et al.*, 2011). Pontin *et al.* (2014) used parametric feedback to stabilize one quadrature without affecting the other and achieved squeezing of  $-7.9$  dB. The realization of quantum squeezing of a quadrature below the zero-point fluctuations was achieved by Lecocq *et al.* (2015), Pirkkalainen, Damskäg *et al.* (2015), and Wollman *et al.* (2015).

Further quantum state protocols such as state transfer and swapping were also shown as multimode optomechanics systems began to be explored. These have a strong indication as quantum information protocols but demonstrate powerful state control capabilities with a myriad of techniques. For example, Weaver *et al.* (2017) demonstrated coherent state swapping between modes of two separate mechanical frequencies in the same cavity.

*Squeezing in levitated optomechanics.* Analogously, the novel capability to control the potential and thus the mechanical frequencies enabled squeezing via nonadiabatic pulses (Rashid *et al.*, 2016). A study by Militaru *et al.* (2022) revealed the signatures of squeezing on the scattered light generated by mechanics oscillators. Recently, squeezed states have been discussed for testing small effects such as those predicted by some form of quantum gravity (Belenchia *et al.*, 2016). Squeezing is an operation to affect the mechanical state: it is one crucial operation in the universal toolbox of Gaussian state preparation but is also discussed as a state-preparation step for achieving non-Gaussian states such as quantum superposition in a levitated mechanical system (Riera-Campen *et al.*, 2023), as well as for generating quantum entanglement between two mechanical systems (Cosco, Pedernales, and Plenio, 2021).

In conclusion, squeezing is one option for generating out-of-equilibrium states of a continuously variable system. Squeezing of large-mass mechanical systems has been experimentally demonstrated. Squeezing generates highly sensitive states exhibiting a peculiar quantum signature (Chowdhury *et al.*, 2020) and is used to control the effect of dynamical nonlinearities.

### 2. Spatial superpositions of mechanical systems

The goal of this section is to review the state-of-the-art mechanical quantum systems and how they approach the regime for testing the overlap between quantum mechanics and gravity. One key aspect of this endeavor is the generation of the prototypical quantum state: the spatial superposition state of sufficiently massive or macroscopic systems. An illustrative example is Feynman’s thought experiment, shown in Fig. 11, where a gravitational field source in a quantum superposition interacts with a test mass, leading to different scenarios depending on whether the field remains in a




FIG. 11. Feynman's thought experiment. (a) A gravitational field source (the blue balls) is placed in a quantum superposition and interacts with a test mass (the green ball). The system can develop into either of two scenarios: (b) If the field is in a quantum superposition, the test mass senses two different gravitational forces and also enters a superposition state. (c) If the field stays classical, the test mass senses a single gravitational force and moves accordingly. Adapted from Pedernales and Plenio, 2023.

superposition or collapses to a classical state (Pedernales and Plenio, 2023). Theoretical proposals for achieving this or similar goals, such as those given by Bose *et al.* (2017) and Hanif *et al.* (2023) were described in Sec. IV.C.

There are competing definitions of what *macroscopic* actually means; the definition strongly depends on what aspects of physics are to be tested. A single photon in a superposition or a pair of photons entangled over 1000 km is arguably a large quantum system, while there are various measures of what quantum coherence at macroscopic scales actually means (Leggett, 1980; Dür, Simon, and Cirac, 2002; Björk and Mana, 2004; Cavalcanti and Reid, 2006; Marquardt, Abel, and von Delft, 2008; Lee and Jeong, 2011). However, for the purpose of testing the overlap between quantum mechanics and gravity, it seems advisable here to choose a macroscopicity measure that includes a set of three parameters about the quantum system and is able to compare a manifold of different physical systems in an objective way, such as the macromasure based on matter-wave superposition put forward by Nimmrichter and Hornberger (2013). The measure  $\mu$  is a function of the mass of the system in a spatial superposition, the spatial size of the superposition, and the time for the spatial superposition to exist. Using  $\mu$ , it becomes evident how wide beam-splitting low-mass atomic fountains (Kovachy *et al.*, 2015), large-mass small zero-point motional optomechanical setups (LIGO) in continuously monitored low-phonon states (Whittle *et al.*, 2021), and levitated mechanical systems compare and why levitated mechanics with mesoscopic masses look most promising for delivering the most macroscopic of superpositions. See Table I for a summary of the values computed for the measure thus far. The current mass record in

matter-wave interferometry was reported by Fein *et al.* (2019) for complex molecules at 28 kDa, which achieves  $\mu > 14$ .

Different ways to generate spatial superpositions have been proposed for mechanical systems, both clamped and levitated, and some have already been experimentally demonstrated. The technical challenge, if formulated in matter-wave language, is to split the matter wave front coherently for a minuscule de Broglie wavelength. This typically requires preparing, by cooling, some sort of coherent initial state, and a subsequent application of a coherence beam-splitter operation. For spatial superpositions the prepared coherence length determines both the spatial resolution of the beam splitter and the extent of the final superposition.

Methods to realize beam splitters inspired by established technology from matter-wave interferometry with electrons, neutrons, atoms, and complex molecules (Cronin, Schmiedmayer, and Pritchard, 2009; Hornberger *et al.*, 2012; Juffmann, Ulbricht, and Arndt, 2013; Arndt and Hornberger, 2014; Millen and Stickler, 2020), the use of optical gratings (Bateman *et al.*, 2014; Geraci and Goldman, 2015), nonlinear interaction in a cavity (Bose, Jacobs, and Knight, 1997, 1999; Mancini, Man'ko, and Tombesi, 1997; Romero-Isart *et al.*, 2011), and levitated magnetomechanical oscillators coupled to magnetic fields (Romero-Isart *et al.*, 2012) are among the proposals, which include magnetic beam splitters using ferromagnetic particles (Rahman, 2019). For mechanical systems ideas also include measurement-based multiple-pulsed schemes addressing the position- and momentum-dependent continuous variables (Vanner *et al.*, 2011), as well as continuous weak measurement protocols (Rossi *et al.*, 2018) or advanced protocols from quantum metrology using dynamical model selection and classical and quantum

TABLE I. Macroscopicity measure summarized by [Schrinski \*et al.\* \(2023\)](#).

Experiment		Year	$\mu$
Mechanical resonators	Bulk acoustic waves ( <a href="#">Bild <i>et al.</i>, 2023</a> ; <a href="#">Schrinski <i>et al.</i>, 2023</a> )	2022	11.3
	Phononic-crystal resonator ( <a href="#">Wollack <i>et al.</i>, 2022</a> )	2022	$\sim 9.0^a$
	Surface acoustic waves ( <a href="#">Satzinger <i>et al.</i>, 2018</a> )	2018	$\sim 8.6^a$
Matter-wave interference	Molecule inteferometry ( <a href="#">Fein <i>et al.</i>, 2019</a> )	2019	14.0
	Atom interferometry ( <a href="#">Xu <i>et al.</i>, 2019</a> )	2019	11.8
	BEC interferometry ( <a href="#">Asenbaum <i>et al.</i>, 2017</a> )	2017	12.4

<sup>a</sup>Estimated by [Schrinski \*et al.\* \(2023\)](#).

hypothesis testing ([McMillen \*et al.\*, 2017](#); [Ralph \*et al.\*, 2018](#); [Schrinski \*et al.\*, 2019](#)), schemes that even go conceptually much beyond the classic scenario for generation of superposition states, as well as in evidencing the appearance of nonclassicalities. However, the same measure has to be consistently applied to rank macroscopicity.

While all of the these methods address external ( $x$  and  $p$ ) degrees of freedom for generating superpositions, there are also promising ideas for addressing internal states such as isolated electron and nuclear spin states. If the coherence of such states can be extended to long enough times and coupled to  $x$  and  $p$  in a coherent fashion, then elegant protocols for state preparation can be transferred from the rich toolbox of atomic two- and few-level physics. Such ideas have been put forward for harmonically bound systems ([Scala \*et al.\*, 2013](#)) as well as for free motion ([Wan \*et al.\*, 2016](#)). Again, the massive and freely evolving quantum state keeps its promise to become the most macroscopic quantum one.

Superpositions of different energy states of the harmonic oscillator using strong coupling in cavity-QED-like (Jaynes-Cummings Hamiltonian; see Sec. III.A.2) systems are a further option and have historically been the first demonstration for a quantum superposition of a massive microwave-driven quantum system ([Teufel \*et al.\*, 2011](#)), for surface acoustic-wave phonons ([Satzinger \*et al.\*, 2018](#)), and when using coupling to a superconducting qubit ([Wollack \*et al.\*, 2022](#)), while the spatial extent of the superposition is on the size of the amplitude of the zero-point motion in those systems, and advanced techniques have to be applied to reach the defined macroscopic. The energy state superposition is mapped onto a motional or vibrational state ([O'Connell \*et al.\*, 2010](#); [Bild \*et al.\*, 2023](#); [Schrinski \*et al.\*, 2023](#)). While large-mass hybrid mechanical systems can be prepared in quantum states ([Sletten \*et al.\*, 2019](#)), the macroscopicity of hybrid quantum superposition states has to be analyzed by a generalized measure such as the one shown in Table I. ([Bild \*et al.\*, 2023](#)) produced the largest spatial superposition state to date among hybrid systems for Planck mass acoustic modes. Hybrid mechanical systems are supreme quantum systems; their relevance for gravity measurements is discussed in Sec. V.A.5.

While the previous discussion is about linear motion, macroscopic superpositions can also be achieved by utilizing rotational mechanical degrees of freedom ([Carlesso \*et al.\*, 2017](#); [Stickler, Hornberger, and Kim, 2021](#)). These approaches are promising since the generated quantum state is potentially more protected from noise and decoherence. In addition, the technology for implementing angular

superpositions is different from those needed for linear superpositions and could be easier to realize.

For the realization of superposition experiments, aspects of decoherence such as those resulting from gas collisions and effects of blackbody photons have to be considered and understood ([Hornberger \*et al.\*, 2003](#); [Hackermüller \*et al.\*, 2004](#); [Romero-Isart, 2011](#)). Decoherence puts severe constraints on any attempt to realize macroscopic quantum states, and each experiment has to focus on those aspects. Theoretical studies of decoherence effects have been carried out using an open quantum system dynamical model on the level of master equations; see Sec. III.B.1 for a more detailed summary.

Another interesting area of research focuses on the superposition of massive electromechanical resonators, which carries significant implications for exploring the effects of general relativity ([Gely and Steele, 2021](#)). To investigate these effects effectively, it is crucial to ensure that the coherence time of the superposition state exceeds the timescale associated with general relativity. In pursuit of this goal, one approach involves integrating clamped mechanical oscillators, such as silicon nitride membranes, with superconducting circuits. This integration allows for the preparation of these resonators in small cat or Fock states, which enables experiments to probe the interplay between quantum mechanics and general relativity ([Albrecht, Retzker, and Plenio, 2014](#); [Liu, Mummery \*et al.\*, 2021](#)).

Furthermore, the coherent coupling of mechanical vibrations in carbon nanotube resonators, controlled by the electronic spin of a nitrogen vacancy, has emerged as an interesting avenue of research ([Qin \*et al.\*, 2019](#)). When these resonators are cooled to their quantum ground state, it becomes possible for the mechanical phonons within them to exhibit both wavelike and particlelike behavior, effectively manifesting the essence of quantum superposition. This line of inquiry not only pushes the boundaries of quantum mechanics but also highlights the potential applications of electromechanical resonators in quantum information processing and quantum metrology.

Additionally, electromechanical systems offer opportunities for experiments involving Paul traps, which can shed light on the interplay between quantum mechanics and general relativity. [Martinez \*et al.\* \(2020\)](#) proposed a Paul trap to trap and cool a single charged nanoparticle to its quantum ground state. Through the controlled application of laser beams and the analysis of the nanoparticle's evolution, they were able to demonstrate the sensitivity of the nanoparticle to gravity and place constraints on non-Newtonian gravitational interactions. Moreover, electrically levitated nanorotors, when coupled to a



superconducting qubit, enable ultrashort timescale interference experiments by achieving a quantum superposition state through controlled rotational and translational motion. These developments represent strong prospects for furthering our understanding of macroscopic quantum phenomena and their implications for our understanding of both quantum mechanics and gravity.

Based on considerations of thermal decoherence effects, a proposal for a superconducting and magnetic version of a superposition experiment has been made. The low-temperature and extreme-high-vacuum setting appear to be the most promising (Pino *et al.*, 2018). The extreme settings where the quantum system is—as much as possible—decoupled from its environments can be analyzed in terms of the duration of free evolution (spreading) of the wave function and demands extremely small amplitudes (typically smaller than the spatial size of the de Broglie wavelength) of all guiding fields and vibrations. This adds another serious technical demand to any experimental realization.

As dictated by Schrödinger dynamics, the free evolution time grows proportionally to the mass of the quantum system and is pushing realistic attempts to surpass the existing mass record well beyond some 100 ms superposition lifetime. All decoherence effects and noises have to be controlled to be smaller than the evolution amplitudes during that same time. Ideally, the wave function is left alone for some seconds, as is at the core of proposals for macroscopic quantum superposition on a dedicated satellite in space (Kaltenbaek *et al.*, 2022).

However, an alternative solution may come from boost or inflation operations, which accelerate the spread of the wave function significantly (Romero-Isart, 2017). The boost has to be coherent, so it does not spatially resolve the position of the particle during the boost. Boosts have been demonstrated by Stern-Gerlach beam splitters for atoms on a chip (Margalit *et al.*, 2021), and it seems possible to translate the same beam-splitting technique to the much more massive NV-defect center diamond nanoparticles and other spin systems.

Another severe experimental challenge is that mechanical experiments, which are usually single-particle experiments, have to be repeated many times to achieve particle number statistics to show unique quantum features (Neumeier *et al.*, 2022). All known measures of quantumness are statistical ensemble measures on the level of the density matrix rather than the wave function directly, as every run of a quantum experiment or operation has a completely random outcome and cannot be predicted by quantum mechanics. As there is no coherent ensemble of massive particles equivalent to, for instance, an atomic BEC, each large-mass single-particle experiment has to be repeated many times (say, at least 1000 times) under the exact same conditions. Nevertheless, many researchers have faced the challenge of working toward the first generation of a truly macroscopic quantum superposition, which will question our understanding of quantum mechanics as well as gravity and will hint at how the two important theories are connected fundamentally.

Romero-Sanchez *et al.* (2018) explored the realm of ultrastrong coupling between a mechanical oscillator and an *LC* resonator, which was achieved through magnetically

induced electromotive force. This interesting approach to coupling magnetomechanical oscillators in the ultrastrong regime has opened up a wealth of possibilities ranging from sensitive weak-force detection to advanced electromechanical state manipulation.

Furthermore, the use of levitated superconducting microrings offers a unique advantage over traditional microspheres by capitalizing on flux conservation within the ring structure (Navau *et al.*, 2021). This innovative approach provides a versatile platform for the design and optimization of magneto-mechanical ring oscillators. Notably, when it comes to generating quantum superpositions through ground-state cooling, the separation between the peaks of the wave function must exceed the physical dimensions of the object. In this regard ring geometries outperform traditional spherical counterparts, making magnetomechanical ring resonators particularly suitable for experiments investigating gravitational interactions. Recent breakthroughs have demonstrated precise control and levitation of high-*Q* superconducting microspheres (Gutierrez Latorre *et al.*, 2023; Hofer *et al.*, 2023). Appealing aspects of these systems include the mechanical frequencies on the order of 100 Hz, the access to trap anharmonicities, and the scalability of mass of the levitated particle, which is of particular relevance for testing macroscopic quantum states and the role of gravity affecting quantum states. One viable approach involves utilizing a static magnetic trap formed by two coils configured in an anti-Helmholtz arrangement to achieve stable levitation, which was first numerically analyzed in detail for an on-chip configuration (Latorre *et al.*, 2020).

In addition, nanomechanical oscillators can be effectively cooled to their ground state when levitated under an inhomogeneous field, a phenomenon facilitated by the Meissner effect (Cirio, Brennen, and Twamley, 2012). When these oscillators are inductively coupled to a flux qubit, a considerable opportunity to create macroscopic entangled states within these sizable objects arises. This breakthrough holds significant promise for practical applications, especially in the precise measurement of force gradients, such as those encountered in the study of gravity (Johnsson, Brennen, and Twamley, 2016).

In conclusion, while quantum superpositions have been achieved in clamped and hybrid mechanical systems, we are still awaiting their experimental demonstration for levitated mechanics. There are plentiful ideas and approaches under active research, and we expect the first levitated superposition within approximately the next five years.

### 3. Entanglement in mechanical systems

The quantum entanglement of massive objects has been a focal point in the intersection of quantum mechanics and gravity research. An understanding of entanglement in macroscopic systems offers insights into the quantum-to-classical transition and theories related to wave-function collapse, as discussed in Sec. IV.B which delves into gravitational decoherence, semiclassical models, self-energy, and gravitationally-induced wave-function collapse. The intricacies of entanglement mediated by gravity are elaborated upon in Sec. IV.C. Various approaches have been explored to create

and manipulate entanglement in massive quantum systems and aim at unraveling the mysteries of quantum mechanics and its connection to gravity. Several approaches have been proposed for entangling massive quantum systems, and some have recently been experimentally demonstrated in clamped optomechanical systems. Quantum entanglement has not yet been demonstrated in levitated mechanical systems but is an active research objective.

In a clamped system [Eichler \*et al.\* \(2014\)](#) used a Bose-Hubbard dimer consisting of two bosonic modes with an on-site interaction strength to experimentally demonstrate quantum-limited amplification and entanglement. They studied how this system responds in different parameter regimes, and by applying a coherent drive field, they were able to generate entangled photon pairs from vacuum input fields demonstrated through the measurement of cumulants. Their work could prove useful for experimental studies related to non-equilibrium many-particle physics in photonic systems and could be applied toward massive resonators, which would be used for exploring wave-function collapse theories at macroscopic scales due to gravitational interactions between them. [Riedinger \*et al.\* \(2018\)](#) demonstrated the generation of distributed entanglement between two nanomechanical phononic-crystal resonators using a three-step protocol that consists of cryogenically cooling the two mechanical resonators, sending a weak pump pulse into a phase-stabilized interferometer, and creating a phonon. The joint state of the two mechanical systems was then entangled and the entanglement was verified by mapping the mechanical state onto an optical field in this measurement-based entanglement scheme. The simultaneous coupling of two nanomechanical resonators to a superconducting qubit in the strong dispersive regime was used to entangle the two nanomechanical devices ([Wollack \*et al.\*, 2022](#)).

Cavity-optomechanical setups have also been demonstrated experimentally for generating entanglement between two massive mechanical oscillators ([Ockeloen-Korppi \*et al.\*, 2018](#)) via a two-mode backaction evading measurement to verify entanglement in the cavity mode. [Thomas \*et al.\* \(2021\)](#) experimentally achieved entanglement between a macroscopic mechanical oscillator and an atomic spin oscillator. This achievement was accomplished using a millimeter-sized dielectric membrane and an ensemble of roughly  $10^9$  atoms within a magnetic field. The entanglement was confirmed by achieving an Einstein-Podolsky-Rosen variance below the separability limit. The process involved manipulating the light that passed through the two spatially separated systems, with the collective atomic spin serving as an effective negative-mass reference, thereby suppressing quantum backaction.

Further theoretical proposals for entanglement schemes making use of access to a discrete variable qubit quantum system and/or cavity modes include concepts like a coherent feedback loop ([Li \*et al.\*, 2017](#)) applied to two macroscopic mechanical resonators that were strongly coupled to a common optical mode. [Asadian and Abdi \(2016\)](#) proposed using a sequence of pulses to periodically flip a qubit synchronized with the resonator frequency. A conditional photon emission is then applied to the qubit to produce a single photon, depending on the state of the qubit prior to the pulse, to create a mechanically entangled coherent state,

namely, a Schrödinger cat state. [Yi \*et al.\* \(2021\)](#) discussed the use of large-scale spatial qubits to explore macroscopic nonclassicality and entanglement generated through the Casimir effect. As mentioned in Sec. V.B.1, [Cosco, Pedernales, and Plenio \(2021\)](#) proposed a protocol for generating entanglement between two weakly interacting massive resonators, such as nanoparticles levitated by optical means or massive pendula tethered to a base. The protocol involves applying a continuously squeezed protocol to the two resonators and removing the squeezing quickly after generating the desired entanglement. They also proposed the reverse protocol to dramatically reduce the decay rate of the entanglement.

Another promising and especially interesting approach for low-frequency mechanical oscillators, as proposed by [Li and Gröblacher \(2020\)](#), involves the preparation of entangled states between a massive membrane and a low-frequency  $LC$  resonator, and [Li and Gröblacher \(2021\)](#) proposed using cavity magnomechanics for entangling vibrational modes of two massive ferromagnetic spheres. [Xu and Blencowe \(2022\)](#) focused on the entanglement dynamics of spatially separated local  $LC$  oscillators coupled to a long, partially metalized elastic strip through the optomechanical interaction, which is proposed to be usable to observe quantum-gravity-induced entanglement at low energies. Electromechanical systems have been considered as well: [Khosla \*et al.\* \(2018\)](#) introduced a novel approach involving the coupling of multiple electromechanical resonators to a common qubit. Such electromechanical systems also allow one to verify entanglement of mechanical oscillators via qubits ([Bose and Agarwal, 2006](#)). This coupling scheme results in the entanglement of these massive oscillators. This entanglement manifests itself in the form of quantum interference patterns observed in the displacement of the resonators.

In conclusion, various protocols have been proposed and some have already been demonstrated, with each offering unique insights into the interplay between quantum mechanics and gravity. These advancements not only contribute to our understanding of fundamental quantum principles but also have the potential to validate or challenge extensions of the Schrödinger equation, furthering our grasp of the complex realm of quantum mechanics in gravitational contexts, including some interesting questions, such as one considering whether gravity can be used to quantum entangle two particles; see Sec. IV.C. We emphasize that large-mass mechanical systems are considered the predominant system for experimental exploration at the quantum-gravity interface.

## VI. OUTLOOK

In this review we have outlined ideas and proposals for probing the interface between quantum mechanics and gravity with massive quantum systems. We emphasize that many unknowns still persist and that numerous questions remain completely open in the field. Gravity appears to behave differently than other forces in that it can be formulated as curvature and allows for an equivalence principle. A fully fledged theory that consistently combines quantum mechanics

and gravity might ultimately manifest in a completely unexpected manner.

To conclude, we offer a few remarks on some of the outstanding questions raised throughout this review. To begin, we have seen that even classical gravity is one of the least precisely tested forces, especially at small length scales. New and better precision tests are needed to further constrain modified gravity theories and verify our current description of gravity. Quantum-enhanced sensors play an important role here. Equally interesting are questions regarding the nature of gravity itself, and whether its interaction with quantum systems results in a decohering or an entangling process. Here tests with masses prepared in quantum superpositions could provide a key step toward establishing the correct theoretical description. Beyond nonrelativistic quantum mechanics, the mathematical formalism of quantum field theory in curved spacetime enables the interplay of quantum and relativistic effects to be studied at low energies. It has, however, not yet been demonstrated in the laboratory, and the formalism itself gives rise to additional tensions, such as the black-hole information paradox.

All of the aforementioned questions warrant further study. Ultimately, to develop a theory that incorporates quantum and general relativistic effects in a consistent way, it is necessary to understand what principles are truly fundamental. Experiments and the development of quantum technologies play a key role here. In particular, increasing the masses and coherence times of quantum systems may allow for some of the proposals outlined in this review to be realized in the laboratory. The results could help guide future research, which thus far has relied mainly on mathematical and theoretical arguments.

In summary, the prospect of using massive quantum systems to explore the interplay between quantum mechanics and gravity opens up a number of novel questions and exciting challenges. The field is ripe for exploration and potentially groundbreaking discoveries. We hope that this review will inspire many future discussions around these topics and look forward to learning about and partaking in potential discoveries in the future.

## ACKNOWLEDGMENTS

We thank Angelo Bassi, Miles P. Blencowe, Caslav Brukner, and Gary Steele for the fruitful discussions. We thank Gerard Higgins and Onus Hosten for their help with compiling data for Fig. 7. We also thank Jack Clarke, Flaminia Giacomini, Simon Haine, Klaus Hornberger, Nathan Inan, Jie Li, Eduardo Martin-Martinez, Anupam Mazumdar, Julian Pedernales, Igor Pikovski, Martin Plenio, Liu Qui, Tejinder Singh, Michael Tobar, Witlief Wieczorek, and Magdalena Zych for their helpful comments. We acknowledge the EPSRC International Quantum Technologies Network (Grant No. EP/W02683X/1) Levitation Network for Advanced Quantum Technologies for their support. This collaboration began as the U.K. Optomechanics Research Network (UniKORN) online seminar series (2020–2022). We thank all speakers, panel members, and audience participants who contributed to the seminars, which inspired us to write this review. I.F. thanks Eugene Jhong, John Moussouris,

Jussi Westergren, and the Emmy Network for support and research funding and acknowledges support from the Leverhulme Trust project MONDMag (Grant No. RPG-2022-57). A.A.G. is supported in part by NSF Grants No. PHY-2110524 and No. PHY-2111544, the Heising-Simons Foundation, the W.M. Keck Foundation, the John Templeton Foundation, and ONR Grant No. N00014-18-1-2370. M.R. acknowledges funding from the EPSRC via Grants No. EP/N031105/1, No. EP/S000267/1, No. EP/W029626/1, and No. EP/S021582/1 and the STFC via Grant No. ST/W006170/1. S.Q. is funded in part by the Wallenberg Initiative on Networks and Quantum Information (WINQ) and in part by the Marie Skłodowska-Curie Action IF program Nonlinear Optomechanics for Verification, Utility, and Sensing (NOVUS) Grant No. 101027183. Nordita is supported in part by NordForsk. M.T. acknowledges funding from the Leverhulme Trust (Grant No. RPG-2020-197). H.U. acknowledges support from the QuantERA grant LEMAQUME, funded by the QuantERA II ERA-NET Cofund in Quantum Technologies implemented within the EU Horizon 2020 program, from the U.K. funding agency EPSRC (Grants No. EP/W007444/1, No. EP/V035975/1, No. EP/V000624/1, and No. EP/X009491/1), the Leverhulme Trust project MONDMag (Grant No. RPG-2022-57), the EU Horizon 2020 FET-Open project TeQ (Grant No. 766900), the EU Horizon Europe EIC Pathfinder project QuCoM (Grant No. 10032223), and the European Space Agency for the ESA Payload Masters project Op-To-Space, as well as the U.K. Space Agency for the IBF project A3S. C.C.W. acknowledges funding from the Winton Programme for the Physics of Sustainability, EPSRC (Grant No. EP/R513180/1), and the European Union's Horizon 2020 research and innovation program under Grant Agreement No. 732894 (FET-Proactive HOT). S.M.K. acknowledges funding from the ERC via Grant Agreement No. 818751. M.T. acknowledges funding from the Slovenian Research and Innovation Agency (ARIS) under Contracts No. N1-0392, No. P1-0416, and No. SNZRD/ 22-27/0510 (RSUL Toroš). The work of S.B. and A.A.G. is funded by the Gordon and Betty Moore Foundation through Grant No. GBMF12328. This material is based upon work of S.B. and A.A.G. supported by the Alfred P. Sloan Foundation under Grant No. G-2023-21130.

## REFERENCES

- Aasi, Junaid, *et al.*, 2013, “Enhanced sensitivity of the LIGO gravitational wave detector by using squeezed states of light,” *Nat. Photonics* **7**, 613–619.
- Abbott, B., *et al.*, 2009, “Observation of a kilogram-scale oscillator near its quantum ground state,” *New J. Phys.* **11**, 073032.
- Abbott, B.P., *et al.*, 2016a, “Observation of Gravitational Waves from a Binary Black Hole Merger,” *Phys. Rev. Lett.* **116**, 061102.
- Abbott, Benjamin P., *et al.*, 2016b, “First targeted search for gravitational-wave bursts from core-collapse supernovae in data of first-generation laser interferometer detectors,” *Phys. Rev. D* **94**, 102001.
- Abe, Mahiro, *et al.*, 2021, “Matter-Wave Atomic Gradiometer Interferometric Sensor (MAGIS-100),” *Quantum Sci. Technol.* **6**, 044003.



- Abele, Hartmut, and Helmut Leeb, 2012, “Gravitation and quantum interference experiments with neutrons,” *New J. Phys.* **14**, 055010.
- Abend, S., *et al.*, 2016, “Atom-Chip Fountain Gravimeter,” *Phys. Rev. Lett.* **117**, 203003.
- Adesso, G., and I. Fuentes, 2009, “Correlation loss and multipartite entanglement across a black hole horizon,” *Quantum Inf. Comput.* **9**, 0657–0665.
- Adler, Stephen L., 2004, *Quantum Theory as an Emergent Phenomenon: The Statistical Mechanics of Matrix Models as the Precursor of Quantum Field Theory* (Cambridge University Press, Cambridge, England).
- Adler, Stephen L., 2007, “Comments on proposed gravitational modifications of Schrödinger dynamics and their experimental implications,” *J. Phys. A* **40**, 755.
- Adler, Stephen L., Angelo Bassi, and Emiliano Ippoliti, 2005, “Towards quantum superpositions of a mirror: An exact open systems analysis—Calculational details,” *J. Phys. A* **38**, 2715.
- Afek, Gadi, Daniel Carney, and David C. Moore, 2022, “Coherent Scattering of Low Mass Dark Matter from Optically Trapped Sensors,” *Phys. Rev. Lett.* **128**, 101301.
- Agafonova, Sofia, Pere Rossello, Manuel Mekonnen, and Onur Hosten, 2024, “Laser cooling a 1-milligram torsional pendulum to 240 microkelvins,” [arXiv:2408.09445](https://arxiv.org/abs/2408.09445).
- Aguilar, Odylio Denys, 2011, “Past, present and future of the resonant-mass gravitational wave detectors,” *Res. Astron. Astrophys.* **11**, 1.
- Ahmadi, Mehdi, David Edward Bruschi, and Ivette Fuentes, 2014, “Quantum metrology for relativistic quantum fields,” *Phys. Rev. D* **89**, 065028.
- Ahmadi, Mehdi, David Edward Bruschi, Carlos Sabín, Gerardo Adesso, and Ivette Fuentes, 2014, “Relativistic quantum metrology: Exploiting relativity to improve quantum measurement technologies,” *Sci. Rep.* **4**, 4996.
- Aimet, Stefan, Hadrien Chevalier, and M. S. Kim, 2022, “Gravity mediated entanglement between light beams as a table-top test of quantum gravity,” [arXiv:2210.12713](https://arxiv.org/abs/2210.12713).
- Akram, Muhammad Javed, Fazal Ghafoor, and Farhan Saif, 2015, “Electromagnetically induced transparency and tunable fano resonances in hybrid optomechanics,” *J. Phys. B* **48**, 065502.
- Al Balushi, Abdulrahim, Wan Cong, and Robert B. Mann, 2018, “Optomechanical quantum Cavendish experiment,” *Phys. Rev. A* **98**, 043811.
- Albers, Henning, *et al.*, 2020, “Quantum test of the universality of free fall using rubidium and potassium,” *Eur. Phys. J. D* **74**, 145.
- Albrecht, Andreas, Alex Retzker, and Martin B. Plenio, 2014, “Testing quantum gravity by nanodiamond interferometry with nitrogen-vacancy centers,” *Phys. Rev. A* **90**, 033834.
- Alexander, Stephon, Lee Samuel Finn, and Nicolas Yunes, 2008, “Gravitational-wave probe of effective quantum gravity,” *Phys. Rev. D* **78**, 066005.
- Almheiri, Ahmed, Thomas Hartman, Juan Maldacena, Edgar Shaghoulian, and Amirhossein Tajdini, 2021, “The entropy of Hawking radiation,” *Rev. Mod. Phys.* **93**, 035002.
- Alsing, Paul M., and Ivette Fuentes, 2012, “Observer-dependent entanglement,” *Classical Quantum Gravity* **29**, 224001.
- Altarev, I. S., *et al.*, 1992, “New method of measuring the electric dipole moment of the neutron,” *Phys. Lett. A* **276**, 242–246.
- Amati, Daniele, Marcello Ciafaloni, and Gabriele Veneziano, 1989, “Can spacetime be probed below the string size?,” *Phys. Lett. B* **216**, 41–47.
- Amelino-Camelia, Giovanni, John Ellis, N. E. Mavromatos, Dimitri V. Nanopoulos, and Subir Sarkar, 1998, “Tests of quantum gravity from observations of  $\gamma$ -ray bursts,” *Nature (London)* **393**, 763–765.
- Anastopoulos, C., 1996, “Quantum theory of nonrelativistic particles interacting with gravity,” *Phys. Rev. D* **54**, 1600.
- Anastopoulos, C., and B. L. Hu, 2013, “A master equation for gravitational decoherence: Probing the textures of spacetime,” *Classical Quantum Gravity* **30**, 165007.
- Anastopoulos, C., and B. L. Hu, 2014a, “Newton-Schrödinger equations are not derivable from general relativity + quantum field theory,” [arXiv:1402.3813](https://arxiv.org/abs/1402.3813).
- Anastopoulos, C., and B. L. Hu, 2014b, “Problems with the Newton-Schrödinger equations,” *New J. Phys.* **16**, 085007.
- Anastopoulos, Charis, and Bei-Lok Hu, 2018, “Equivalence principle for quantum systems: Dephasing and phase shift of free-falling particles,” *Classical Quantum Gravity* **35**, 035011.
- Anastopoulos, Charis, and Bei-Lok Hu, 2020, “Quantum superposition of two gravitational cat states,” *Classical Quantum Gravity* **37**, 235012.
- Anastopoulos, Charis, and Bei-Lok Hu, 2022, “Gravitational decoherence: A thematic overview,” *AVS Quantum Sci.* **4**, 015602.
- Arcizet, O., T. Briant, A. Heidmann, and M. Pinard, 2006, “Beating quantum limits in an optomechanical sensor by cavity detuning,” *Phys. Rev. A* **73**, 033819.
- Arcizet, Olivier, Vincent Jacques, Alessandro Siria, Philippe Poncharal, Pascal Vincent, and Signe Seidelin, 2011, “A single nitrogen-vacancy defect coupled to a nanomechanical oscillator,” *Nat. Phys.* **7**, 879–883.
- Arkani-Hamed, Nima, Savas Dimopoulos, and Gia Dvali, 1998, “The hierarchy problem and new dimensions at a millimeter,” *Phys. Lett. B* **429**, 263–272.
- Armano, Michele, *et al.*, 2016, “Sub-Femto- $g$  Free Fall for Space-Based Gravitational Wave Observatories: LISA Pathfinder Results,” *Phys. Rev. Lett.* **116**, 231101.
- Armata, Federico, Ludovico Latmiral, Igor Pikovski, Michael R. Vanner, Časlav Brukner, and M. S. Kim, 2016, “Quantum and classical phases in optomechanics,” *Phys. Rev. A* **93**, 063862.
- Armata, Federico, Ludovico Latmiral, ADK Plato, and M. S. Kim, 2017, “Quantum limits to gravity estimation with optomechanics,” *Phys. Rev. A* **96**, 043824.
- Arndt, Markus, and Klaus Hornberger, 2014, “Testing the limits of quantum mechanical superpositions,” *Nat. Phys.* **10**, 271–277.
- Arnowitz, Richard, Stanley Deser, and Charles W. Misner, 1959, “Dynamical structure and definition of energy in general relativity,” *Phys. Rev.* **116**, 1322.
- Arnquist, I. J., *et al.*, 2022, “Search for Spontaneous Radiation from Wave Function Collapse in the Majorana Demonstrator,” *Phys. Rev. Lett.* **129**, 080401.
- Arrangoiz-Arriola, Patricio, E. Alex Wollack, Zhaoyou Wang, Marek Pechal, Wentao Jiang, Timothy P. McKenna, Jeremy D. Witmer, Raphaël Van Laer, and Amir H. Safavi-Naeini, 2019, “Resolving the energy levels of a nanomechanical oscillator,” *Nature (London)* **571**, 537–540.
- Arvidsson, Elisabet K., Ermes Scarano, August K. Roos, Sofia Qvarfort, and David B. Haviland, 2024, “Sensing force gradients with cavity optomechanics while evading backaction,” [arXiv:2405.06589](https://arxiv.org/abs/2405.06589).
- Asadian, Ali, and Mehdi Abdi, 2016, “Heralded entangled coherent states between spatially separated massive resonators,” *Phys. Rev. A* **93**, 052315.
- Asenbaum, Peter, Chris Overstreet, Minjeong Kim, Joseph Curti, and Mark A. Kasevich, 2020, “Atom-Interferometric Test of the Equivalence Principle at the  $10^{-12}$  Level,” *Phys. Rev. Lett.* **125**, 191101.
- Asenbaum, Peter, Chris Overstreet, Tim Kovachy, Daniel D. Brown, Jason M. Hogan, and Mark A. Kasevich, 2017, “Phase Shift in an

- Atom Interferometer due to Spacetime Curvature across Its Wave Function,” *Phys. Rev. Lett.* **118**, 183602.
- Ashkin, Arthur, and J. M. Dziedzic, 1971, “Optical levitation by radiation pressure,” *Appl. Phys. Lett.* **19**, 283–285.
- Ashtekar, Abhay, and Eugenio Bianchi, 2021, “A short review of loop quantum gravity,” *Rep. Prog. Phys.* **84**, 042001.
- Aspelmeyer, Markus, 2022, “When Zeh meets Feynman: How to avoid the appearance of a classical world in gravity experiments,” in *From Quantum to Classical: Essays in Honour of H.-Dieter Zeh*, Fundamental Theories of Physics Vol. 204, edited by Claus Kiefer (Springer, New York), pp. 85–95.
- Aspelmeyer, Markus, Tobias J. Kippenberg, and Florian Marquardt, 2014, “Cavity optomechanics,” *Rev. Mod. Phys.* **86**, 1391.
- Asprea, L., A. Bassi, H. Ulbricht, and G. Gasbarri, 2021, “Gravitational Decoherence and the Possibility of Its Interferometric Detection,” *Phys. Rev. Lett.* **126**, 200403.
- Audretsch, J., and C. Lammerzahl, 1983, “Neutron interference: General theory of the influence of gravity, inertia and space-time torsion,” *J. Phys. A* **16**, 2457.
- Bachtold, Adrian, Joel Moser, and M. I. Dykman, 2022, “Mesoscopic physics of nanomechanical systems,” *Rev. Mod. Phys.* **94**, 045005.
- Badurina, L., *et al.*, 2020, “AION: An atom interferometer observatory and network,” *J. Cosmol. Astropart. Phys.* **05**, 011.
- Bahrami, Mohammad, André Großardt, Sandro Donadi, and Angelo Bassi, 2014, “The Schrödinger-Newton equation and its foundations,” *New J. Phys.* **16**, 115007.
- Bahrami, Mohammad, Andrea Smirne, and Angelo Bassi, 2014, “Role of gravity in the collapse of a wave function: A probe into the Diósi-Penrose model,” *Phys. Rev. A* **90**, 062105.
- Ball, Jonathan L., Ivette Fuentes-Schuller, and Frederic P. Schuller, 2006, “Entanglement in an expanding spacetime,” *Phys. Lett. A* **359**, 550–554.
- Barausse, Enrico, *et al.*, 2020, “Prospects for fundamental physics with LISA,” *Gen. Relativ. Gravit.* **52**, 81.
- Barchielli, Alberto, and Bassano Vacchini, 2015, “Quantum Langevin equations for optomechanical systems,” *New J. Phys.* **17**, 083004.
- Bar-Gill, Nir, Linh M. Pham, Andrejs Jarmola, Dmitry Budker, and Ronald L. Walsworth, 2013, “Solid-state electronic spin coherence time approaching one second,” *Nat. Commun.* **4**, 1743.
- Barker, P. F., and M. N. Shneider, 2010, “Cavity cooling of an optically trapped nanoparticle,” *Phys. Rev. A* **81**, 023826.
- Barman, Dipankar, Angshuman Choudhury, Bhushan Kad, and Bibhas Ranjan Majhi, 2023, “Spontaneous entanglement leakage of two static entangled Unruh-Dewitt detectors,” *Phys. Rev. D* **107**, 045001.
- Barzanjeh, Shabir, André Xuereb, Simon Gröblacher, Mauro Paternostro, Cindy A. Regal, and Eva M. Weig, 2022, “Optomechanics for quantum technologies,” *Nat. Phys.* **18**, 15–24.
- Bassi, Angelo, and GianCarlo Ghirardi, 2003, “Dynamical reduction models,” *Phys. Rep.* **379**, 257–426.
- Bassi, Angelo, André Großardt, and Hendrik Ulbricht, 2017, “Gravitational decoherence,” *Classical Quantum Gravity* **34**, 193002.
- Bassi, Angelo, Emiliano Ippoliti, and Stephen L. Adler, 2005, “Towards Quantum Superpositions of a Mirror: An Exact Open Systems Analysis,” *Phys. Rev. Lett.* **94**, 030401.
- Bassi, Angelo, Kinjalk Lochan, Seema Satin, Tejinder P. Singh, and Hendrik Ulbricht, 2013, “Models of wave-function collapse, underlying theories, and experimental tests,” *Rev. Mod. Phys.* **85**, 471.
- Bateman, James, Stefan Nimmrichter, Klaus Hornberger, and Hendrik Ulbricht, 2014, “Near-field interferometry of a free-falling nanoparticle from a point-like source,” *Nat. Commun.* **5**, 4788.
- Bawaj, Mateusz, *et al.*, 2015, “Probing deformed commutators with macroscopic harmonic oscillators,” *Nat. Commun.* **6**, 7503.
- Beckerling Vinckers, Ulrich K., Álvaro de la Cruz-Dombriz, and Anupam Mazumdar, 2023, “Quantum entanglement of masses with nonlocal gravitational interaction,” *Phys. Rev. D* **107**, 124036.
- Bedingham, Daniel, Detlef Dürr, GianCarlo Ghirardi, Sheldon Goldstein, Roderich Tumulka, and Nino Zanghi, 2014, “Matter density and relativistic models of wave function collapse,” *J. Stat. Phys.* **154**, 623–631.
- Belenchia, Alessio, Dionigi M. T. Benincasa, Stefano Liberati, Francesco Marin, Francesco Marino, and Antonello Ortolan, 2016, “Testing Quantum Gravity Induced Nonlocality via Optomechanical Quantum Oscillators,” *Phys. Rev. Lett.* **116**, 161303.
- Belenchia, Alessio, Robert M. Wald, Flaminia Giacomini, Esteban Castro-Ruiz, Časlav Brukner, and Markus Aspelmeyer, 2018, “Quantum superposition of massive objects and the quantization of gravity,” *Phys. Rev. D* **98**, 126009.
- Belenchia, Alessio, *et al.*, 2022, “Quantum physics in space,” *Phys. Rep.* **951**, 1–70.
- Bellei, G., P. Droll, F. Delhaise, I. Harrison, and D. Amend, 2017, “Laser Interferometer Space Antenna (LISA) Pathfinder: New methods for acquisition of signal after large apogee-raising maneuvers,” in *Space Operations: Contributions from the Global Community*, edited by C. Cruzen, M. Schmidhuber, Y. Lee, and B. Kim (Springer, Cham, Switzerland), pp. 567–583, 10.1007/978-3-319-51941-8\_25.
- Bennett, James S., Kiran Khosla, Lars S. Madsen, Michael R. Vanner, Halina Rubinsztein-Dunlop, and Warwick P. Bowen, 2016, “A quantum optomechanical interface beyond the resolved sideband limit,” *New J. Phys.* **18**, 053030.
- Bennett, Steven D., Lynda Cockins, Yoichi Miyahara, Peter Grütter, and Aashish A. Clerk, 2010, “Strong Electromechanical Coupling of an Atomic Force Microscope Cantilever to a Quantum Dot,” *Phys. Rev. Lett.* **104**, 017203.
- Bera, Sayantani, Ravi Mohan, and Tejinder P. Singh, 2015, “Stochastic modification of the Schrödinger-Newton equation,” *Phys. Rev. D* **92**, 025054.
- Berglund, Per, Andrew Geraci, Tristan Hübsch, David Mattingly, and Djordje Minic, 2023, “Triple interference, non-linear Talbot effect and gravitization of the quantum,” *Classical Quantum Gravity* **40**, 155008.
- Berglund, Per, Tristan Hübsch, David Mattingly, and Djordje Minic, 2022, “Gravitizing the quantum,” *Int. J. Mod. Phys. D* **31**, 2242024.
- Bernád, József Zsolt, Lajos Diósi, and Tamás Geszti, 2006, “Quest for Quantum Superpositions of a Mirror: High and Moderately Low Temperatures,” *Phys. Rev. Lett.* **97**, 250404.
- Bertlmann, Reinhold A., and Beatrix C. Hiesmayr, 2001, “Bell inequalities for entangled kaons and their unitary time evolution,” *Phys. Rev. A* **63**, 062112.
- Betz, J., J. Manley, E. M. Wright, D. Grin, and S. Singh, 2022, “Searching for Chameleon Dark Energy with Mechanical Systems,” *Phys. Rev. Lett.* **129**, 131302.
- Bhattacharyya, Sukanta, Sunandan Gangopadhyay, and Anirban Saha, 2020, “Generalized uncertainty principle in resonant detectors of gravitational waves,” *Classical Quantum Gravity* **37**, 195006.
- Bild, Marius, Matteo Fadel, Y. u. Yang, Uwe von Lüpke, Phillip Martin, Alessandro Bruno, and Yiwen Chu, 2023, “Schrödinger cat states of a 16-microgram mechanical oscillator,” *Science* **380**, 274–278.
- Birrell, N., and P. Davies, 1982, *Quantum Fields in Curved Space*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, England).

- Bishop, Michael, Jaeyeon Lee, and Douglas Singleton, 2020, "Modified commutators are not sufficient to determine a quantum gravity minimal length scale," *Phys. Lett. B* **802**, 135209.
- Biswas, Dripto, Sougato Bose, Anupam Mazumdar, and Marko Toroš, 2023, "Gravitational optomechanics: Photon-matter entanglement via graviton exchange," *Phys. Rev. D* **108**, 064023.
- Bjerrum-Bohr, N. E. J., John F. Donoghue, and Barry R. Holstein, 2003, "Quantum gravitational corrections to the nonrelativistic scattering potential of two masses," *Phys. Rev. D* **67**, 084033.
- Bjerrum-Bohr, N. E. J., John F. Donoghue, Barry R. Holstein, Ludovic Planté, and Pierre Vanhove, 2015, "Bending of Light in Quantum Gravity," *Phys. Rev. Lett.* **114**, 061301.
- Bjerrum-Bohr, N. E. J., John F. Donoghue, and Pierre Vanhove, 2014, "On-shell techniques and universal results in quantum gravity," *J. High Energy Phys.* **02**, 111.
- Björk, Gunnar, and Piero G. Luca Mana, 2004, "A size criterion for macroscopic superposition states," *J. Opt. B* **6**, 429.
- Blakemore, Charles P., Alexander Fieguth, Akio Kawasaki, Nadav Priel, Denzal Martin, Alexander D. Rider, Qidong Wang, and Giorgio Gratta, 2021, "Search for non-Newtonian interactions at micrometer scale with a levitated test mass," *Phys. Rev. D* **104**, L061101.
- Blencowe, M. P., 2013, "Effective Field Theory Approach to Gravitationally Induced Decoherence," *Phys. Rev. Lett.* **111**, 021302.
- Bleuler, K., 1950, "Eine neue methode zur behandlung der longitudinalen und skalaren photonen," [10.5169/SEALS-112124](https://arxiv.org/abs/10.5169/SEALS-112124).
- Bollinger, John J., Wayne M. Itano, David J. Wineland, and Daniel J. Heinzen, 1996, "Optimal frequency measurements with maximally correlated states," *Phys. Rev. A* **54**, R4649.
- Bose, S., and G. S. Agarwal, 2006, "Entangling pairs of nanocantilevers, Cooper-pair boxes and mesoscopic teleportation," *New J. Phys.* **8**, 34.
- Bose, S., K. Jacobs, and P. L. Knight, 1997, "Preparation of non-classical states in cavities with a moving mirror," *Phys. Rev. A* **56**, 4175.
- Bose, Sougato, 2006, "Qubit Assisted Probing of Coherence between Mesoscopic States of an Apparatus," *Phys. Rev. Lett.* **96**, 060402.
- Bose, Sougato, 2016, "Matter wave Ramsey interferometry and the quantum nature of gravity," lecture, Fundamental Problems of Quantum Physics, International Centre for Theoretical Sciences, Bangalore, India, 2016, [https://www.youtube.com/watch?v=0Fv-0k13s\\_k](https://www.youtube.com/watch?v=0Fv-0k13s_k).
- Bose, Sougato, Kurt Jacobs, and Peter L. Knight, 1999, "Scheme to probe the decoherence of a macroscopic object," *Phys. Rev. A* **59**, 3204.
- Bose, Sougato, Anupam Mazumdar, Gavin W. Morley, Hendrik Ulbricht, Marko Toroš, Mauro Paternostro, Andrew A. Geraci, Peter F. Barker, M. S. Kim, and Gerard Milburn, 2017, "Spin Entanglement Witness for Quantum Gravity," *Phys. Rev. Lett.* **119**, 240401.
- Bose, Sougato, Anupam Mazumdar, Martine Schut, and Marko Toroš, 2022, "Mechanism for the quantum natured gravitons to entangle masses," *Phys. Rev. D* **105**, 106028.
- Bowen, Warwick P., and Gerard J. Milburn, 2015, *Quantum Optomechanics* (CRC Press, Boca Raton).
- Brady, Anthony J., Xin Chen, Kewen Xiao, Y. i. Xia, Zhen Liu, Roni Harnik, Dalziel J. Wilson, Zheshen Zhang, and Quntao Zhuang, 2022, "Entanglement-enhanced optomechanical sensor array for dark matter searches," [arXiv:2210.07291](https://arxiv.org/abs/2210.07291).
- Braginsky, V. B., and F. Ya. Khalili, 1996, "Quantum nondemolition measurements: The route from toys to tools," *Rev. Mod. Phys.* **68**, 1–11.
- Braginsky, Vladimir B., and Farid Ya. Khalili, 1995, *Quantum Measurement* (Cambridge University Press, Cambridge, England).
- Braginsky, Vladimir B., Yuri I. Vorontsov, and Kip S. Thorne, 1980, "Quantum nondemolition measurements," *Science* **209**, 547–557.
- Branford, Dominic, Haixing Miao, and Animesh Datta, 2018, "Fundamental Quantum Limits of Multicarrier Optomechanical Sensors," *Phys. Rev. Lett.* **121**, 110505.
- Braunstein, Samuel L., and Carlton M. Caves, 1994, "Statistical Distance and the Geometry of Quantum States," *Phys. Rev. Lett.* **72**, 3439–3443.
- Braunstein, Samuel L., Carlton M. Caves, and G. J. Milburn, 1996, "Generalized uncertainty relations: Theory, examples, and Lorentz invariance," *Ann. Phys. (N.Y.)* **247**, 135–173.
- Bravo, Tupac, Dennis Rätzel, and Ivette Fuentes, 2020, "Phononic gravity gradiometry with Bose-Einstein condensates," [arXiv:2001.10104v2](https://arxiv.org/abs/2001.10104v2).
- Brax, Philippe, Carsten van de Bruck, Anne-Christine Davis, Justin Khoury, and Amanda Weltman, 2004, "Detecting dark energy in orbit: The cosmological chameleon," *Phys. Rev. D* **70**, 123518.
- Breuer, H. P., P. I. H. P. Breuer, F. Petruccione, and S. P. A. P. F. Petruccione, 2002, *The Theory of Open Quantum Systems* (Oxford University Press, New York).
- Breuer, Heinz-Peter, Ertan Göklü, and Claus Lämmerzahl, 2009, "Metric fluctuations and decoherence," *Classical Quantum Gravity* **26**, 105012.
- Briegel, Hans-Jürgen, and Berthold-Georg Englert, 1993, "Quantum optical master equations: The use of damping bases," *Phys. Rev. A* **47**, 3311–3329.
- Brodeur, M., *et al.*, 2023, "Nuclear  $\beta$  decay as a probe for physics beyond the standard model," [arXiv:2301.03975](https://arxiv.org/abs/2301.03975).
- Bronstein, M., 1933, "K voprosu o vozmoznoy teorii mira kak tselogo [On the question of a possible theory of the world as a whole]," *Usp. Astron. Nauk. Sb.* **3**, 3–30.
- Brunelli, Matteo, Daniel Malz, and Andreas Nunnenkamp, 2019, "Conditional Dynamics of Optomechanical Two-Tone Backaction-Evading Measurements," *Phys. Rev. Lett.* **123**, 093602.
- Brunelli, Matteo, Daniel Malz, Albert Schliesser, and Andreas Nunnenkamp, 2020, "Stroboscopic quantum optomechanics," *Phys. Rev. Res.* **2**, 023241.
- Bruschi, David Edward, Animesh Datta, Rupert Ursin, Timothy C. Ralph, and Ivette Fuentes, 2014, "Quantum estimation of the Schwarzschild spacetime parameters of the Earth," *Phys. Rev. D* **90**, 124001.
- Bruschi, David Edward, Ivette Fuentes, and Jorma Louko, 2012, "Voyage to Alpha Centauri: Entanglement degradation of cavity modes due to motion," *Phys. Rev. D* **85**, 061701.
- Bruschi, David Edward, Benjamin Morris, and Ivette Fuentes, 2020, "Thermodynamics of relativistic quantum fields confined in cavities," *Phys. Lett. A* **384**, 126601.
- Bruschi, David Edward, Timothy C. Ralph, Ivette Fuentes, Thomas Jennewein, and Mohsen Razavi, 2014, "Spacetime effects on satellite-based quantum communications," *Phys. Rev. D* **90**, 045041.
- Bruschi, David Edward, Carlos Sabín, Angela White, Valentina Baccetti, Daniel K. L. Oi, and Ivette Fuentes, 2014, "Testing the effects of gravity and motion on quantum entanglement in space-based experiments," *New J. Phys.* **16**, 053041.
- Bub, Mathew W., Yanbei Chen, Yufeng Du, Dongjun Li, Yiwen Zhang, and Kathryn M. Zurek, 2023, "Quantum gravity background in next-generation gravitational wave detectors," [arXiv:2305.11224](https://arxiv.org/abs/2305.11224).



- Bubuianu, Iuliana, Sergiu I. Vacaru, and Elsen Veli Veliev, 2021, “Kaluza-Klein gravity and cosmology emerging from G. Perelman’s entropy functionals and quantum geometric information flows,” *Eur. Phys. J. Plus* **136**, 149.
- Buikema, Aaron, *et al.*, 2020, “Sensitivity and performance of the Advanced LIGO detectors in the third observing run,” *Phys. Rev. D* **102**, 062003.
- Bures, Donald, 1969, “An extension of Kakutani’s theorem on infinite product measures to the tensor product of semifinite  $w^*$ -algebras,” *Trans. Am. Math. Soc.* **135**, 199–212, <https://community.ams.org/journals/tran/1969-135-00/S0002-9947-1969-0236719-2/S0002-9947-1969-0236719-2.pdf>.
- Burrage, Clare, and Jeremy Sakstein, 2018, “Tests of chameleon gravity,” *Living Rev. Relativity* **21**, 1–58.
- Bushev, P. A., J. Bourhill, Maxim Goryachev, N. Kukharchyk, E. Ivanov, Serge Galliou, M. E. Tobar, and S. Danilishin, 2019, “Testing the generalized uncertainty principle with macroscopic mechanical oscillators and pendulums,” *Phys. Rev. D* **100**, 066020.
- Caldeira, Amir O., and Anthony J. Leggett, 1983a, “Path integral approach to quantum Brownian motion,” *Physica (Amsterdam)* **121A**, 587–616.
- Caldeira, Amir O., and Anthony J. Leggett, 1983b, “Quantum tunnelling in a dissipative system,” *Ann. Phys. (N.Y.)* **149**, 374–456.
- Camerer, Stephan, Maria Korppi, Andreas Jöckel, David Hunger, Theodor W. Hänsch, and Philipp Treutlein, 2011, “Realization of an Optomechanical Interface between Ultracold Atoms and a Membrane,” *Phys. Rev. Lett.* **107**, 223001.
- Campaioli, Francesco, Jared H. Cole, and Harini Hapuarachchi, 2023, “A tutorial on quantum master equations: Tips and tricks for quantum optics, quantum computing and beyond,” *arXiv:2303.16449*.
- Campbell, William M., Michael E. Tobar, Maxim Goryachev, and Serge Galliou, 2023, “Improved constraints on minimum length models with a macroscopic low loss phonon cavity,” *Phys. Rev. D* **108**, 102006.
- Cano, Daniel, B. Kasch, Helge Hattermann, Reinhold Kleiner, Claus Zimmermann, Dieter Koelle, and József Fortágh, 2008, “Meissner Effect in Superconducting Microtraps,” *Phys. Rev. Lett.* **101**, 183006.
- Canuel, B., *et al.*, 2018, “Exploring gravity with the MIGA large scale atom interferometer,” *Sci. Rep.* **8**, 14064.
- Carlesso, Matteo, Angelo Bassi, Paolo Falferi, and Andrea Vinante, 2016, “Experimental bounds on collapse models from gravitational wave detectors,” *Phys. Rev. D* **94**, 124036.
- Carlesso, Matteo, Sandro Donadi, Luca Ferialdi, Mauro Paternostro, Hendrik Ulbricht, and Angelo Bassi, 2022, “Present status and future challenges of non-interferometric tests of collapse models,” *Nat. Phys.* **18**, 243–250.
- Carlesso, Matteo, Mauro Paternostro, Hendrik Ulbricht, and Angelo Bassi, 2017, “When Cavendish meets Feynman: A quantum torsion balance for testing the quantumness of gravity,” *arXiv:1710.08695*.
- Carlip, Steve, 2008, “Is quantum gravity necessary?,” *Classical Quantum Gravity* **25**, 154010.
- Carney, D., *et al.*, 2021, “Mechanical quantum sensing in the search for dark matter,” *Quantum Sci. Technol.* **6**, 024002.
- Carney, Daniel, 2022, “Newton, entanglement, and the graviton,” *Phys. Rev. D* **105**, 024029.
- Carney, Daniel, Anson Hook, Zhen Liu, Jacob M. Taylor, and Yue Zhao, 2021, “Ultralight dark matter detection with mechanical quantum sensors,” *New J. Phys.* **23**, 023041.
- Carney, Daniel, Holger Müller, and Jacob M. Taylor, 2021, “Using an atom interferometer to infer gravitational entanglement generation,” *PRX Quantum* **2**, 030330.
- Carney, Daniel, Philip C.E. Stamp, and Jacob M. Taylor, 2019, “Tabletop experiments for quantum gravity: A user’s manual,” *Classical Quantum Gravity* **36**, 034001.
- Carusotto, Iacopo, Serena Fagnocchi, Alessio Recati, Roberto Balbinot, and Alessandro Fabbri, 2008, “Numerical observation of Hawking radiation from acoustic black holes in atomic Bose-Einstein condensates,” *New J. Phys.* **10**, 103001.
- Cattiaux, D., *et al.*, 2021, “A macroscopic object passively cooled into its quantum ground state of motion beyond single-mode cooling,” *Nat. Commun.* **12**, 6182.
- Cavalcanti, E. G., and M. D. Reid, 2006, “Signatures for Generalized Macroscopic Superpositions,” *Phys. Rev. Lett.* **97**, 170405.
- Caves, Carlton M., 1979, “Microwave cavity gravitational radiation detectors,” *Phys. Lett.* **80B**, 323–326.
- Caves, Carlton M., 1982, “Quantum limits on noise in linear amplifiers,” *Phys. Rev. D* **26**, 1817–1839.
- Caves, Carlton M., Joshua Combes, Zhang Jiang, and Shashank Pandey, 2012, “Quantum limits on phase-preserving linear amplifiers,” *Phys. Rev. A* **86**, 063802.
- Caves, Carlton M., and G. J. Milburn, 1987, “Quantum-mechanical model for continuous position measurements,” *Phys. Rev. A* **36**, 5543–5555.
- Caves, Carlton M., Kip S. Thorne, Ronald W. P. Drever, Vernon D. Sandberg, and Mark Zimmermann, 1980, “On the measurement of a weak classical force coupled to a quantum-mechanical oscillator. I. Issues of principle,” *Rev. Mod. Phys.* **52**, 341–392.
- Chakraborty, Sumanta, Anupam Mazumdar, and Ritapriya Pradhan, 2023, “Distinguishing Jordan and Einstein frames in gravity through entanglement,” *arXiv:2310.06899*.
- Chan, Jasper, T. P. Mayer Alegre, Amir H. Safavi-Naeini, Jeff T. Hill, Alex Krause, Simon Gröblacher, Markus Aspelmeyer, and Oskar Painter, 2011, “Laser cooling of a nanomechanical oscillator into its quantum ground state,” *Nature (London)* **478**, 89–92.
- Chang, Darrick E., C. A. Regal, S. B. Papp, D. J. Wilson, J. Ye, O. Painter, H. Jeff Kimble, and P. Zoller, 2010, “Cavity optomechanics using an optically levitated nanosphere,” *Proc. Natl. Acad. Sci. U.S.A.* **107**, 1005–1010.
- Chang, Lay Nam, Djordje Minic, Naotoshi Okamura, and Tatsuo Takeuchi, 2002, “Exact solution of the harmonic oscillator in arbitrary dimensions with minimal length uncertainty relations,” *Phys. Rev. D* **65**, 125027.
- Chatterjee, Riddhi, Sunandan Gangopadhyay, and A. S. Majumdar, 2021, “Violation of equivalence in an accelerating atom-mirror system in the generalized uncertainty principle framework,” *Phys. Rev. D* **104**, 124001.
- Chen, Lei, Jian Liu, and Ka-Di Zhu, 2022, “Constraining the axion-nucleon coupling and non-Newtonian gravity with a levitated optomechanical device,” *Phys. Rev. D* **106**, 095007.
- Chen, Lin-Qing, Flaminia Giacomini, and Carlo Rovelli, 2023, “Quantum states of fields for quantum split sources,” *Quantum* **7**, 958.
- Chen, Xi, Yong-Chun Liu, Pai Peng, Yanyan Zhi, and Yun-Feng Xiao, 2015, “Cooling of macroscopic mechanical resonators in hybrid atom-optomechanical systems,” *Phys. Rev. A* **92**, 033841.
- Chevalier, Hadrien, A. J. Paige, and M. S. Kim, 2020, “Witnessing the nonclassical nature of gravity in the presence of unknown interactions,” *Phys. Rev. A* **102**, 022428.
- Chiao, R. Y., N. A. Inan, M. Scheibner, J. Sharping, D. A. Singleton, and M. E. Tobar, 2023, “Gravitational Aharonov-Bohm effect,” *arXiv:2311.07764*.
- Chiao, Raymond Y., Robert W. Haun, Nader A. Inan, Bong-Soo Kang, Luis A. Martinez, Stephen J. Minter, Gerardo A. Munoz, and Douglas A. Singleton, 2014, “A gravitational Aharonov-Bohm

- effect, and its connection to parametric oscillators and gravitational radiation,” in *Quantum Theory: A Two-Time Success Story—Yakir Aharonov Festschrift*, edited by Daniele C. Struppa and Jeffrey M. Tollaksen (Springer, New York), pp. 213–246.
- Chiaverini, J., S. J. Smullin, A. A. Geraci, D. M. Weld, and A. Kapitulnik, 2003, “New Experimental Constraints on Non-Newtonian Forces below 100  $\mu\text{m}$ ,” *Phys. Rev. Lett.* **90**, 151101.
- Chiu, Sheng-wei, Tim Kovachy, Hui-Chun Chien, and Mark A. Kasevich, 2011, “102 $hk$  Large Area Atom Interferometers,” *Phys. Rev. Lett.* **107**, 130403.
- Chowdhury, A., P. Vezio, M. Bonaldi, A. Borrielli, F. Marino, Bruno Morana, G. A. Prodi, Pasqualina M. Sarro, Enrico Serra, and F. Marin, 2020, “Quantum Signature of a Squeezed Mechanical Oscillator,” *Phys. Rev. Lett.* **124**, 023601.
- Christian, Joy, 1997, “Exactly soluble sector of quantum gravity,” *Phys. Rev. D* **56**, 4844.
- Christodoulou, Marios, Andrea Di Biagio, Markus Aspelmeyer, Časlav Brukner, Carlo Rovelli, and Richard Howl, 2023, “Locally Mediated Entanglement in Linearized Quantum Gravity,” *Phys. Rev. Lett.* **130**, 100202.
- Christodoulou, Marios, and Carlo Rovelli, 2019, “On the possibility of laboratory evidence for quantum superposition of geometries,” *Phys. Lett. B* **792**, 64–68.
- Chu, Yiwen, and Simon Gröblacher, 2020, “A perspective on hybrid quantum opto- and electromechanical systems,” *Appl. Phys. Lett.* **117**, 150503.
- Chu, Yiwen, Prashanta Kharel, William H. Renninger, Luke D. Burkhardt, Luigi Frunzio, Peter T. Rakich, and Robert J. Schoelkopf, 2017, “Quantum acoustics with superconducting qubits,” *Science* **358**, 199–202.
- Chu, Yiwen, Prashanta Kharel, Taekwan Yoon, Luigi Frunzio, Peter T. Rakich, and Robert J. Schoelkopf, 2018, “Creation and control of multi-phonon Fock states in a bulk acoustic-wave resonator,” *Nature (London)* **563**, 666–670.
- Cirac, Juan I., and Peter Zoller, 1995, “Quantum Computations with Cold Trapped Ions,” *Phys. Rev. Lett.* **74**, 4091.
- Cirio, M., G. K. Brennen, and J. Twamley, 2012, “Quantum Magnetomechanics: Ultrahigh- $q$ -Levitated Mechanical Oscillators,” *Phys. Rev. Lett.* **109**, 147206.
- Clark, Jeremy B., Florent Lecocq, Raymond W. Simmonds, José Aumentado, and John D. Teufel, 2017, “Sideband cooling beyond the quantum backaction limit with squeezed light,” *Nature (London)* **541**, 191–195.
- Clarke, J., P. Sahium, K. E. Khosla, Igor Pikovski, M. S. Kim, and M. R. Vanner, 2020, “Generating mechanical and optomechanical entanglement via pulsed interaction and measurement,” *New J. Phys.* **22**, 063001.
- Clarke, Jack, Pascal Neveu, Kiran E. Khosla, Ewold Verhagen, and Michael R. Vanner, 2023, “Cavity Quantum Optomechanical Nonlinearities and Position Measurement beyond the Breakdown of the Linearized Approximation,” *Phys. Rev. Lett.* **131**, 053601.
- Clarke, Jack, and Michael R. Vanner, 2018, “Growing macroscopic superposition states via cavity quantum optomechanics,” *Quantum Sci. Technol.* **4**, 014003.
- Clerk, A. A., M. H. Devoret, S. M. Girvin, Florian Marquardt, and R. J. Schoelkopf, 2010, “Introduction to quantum noise, measurement, and amplification,” *Rev. Mod. Phys.* **82**, 1155–1208.
- Clerk, A. A., F. Marquardt, and K. Jacobs, 2008, “Back-action evasion and squeezing of a mechanical resonator using a cavity detector,” *New J. Phys.* **10**, 095010.
- Clerk, Aashish A., 2020, “Optomechanics and quantum measurement,” in *Quantum Optomechanics and Nanomechanics*, Proceedings of the Les Houches Summer School, Session CV, edited by Pierre-Francois Cohadon, Jack Harris, Florian Marquardt, and Leticia Cugliandolo (Oxford University Press, New York).
- Clifton, Timothy, Pedro G. Ferreira, Antonio Padilla, and Constantinos Skordis, 2012, “Modified gravity and cosmology,” *Phys. Rep.* **513**, 1–189.
- Colella, R., A. W. Overhauser, and S. A. Werner, 1975, “Observation of Gravitationally Induced Quantum Interference,” *Phys. Rev. Lett.* **34**, 1472–1474.
- Colin, Samuel, Thomas Durt, and Ralph Willox, 2016, “Crucial tests of macrorealist and semiclassical gravity models with freely falling mesoscopic nanospheres,” *Phys. Rev. A* **93**, 062102.
- Cosco, F., J. S. Pedernales, and Martin B. Plenio, 2021, “Enhanced force sensitivity and entanglement in periodically driven optomechanics,” *Phys. Rev. A* **103**, L061501.
- Cronin, Alexander D., Jörg Schmiedmayer, and David E. Pritchard, 2009, “Optics and interferometry with atoms and molecules,” *Rev. Mod. Phys.* **81**, 1051–1129.
- Cui, Dianzhen, T. Li, Jianning Li, and Xuexi Yi, 2021, “Detecting deformed commutators with exceptional points in optomechanical sensors,” *New J. Phys.* **23**, 123037.
- Dahleh, M., A. P. Peirce, and H. Rabitz, 1990, “Optimal control of uncertain quantum systems,” *Phys. Rev. A* **42**, 1065–1079.
- Damour, Thibault, 2012, “Theoretical aspects of the equivalence principle,” *Classical Quantum Gravity* **29**, 184001.
- Danielson, Daine L., Gautam Satishchandran, and Robert M. Wald, 2022, “Gravitationally mediated entanglement: Newtonian field versus gravitons,” *Phys. Rev. D* **105**, 086001.
- Danilishin, Stefan L., and Farid Ya. Khalili, 2012, “Quantum measurement theory in gravitational-wave detectors,” *Living Rev. Relativity* **15**, 1–147.
- Das, Ashmita, Saurya Das, Noor R. Mansour, and Elias C. Vagenas, 2021, “Bounds on GUP parameters from GW150914 and GW190521,” *Phys. Lett. B* **819**, 136429.
- Das, Saurya, and Elias C. Vagenas, 2008, “Universality of Quantum Gravity Corrections,” *Phys. Rev. Lett.* **101**, 221301.
- Datta, Animesh, and Haixing Miao, 2021, “Signatures of the quantum nature of gravity in the differential motion of two masses,” *Quantum Sci. Technol.* **6**, 045014.
- Davies, P. C. W., and P. G. Falkowski, 1982, “Quantum theory and the equivalence principle,” *Proc. R. Soc. A* **381**, 469–478.
- Davies, Paul C. W., 1975, “Scalar production in Schwarzschild and Rindler metrics,” *J. Phys. A* **8**, 609.
- De Bonis, S. L., C. Urgell, W. Yang, C. Samanta, Adrien Noury, J. Vergara-Cruz, Q. Dong, Y. Jin, and Adrian Bachtold, 2018, “Ultrasensitive displacement noise measurement of carbon nanotube mechanical resonators,” *Nano Lett.* **18**, 5324–5328.
- de la Hamette, Anne-Catherine, Viktoria Kabel, Marios Christodoulou, and Časlav Brukner, 2022, “Quantum diffeomorphisms cannot make indefinite causal order definite,” *arXiv*: 2211.15685.
- de Lépinay, Laure Mercier, Caspar F. Ockeloen-Korppi, Daniel Malz, and Mika A. Sillanpää, 2020, “Nonreciprocal Transport Based on Cavity Floquet Modes in Optomechanics,” *Phys. Rev. Lett.* **125**, 023603.
- de Lépinay, Laure Mercier, Caspar F. Ockeloen-Korppi, Matthew J. Woolley, and Mika A. Sillanpää, 2021, “Quantum mechanics-free subsystem with mechanical oscillators,” *Science* **372**, 625–629.
- Delić, Uroš, Manuel Reisenbauer, Kahan Dare, David Grass, Vladan Vuletić, Nikolai Kiesel, and Markus Aspelmeyer, 2020, “Cooling of a levitated nanoparticle to the motional quantum ground state,” *Science* **367**, 892–895.

- Dellantonio, Luca, Oleksandr Kyriienko, Florian Marquardt, and Anders S. Sørensen, 2018, “Quantum nondemolition measurement of mechanical motion quanta,” *Nat. Commun.* **9**, 3621.
- De Lorenzo, L. A., and K. C. Schwab, 2014, “Superfluid optomechanics: Coupling of a superfluid to a superconducting condensate,” *New J. Phys.* **16**, 113020.
- del Pino, Javier, Jesse J. Slim, and Ewold Verhagen, 2022, “Non-Hermitian chiral phononics through optomechanically induced squeezing,” *Nature (London)* **606**, 82–87.
- Dias, Mafalda, Jonathan Frazer, Ander Retolaza, and Alexander Westphal, 2019, “Primordial gravitational waves and the swampland,” *Fortschr. Phys.* **67**, 1800063.
- Dienes, Keith R., 1997, “String theory and the path to unification: A review of recent developments,” *Phys. Rep.* **287**, 447–525.
- Dimopoulos, Savas, and Andrew A. Geraci, 2003, “Probing sub-micron forces by interferometry of Bose-Einstein condensed atoms,” *Phys. Rev. D* **68**, 124021.
- Dimopoulos, Savas, Peter W. Graham, Jason M. Hogan, and Mark A. Kasevich, 2007, “Testing General Relativity with Atom Interferometry,” *Phys. Rev. Lett.* **98**, 111102.
- Dimopoulos, Savas, Peter W. Graham, Jason M. Hogan, and Mark A. Kasevich, 2008, “General relativistic effects in atom interferometry,” *Phys. Rev. D* **78**, 042003.
- Dimopoulos, Savas, and Greg Landsberg, 2001, “Black Holes at the Large Hadron Collider,” *Phys. Rev. Lett.* **87**, 161602.
- Diósi, L., 1984, “Gravitation and quantum-mechanical localization of macro-objects,” *Phys. Lett.* **105A**, 199–202.
- Diósi, L., 1987, “A universal master equation for the gravitational violation of quantum mechanics,” *Phys. Lett. A* **120**, 377–381.
- Diósi, L., and B. Lukács, 1993, “Calculation of x-ray signals from Károlyházy hazy space-time,” *Phys. Lett. A* **181**, 366–368.
- Diósi, Lajos, 1989, “Models for universal reduction of macroscopic quantum fluctuations,” *Phys. Rev. A* **40**, 1165.
- Diósi, Lajos, 2005, “Intrinsic time-uncertainties and decoherence: Comparison of 4 models,” *Braz. J. Phys.* **35**, 260–265.
- Diósi, Lajos, 2007, “Notes on certain Newton gravity mechanisms of wavefunction localization and decoherence,” *J. Phys. A* **40**, 2989–2995.
- Diósi, Lajos, 2018, “How to teach and think about spontaneous wave function collapse theories: Not like before,” in *Collapse of the Wave Function: Models, Ontology, Origin, and Implications*, edited by Shan Gao (Cambridge University Press, Cambridge, England), pp. 3–11.
- Diósi, Lajos, and Jonathan J. Halliwell, 1998, “Coupling Classical and Quantum Variables Using Continuous Quantum Measurement Theory,” *Phys. Rev. Lett.* **81**, 2846.
- Doherty, Andrew C., Salman Habib, Kurt Jacobs, Hideo Mabuchi, and Sze M. Tan, 2000, “Quantum feedback control and classical control theory,” *Phys. Rev. A* **62**, 012105.
- Donadi, Sandro, Kristian Piscicchia, Catalina Curceanu, Lajos Diósi, Matthias Laubenstein, and Angelo Bassi, 2021, “Underground test of gravity-related wave function collapse,” *Nat. Phys.* **17**, 74–78.
- Donoghue, John F., 1994, “General relativity as an effective field theory: The leading quantum corrections,” *Phys. Rev. D* **50**, 3874.
- Dowling, Jonathan P., 1998, “Correlated input-port, matter-wave interferometer: Quantum-noise limits to the atom-laser gyroscope,” *Phys. Rev. A* **57**, 4736.
- Drinkwater, Mark, Rune Floberghagen, R. Haagsmans, D. Muzi, and A. Popescu, 2003, “GOCE: ESA’s first Earth Explorer core mission,” *Space Sci. Rev.* **108**, 419–432.
- Du, Haoxing, and Robert B. Mann, 2021, “Fisher information as a probe of spacetime structure: Relativistic quantum metrology in (A)dS,” *J. High Energy Phys.* **05**, 112.
- Duan, Xiao-Chun, Xiao-Bing Deng, Min-Kang Zhou, K. e. Zhang, Wen-Jie Xu, Feng Xiong, Yao-Yao Xu, Cheng-Gang Shao, Jun Luo, and Zhong-Kun Hu, 2016, “Test of the Universality of Free Fall with Atoms in Different Spin Orientations,” *Phys. Rev. Lett.* **117**, 023001.
- Dür, W., M. Skotiniotis, F. Fröwis, and B. Kraus, 2014, “Improved Quantum Metrology Using Quantum Error Correction,” *Phys. Rev. Lett.* **112**, 080801.
- Dür, Wolfgang, Christoph Simon, and J. Ignacio Cirac, 2002, “Effective Size of Certain Macroscopic Quantum Superpositions,” *Phys. Rev. Lett.* **89**, 210402.
- Eichler, C., Y. Salathe, J. Mlynek, S. Schmidt, and A. Wallraff, 2014, “Quantum-Limited Amplification and Entanglement in Coupled Nonlinear Resonators,” *Phys. Rev. Lett.* **113**, 110502.
- Elahi, Shafaq Gulzar, and Anupam Mazumdar, 2023, “Probing massless and massive gravitons via entanglement in a warped extra dimension,” *arXiv:2303.07371*.
- Elste, Florian, S. M. Girvin, and A. A. Clerk, 2009, “Quantum Noise Interference and Backaction Cooling in Cavity Nanomechanics,” *Phys. Rev. Lett.* **102**, 207209.
- Ernzer, Maryse, Manel Bosch Aguilera, Matteo Brunelli, Gian-Luca Schmid, Christoph Bruder, Patrick P. Potts, and Philipp Treutlein, 2023, “Optical Coherent Feedback Control of a Mechanical Oscillator,” *Phys. Rev. X* **13**, 021023.
- Escher, B. M., R. L. de Matos Filho, and Luiz Davidovich, 2011, “General framework for estimating the ultimate precision limit in noisy quantum-enhanced metrology,” *Nat. Phys.* **7**, 406–411.
- Esteve, J., C. Gross, A. Weller, S. Giovanazzi, and M. K. Oberthaler, 2008, “Squeezing and entanglement in a Bose-Einstein condensate,” *Nature (London)* **455**, 1216–1219.
- Etezad-Razavi, Saba, and Lucien Hardy, 2023, “Paradox with phase-coupled interferometers,” *arXiv:2305.14241*.
- Everitt, C. W. Francis, *et al.*, 2011, “Gravity Probe B: Final Results of a Space Experiment to Test General Relativity,” *Phys. Rev. Lett.* **106**, 221101.
- Fagnocchi, S., S. Finazzi, S. Liberati, M. Kormos, and A. Trombettoni, 2010, “Relativistic Bose-Einstein condensates: A new system for analogue models of gravity,” *New J. Phys.* **12**, 095012.
- Famaey, Benoît, and Stacy S. McGaugh, 2012, “Modified Newtonian dynamics (MOND): Observational phenomenology and relativistic extensions,” *Living Rev. Relativity* **15**, 1–159.
- Fein, Yaakov Y., Philipp Geyer, Patrick Zwick, Filip Kiałka, Sebastian Pedalino, Marcel Mayor, Stefan Gerlich, and Markus Arndt, 2019, “Quantum superposition of molecules beyond 25 kDa,” *Nat. Phys.* **15**, 1242–1245.
- Fermi, Enrico, *et al.*, 1936, “Sul moto dei neutroni nelle sostanze idrogenate [Motion of neutrons in hydrogenous substances],” *Ric. Sci.* **7**, 13–52.
- Fernandez-Melendez, Hector A., Alexander Belyaev, Vahe Gurzadyan, and Ivette Fuentes, 2024, “Probing Lambda-gravity with Bose-Einstein condensate,” *arXiv:2409.19755*.
- Ferrari, G., N. Poli, F. Sorrentino, and G. M. Tino, 2006, “Long-Lived Bloch Oscillations with Bosonic Sr Atoms and Application to Gravity Measurement at the Micrometer Scale,” *Phys. Rev. Lett.* **97**, 060402.
- Ferrari, V., G. Pizzella, M. Lee, and J. Weber, 1982, “Search for correlations between the University of Maryland and the University of Rome gravitational radiation antennas,” *Phys. Rev. D* **25**, 2471–2486.
- Fewster, Christopher J., and Rainer Verch, 2020, “Quantum fields and local measurements,” *Commun. Math. Phys.* **378**, 851–889.



- Feynman, R. P., 2011, in *The Role of Gravitation in Physics: Report from the 1957 Chapel Hill Conference*, edited by Cécile M. DeWitt and Dean Rickles (Max-Planck-Gesellschaft zur Förderung der Wissenschaften, Berlin).
- Forstner, Stefan, Magdalena Zych, Sahar Basiri-Esfahani, Kiran E. Khosla, and Warwick P. Bowen, 2020, “Nanomechanical test of quantum linearity,” *Optica* **7**, 1427–1434.
- Fragkos, Vasileios, Michael Kopp, and Igor Pikovski, 2022, “On inference of quantization from gravitationally induced entanglement,” *AVS Quantum Sci.* **4**, 045601.
- Fragolino, Paolo, Martine Schut, Marko Toroš, Sougato Bose, and Anupam Mazumdar, 2023, “Decoherence of a matter-wave interferometer due to dipole-dipole interactions,” *arXiv:2307.07001*.
- Friis, N., A. R. Lee, K. Truong, C. Sabín, E. Solano, G. Johansson, and I. Fuentes, 2013, “Relativistic Quantum Teleportation with Superconducting Circuits,” *Phys. Rev. Lett.* **110**, 113602.
- Frimmer, Martin, Karol Luszcz, Sandra Ferreiro, Vijay Jain, Erik Hebestreit, and Lukas Novotny, 2017, “Controlling the net charge on a nanoparticle optically levitated in vacuum,” *Phys. Rev. A* **95**, 061801.
- Fuchs, Tim M., Dennis G. Uitenbroek, Jaimy Plugge, Noud van Halteren, Jean-Paul van Soest, Andrea Vinante, Hendrik Ulbricht, and Tjerk H. Oosterkamp, 2024, “Measuring gravity with milligram levitated masses,” *Sci. Adv.* **10**, eadk2949.
- Fuentes, Ivette, Robert B. Mann, Eduardo Martín-Martínez, and Shahpoor Moradi, 2010, “Entanglement of Dirac fields in an expanding spacetime,” *Phys. Rev. D* **82**, 045030.
- Fuentes, Ivette, and Roger Penrose, 2018, “Quantum state reduction via gravity, and possible tests using Bose-Einstein condensates,” in *Collapse of the Wave Function*, edited by Shan Gao (Cambridge University Press, Cambridge, England), pp. 187–206.
- Fuentes, Ivette, Dennis Rätzel, Tupac Bravo Ibarra, and David E. Bruschi, 2019, “Quantum gravimeters and gradiometers,” European Patent No. 20734280.9 and U.S. Patent No. US-2022-0171089-A1.
- Fuentes-Schuller, Ivette, and Robert B. Mann, 2005, “Alice Falls into a Black Hole: Entanglement in Noninertial Frames,” *Phys. Rev. Lett.* **95**, 120404.
- Fulling, S. A., and P. C. W. Davies, 1976, “Radiation from a moving mirror in two dimensional space-time: Conformal anomaly,” *Proc. R. Soc. A* **348**, 393–414.
- Fulling, Stephen A., 1973, “Nonuniqueness of canonical field quantization in Riemannian space-time,” *Phys. Rev. D* **7**, 2850.
- Fulling, Stephen A., 1989, *Aspects of Quantum Field Theory in Curved Spacetime*, London Mathematical Society Student Texts Vol. 17 (Cambridge University Press, Cambridge, England).
- Gallerati, A., G. Modanese, and G. A. Ummary, 2022, “Interaction between macroscopic quantum systems and gravity,” *Front. Phys.* **10**, 941858.
- Galley, Thomas D., Flaminia Giacomini, and John H. Selby, 2022, “A no-go theorem on the nature of the gravitational field beyond quantum theory,” *Quantum* **6**, 779.
- Galley, Thomas D., Flaminia Giacomini, and John H. Selby, 2023, “Any consistent coupling between classical gravity and quantum matter is fundamentally irreversible,” *arXiv:2301.10261*.
- Garay, Luis J., 1995, “Quantum gravity and minimum length,” *Int. J. Mod. Phys. A* **10**, 145–165.
- Gardiner, C. W., and M. J. Collett, 1985, “Input and output in damped quantum systems: Quantum stochastic differential equations and the master equation,” *Phys. Rev. A* **31**, 3761–3774.
- Gardiner, C. W., and P. Zoller, 2000, *Quantum Noise* (Springer, New York).
- Gasbarri, Giulio, Marko Toroš, Sandro Donadi, and Angelo Bassi, 2017, “Gravity induced wave function collapse,” *Phys. Rev. D* **96**, 104013.
- Gely, Mario F., and Gary A. Steele, 2021, “Superconducting electro-mechanics to test Diósi-Penrose effects of general relativity in massive superpositions,” *AVS Quantum Sci.* **3**, 035601.
- Genoni, Marco G., Jinglei Zhang, James Millen, Peter F. Barker, and Alessio Serafini, 2015, “Quantum cooling and squeezing of a levitating nanosphere via time-continuous measurements,” *New J. Phys.* **17**, 073019.
- Geraci, Andrew, and Hart Goldman, 2015, “Sensing short range forces with a nanosphere matter-wave interferometer,” *Phys. Rev. D* **92**, 062002.
- Geraci, Andrew A., Scott B. Papp, and John Kitching, 2010, “Short-Range Force Detection Using Optically Cooled Levitated Microspheres,” *Phys. Rev. Lett.* **105**, 101101.
- Geraci, Andrew A., Sylvia J. Smullin, David M. Weld, John Chiaverini, and Aharon Kapitlnik, 2008, “Improved constraints on non-Newtonian forces at 10 microns,” *Phys. Rev. D* **78**, 022002.
- Ghirardi, Gian Carlo, Philip Pearle, and Alberto Rimini, 1990, “Markov processes in Hilbert space and continuous spontaneous localization of systems of identical particles,” *Phys. Rev. A* **42**, 78.
- Ghirardi, Gian Carlo, Alberto Rimini, and Tullio Weber, 1986, “Unified dynamics for microscopic and macroscopic systems,” *Phys. Rev. D* **34**, 470.
- Ghirardi, GianCarlo, Renata Grassi, and Alberto Rimini, 1990, “Continuous-spontaneous-reduction model involving gravity,” *Phys. Rev. A* **42**, 1057.
- Ghosal, Pratik, Arkaprabha Ghosal, and Somshubhro Bandyopadhyay, 2023, “Distribution of quantum gravity induced entanglement in many-body systems,” *arXiv:2311.08291*.
- Giacomini, Flaminia, and Časlav Brukner, 2022, “Quantum superposition of spacetimes obeys Einstein’s equivalence principle,” *AVS Quantum Sci.* **4**, 015601.
- Giacomini, Flaminia, Esteban Castro-Ruiz, and Časlav Brukner, 2019, “Quantum mechanics and the covariance of physical laws in quantum reference frames,” *Nat. Commun.* **10**, 494.
- Giovannetti, Vittorio, Seth Lloyd, and Lorenzo Maccone, 2011, “Advances in quantum metrology,” *Nat. Photonics* **5**, 222–229.
- Giovannetti, Vittorio, Seth Lloyd, and Lorenzo Maccone, 2023, “Geometric event-based quantum mechanics,” *New J. Phys.* **25**, 023027.
- Girdhar, Parth, and Andrew C. Doherty, 2020, “Testing generalised uncertainty principles through quantum noise,” *New J. Phys.* **22**, 093073.
- Gisin, Nicolas, 1989, “Stochastic quantum dynamics and relativity,” *Helv. Phys. Acta* **62**, 363–371, <https://www.osti.gov/etdeweb/biblio/5663927>.
- Giulini, Domenico, 2000, “Decoherence: A dynamical approach to superselection rules?,” in *Relativistic Quantum Measurement and Decoherence*, Lecture Notes in Physics Vol. 559, edited by Heinz-Peter Breuer and Francesco Petruccione (Springer, New York), pp. 67–92.
- Giulini, Domenico, and André Großardt, 2011, “Gravitationally induced inhibitions of dispersion according to the Schrödinger-Newton equation,” *Classical Quantum Gravity* **28**, 195026.
- Giulini, Domenico, and André Großardt, 2012, “The Schrödinger-Newton equation as a non-relativistic limit of self-gravitating Klein-Gordon and Dirac fields,” *Classical Quantum Gravity* **29**, 215010.
- Goodkind, John M., 1999, “The superconducting gravimeter,” *Rev. Sci. Instrum.* **70**, 4131–4152.

- Gorini, Vittorio, Andrzej Kossakowski, and Ennackal Chandy George Sudarshan, 1976, “Completely positive dynamical semi-groups of  $N$ -level systems,” *Rep. Math. Phys.* **17**, 821–825.
- Gosling, Jonathan M. H., Markus Rademacher, Jence T. Mulder, Arjan J. Houtepen, Marko Toroš, A. T. M. Anishur Rahman, Antonio Pontin, and P. F. Barker, 2024, “Levitodynamic spectroscopy for single nanoparticle characterisation,” [arXiv:2401.11551](#).
- Greenberger, Daniel, 1968, “The role of equivalence in quantum mechanics,” *Ann. Phys. (N.Y.)* **47**, 116–126.
- Greenberger, Daniel M., 1983, “The neutron interferometer as a device for illustrating the strange behavior of quantum systems,” *Rev. Mod. Phys.* **55**, 875–905.
- Griffiths, David J., and Darrell F. Schroeter, 2018, *Introduction to Quantum Mechanics* (Cambridge University Press, Cambridge, England).
- Gröblacher, Simon, Klemens Hammerer, Michael R. Vanner, and Markus Aspelmeyer, 2009, “Observation of strong coupling between a micromechanical resonator and an optical cavity field,” *Nature (London)* **460**, 724–727.
- Gröblacher, Simon, Jared B. Hertzberg, Michael R. Vanner, Garrett D. Cole, Sylvain Gigan, K. C. Schwab, and Markus Aspelmeyer, 2009, “Demonstration of an ultracold micro-optomechanical oscillator in a cryogenic cavity,” *Nat. Phys.* **5**, 485–488.
- Gross, Eugene P., 1961, “Structure of a quantized vortex in boson systems,” *Nuovo Cimento* **20**, 454–477.
- Großardt, André, James Bateman, Hendrik Ulbricht, and Angelo Bassi, 2016, “Optomechanical test of the Schrödinger-Newton equation,” *Phys. Rev. D* **93**, 096003.
- Guérin, Philippe Allard, and Časlav Brukner, 2018, “Observer-dependent locality of quantum events,” *New J. Phys.* **20**, 103031.
- Guff, Thomas, Nicolas Boulle, and Igor Pikovski, 2022, “Optimal fidelity witnesses for gravitational entanglement,” *Phys. Rev. A* **105**, 022444.
- Gundlach, Jens H., and Stephen M. Merkowitz, 2000, “Measurement of Newton’s Constant Using a Torsion Balance with Angular Acceleration Feedback,” *Phys. Rev. Lett.* **85**, 2869–2872.
- Gunnink, Fabian, Anupam Mazumdar, Martine Schut, and Marko Toroš, 2022, “Gravitational decoherence by the apparatus in the quantum-gravity induced entanglement of masses,” [arXiv:2210.16919](#).
- Guo, Jingkun, and Simon Gröblacher, 2022, “Coherent feedback in optomechanical systems in the sideband-unresolved regime,” *Quantum* **6**, 848.
- Guo, Jingkun, Richard Norte, and Simon Gröblacher, 2019, “Feedback Cooling of a Room Temperature Mechanical Oscillator Close to Its Motional Ground State,” *Phys. Rev. Lett.* **123**, 223602.
- Gupta, Suraj N., 1950, “Theory of longitudinal photons in quantum electrodynamics,” *Proc. Phys. Soc. London Sect. A* **63**, 681–691.
- Gupta, Suraj N., 1952a, “Quantization of Einstein’s gravitational field: General treatment,” *Proc. Phys. Soc. London Sect. A* **65**, 608.
- Gupta, Suraj N., 1952b, “Quantization of Einstein’s gravitational field: Linear approximation,” *Proc. Phys. Soc. London Sect. A* **65**, 161.
- Gutierrez Latorre, Martí, Gerard Higgins, Achintya Paradkar, Thilo Bauch, and Witlef Wieczorek, 2023, “Superconducting Microsphere Magnetically Levitated in an Anharmonic Potential with Integrated Magnetic Readout,” *Phys. Rev. Appl.* **19**, 054047.
- Hackermüller, Lucia, Klaus Hornberger, Björn Brezger, Anton Zeilinger, and Markus Arndt, 2004, “Decoherence of matter waves by thermal emission of radiation,” *Nature (London)* **427**, 711–714.
- Haddock, Christopher C., *et al.*, 2018, “Search for deviations from the inverse square law of gravity at nm range using a pulsed neutron beam,” *Phys. Rev. D* **97**, 062002.
- Hadjar, Y., P. F. Cohadon, C. G. Aminoff, M. Pinard, and A. Heidmann, 1999, “High-sensitivity optical measurement of mechanical Brownian motion,” *Europhys. Lett.* **47**, 545.
- Haine, Simon A., 2021, “Searching for signatures of quantum gravity in quantum gases,” *New J. Phys.* **23**, 033020.
- Hamerly, Ryan, and Hideo Mabuchi, 2012, “Advantages of Coherent Feedback for Cooling Quantum Oscillators,” *Phys. Rev. Lett.* **109**, 173602.
- Hamerly, Ryan, and Hideo Mabuchi, 2013, “Coherent controllers for optical-feedback cooling of quantum oscillators,” *Phys. Rev. A* **87**, 013815.
- Hanif, Farhan, Debarshi Das, Jonathan Halliwell, Dipankar Home, Anupam Mazumdar, Hendrik Ulbricht, and Sougato Bose, 2023, “Testing whether gravity acts as a quantum entity when measured,” [arXiv:2307.08133](#).
- Hao, Xiang, and Yinzong Wu, 2016, “Quantum parameter estimation in the Unruh-DeWitt detector model,” *Ann. Phys. (Amsterdam)* **372**, 110–118.
- Harber, D. M., J. M. Obrecht, J. M. McGuirk, and E. A. Cornell, 2005, “Measurement of the Casimir-Polder force through center-of-mass oscillations of a Bose-Einstein condensate,” *Phys. Rev. A* **72**, 033610.
- Hardman, K. S., P. J. Everitt, G. D. McDonald, P. Manju, P. B. Wigley, M. A. Sooriyabandara, C. C. N. Kuhn, J. E. Debs, J. D. Close, and N. P. Robins, 2016, “Simultaneous Precision Gravimetry and Magnetic Gradiometry with a Bose-Einstein Condensate: A High Precision, Quantum Sensor,” *Phys. Rev. Lett.* **117**, 138501.
- Hardy, Lucien, 2018, “The construction interpretation: Conceptual roads to quantum gravity,” [arXiv:1807.10980](#).
- Hartley, Daniel, Christian Käding, Richard Howl, and Ivette Fuentes, 2019a, “Quantum simulation of dark energy candidates,” *Phys. Rev. D* **99**, 105002.
- Hartley, Daniel, Christian Käding, Richard Howl, and Ivette Fuentes, 2024b, “Quantum-enhanced screened dark energy detection,” *Eur. Phys. J. C* **84**, 49.
- Harwood, Alfred, Matteo Brunelli, and Alessio Serafini, 2021, “Cavity optomechanics assisted by optical coherent feedback,” *Phys. Rev. A* **103**, 023509.
- Hauss, Julian, Arkady Fedorov, Stephan André, Valentina Brosco, Carsten Hutter, Robin Kothari, Sunil Yeshwanth, Alexander Shnirman, and Gerd Schön, 2008, “Dissipation in circuit quantum electrodynamics: Lasing and cooling of a low-frequency oscillator,” *New J. Phys.* **10**, 095018.
- Hawking, Stephen W., 1974, “Black hole explosions?,” *Nature (London)* **248**, 30–31.
- Hawking, Stephen W., 1975, “Particle creation by black holes,” *Commun. Math. Phys.* **43**, 199–220.
- Hees, Aurélien, Olivier Minazzoli, Etienne Savalle, Yevgeny V. Stadnik, and Peter Wolf, 2018, “Violation of the equivalence principle from light scalar dark matter,” *Phys. Rev. D* **98**, 064051.
- Heikkilä, Tero T., Francesco Massel, Jani Tuorila, Raphaël Khan, and Mika A. Sillanpää, 2014, “Enhancing Optomechanical Coupling via the Josephson Effect,” *Phys. Rev. Lett.* **112**, 203603.
- Helou, Bassam, B. J. J. Slagmolen, David E. McClelland, and Yanbei Chen, 2017, “LISA Pathfinder appreciably constrains collapse models,” *Phys. Rev. D* **95**, 084054.
- Helstrom, Carl W., 1967, “Minimum mean-squared error of estimates in quantum statistics,” *Phys. Lett.* **25A**, 101–102.
- Helstrom, Carl W., 1969, “Quantum detection and estimation theory,” *J. Stat. Phys.* **1**, 231–252.
- Hempston, David, Jamie Vovrosh, Marko Toroš, George Winstone, Muddassar Rashid, and Hendrik Ulbricht, 2017, “Force sensing

- with an optically levitated charged nanoparticle,” *Appl. Phys. Lett.* **111**, 133111.
- Henkel, Carsten, and Ron Folman, 2022, “Internal decoherence in nano-object interferometry due to phonons,” *AVS Quantum Sci.* **4**, 025602.
- Henkel, Carsten, and Ron Folman, 2023, “Universal limit on spatial quantum superpositions with massive objects due to phonons,” *arXiv:2305.15230*.
- Hensen, Bas, *et al.*, 2015, “Loophole-free Bell inequality violation using electron spins separated by 1.3 kilometres,” *Nature (London)* **526**, 682–686.
- Hoang, Thai M., Jonghoon Ahn, Jaehoon Bang, and Tongcang Li, 2016, “Electron spin control of optically levitated nanodiamonds in vacuum,” *Nat. Commun.* **7**, 12250.
- Hofer, J., *et al.*, 2023, “High- $q$  Magnetic Levitation and Control of Superconducting Microspheres at Millikelvin Temperatures,” *Phys. Rev. Lett.* **131**, 043603.
- Hoff, Ulrich B., Johann Kollath-Bönig, Jonas S. Neergaard-Nielsen, and Ulrik L. Andersen, 2016, “Measurement-Induced Macroscopic Superposition States in Cavity Optomechanics,” *Phys. Rev. Lett.* **117**, 143601.
- Hohensee, Michael A., Brian Estey, Paul Hamilton, Anton Zeilinger, and Holger Müller, 2012, “Force-Free Gravitational Redshift: Proposed Gravitational Aharonov-Bohm Experiment,” *Phys. Rev. Lett.* **108**, 230404.
- Holevo, Alexander S., 2011, *Probabilistic and Statistical Aspects of Quantum Theory*, Vol. 1 (Springer, New York).
- Hollands, Stefan, and Robert M. Wald, 2015, “Quantum fields in curved spacetime,” *Phys. Rep.* **574**, 1–35.
- Holstein, Barry R., 2017, “Analytical on-shell calculation of low energy higher order scattering,” *J. Phys. G* **44**, 01LT01.
- Holstein, Barry R., and Andreas Ross, 2008, “Spin effects in long range gravitational scattering,” *arXiv:0802.0716*.
- Hornberger, Klaus, Stefan Gerlich, Philipp Haslinger, Stefan Nimmrichter, and Markus Arndt, 2012, “Colloquium: Quantum interference of clusters and molecules,” *Rev. Mod. Phys.* **84**, 157–173.
- Hornberger, Klaus, Stefan Uttenthaler, Björn Brezger, Lucia Hackermüller, Markus Arndt, and Anton Zeilinger, 2003, “Collisional Decoherence Observed in Matter Wave Interferometry,” *Phys. Rev. Lett.* **90**, 160401.
- Hossenfelder, Sabine, 2013, “Minimal length scale scenarios for quantum gravity,” *Living Rev. Relativity* **16**, 1–90.
- Hosten, Onur, 2022, “Constraints on probing quantum coherence to infer gravitational entanglement,” *Phys. Rev. Res.* **4**, 013023.
- Howl, R., and I. Fuentes, 2023, “Quantum frequency interferometry: With applications ranging from gravitational wave detection to dark matter searches,” *AVS Quantum Sci.* **5**, 014402.
- Howl, Richard, Roger Penrose, and Ivette Fuentes, 2019, “Exploring the unification of quantum theory and general relativity with a Bose-Einstein condensate,” *New J. Phys.* **21**, 043047.
- Howl, Richard, Vlatko Vedral, Devang Naik, Marios Christodoulou, Carlo Rovelli, and Aditya Iyer, 2021, “Non-Gaussianity as a signature of a quantum theory of gravity,” *PRX Quantum* **2**, 010325.
- Hu, Bei Lok, Juan Pablo Paz, and Yuhong Zhang, 1992, “Quantum Brownian motion in a general environment: Exact master equation with nonlocal dissipation and colored noise,” *Phys. Rev. D* **45**, 2843.
- Hu, Bei Lok, and Enric Verdager, 2008, “Stochastic gravity: Theory and applications,” *Living Rev. Relativity* **11**, 1–112.
- Hu, Dan, Shang-Yu Huang, Jie-Qiao Liao, Lin Tian, and Hsi-Sheng Goan, 2015, “Quantum coherence in ultrastrong optomechanics,” *Phys. Rev. A* **91**, 013812.
- Huang, Sumei, and G. S. Agarwal, 2017, “Robust force sensing for a free particle in a dissipative optomechanical system with a parametric amplifier,” *Phys. Rev. A* **95**, 023844.
- Hui, Lam, Alberto Nicolis, and Christopher W. Stubbs, 2009, “Equivalence principle implications of modified gravity models,” *Phys. Rev. D* **80**, 104002.
- Huimann, Stephan, 2020, “The quantum harmonic oscillator under the influence of gravity,” master’s thesis (University of Vienna).
- Iakovleva, Tatiana, Bijita Sarma, and Jason Twamley, 2023, “Zeptometer displacement sensing using cavity opto-magneto-mechanics,” *arXiv:2302.06795*.
- Ivanov, A. N., M. Wellenzohn, and H. Abele, 2021, “Quantum gravitational states of ultracold neutrons as a tool for probing of beyond-Riemann gravity,” *Phys. Lett. B* **822**, 136640.
- Jacobs, Kurt, 2014, *Quantum Measurement Theory and Its Applications* (Cambridge University Press, Cambridge, England).
- Jacobs, Kurt, and Daniel A. Steck, 2006, “A straightforward introduction to continuous quantum measurement,” *Contemp. Phys.* **47**, 279–303.
- Jacobs, Kurt, Xiaoting Wang, and Howard M. Wiseman, 2014, “Coherent feedback that beats all measurement-based feedback protocols,” *New J. Phys.* **16**, 073036.
- Jaehne, Konstanze, Klemens Hammerer, and Margareta Wallquist, 2008, “Ground-state cooling of a nanomechanical resonator via a Cooper-pair box qubit,” *New J. Phys.* **10**, 095019.
- Jaffe, Matt, Philipp Haslinger, Victoria Xu, Paul Hamilton, Amol Upadhye, Benjamin Elder, Justin Khoury, and Holger Müller, 2017, “Testing sub-gravitational forces on atoms from a miniature in-vacuum source mass,” *Nat. Phys.* **13**, 938–942.
- Jain, Vijay, Jan Gieseler, Clemens Moritz, Christoph Dellago, Romain Quidant, and Lukas Novotny, 2016, “Direct Measurement of Photon Recoil from a Levitated Nanoparticle,” *Phys. Rev. Lett.* **116**, 243601.
- Japha, Yonathan, and Ron Folman, 2022, “Role of rotations in Stern-Gerlach interferometry with massive objects,” *arXiv:2202.10535*.
- Jenke, T., *et al.*, 2014, “Gravity Resonance Spectroscopy Constrains Dark Energy and Dark Matter Scenarios,” *Phys. Rev. Lett.* **112**, 151105.
- Jenke, Tobias, David Stadler, Hartmut Abele, and Peter Geltenbort, 2009, “Q-BOUNCE—Experiments with quantum bouncing ultracold neutrons,” *Nucl. Instrum. Methods Phys. Res., Sect. A* **611**, 318–321.
- Jenke, Tobias, *et al.*, 2019, “Testing gravity at short distances: Gravity resonance spectroscopy with  $q$ BOUNCE,” *EPJ Web Conf.* **219**, 05003.
- Jing, Limei, and Jiliang Jing, 2023, “Bosonic entanglement generated by projective measurements in Einstein-Gauss-Bonnet black hole,” *Quantum Inf. Process.* **22**, 371.
- Jo, G.-B., Y. Shin, S. Will, T. A. Pasquini, M. Saba, W. Ketterle, D. E. Pritchard, M. Vengalattore, and M. Prentiss, 2007, “Long Phase Coherence Time and Number Squeezing of Two Bose-Einstein Condensates on an Atom Chip,” *Phys. Rev. Lett.* **98**, 030407.
- Johansson, J. Robert, Paul D. Nation, and Franco Nori, 2012, “QuTiP: An open-source Python framework for the dynamics of open quantum systems,” *Comput. Phys. Commun.* **183**, 1760–1772.
- Johansson, Mattias T., Gavin K. Brennen, and Jason Twamley, 2016, “Macroscopic superpositions and gravimetry with quantum magnetomechanics,” *Sci. Rep.* **6**, 1–13.



- Judson, Richard S., and Herschel Rabitz, 1992, “Teaching Lasers to Control Molecules,” *Phys. Rev. Lett.* **68**, 1500–1503.
- Juffmann, Thomas, Hendrik Ulbricht, and Markus Arndt, 2013, “Experimental methods of molecular matter-wave optics,” *Rep. Prog. Phys.* **76**, 086402.
- Kafri, D., J. M. Taylor, and G. J. Milburn, 2014, “A classical channel model for gravitational decoherence,” *New J. Phys.* **16**, 065020.
- Kafri, Dvir, and J. M. Taylor, 2013, “A noise inequality for classical forces,” [arXiv:1311.4558](#).
- Kaltenbaek, Rainer, *et al.*, 2022, “MAQRO—BPS 2023 research campaign whitepaper,” [arXiv:2202.01535](#).
- Kamba, M., H. Kiuchi, T. Yotsuya, and K. Aikawa, 2021, “Recoil-limited feedback cooling of single nanoparticles near the ground state in an optical lattice,” *Phys. Rev. A* **103**, L051701.
- Kamba, M., R. Shimizu, and K. Aikawa, 2023, “Nanoscale feedback control of six degrees of freedom of a near-sphere,” [arXiv:2303.02831](#).
- Kamba, Mitsuyoshi, Ryoga Shimizu, and Kiyotaka Aikawa, 2022, “Optical cold damping of neutral nanoparticles near the ground state in an optical lattice,” *Opt. Express* **30**, 26716–26727.
- Kamionkowski, Marc, and Ely D. Kovetz, 2016, “The quest for  $B$  modes from inflationary gravitational waves,” *Annu. Rev. Astron. Astrophys.* **54**, 227–269.
- Kamiya, Y., K. Itagaki, M. Tani, G. N. Kim, and S. Komamiya, 2015, “Constraints on New Gravitylike Forces in the Nanometer Range,” *Phys. Rev. Lett.* **114**, 161101.
- Kanari-Naish, Lydia A., Jack Clarke, Sofia Qvarfort, and Michael R. Vanner, 2022, “Two-mode Schrödinger-cat states with nonlinear optomechanics: Generation and verification of non-Gaussian mechanical entanglement,” *Quantum Sci. Technol.* **7**, 035012.
- Kanno, Sugumi, Jiro Soda, and Junsei Tokuda, 2021, “Noise and decoherence induced by gravitons,” *Phys. Rev. D* **103**, 044017.
- Kapner, D. J., T. S. Cook, E. G. Adelberger, J. H. Gundlach, B. R. Heckel, C. D. Hoyle, and H. E. Swanson, 2007, “Tests of the Gravitational Inverse-Square Law below the Dark-Energy Length Scale,” *Phys. Rev. Lett.* **98**, 021101.
- Károlyházy, Frederick, 1966, “Gravitation and quantum mechanics of macroscopic objects,” *Nuovo Cimento A* **42**, 390–402.
- Kasevich, Mark, and Steven Chu, 1991, “Atomic Interferometry Using Stimulated Raman Transitions,” *Phys. Rev. Lett.* **67**, 181–184.
- Kawana, Kiyoharu, and Daiki Ueda, 2019, “Amplification of gravitational motion via quantum weak measurement,” *Prog. Theor. Exp. Phys.* 041A01.
- Kempf, Achim, Gianpiero Mangano, and Robert B. Mann, 1995, “Hilbert space representation of the minimal length uncertainty relation,” *Phys. Rev. D* **52**, 1108–1118.
- Kent, Adrian, and Damián Pitalúa-García, 2021, “Testing the non-classicality of spacetime: What can we learn from Bell–Bose *et al.*-Marletto-Vedral experiments?” *Phys. Rev. D* **104**, 126030.
- Kessler, E. M., I. Lovchinsky, A. O. Sushkov, and M. D. Lukin, 2014, “Quantum Error Correction for Metrology,” *Phys. Rev. Lett.* **112**, 150802.
- Khintchine, A., 1934, “Korrelationstheorie der stationären stochastischen Prozesse,” *Math. Ann.* **109**, 604–615.
- Khosla, K. E., M. R. Vanner, W. P. Bowen, and G. J. Milburn, 2013, “Quantum state preparation of a mechanical resonator using an optomechanical geometric phase,” *New J. Phys.* **15**, 043025.
- Khosla, Kiran E., Michael R. Vanner, Natalia Ares, and Edward Alexander Laird, 2018, “Displacement Electromechanics: How to Detect Quantum Interference in a Nanomechanical Resonator,” *Phys. Rev. X* **8**, 021052.
- Khoury, Justin, and Amanda Weltman, 2004, “Chameleon cosmology,” *Phys. Rev. D* **69**, 044026.
- Khoury, Justin, and Amanda Weltman, 2004b, “Chameleon Fields: Awaiting Surprises for Tests of Gravity in Space,” *Phys. Rev. Lett.* **93**, 171104.
- Kibble, T. W. B., 1981, “Is a semi-classical theory of gravity viable?,” *Quantum Gravity 2: A Second Oxford Symposium*, edited by C. J. Isham, R. Penrose, and D. W. Sciama (Oxford University Press, New York), p. 63.
- Kilian, Eva, Marko Toroš, Frank F. Deppisch, Ruben Saakyan, and Sougato Bose, 2023, “Requirements on quantum superpositions of macro-objects for sensing neutrinos,” *Phys. Rev. Res.* **5**, 023012.
- Kilian, Eva, *et al.*, 2024, “Dark matter searches with levitated sensors,” *AVS Quantum Sci.* **6**, 030503.
- Kirilil, G. G., and I. B. Khriplovich, 2002, “Quantum power correction to the Newton law,” *J. Exp. Theor. Phys.* **95**, 981–986.
- Kirsten-Siemß, J.-N., F. Fitzek, C. Schubert, E. M. Rasel, N. Gaaloul, and K. Hammerer, 2023, “Large-Momentum-Transfer Atom Interferometers with  $\mu$ rad-Accuracy Using Bragg Diffraction,” *Phys. Rev. Lett.* **131**, 033602.
- Kleckner, Dustin, Igor Pikovski, Evan Jeffrey, Luuk Ament, Eric Eliel, Jeroen van den Brink, and Dirk Bouwmeester, 2008, “Creating and verifying a quantum superposition in a micro-optomechanical system,” *New J. Phys.* **10**, 095020.
- Knee, George C., G. Andrew D. Briggs, Simon C. Benjamin, and Erik M. Gauger, 2013, “Quantum sensors based on weak-value amplification cannot overcome decoherence,” *Phys. Rev. A* **87**, 012115.
- Koch, Florian, and Jan Carl Budich, 2022, “Quantum non-Hermitian topological sensors,” *Phys. Rev. Res.* **4**, 013113.
- Kohen, D., C. C. Marston, and D. J. Tannor, 1997, “Phase space approach to theories of quantum dissipation,” *J. Chem. Phys.* **107**, 5236–5253.
- Kohlrus, Jan, David Edward Bruschi, and Ivette Fuentes, 2019, “Quantum-metrology estimation of spacetime parameters of the Earth outperforming classical precision,” *Phys. Rev. A* **99**, 032350.
- Kohlrus, Jan, David Edward Bruschi, Jorma Louko, and Ivette Fuentes, 2017, “Quantum communications and quantum metrology in the spacetime of a rotating planet,” *Eur. Phys. J. Quantum Technol.* **4**, 7.
- Kok, Pieter, and Ulvi Yurtsever, 2003, “Gravitational decoherence,” *Phys. Rev. D* **68**, 085006.
- Kolkowitz, Shimon, Ania C. Bleszynski Jayich, Quirin P. Unterreithmeier, Steven D. Bennett, Peter Rabl, J. G. E. Harris, and Mikhail D. Lukin, 2012, “Coherent sensing of a mechanical resonator with a single-spin qubit,” *Science* **335**, 1603–1606.
- Komatsu, Eiichiro, 2010, “Hunting for primordial non-Gaussianity in the cosmic microwave background,” *Classical Quantum Gravity* **27**, 124010.
- Kononchuk, Rodion, Jizhe Cai, Fred Ellis, Ramathanas Thevamaran, and Tsampikos Kottos, 2022, “Exceptional-point-based accelerometers with enhanced signal-to-noise ratio,” *Nature (London)* **607**, 697–702.
- Kounalakis, Marios, Yaroslav M. Blanter, and Gary A. Steele, 2020, “Flux-mediated optomechanics with a transmon qubit in the single-photon ultrastrong-coupling regime,” *Phys. Rev. Res.* **2**, 023335.
- Kovachy, T., P. Asenbaum, C. Overstreet, C. A. Donnelly, S. M. Dickerson, A. Sugarbaker, J. M. Hogan, and M. A. Kasevich, 2015, “Quantum superposition at the half-metre scale,” *Nature (London)* **528**, 530–533.
- Kreuz, M., *et al.*, 2009, “A method to measure the resonance transitions between the gravitationally bound quantum states of

- neutrons in the GRANIT spectrometer,” *Nucl. Instrum. Methods Phys. Res., Sect. A* **611**, 326–330.
- Krisnanda, Tanjung, Guo Yao Tham, Mauro Paternostro, and Tomasz Paterek, 2020, “Observable quantum entanglement due to gravity,” *npj Quantum Inf.* **6**, 1–6.
- Krisnanda, Tanjung, Margherita Zuppardo, Mauro Paternostro, and Tomasz Paterek, 2017, “Revealing Nonclassicality of Inaccessible Objects,” *Phys. Rev. Lett.* **119**, 120402.
- Kumar, Shreya P., and Martin B. Plenio, 2018, “Quantum-optical tests of Planck-scale physics,” *Phys. Rev. A* **97**, 063855.
- Kumar, Shreya P., and Martin B. Plenio, 2020, “On quantum gravity tests with composite particles,” *Nat. Commun.* **11**, 3900.
- Kwiat, Paul G., Klaus Mattle, Harald Weinfurter, Anton Zeilinger, Alexander V. Sergienko, and Yanhua Shih, 1995, “New High-Intensity Source of Polarization-Entangled Photon Pairs,” *Phys. Rev. Lett.* **75**, 4337–4341.
- Lagouvardos, Michalis, and Charis Anastopoulos, 2021, “Gravitational decoherence of photons,” *Classical Quantum Gravity* **38**, 115012.
- Lahav, Oren, Amir Itah, Alex Blumkin, Carmit Gordon, Shahar Rinott, Alona Zayats, and Jeff Steinhauer, 2010, “Realization of a Sonic Black Hole Analog in a Bose-Einstein Condensate,” *Phys. Rev. Lett.* **105**, 240401.
- Lami, Ludovico, Julien S. Pedernales, and Martin B. Plenio, 2023, “Testing the quantumness of gravity without entanglement,” [arXiv:2302.03075](https://arxiv.org/abs/2302.03075).
- Lamine, B., R. Hervé, A. Lambrecht, and S. Reynaud, 2006, “Decoherence induced by stochastic background of gravitational waves on matter-wave interferometers,” *Appl. Phys. B* **84**, 575–578.
- Lämmerzahl, Claus, 1996, “On the equivalence principle in quantum theory,” *Gen. Relativ. Gravit.* **28**, 1043–1070.
- Lämmerzahl, Claus, 1998, “Minimal coupling and the equivalence principle in quantum mechanics,” *Acta Phys. Pol. B* **29**, 1057, <https://www.actaphys.uj.edu.pl/R/29/4/1057/pdf>.
- Lamporesi, G., A. Bertoldi, A. Cecchetti, B. Duhlach, M. Fattori, A. Malengo, S. Pettoroso, M. Prevedelli, and G.M. Tino, 2007, “Source mass and positioning system for an accurate measurement of  $G$ ,” *Rev. Sci. Instrum.* **78**, 075109.
- Latorre, Martí Gutierrez, Joachim Hofer, Matthias Rudolph, and Witlief Wieczorek, 2020, “Chip-based superconducting traps for levitation of micrometer-sized particles in the Meissner state,” *Supercond. Sci. Technol.* **33**, 105002.
- Lau, Hoi-Kwan, and Aashish A. Clerk, 2018, “Fundamental limits and non-reciprocal approaches in non-Hermitian quantum sensing,” *Nat. Commun.* **9**, 4320.
- Law, C.K., 1995, “Interaction between a moving mirror and radiation pressure: A Hamiltonian formulation,” *Phys. Rev. A* **51**, 2537.
- Lecocq, Florent, Jeremy B. Clark, Raymond W. Simmonds, Jose Aumentado, and John D. Teufel, 2015, “Quantum Nondemolition Measurement of a Nonclassical State of a Massive Object,” *Phys. Rev. X* **5**, 041037.
- Lee, Chang-Woo, and Hyunseok Jeong, 2011, “Quantification of Macroscopic Quantum Superpositions within Phase Space,” *Phys. Rev. Lett.* **106**, 220401.
- Lee, Chang-Woo, Jae Hoon Lee, Jaewoo Joo, and Hyojun Seok, 2022, “Quantum Fisher information of an optomechanical force sensor driven by a squeezed vacuum field,” *Opt. Express* **30**, 25249–25261.
- Lee, J.G., E. G. Adelberger, T. S. Cook, S. M. Fleischer, and B. R. Heckel, 2020, “New Test of the Gravitational  $1/r^2$  Law at Separations down to 52  $\mu\text{m}$ ,” *Phys. Rev. Lett.* **124**, 101101.
- Leggett, Anthony J., 1980, “Macroscopic quantum systems and the quantum theory of measurement,” *Prog. Theor. Phys. Suppl.* **69**, 80–100.
- Lei, C. U., A. J. Weinstein, J. Suh, E. E. Wollman, A. Kronwald, F. Marquardt, A. A. Clerk, and K. C. Schwab, 2016, “Quantum Nondemolition Measurement of a Quantum Squeezed State beyond the 3 dB Limit,” *Phys. Rev. Lett.* **117**, 100801.
- Leibfried, D., M. D. Barrett, T. Schaetz, J. Britton, J. Chiaverini, W. M. Itano, J. D. Jost, C. Langer, and D. J. Wineland, 2004, “Toward Heisenberg-limited spectroscopy with multiparticle entangled states,” *Science* **304**, 1476–1478.
- Leibfried, Dietrich, Rainer Blatt, Christopher Monroe, and David Wineland, 2003, “Quantum dynamics of single trapped ions,” *Rev. Mod. Phys.* **75**, 281.
- Lewandowski, Charles W., Tyler D. Knowles, Zachariah B. Etienne, and Brian D’Urso, 2021, “High-Sensitivity Accelerometry with a Feedback-Cooled Magnetically Levitated Microsphere,” *Phys. Rev. Appl.* **15**, 014050.
- Lewis, Zachary, and Tatsu Takeuchi, 2011, “Position and momentum uncertainties of the normal and inverted harmonic oscillators under the minimal length uncertainty relation,” *Phys. Rev. D* **84**, 105029.
- Li, Bei-Bei, Lingfeng Ou, Yuechen Lei, and Yong-Chun Liu, 2021, “Cavity optomechanical sensing,” *Nanophotonics* **10**, 2799–2832.
- Li, Jie, and Simon Gröblacher, 2020, “Stationary quantum entanglement between a massive mechanical membrane and a low frequency  $LC$  circuit,” *New J. Phys.* **22**, 063041.
- Li, Jie, and Simon Gröblacher, 2021, “Entangling the vibrational modes of two massive ferromagnetic spheres using cavity magnomechanics,” *Quantum Sci. Technol.* **6**, 024005.
- Li, Jie, Gang Li, Stefano Zippilli, David Vitali, and Tiancai Zhang, 2017, “Enhanced entanglement of two different mechanical resonators via coherent feedback,” *Phys. Rev. A* **95**, 043819.
- Li, Tongcang, Simon Kheifets, and Mark G. Raizen, 2011, “Millikelvin cooling of an optically trapped microsphere in vacuum,” *Nat. Phys.* **7**, 527–530.
- Libbrecht, Kenneth G., and Eric D. Black, 2004, “Toward quantum-limited position measurements using optically levitated microspheres,” *Phys. Lett. A* **321**, 99–102.
- Lieu, Richard, and Lloyd W. Hillman, 2003, “The phase coherence of light from extragalactic sources: Direct evidence against first-order Planck-scale fluctuations in time and space,” *Astrophys. J.* **585**, L77.
- LIGO Scientific Collaboration, 2011, “A gravitational wave observatory operating beyond the quantum shot-noise limit,” *Nat. Phys.* **7**, 962–965.
- Lindblad, Goran, 1976, “On the generators of quantum dynamical semigroups,” *Commun. Math. Phys.* **48**, 119–130.
- Lindner, Netanel H., and Asher Peres, 2005, “Testing quantum superpositions of the gravitational field with Bose-Einstein condensates,” *Phys. Rev. A* **71**, 024101.
- Liu, Jing, Xiao-Xing Jing, Wei Zhong, and Xiao-Guang Wang, 2014, “Quantum Fisher information for density matrices with arbitrary ranks,” *Commun. Theor. Phys.* **61**, 45.
- Liu, Jing, Haidong Yuan, Xiao-Ming Lu, and Xiaoguang Wang, 2020, “Quantum Fisher information matrix and multiparameter estimation,” *J. Phys. A* **53**, 023001.
- Liu, Shaopeng, B. o. Liu, Junfeng Wang, Lilong Zhao, and Wen-Xing Yang, 2021, “Gain-type optomechanically induced absorption and precise mass sensor in a hybrid optomechanical system,” *J. Appl. Phys.* **129**, 084504.
- Liu, Xinmiao, Weixin Liu, Zhihao Ren, Yiming Ma, Bowei Dong, Guangya Zhou, and Chengkuo Lee, 2021, “Progress of

- optomechanical micro/nano sensors: A review,” *Int. J. Optomechanics* **15**, 120–159.
- Liu, Yong-Chun, Yu-Wen Hu, Chee Wei Wong, and Yun-Feng Xiao, 2013, “Review of cavity optomechanical cooling,” *Chin. Phys. B* **22**, 114213.
- Liu, Yulong, Jay Mummery, Jingwei Zhou, and Mika A Sillanpää, 2021, “Gravitational Forces between Nonclassical Mechanical Oscillators,” *Phys. Rev. Appl.* **15**, 034004.
- Lloyd, Seth, 2000, “Coherent quantum feedback,” *Phys. Rev. A* **62**, 022108.
- Lock, Maximilian P.E., and Ivette Fuentes, 2017, “Dynamical Casimir effect in curved spacetime,” *New J. Phys.* **19**, 073005.
- Loll, Renate, 2020, “Quantum gravity from causal dynamical triangulations: A review,” *Classical Quantum Gravity* **37**, 013002.
- Long, Joshua C., Hilton W. Chan, Allison B. Churnside, Eric A. Gulbis, Michael C.M. Varney, and John C. Price, 2003, “Upper limits to submillimetre-range forces from extra space-time dimensions,” *Nature (London)* **421**, 922–925.
- Lopp, Richard, and Eduardo Martín-Martínez, 2018, “Light, matter, and quantum randomness generation: A relativistic quantum information perspective,” *Opt. Commun.* **423**, 29–47.
- Lushikov, I., *et al.*, 1969, “Observation of ultracold neutrons,” *JETP Lett.* **9**, 23, <http://jetpletters.ru/ps/1639/article%5F25024.shtml>.
- Lynden-Bell, D., 1961, “Stellar dynamics: Exact solution of the self-gravitation equation,” *Mon. Not. R. Astron. Soc.* **123**, 447.
- Ma, Yiqiu, Haixing Miao, Belinda Heyun Pang, Matthew Evans, Chunnong Zhao, Jan Harms, Roman Schnabel, and Yanbei Chen, 2017, “Proposal for gravitational-wave detection beyond the standard quantum limit through EPR entanglement,” *Nat. Phys.* **13**, 776–780.
- Ma, Yue, Thomas Guff, Gavin W. Morley, Igor Pikovski, and M. S. Kim, 2022, “Limits on inference of gravitational entanglement,” *Phys. Rev. Res.* **4**, 013024.
- Maggiore, Michele, 1993a, “A generalized uncertainty principle in quantum gravity,” *Phys. Lett. B* **304**, 65–69.
- Maggiore, Michele, 1993b, “The algebraic structure of the generalized uncertainty principle,” *Phys. Lett. B* **319**, 83–86.
- Magrini, Lorenzo, Philipp Rosenzweig, Constanze Bach, Andreas Deutschmann-Olek, Sebastian G. Hofer, Sungkun Hong, Nikolai Kiesel, Andreas Kugi, and Markus Aspelmeyer, 2021, “Real-time optimal quantum control of mechanical motion at room temperature,” *Nature (London)* **595**, 373–377.
- Mancini, S., V.I. Man’ko, and P. Tombesi, 1997, “Ponderomotive control of quantum macroscopic coherence,” *Phys. Rev. A* **55**, 3042.
- Manley, Jack, Mitul Dey Chowdhury, Daniel Grin, Swati Singh, and Dalziel J. Wilson, 2021, “Searching for Vector Dark Matter with an Optomechanical Accelerometer,” *Phys. Rev. Lett.* **126**, 061301.
- Mann, Robert B., and Timothy C. Ralph, 2012, “Relativistic quantum information,” *Classical Quantum Gravity* **29**, 220301.
- Mansouri, Daryoosh, Behrooz Rezaie, Abolfazl Ranjbar N, and Abolghasem Daeichian, 2022, “Cavity-assisted coherent feedback cooling of a mechanical resonator to the ground-state in the unresolved sideband regime,” *J. Phys. B* **55**, 165501.
- March, James E., *et al.*, 2023, “Long Spin Coherence and Relaxation Times in Nanodiamonds Milled from Polycrystalline  $^{12}\text{C}$  Diamond,” *Phys. Rev. Appl.* **20**, 044045.
- Margalit, Yair, *et al.*, 2021, “Realization of a complete Stern-Gerlach interferometer: Toward a test of quantum gravity,” *Sci. Adv.* **7**, eabg2879.
- Marletto, Chiara, and Vlatko Vedral, 2017, “Gravitationally Induced Entanglement between Two Massive Particles Is Sufficient Evidence of Quantum Effects in Gravity,” *Phys. Rev. Lett.* **119**, 240402.
- Marquardt, Florian, Benjamin Abel, and Jan von Delft, 2008, “Measuring the size of a quantum superposition of many-body states,” *Phys. Rev. A* **78**, 012109.
- Marquardt, Florian, Joe P. Chen, A. A. Clerk, and S. M. Girvin, 2007, “Quantum Theory of Cavity-Assisted Sideband Cooling of Mechanical Motion,” *Phys. Rev. Lett.* **99**, 093902.
- Marshall, William, Christoph Simon, Roger Penrose, and Dik Bouwmeester, 2003, “Towards Quantum Superpositions of a Mirror,” *Phys. Rev. Lett.* **91**, 130401.
- Marshman, Ryan J., Anupam Mazumdar, and Sougato Bose, 2020, “Locality and entanglement in table-top testing of the quantum nature of linearized gravity,” *Phys. Rev. A* **101**, 052110.
- Marshman, Ryan J., Anupam Mazumdar, Ron Folman, and Sougato Bose, 2021, “Large splitting massive Schrödinger kittens,” *arXiv:2105.01094*.
- Marshman, Ryan J., Anupam Mazumdar, Gavin W. Morley, Peter F. Barker, Steven Hoekstra, and Sougato Bose, 2020, “Mesoscopic interference for metric and curvature & gravitational wave detection,” *New J. Phys.* **22**, 083012.
- Martin, Ivar, Alexander Shnirman, Lin Tian, and Peter Zoller, 2004, “Ground-state cooling of mechanical resonators,” *Phys. Rev. B* **69**, 125339.
- Martinez, Lukas, Klaus Hornberger, James Millen, M. S. Kim, and Benjamin A. Stickler, 2020, “Quantum electromechanics with levitated nanoparticles,” *npj Quantum Inf.* **6**, 101.
- Martín-Martínez, Eduardo, and T. Rick Perche, 2023, “What gravity mediated entanglement can really tell us about quantum gravity,” *Phys. Rev. D* **108**, L101702.
- Martynov, D. V., *et al.*, 2016, “Sensitivity of the Advanced LIGO detectors at the beginning of gravitational wave astronomy,” *Phys. Rev. D* **93**, 112004.
- Marzlin, Karl-Peter, 1995, “Dipole coupling of atoms and light in gravitational fields,” *Phys. Rev. A* **51**, 625–631.
- Matsumura, Akira, and Kazuhiro Yamamoto, 2020, “Gravity-induced entanglement in optomechanical systems,” *Phys. Rev. D* **102**, 106021.
- McDonald, A., T. Pereg-Barnea, and A. A. Clerk, 2018, “Phase-Dependent Chiral Transport and Effective Non-Hermitian Dynamics in a Bosonic Kitaev-Majorana Chain,” *Phys. Rev. X* **8**, 041031.
- McDonald, Alexander, and Aashish A. Clerk, 2020, “Exponentially-enhanced quantum sensing with non-Hermitian lattice dynamics,” *Nat. Commun.* **11**, 5382.
- McMillen, Sam, Matteo Brunelli, Matteo Carlesso, Angelo Bassi, Hendrik Ulbricht, Matteo GA Paris, and Mauro Paternostro, 2017, “Quantum-limited estimation of continuous spontaneous localization,” *Phys. Rev. A* **95**, 012132.
- Mehdi, Zain, Joseph J. Hope, and Simon A. Haine, 2023, “Signatures of Quantum Gravity in the Gravitational Self-Interaction of Photons,” *Phys. Rev. Lett.* **130**, 240203.
- Merali, Zeeya, 2013, “Astrophysics: Fire in the hole!,” *Nature (London)* **496**, 20–23.
- Metelmann, A., and A. A. Clerk, 2015, “Nonreciprocal Photon Transmission and Amplification via Reservoir Engineering,” *Phys. Rev. X* **5**, 021025.
- Metelmann, A., and A. A. Clerk, 2017, “Nonreciprocal quantum interactions and devices via autonomous feedforward,” *Phys. Rev. A* **95**, 013837.
- Miao, Haixing, Denis Martynov, Huan Yang, and Animesh Datta, 2020, “Quantum correlations of light mediated by gravity,” *Phys. Rev. A* **101**, 063804.



- Mielnik, Bogdan, 1974, “Generalized quantum mechanics,” *Commun. Math. Phys.* **37**, 221–256.
- Mielnik, Bogdan, 1980, “Mobility of nonlinear systems,” *J. Math. Phys. (N.Y.)* **21**, 44–54.
- Miki, Daisuke, Akira Matsumura, and Kazuhiro Yamamoto, 2021, “Entanglement and decoherence of massive particles due to gravity,” *Phys. Rev. D* **103**, 026017.
- Milburn, G. J., S. Schneider, and D. F. V. James, 2000, “Ion trap quantum computing with warm ions,” *Fortschr. Phys.* **48**, 801–810.
- Milburn, T. J., M. S. Kim, and M. R. Vanner, 2016, “Nonclassical-state generation in macroscopic systems via hybrid discrete-continuous quantum measurements,” *Phys. Rev. A* **93**, 053818.
- Militaru, Andrei, Massimiliano Rossi, Felix Tebbenjohanns, Oriol Romero-Isart, Martin Frimmer, and Lukas Novotny, 2022, “Ponderomotive Squeezing of Light by a Levitated Nanoparticle in Free Space,” *Phys. Rev. Lett.* **129**, 053602.
- Millen, James, and Benjamin A. Stickler, 2020, “Quantum experiments with microscale particles,” *Contemp. Phys.* **61**, 155–168.
- Misner, Charles W., Kip S. Thorne, and John Archibald Wheeler, 1973, *Gravitation* (W. H. Freeman, New York).
- Mohageg, Makan, *et al.*, 2022, “The deep space quantum link: Prospective fundamental physics experiments using long-baseline quantum optics,” *Eur. Phys. J. Quantum Technol.* **9**, 25.
- Møller, Christian, *et al.*, 1962, “The energy-momentum complex in general relativity and related problems,” in *Les Théories Relativistes de la Gravitation-Colloques Internationaux*, CNRS Vol. 91, edited by A. Lichnerowicz and M. A. Tonnelat (CNRS, Paris), pp. 15–29.
- Møller, Christoffer B., Rodrigo A. Thomas, Georgios Vasilakis, Emil Zeuthen, Yeghishe Tsaturyan, Mikhail Balabas, Kasper Jensen, Albert Schliesser, Klemens Hammerer, and Eugene S. Polzik, 2017, “Quantum back-action-evading measurement of motion in a negative mass reference frame,” *Nature (London)* **547**, 191–195.
- Mollow, B. R., and R. J. Glauber, 1967, “Quantum theory of parametric amplification. I,” *Phys. Rev.* **160**, 1076–1096.
- Mølmer, Klaus, and Anders Sørensen, 1999, “Multiparticle Entanglement of Hot Trapped Ions,” *Phys. Rev. Lett.* **82**, 1835.
- Monteiro, Fernando, Sumita Ghosh, Adam Getzels Fine, and David C. Moore, 2017, “Optical levitation of 10-ng spheres with nano-*g* acceleration sensitivity,” *Phys. Rev. A* **96**, 063841.
- Monteiro, Fernando, Wenqiang Li, Gadi Afek, Chang-ling Li, Michael Mossman, and David C. Moore, 2020, “Force and acceleration sensing with optically levitated nanogram masses at microkelvin temperatures,” *Phys. Rev. A* **101**, 053835.
- Moody, M. Vol, Ho. Jung Paik, and Edgar R. Canavan, 2002, “Three-axis superconducting gravity gradiometer for sensitive gravity experiments,” *Rev. Sci. Instrum.* **73**, 3957–3974.
- Moore, David C., and Andrew A. Geraci, 2021, “Searching for new physics using optically levitated sensors,” *Quantum Sci. Technol.* **6**, 014008.
- Moore, Gerald T., 1970, “Quantum theory of the electromagnetic field in a variable-length one-dimensional cavity,” *J. Math. Phys. (N.Y.)* **11**, 2679–2691.
- Moroz, Irene M., Roger Penrose, and Paul Tod, 1998, “Spherically-symmetric solutions of the Schrödinger-Newton equations,” *Classical Quantum Gravity* **15**, 2733.
- Moser, Joel, Alexander Eichler, Johannes Güttinger, Mark I. Dykman, and Adrian Bachtold, 2014, “Nanotube mechanical resonators with quality factors of up to 5 million,” *Nat. Nanotechnol.* **9**, 1007–1011.
- Motazedifard, Ali, F. Bemani, M. H. Naderi, R. Roknizadeh, and D. Vitali, 2016, “Force sensing based on coherent quantum noise cancellation in a hybrid optomechanical cavity with squeezed-vacuum injection,” *New J. Phys.* **18**, 073040.
- Motazedifard, Ali, A. Dalafi, F. Bemani, and M. H. Naderi, 2019, “Force sensing in hybrid Bose-Einstein-condensate optomechanics based on parametric amplification,” *Phys. Rev. A* **100**, 023815.
- Motazedifard, Ali, A. Dalafi, and M. H. Naderi, 2021, “Ultraprecision quantum sensing and measurement based on nonlinear hybrid optomechanical systems containing ultracold atoms or atomic Bose-Einstein condensate,” *AVS Quantum Sci.* **3**, 024701.
- Müller, Holger, Achim Peters, and Steven Chu, 2010, “A precision measurement of the gravitational redshift by the interference of matter waves,” *Nature (London)* **463**, 926–929.
- Müntinga, H., *et al.*, 2013, “Interferometry with Bose-Einstein Condensates in Microgravity,” *Phys. Rev. Lett.* **110**, 093602.
- Murata, Jiro, and Saki Tanaka, 2015, “A review of short-range gravity experiments in the LHC era,” *Classical Quantum Gravity* **32**, 033001.
- Naikoo, Javid, Ashutosh Kumar Alok, Subhashish Banerjee, S. Uma Sankar, Giacomo Guarnieri, Christiane Schultze, and Beatrix C. Hiesmayr, 2020, “A quantum information theoretic quantity sensitive to the neutrino mass-hierarchy,” *Nucl. Phys.* **B951**, 114872.
- Navau, Carles, Stefan Minniberger, Michael Trupke, and Alvaro Sanchez, 2021, “Levitation of superconducting microrings for quantum magnetomechanics,” *Phys. Rev. B* **103**, 174436.
- Nelson, Richard J., Yaakov Weinstein, David Cory, and Seth Lloyd, 2000, “Experimental Demonstration of Fully Coherent Quantum Feedback,” *Phys. Rev. Lett.* **85**, 3045–3048.
- Nesvizhevsky, Valery V., *et al.*, 2002, “Quantum states of neutrons in the Earth’s gravitational field,” *Nature (London)* **415**, 297–299.
- Neukirch, Levi P., Eva von Haartman, Jessica M. Rosenholm, and A. Nick Vamivakas, 2015, “Multi-dimensional single-spin nano-optomechanics with a levitated nanodiamond,” *Nat. Photonics* **9**, 653–657.
- Neumeier, Lukas, Mario A. Ciampini, Oriol Romero-Isart, Markus Aspelmeyer, and Nikolai Kiesel, 2022, “Fast quantum interference of a nanoparticle via optical potential control,” *arXiv:2207.12539*.
- Neveu, Pascal, Jack Clarke, Michael R. Vanner, and Ewold Verhagen, 2021, “Preparation and verification of two-mode mechanical entanglement through pulsed optomechanical measurements,” *New J. Phys.* **23**, 023026.
- Newton, S. I., M. P. Bouguer, and H. Cavendish, 1900, “Experiments to determine the density of the Earth,” in *The Laws of Gravitation: Memoirs by Newton, Bouguer and Cavendish, Together with Abstracts of Other Important Memoirs*, edited by A. S. MacKenzie (American Book Co., New York), pp. 57–107.
- Ng, Keith K., Chen Zhang, Jorma Louko, and Robert B. Mann, 2022, “A little excitement across the horizon,” *New J. Phys.* **24**, 103018.
- Niedermaier, M., 2007, “The asymptotic safety scenario in quantum gravity: An introduction,” *Classical Quantum Gravity* **24**, R171–R230.
- Nimmrichter, Stefan, and Klaus Hornberger, 2013, “Macroscopicity of Mechanical Quantum Superposition States,” *Phys. Rev. Lett.* **110**, 160403.
- Nongthombam, Roson, Ambareesh Sahoo, and Amarendra K. Sarma, 2021, “Ground-state cooling of a mechanical oscillator via a hybrid electro-optomechanical system,” *Phys. Rev. A* **104**, 023509.
- Norton, J. D., 1993, *Rep. Prog. Phys.* **56**, 791–858.
- Nunnenkamp, Andreas, Kjetil Børkje, and Steven M. Girvin, 2011, “Single-photon optomechanics,” *Phys. Rev. Lett.* **107**, 063602.
- Nurdin, Hendra I., Matthew R. James, and Ian R. Petersen, 2009, “Coherent quantum LQG control,” *Automatica* **45**, 1837–1846.

- Ockeloen-Korppi, C. F., E. Damskäg, J.-M. Pirkkalainen, M. Asjad, A. A. Clerk, F. Massel, M. J. Woolley, and M. A. Sillanpää, 2018, “Stabilized entanglement of massive mechanical oscillators,” *Nature (London)* **556**, 478–482.
- Ockeloen-Korppi, C. F., E. Damskäg, J.-M. Pirkkalainen, A. A. Clerk, M. J. Woolley, and M. A. Sillanpää, 2016, “Quantum Backaction Evading Measurement of Collective Mechanical Modes,” *Phys. Rev. Lett.* **117**, 140401.
- O’Connell, Aaron D., *et al.*, 2010, “Quantum ground state and single-phonon control of a mechanical resonator,” *Nature (London)* **464**, 697–703.
- Oniga, Teodora, and Charles H.-T. Wang, 2016, “Quantum gravitational decoherence of light and matter,” *Phys. Rev. D* **93**, 044027.
- Oniga, Teodora, and Charles H.-T. Wang, 2017, “Quantum coherence, radiance, and resistance of gravitational systems,” *Phys. Rev. D* **96**, 084014.
- Onofrio, Roberto, 2006, “Casimir forces and non-Newtonian gravitation,” *New J. Phys.* **8**, 237.
- Oppenheim, Jonathan, 2023, “A Postquantum Theory of Classical Gravity?,” *Phys. Rev. X* **13**, 041040.
- Oppenheim, Jonathan, Carlo Sparaciari, Barbara Šoda, and Zachary Weller-Davies, 2022, “Gravitationally induced decoherence vs space-time diffusion: Testing the quantum nature of gravity,” *arXiv*: 2203.01982.
- Orlando, Patrick J., Robert B. Mann, Kavan Modi, and Felix A. Pollock, 2016, “A test of the equivalence principle(s) for quantum superpositions,” *Classical Quantum Gravity* **33**, 19LT01.
- Overhauser, A. W., and R. Colella, 1974, “Experimental Test of Gravitationally Induced Quantum Interference,” *Phys. Rev. Lett.* **33**, 1237.
- Overstreet, Chris, Peter Asenbaum, Joseph Curti, Minjeong Kim, and Mark A. Kasevich, 2022, “Observation of a gravitational Aharonov-Bohm effect,” *Science* **375**, 226–229.
- Overstreet, Chris, Peter Asenbaum, Tim Kovachy, Remy Notermans, Jason M. Hogan, and Mark A. Kasevich, 2018, “Effective Inertial Frame in an Atom Interferometric Test of the Equivalence Principle,” *Phys. Rev. Lett.* **120**, 183604.
- Overstreet, Chris, Joseph Curti, Minjeong Kim, Peter Asenbaum, Mark A. Kasevich, and Flaminia Giacomini, 2023, “Inference of gravitational field superposition from quantum measurements,” *Phys. Rev. D* **108**, 084038.
- Padilla, Antonio, 2015, “Lectures on the cosmological constant problem,” *arXiv*:1502.05296.
- Panda, C. D., M. Tao, J. Eggelhof, M. Ceja, A. Reynoso, V. Xu, and H. Müller, 2023, “Probing gravity for one minute with an optical-lattice atom interferometer,” in *Proceedings of the Ninth Meeting on CPT and Lorentz Symmetry, Bloomington, IN, 2022*, edited by R. Lehnert (World Scientific, Singapore), pp. 136–139.
- Pang, Shengshi, and Todd A. Brun, 2014, “Quantum metrology for a general Hamiltonian parameter,” *Phys. Rev. A* **90**, 022117.
- Parikh, Maulik, Frank Wilczek, and George Zahariade, 2020, “The noise of gravitons,” *Int. J. Mod. Phys. D* **29**, 2042001.
- Parikh, Maulik, Frank Wilczek, and George Zahariade, 2021a, “Quantum Mechanics of Gravitational Waves,” *Phys. Rev. Lett.* **127**, 081602.
- Parikh, Maulik, Frank Wilczek, and George Zahariade, 2021b, “Signatures of the quantization of gravity at gravitational wave detectors,” *Phys. Rev. D* **104**, 046021.
- Paris, Matteo G. A., 2009, “Quantum estimation for quantum technology,” *Int. J. Quantum Inf.* **07**, 125–137.
- Parker, Leonard, and David Toms, 2009, *Quantum Field Theory in Curved Spacetime: Quantized Fields and Gravity*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, England).
- Paul, Wolfgang, 1990, “Electromagnetic traps for charged and neutral particles,” *Angew. Chem., Int. Ed. Engl.* **29**, 739–748.
- Peano, V., H. G. L. Schwefel, Ch. Marquardt, and F. Marquardt, 2015, “Intracavity Squeezing Can Enhance Quantum-Limited Optomechanical Position Detection through Deamplification,” *Phys. Rev. Lett.* **115**, 243603.
- Pedernales, Julien S., Gavin W. Morley, and Martin B. Plenio, 2020, “Motional Dynamical Decoupling for Interferometry with Macroscopic Particles,” *Phys. Rev. Lett.* **125**, 023602.
- Pedernales, Julien S., and Martin B. Plenio, 2023, “On the origin of force sensitivity in tests of quantum gravity with delocalised mechanical systems,” *Contemp. Phys.* **64**, 147–163.
- Pedernales, Julien S., Kirill Streletsov, and Martin B. Plenio, 2022, “Enhancing Gravitational Interaction between Quantum Systems by a Massive Mediator,” *Phys. Rev. Lett.* **128**, 110401.
- Peirce, Anthony P., Mohammed A. Dahleh, and Herschel Rabitz, 1988, “Optimal control of quantum-mechanical systems: Existence, numerical approximation, and applications,” *Phys. Rev. A* **37**, 4950–4964.
- Pendlebury, J. M., *et al.*, 2015, “Revised experimental upper limit on the electric dipole moment of the neutron,” *Phys. Rev. D* **92**, 092003.
- Penington, Geoffrey, 2020, “Entanglement wedge reconstruction and the information paradox,” *J. High Energy Phys.* **09**, 002.
- Penrose, Roger, 1986, “Gravity and state vector reduction,” in *Quantum Concepts in Space and Time*, edited by Roger Penrose and C. J. Isham (Oxford University Press, New York), pp. 1–129.
- Penrose, Roger, 1996, “On gravity’s role in quantum state reduction,” *Gen. Relativ. Gravit.* **28**, 581–600.
- Penrose, Roger, 1998, “Quantum computation, entanglement and state reduction,” *Phil. Trans. R. Soc. A* **356**, 1927–1939.
- Penrose, Roger, 2014, “On the gravitization of quantum mechanics 1: Quantum state reduction,” *Found. Phys.* **44**, 557–575.
- Pereira dos Santos, F., X. Alauze, C. Solaro, and A. Bonnin, 2017, “Trapped atom interferometry for the study of Casimir forces and gravitation at short range,” in *Proceedings of the 52nd Rencontres de Moriond on Gravitation, La Thuile, Italy, 2017*, edited by E. Auge, J. Dumarchez, and J. Tran Thanh Van (ARISF, Paris), pp. 185–190.
- Peres, A., 1993, *Quantum Theory: Concepts and Methods* (Kluwer, Dordrecht).
- Peters, Achim, Keng Yeow Chung, and Steven Chu, 1999, “Measurement of gravitational acceleration by dropping atoms,” *Nature (London)* **400**, 849–852.
- Peters, Achim, Keng Yeow Chung, and Steven Chu, 2001, “High-precision gravity measurements using atom interferometry,” *Metrologia* **38**, 25.
- Petruzzello, Luciano, 2021, “Generalized uncertainty principle with maximal observable momentum and no minimal length indeterminacy,” *Classical Quantum Gravity* **38**, 135005.
- Pfister, Corsin, Jed Kaniewski, M. Tomamichel, A. Mantri, R. Schmucker, N. McMahon, G. Milburn, and Stephanie Wehner, 2016, “A universal test for gravitational decoherence,” *Nat. Commun.* **7**, 13022.
- Philbin, Thomas G., Chris Kuklewicz, Scott Robertson, Stephen Hill, Friedrich König, and Ulf Leonhardt, 2008, “Fiber-optical analog of the event horizon,” *Science* **319**, 1367–1370.
- Pike, W. T., I. M. Standley, S. B. Calcutt, and A. G. Mukherjee, 2018, “A broad-band silicon microseismometer with 0.25 NG/rtHz performance,” in *Proceedings of the 31st IEEE International*

- Conference on Micro Electro Mechanical Systems (MEMS)*, *Belfast, 2018* (IEEE, New York), pp. 113–116.
- Pikovski, Igor, Michael R. Vanner, Markus Aspelmeyer, M. S. Kim, and Časlav Brukner, 2012, “Probing Planck-scale physics with quantum optics,” *Nat. Phys.* **8**, 393–397.
- Pikovski, Igor, Magdalena Zych, Fabio Costa, and Časlav Brukner, 2015, “Universal decoherence due to gravitational time dilation,” *Nat. Phys.* **11**, 668–672.
- Pikovski, Igor, Magdalena Zych, Fabio Costa, and Časlav Brukner, 2017, “Time dilation in quantum systems and decoherence,” *New J. Phys.* **19**, 025011.
- Pinel, O., P. Jian, N. Treps, C. Fabre, and D. Braun, 2013, “Quantum parameter estimation using general single-mode Gaussian states,” *Phys. Rev. A* **88**, 040102.
- Pino, H., J. Prat-Camps, K. Sinha, B. Prasanna Venkatesh, and O. Romero-Isart, 2018, “On-chip quantum interference of a superconducting microsphere,” *Quantum Sci. Technol.* **3**, 025001.
- Piotrowski, Johannes, Dominik Windey, Jayadev Vijayan, Carlos Gonzalez-Ballester, Andrés de los Ríos Sommer, Nadine Meyer, Romain Quidant, Oriol Romero-Isart, René Reimann, and Lukas Novotny, 2023, “Simultaneous ground-state cooling of two mechanical modes of a levitated nanoparticle,” *Nat. Phys.* **19**, 1009–1013.
- Pirkkalainen, J.-M., S. U. Cho, Francesco Massel, J. Tuorila, T. T. Heikkilä, P. J. Hakonen, and M. A. Sillanpää, 2015, “Cavity optomechanics mediated by a quantum two-level system,” *Nat. Commun.* **6**, 6981.
- Pirkkalainen, J.-M., Erno Damskägg, Matthias Brandt, Francesco Massel, and Mika A. Sillanpää, 2015, “Squeezing of Quantum Noise of Motion in a Micromechanical Resonator,” *Phys. Rev. Lett.* **115**, 243601.
- Pitaevskii, Lev P., 1961, “Vortex lines in an imperfect Bose gas,” *Sov. Phys. JETP* **13**, 451–454.
- Plotkin-Swing, Benjamin, Daniel Gochner, Katherine E. McAlpine, Eric S. Cooper, Alan O. Jamison, and Subhadeep Gupta, 2018, “Three-Path Atom Interferometry with Large Momentum Separation,” *Phys. Rev. Lett.* **121**, 133201.
- Poggio, M., C. L. Degen, H. J. Mamin, and D. Rugar, 2007, “Feedback Cooling of a Cantilever’s Fundamental Mode below 5 mK,” *Phys. Rev. Lett.* **99**, 017201.
- Poisson, Eric, 2004, *A Relativist’s Toolkit: The Mathematics of Black-Hole Mechanics* (Cambridge University Press, Cambridge, England).
- Pokotilovski, Yu. N., 2018, “Experiments with ultracold neutrons—First 50 years,” *arXiv:1805.05292*.
- Polarski, David, and Alexei A. Starobinsky, 1996, “Semiclassicality and decoherence of cosmological perturbations,” *Classical Quantum Gravity* **13**, 377.
- Polchinski, Joseph, 1991, “Weinberg’s Nonlinear Quantum Mechanics and the Einstein-Podolsky-Rosen Paradox,” *Phys. Rev. Lett.* **66**, 397.
- Pontin, A., Michele Bonaldi, A. Borrielli, F. S. Cataliotti, F. Marino, G. A. Prodi, Enrico Serra, and F. Marin, 2014, “Squeezing a Thermal Mechanical Oscillator by Stabilized Parametric Effect on the Optical Spring,” *Phys. Rev. Lett.* **112**, 023601.
- Pontin, A., N. P. Bullier, M. Toroš, and P. F. Barker, 2020, “Ultra-narrow-linewidth levitated nano-oscillator for testing dissipative wave-function collapse,” *Phys. Rev. Res.* **2**, 023349.
- Pontin, A., H. Fu, M. Toroš, T. S. Monteiro, and P. F. Barker, 2023, “Simultaneous cavity cooling of all six degrees of freedom of a levitated nanoparticle,” *Nat. Phys.* **19**, 1003–1008.
- Poot, Menno, and Herre S. J. van der Zant, 2012, “Mechanical systems in the quantum regime,” *Phys. Rep.* **511**, 273–335.
- Priel, Nadav, Alexander Fieguth, Charles P. Blakemore, Emmett Hough, Akio Kawasaki, Denzel Martin, Gautam Venugopalan, and Giorgio Gratta, 2022, “Dipole moment background measurement and suppression for levitated charge sensors,” *Sci. Adv.* **8**, eabo2361.
- Purdy, Tom P., Robert W. Peterson, and C. A. Regal, 2013, “Observation of radiation pressure shot noise on a macroscopic object,” *Science* **339**, 801–804.
- Qin, Wei, Adam Miranowicz, Guilu Long, J. Q. You, and Franco Nori, 2019, “Proposal to test quantum wave-particle superposition on massive mechanical resonators,” *npj Quantum Inf.* **5**, 1–8.
- Qiu, Liu, Itay Shomroni, Paul Seidler, and Tobias J. Kippenberg, 2020, “Laser Cooling of a Nanomechanical Oscillator to Its Zero-Point Energy,” *Phys. Rev. Lett.* **124**, 173601.
- Qvarfort, Sofia, 2023, “Enhanced optomechanical nonlinearity through non-Markovian mechanical noise,” *arXiv:2308.01115*.
- Qvarfort, Sofia, Sougato Bose, and Alessio Serafini, 2020, “Mesoscopic entanglement through central-potential interactions,” *J. Phys. B* **53**, 235501.
- Qvarfort, Sofia, A. Douglas K. Plato, David Edward Bruschi, Fabienne Schneiter, Daniel Braun, Alessio Serafini, and Dennis Rätzel, 2021, “Optimal estimation of time-dependent gravitational fields with quantum optomechanical systems,” *Phys. Rev. Res.* **3**, 013159.
- Qvarfort, Sofia, and Igor Pikovski, 2022, “Solving quantum dynamics with a Lie algebra decoupling method,” *arXiv:2210.11894*.
- Qvarfort, Sofia, Dennis Rätzel, and Stephen Stopyra, 2022, “Constraining modified gravity with quantum optomechanics,” *New J. Phys.* **24**, 033009.
- Qvarfort, Sofia, Alessio Serafini, Peter F. Barker, and Sougato Bose, 2018, “Gravimetry through non-linear optomechanics,” *Nat. Commun.* **9**, 3690.
- Qvarfort, Sofia, Alessio Serafini, André Xuereb, Dennis Rätzel, and David Edward Bruschi, 2019, “Enhanced continuous generation of non-Gaussianity through optomechanical modulation,” *New J. Phys.* **21**, 055004.
- Qvarfort, Sofia, Michael R. Vanner, Peter F. Barker, and David Edward Bruschi, 2021, “Master-equation treatment of nonlinear optomechanical systems with optical loss,” *Phys. Rev. A* **104**, 013501.
- Rabl, Peter, 2011, “Photon Blockade Effect in Optomechanical Systems,” *Phys. Rev. Lett.* **107**, 063601.
- Rademacher, Markus, Jonathan Gosling, Antonio Pontin, Marko Toroš, Jence T. Mulder, Arjan J. Houtepen, and P. F. Barker, 2022, “Measurement of single nanoparticle anisotropy by laser induced optical alignment and Rayleigh scattering for determining particle morphology,” *Appl. Phys. Lett.* **121**, 221102.
- Rademacher, Markus, Michael Konopik, Maxime Debiossac, David Grass, Eric Lutz, and Nikolai Kiesel, 2022, “Nonequilibrium Control of Thermal and Mechanical Changes in a Levitated System,” *Phys. Rev. Lett.* **128**, 070601.
- Rademacher, Markus, James Millen, and Ying Lia Li, 2020, “Quantum sensing with nanoparticles for gravimetry: When bigger is better,” *Adv. Opt. Technol.* **9**, 227–239.
- Ragazzoni, Roberto, Massimo Turatto, and Wolfgang Gaessler, 2003, “The lack of observational evidence for the quantum structure of spacetime at Planck scales,” *Astrophys. J.* **587**, L1.
- Rahman, A. T. M. Anishur, 2019, “Large spatial Schrödinger cat state using a levitated ferrimagnetic nanoparticle,” *New J. Phys.* **21**, 113011.
- Raju, Suvrat, 2022, “Lessons from the information paradox,” *Phys. Rep.* **943**, 1–80.
- Ralph, Jason F., Marko Toroš, Simon Maskell, Kurt Jacobs, Muddassar Rashid, Ashley J. Setter, and Hendrik Ulbricht,



- 2018, “Dynamical model selection near the quantum-classical boundary,” *Phys. Rev. A* **98**, 010102.
- Ranfagni, A., K. Børkje, F. Marino, and F. Marin, 2022, “Two-dimensional quantum motion of a levitated nanosphere,” *Phys. Rev. Res.* **4**, 033051.
- Ranjit, Gambhir, Mark Cunningham, Kirsten Casey, and Andrew A. Geraci, 2016, “Zeptonewton force sensing with nanospheres in an optical lattice,” *Phys. Rev. A* **93**, 053801.
- Rashid, Muddassar, Tommaso Tufarelli, James Bateman, Jamie Vovrosh, David Hempston, M. S. Kim, and Hendrik Ulbricht, 2016, “Experimental Realization of a Thermal Squeezed State of Levitated Optomechanics,” *Phys. Rev. Lett.* **117**, 273601.
- Rätzel, Dennis, Richard Howl, Joel Lindkvist, and Ivette Fuentes, 2018, “Dynamical response of Bose-Einstein condensates to oscillating gravitational fields,” *New J. Phys.* **20**, 073044.
- Rauch, H., A. Zeilinger, G. Badurek, A. Wilfing, W. Bauspiess, and U. Bonse, 1975, “Verification of coherent spinor rotation of fermions,” *Phys. Lett.* **54A**, 425–427.
- Rauch, Helmut, and Samuel A. Werner, 2015, *Neutron Interferometry: Lessons in Experimental Quantum Mechanics, Wave-Particle Duality, and Entanglement* (Oxford University Press, Oxford).
- Regal, C. A., and K. W. Lehnert, 2011, “From cavity electromechanics to cavity optomechanics,” *J. Phys. Conf. Ser.* **264**, 012025.
- Remus, Laura G., Miles P. Blencowe, and Yukihiro Tanaka, 2009, “Damping and decoherence of a nanomechanical resonator due to a few two-level systems,” *Phys. Rev. B* **80**, 174103.
- Reuter, Martin, and Frank Saueressig, 2010, “Functional renormalization group equations, asymptotic safety, and quantum Einstein gravity,” in *Geometric and Topological Methods for Quantum Field Theory*, edited by Hernan Ocampo, Eddy Pariguan, and Sylvie Paycha (Cambridge University Press, Cambridge, England), pp. 288–329, [10.1017/CBO9780511712135.008](https://doi.org/10.1017/CBO9780511712135.008).
- Reznik, Benni, 2003, “Entanglement from the vacuum,” *Found. Phys.* **33**, 167–176.
- Riedinger, Ralf, Andreas Wallucks, Igor Marinković, Clemens Löschnauer, Markus Aspelmeyer, Sungkun Hong, and Simon Gröblacher, 2018, “Remote quantum entanglement between two micromechanical oscillators,” *Nature (London)* **556**, 473–477.
- Riera-Campeny, Andreu, Marc Roda-Llodes, Piotr T. Grochowski, and Oriol Romero-Isart, 2023, “Wigner analysis of particle dynamics in wide nonharmonic potentials,” [arXiv:2307.14106](https://arxiv.org/abs/2307.14106).
- Rijavec, Simone, Matteo Carlesso, Angelo Bassi, Vlatko Vedral, and Chiara Marletto, 2021, “Decoherence effects in non-classicality tests of gravity,” *New J. Phys.* **23**, 043040.
- Ringbauer, Martin, Till J. Weinhold, L. A. Howard, A. G. White, and M. R. Vanner, 2018, “Generation of mechanical interference fringes by multi-photon counting,” *New J. Phys.* **20**, 053042.
- Riviere, Remi, Samuel Deleglise, Stefan Weis, Emanuel Gavartin, Olivier Arcizet, Albert Schliesser, and Tobias J. Kippenberg, 2011, “Optomechanical sideband cooling of a micromechanical oscillator close to the quantum ground state,” *Phys. Rev. A* **83**, 063835.
- Roati, G., E. de Mirandes, F. Ferlaino, H. Ott, G. Modugno, and M. Inguscio, 2004, “Atom Interferometry with Trapped Fermi Gases,” *Phys. Rev. Lett.* **92**, 230402.
- Rogers, Benjamin, N. Lo Gullo, Gabriele De Chiara, G. Massimo Palma, and Mauro Paternostro, 2014, “Hybrid optomechanics for quantum technologies,” *Quantum Meas. Quantum Metrol.* **2**, 11–43.
- Romero-Isart, O., L. Clemente, C. Navau, A. Sanchez, and J. I. Cirac, 2012, “Quantum Magnetomechanics with Levitating Superconducting Microspheres,” *Phys. Rev. Lett.* **109**, 147205.
- Romero-Isart, Oriol, 2011, “Quantum superposition of massive objects and collapse models,” *Phys. Rev. A* **84**, 052121.
- Romero-Isart, Oriol, 2017, “Coherent inflation for large quantum superpositions of levitated microspheres,” *New J. Phys.* **19**, 123029.
- Romero-Isart, Oriol, Mathieu L. Juan, Romain Quidant, and J. Ignacio Cirac, 2010, “Toward quantum superposition of living organisms,” *New J. Phys.* **12**, 033015.
- Romero-Isart, Oriol, Anika C. Pflanzner, Mathieu L. Juan, Romain Quidant, Nikolai Kiesel, Markus Aspelmeyer, and J. Ignacio Cirac, 2011, “Optically levitating dielectrics in the quantum regime: Theory and protocols,” *Phys. Rev. A* **83**, 013803.
- Romero-Sanchez, Erick, Warwick P. Bowen, Michael R. Vanner, K. Xia, and Jason Twamley, 2018, “Quantum magnetomechanics: Towards the ultrastrong coupling regime,” *Phys. Rev. B* **97**, 024109.
- Roos, C. F., M. Chwalla, K. Kim, M. Riebe, and R. Blatt, 2006, “‘Designer atoms’ for quantum metrology,” *Nature (London)* **443**, 316–319.
- Rosenfeld, Leon, 1963, “On quantization of fields,” *Nucl. Phys.* **40**, 353–356.
- Rosi, G., G. D’Amico, L. Cacciapuoti, F. Sorrentino, M. Prevedelli, M. Zych, Č. Brukner, and G. M. Tino, 2017, “Quantum test of the equivalence principle for atoms in coherent superposition of internal energy states,” *Nat. Commun.* **8**, 15529.
- Rosi, G., F. Sorrentino, L. Cacciapuoti, M. Prevedelli, and G. M. Tino, 2014, “Precision measurement of the Newtonian gravitational constant using cold atoms,” *Nature (London)* **510**, 518.
- Rossi, Massimiliano, David Mason, Junxin Chen, Yeghishe Tsaturyan, and Albert Schliesser, 2018, “Measurement-based quantum control of mechanical motion,” *Nature (London)* **563**, 53–58.
- Rothleitner, Christian, and Stephan Schlamminger, 2017, “Invited review article: Measurements of the Newtonian constant of gravitation,  $G$ ,” *Rev. Sci. Instrum.* **88**, 111101.
- Roura, Albert, 2022, “Quantum probe of space-time curvature,” *Science* **375**, 142–143.
- Rovelli, Carlo, 1998, “Loop quantum gravity,” *Living Rev. Relativity* **1**, 1.
- Rudolph, Henning and Uroš Delić, Markus Aspelmeyer, Klaus Hornberger, and Benjamin A. Stickler, 2022, “Force-Gradient Sensing and Entanglement via Feedback Cooling of Interacting Nanoparticles,” *Phys. Rev. Lett.* **129**, 193602.
- Rudolph, Jan, Thomas Wilkason, Megan Nantel, Hunter Swan, Connor M. Holland, Yijun Jiang, Benjamin E. Garber, Samuel P. Carman, and Jason M. Hogan, 2020, “Large Momentum Transfer Clock Atom Interferometry on the 689 nm Intercombination Line of Strontium,” *Phys. Rev. Lett.* **124**, 083604.
- Ruffini, Remo, and Silvano Bonazzola, 1969, “Systems of self-gravitating particles in general relativity and the concept of an equation of state,” *Phys. Rev.* **187**, 1767.
- Rugar, D., and P. Grütter, 1991, “Mechanical Parametric Amplification and Thermomechanical Noise Squeezing,” *Phys. Rev. Lett.* **67**, 699.
- Sabín, Carlos, David Edward Bruschi, Mehdi Ahmadi, and Ivette Fuentes, 2014, “Phonon creation by gravitational waves,” *New J. Phys.* **16**, 085003.
- Sabulsky, D. O., I. Dutta, E. A. Hinds, B. Elder, C. Burrage, and Edmund J. Copeland, 2019, “Experiment to Detect Dark Energy Forces Using Atom Interferometry,” *Phys. Rev. Lett.* **123**, 061102.
- Sakurai, Jun John, and Eugene D. Commins, 1995, *Modern Quantum Mechanics*, revised ed. (Addison-Wesley, Reading, MA).
- Satzinger, Kevin Joseph, *et al.*, 2018, “Quantum control of surface acoustic-wave phonons,” *Nature (London)* **563**, 661–665.
- Scadron, Michael D., 2006, *Advanced Quantum Theory* (World Scientific, Singapore).

- Scala, Matteo, M. S. Kim, G. W. Morley, P. F. Barker, and S. Bose, 2013, “Matter-Wave Interferometry of a Levitated Thermal Nano-Oscillator Induced and Probed by a Spin,” *Phys. Rev. Lett.* **111**, 180403.
- Scardigli, Fabio, 1999, “Generalized uncertainty principle in quantum gravity from micro-black hole gedanken experiment,” *Phys. Lett. B* **452**, 39–44.
- Scardigli, Fabio, and Roberto Casadio, 2015, “Gravitational tests of the generalized uncertainty principle,” *Eur. Phys. J. C* **75**, 1–12.
- Scharpf, Philipp, 2017, “Simulation and visualization of gravitational waves from binary black holes,” Ph.D. thesis (Universität Stuttgart).
- Schleier-Smith, Monika H., Ian D. Leroux, Hao Zhang, Mackenzie A. Van Camp, and Vladan Vuletić, 2011, “Optomechanical Cavity Cooling of an Atomic Ensemble,” *Phys. Rev. Lett.* **107**, 143005.
- Schliesser, Albert, Olivier Arcizet, Rémi Rivière, Georg Anetsberger, and Tobias J. Kippenberg, 2009, “Resolved-sideband cooling and position measurement of a micromechanical oscillator close to the Heisenberg uncertainty limit,” *Nat. Phys.* **5**, 509–514.
- Schlippert, Dennis, Jonas Hartwig, Henning Albers, Logan L. Richardson, Christian Schubert, Albert Roura, Wolfgang P. Schleich, Wolfgang Ertmer, and Ernst M. Rasel, 2014, “Quantum Test of the Universality of Free Fall,” *Phys. Rev. Lett.* **112**, 203002.
- Schmölle, Jonas, Mathias Dragosits, Hans Hepach, and Markus Aspelmeyer, 2016, “A micromechanical proof-of-principle experiment for measuring the gravitational force of milligram masses,” *Classical Quantum Gravity* **33**, 125031.
- Schnabel, Roman, 2017, “Squeezed states of light and their applications in laser interferometers,” *Phys. Rep.* **684**, 1–51.
- Schneider, Ch., Martin Enderlein, Thomas Huber, and Tobias Schätz, 2010, “Optical trapping of an ion,” *Nat. Photonics* **4**, 772–775.
- Schinski, Birthe, Benjamin A. Stickler, and Klaus Hornberger, 2022, “Interferometric control of nanorotor alignment,” *Phys. Rev. A* **105**, L021502.
- Schinski, Björn, Stefan Nimmrichter, Benjamin A. Stickler, and Klaus Hornberger, 2019, “Macroscopicity of quantum mechanical superposition tests via hypothesis falsification,” *Phys. Rev. A* **100**, 032111.
- Schinski, Björn, Yu. Yang, Uwe von Lüpke, Marius Bild, Yiwen Chu, Klaus Hornberger, Stefan Nimmrichter, and Matteo Fadel, 2023, “Macroscopic Quantum Test with Bulk Acoustic Wave Resonators,” *Phys. Rev. Lett.* **130**, 133604.
- Schut, Martine, Andrew Geraci, Sougato Bose, and Anupam Mazumdar, 2023, “Micron-size spatial superpositions for the QGEM-protocol via screening and trapping,” *arXiv:2307.15743*.
- Schut, Martine, Alexey Grinin, Andrew Dana, Sougato Bose, Andrew Geraci, and Anupam Mazumdar, 2023, “Relaxation of experimental parameters in a quantum-gravity-induced entanglement of masses protocol using electromagnetic screening,” *Phys. Rev. Res.* **5**, 043170.
- Schut, Martine, Jules Tilly, Ryan J. Marshman, Sougato Bose, and Anupam Mazumdar, 2022, “Improving resilience of quantum-gravity-induced entanglement of masses to decoherence using three superpositions,” *Phys. Rev. A* **105**, 032411.
- Schwartz, Philip K., and Domenico Giulini, 2019a, “Post-Newtonian corrections to Schrödinger equations in gravitational fields,” *Classical Quantum Gravity* **36**, 095016.
- Schwartz, Philip K., and Domenico Giulini, 2019b, “Post-Newtonian Hamiltonian description of an atom in a weak gravitational field,” *Phys. Rev. A* **100**, 052116.
- Schwarz, John H., and Nathan Seiberg, 1999, “String theory, supersymmetry, unification, and all that,” *Rev. Mod. Phys.* **71**, S112.
- Schweber, Silvan S., 2005, *An Introduction to Relativistic Quantum Field Theory* (Dover Publications, Mineola, NY).
- Seis, Yannick, Thibault Capelle, Eric Langman, Sampo Saarinen, Eric Planz, and Albert Schliesser, 2022, “Ground state cooling of an ultracoherent electromechanical system,” *Nat. Commun.* **13**, 1507.
- Sen, Soham, Sukanta Bhattacharyya, and Sunandan Gangopadhyay, 2022, “Probing the generalized uncertainty principle through quantum noises in optomechanical systems,” *Classical Quantum Gravity* **39**, 075020.
- Seoane, P. Amaro, *et al.*, 2013, “The gravitational universe,” *arXiv:1305.5720*.
- Setter, Ashley, Marko Toroš, Jason F. Ralph, and Hendrik Ulbricht, 2018, “Real-time Kalman filter: Cooling of an optically levitated nanoparticle,” *Phys. Rev. A* **97**, 033822.
- Shaw, E. A., M. P. Ross, C. A. Hagedorn, E. G. Adelberger, and J. H. Gundlach, 2022, “Torsion-balance search for ultralow-mass bosonic dark matter,” *Phys. Rev. D* **105**, 042007.
- Shomer, Assaf, 2007, “A pedagogical explanation for the non-renormalizability of gravity,” *arXiv:0709.3555*.
- Shomroni, Itay, Liu Qiu, Daniel Malz, Andreas Nunnenkamp, and Tobias J. Kippenberg, 2019, “Optical backaction-evading measurement of a mechanical oscillator,” *Nat. Commun.* **10**, 2086.
- Siddiqi, Irfan, 2021, “Engineering high-coherence superconducting qubits,” *Nat. Rev. Mater.* **6**, 875–891.
- Sidhu, Jasminder S., *et al.*, 2021, “Advances in space quantum communications,” *IET Quantum Commun.* **2**, 182–217.
- Singh, S., L. A. De Lorenzo, I. Pikovski, and K. C. Schwab, 2017, “Detecting continuous gravitational waves with superfluid  $4\text{He}$ ,” *New J. Phys.* **19**, 073023.
- Sletten, Lucas R., Bradley A. Moores, Jeremie J. Vienne, and Konrad W. Lehnert, 2019, “Resolving Phonon Fock States in a Multimode Cavity with a Double-Slit Qubit,” *Phys. Rev. X* **9**, 021056.
- Slim, Jesse J., Clara C. Wanjura, Matteo Brunelli, Javier del Pino, Andreas Nunnenkamp, and Ewold Verhagen, 2024, “Optomechanical realization of the bosonic Kitaev-Majorana chain,” *Nature (London)* **627**, 767–771.
- Solano, Enrique, Ruynet Lima de Matos Filho, and Nicim Zagury, 1999, “Deterministic Bell states and measurement of the motional state of two trapped ions,” *Phys. Rev. A* **59**, R2539(R).
- Sonnleitner, Matthias, and Stephen M. Barnett, 2018, “Mass-energy and anomalous friction in quantum optics,” *Phys. Rev. A* **98**, 042106.
- Sorkin, Rafael D., 1993, “Impossible measurements on quantum fields,” in *Directions in General Relativity: Proceedings of the 1993 International Symposium, Vol. 2—Papers in Honor of Dieter Brill*, edited by Bei-Lok Hu and T. A. Jacobson (Cambridge University Press, Cambridge, England), pp. 293–305.
- Sorkin, Rafael D., 1994, “Quantum mechanics as quantum measure theory,” *Mod. Phys. Lett. A* **09**, 3119–3127.
- Sorrentino, F., Y.-H. Lien, G. Rosi, L. Cacciapuoti, M. Prevedelli, and G. M. Tino, 2010, “Sensitive gravity-gradiometry with atom interferometry: Progress towards an improved determination of the gravitational constant,” *New J. Phys.* **12**, 095009.
- Spaventa, Giovanni, Ludovico Lami, and Martin B. Plenio, 2023, “On tests of the quantum nature of gravitational interactions in presence of non-linear corrections to quantum mechanics,” *Quantum* **7**, 1157.
- Steinhauer, Jeff, 2014, “Observation of self-amplifying Hawking radiation in an analogue black-hole laser,” *Nat. Phys.* **10**, 864–869.
- Steinhauer, Jeff, 2022, “Confirmation of stimulated Hawking radiation, but not of black hole lasing,” *Phys. Rev. D* **106**, 102007.

- Steinhauer, Jeff, Murad Abuzarli, Tangui Aladjidi, Tom Bienaimé, Clara Piekarski, Wei Liu, Elisabeth Giacobino, Alberto Bramati, and Quentin Glorieux, 2022, “Analogue cosmological particle creation in an ultracold quantum fluid of light,” *Nat. Commun.* **13**, 2890.
- Steyerl, A., 1969, “Measurements of total cross sections for very slow neutrons with velocities from 100 m/sec to 5 m/sec,” *Phys. Lett.* **29B**, 33–35.
- Steyerl, A., J. M. Pendlebury, C. Kaufman, S. S. Malik, and A. M. Desai, 2012, “Quasielastic scattering in the interaction of ultracold neutrons with a liquid wall and application in a reanalysis of the Mambo I neutron-lifetime experiment,” *Phys. Rev. C* **85**, 065503.
- Stickler, Benjamin A., Klaus Hornberger, and M. S. Kim, 2021, “Quantum rotations of nanoparticles,” *Nat. Rev. Phys.* **3**, 589–597.
- Stickler, Benjamin A., Birthe Papendell, Stefan Kuhn, Björn Schrämski, James Millen, Markus Arndt, and Klaus Hornberger, 2018, “Probing macroscopic quantum superpositions with nanorotors,” *New J. Phys.* **20**, 122001.
- Streed, Erik W., Ananth P. Chikkatur, Todd L. Gustavson, Micah Boyd, Yoshio Torii, Dominik Schneble, Gretchen K. Campbell, David E. Pritchard, and Wolfgang Ketterle, 2006, “Large atom number Bose-Einstein condensate machines,” *Rev. Sci. Instrum.* **77**, 023106.
- Streltsov, Kirill, Julen Simon Pedernales, and Martin Bodo Plenio, 2022, “On the significance of interferometric revivals for the fundamental description of gravity,” *Universe* **8**, 58.
- Suh, J., A. J. Weinstein, C. U. Lei, E. E. Wollman, S. K. Steinke, P. Meystre, A. A. Clerk, and K. C. Schwab, 2014, “Mechanically detecting and avoiding the quantum fluctuations of a microwave field,” *Science* **344**, 1262–1265.
- Suzuki, Fumika, and Friedemann Queisser, 2015, “Environmental gravitational decoherence and a tensor noise model,” *J. Phys. Conf. Ser.* **626**, 012039.
- Szigeti, Stuart S., Onur Hosten, and Simon A. Haine, 2021, “Improving cold-atom sensors with quantum entanglement: Prospects and challenges,” *Appl. Phys. Lett.* **118**, 140501.
- Szigeti, Stuart S., Samuel P. Nolan, John D. Close, and Simon A. Haine, 2020, “High-Precision Quantum-Enhanced Gravimetry with a Bose-Einstein Condensate,” *Phys. Rev. Lett.* **125**, 100402.
- Szorkovszky, Alex, Andrew C. Doherty, Glen I. Harris, and Warwick P. Bowen, 2011, “Mechanical Squeezing via Parametric Amplification and Weak Measurement,” *Phys. Rev. Lett.* **107**, 213603.
- Tapley, Byron D., S. Bettadpur, M. Watkins, and C. Reigber, 2004, “The gravity recovery and climate experiment: Mission overview and early results,” *Geophys. Res. Lett.* **31**, L09607.
- Tavernarakis, A., A. Stavriniadis, A. Nowak, I. Tsioutsios, Adrian Bachtold, and Pierre Verlot, 2018, “Optomechanics with a hybrid carbon nanotube resonator,” *Nat. Commun.* **9**, 662.
- Tebbenjohanns, Felix, Martin Frimmer, Vijay Jain, Dominik Windey, and Lukas Novotny, 2020, “Motional Sideband Asymmetry of a Nanoparticle Optically Levitated in Free Space,” *Phys. Rev. Lett.* **124**, 013603.
- Tebbenjohanns, Felix, M. Luisa Mattana, Massimiliano Rossi, Martin Frimmer, and Lukas Novotny, 2021, “Quantum control of a nanoparticle optically levitated in cryogenic free space,” *Nature (London)* **595**, 378–382.
- Terrano, W. A., E. G. Adelberger, J. G. Lee, and B. R. Heckel, 2015, “Short-Range, Spin-Dependent Interactions of Electrons: A Probe for Exotic Pseudo-Goldstone Bosons,” *Phys. Rev. Lett.* **115**, 201801.
- Teufel, J. D., J. W. Harlow, C. A. Regal, and K. W. Lehnert, 2008, “Dynamical Backaction of Microwave Fields on a Nanomechanical Oscillator,” *Phys. Rev. Lett.* **101**, 197203.
- Teufel, John D., Tobias Donner, Dale Li, Jennifer W. Harlow, M. S. Allman, Katarina Cicak, Adam J. Sirois, Jed D. Whittaker, Konrad W. Lehnert, and Raymond W. Simmonds, 2011, “Sideband cooling of micromechanical motion to the quantum ground state,” *Nature (London)* **475**, 359–363.
- Thomas, Rodrigo A., Michał Parniak, Christoffer Østfeldt, Christoffer B. Møller, Christian Bærentsen, Yeghishe Tsaturyan, Albert Schliesser, Jürgen Appel, Emil Zeuthen, and Eugene S. Polzik, 2021, “Entanglement between distant macroscopic mechanical and spin systems,” *Nat. Phys.* **17**, 228–233.
- Thorne, Kip S., Ronald W. P. Drever, Carlton M. Caves, Mark Zimmermann, and Vernon D. Sandberg, 1978, “Quantum Non-demolition Measurements of Harmonic Oscillators,” *Phys. Rev. Lett.* **40**, 667–671.
- Tian, Zehua, Jieci Wang, Heng Fan, and Jiliang Jing, 2015, “Relativistic quantum metrology in open system dynamics,” *Sci. Rep.* **5**, 7946.
- Tiesinga, Eite, Peter J. Mohr, David B. Newell, and Barry N. Taylor, 2021, “CODATA recommended values of the fundamental physical constants: 2018,” *J. Phys. Chem. Ref. Data* **50**, 033105.
- Tilloy, Antoine, and Lajos Diósi, 2016, “Sourcing semiclassical gravity from spontaneously localized quantum matter,” *Phys. Rev. D* **93**, 024026.
- Tilly, Jules, Ryan J. Marshman, Anupam Mazumdar, and Sougato Bose, 2021, “Qudits for witnessing quantum-gravity-induced entanglement of masses under decoherence,” *Phys. Rev. A* **104**, 052416.
- Timberlake, Chris, Andrea Vinante, Francesco Shankar, Andrea Lapi, and Hendrik Ulbricht, 2021, “Probing modified gravity with magnetically levitated resonators,” *Phys. Rev. D* **104**, L101101.
- Tino, Guglielmo M., 2021, “Testing gravity with cold atom interferometry: Results and prospects,” *Quantum Sci. Technol.* **6**, 024014.
- Tobar, Germain, Sreenath K. Manikandan, Thomas Beitel, and Igor Pikovski, 2023, “Detecting single gravitons with quantum sensing,” *arXiv:2308.15440*.
- Tobar, Germain, Igor Pikovski, and Michael Edmund Tobar, 2024, “Detecting kHz gravitons from a neutron star merger with a multi-mode resonant bar,” *arXiv:2406.16898*.
- Tod, Paul, and Irene M. Moroz, 1999, “An analytical approach to the Schrödinger-Newton equations,” *Nonlinearity* **12**, 201.
- Tolra, B. Laburthe, K. M. O’Hara, J. H. Huckans, W. D. Phillips, S. L. Rolston, and J. V. Porto, 2004, “Observation of Reduced Three-Body Recombination in a Correlated 1D Degenerate Bose Gas,” *Phys. Rev. Lett.* **92**, 190401.
- Toroš, Marko, and Angelo Bassi, 2018, “Bounds on quantum collapse models from matter-wave interferometry: Computational details,” *J. Phys. A* **51**, 115302.
- Toroš, Marko, Thomas W. van de Kamp, Ryan J. Marshman, M. S. Kim, Anupam Mazumdar, and Sougato Bose, 2021, “Relative acceleration noise mitigation for nanocrystal matter-wave interferometry: Applications to entangling masses via quantum gravity,” *Phys. Rev. Res.* **3**, 023178.
- Toroš, Marko, Anupam Mazumdar, and Sougato Bose, 2024, “Loss of coherence and coherence protection from a graviton bath,” *Phys. Rev. D* **109**, 084050.
- Torres, Juan Mauricio, Ralf Betzholz, and Marc Bienert, 2019, “Optomechanical damping basis,” *J. Phys. A* **52**, 08LT02.
- Tóth, Géza, and Iagoba Apellaniz, 2014, “Quantum metrology from a quantum information science perspective,” *J. Phys. A* **47**, 424006.



- Touboul, Pierre, *et al.*, 2017, “MICROSCOPE Mission: First Results of a Space Test of the Equivalence Principle,” *Phys. Rev. Lett.* **119**, 231101.
- Touboul, Pierre, *et al.* (MICROSCOPE Collaboration), 2022, “MICROSCOPE Mission: Final Results of the Test of the Equivalence Principle,” *Phys. Rev. Lett.* **129**, 121102.
- Treutlein, Philipp, Claudiu Genes, Klemens Hammerer, Martino Poggio, and Peter Rabl, 2014, “Hybrid mechanical systems,” in *Cavity Optomechanics: Nano- and Micromechanical Resonators Interacting with Light*, Quantum Science and Technology, edited by Markus Aspelmeyer, Tobias J. Kippenberg, and Florian Marquardt (Springer, Berlin), pp. 327–351.
- Triana, Johan F., Andrés F. Estrada, and Leonardo A. Pachón, 2016, “Ultrafast Optimal Sideband Cooling under Non-Markovian Evolution,” *Phys. Rev. Lett.* **116**, 183602.
- Tsang, Mankei, and Carlton M. Caves, 2010, “Coherent Quantum-Noise Cancellation for Optomechanical Sensors,” *Phys. Rev. Lett.* **105**, 123601.
- Tsang, Mankei, and Carlton M. Caves, 2012, “Evading Quantum Mechanics: Engineering a Classical Subsystem within a Quantum Environment,” *Phys. Rev. X* **2**, 031016.
- Ulbricht, Hendrik, 2021, “Testing fundamental physics by using levitated mechanical systems,” in *Molecular Beams in Physics and Chemistry: From Otto Stern’s Pioneering Exploits to Present-Day Feats*, edited by Bretislav Friedrich and Horst Schmidt-Böcking (Springer, Cham, Switzerland), pp. 303–332.
- Unruh, William G., 1976, “Notes on black-hole evaporation,” *Phys. Rev. D* **14**, 870.
- Unruh, William George, 1981, “Experimental Black-Hole Evaporation?,” *Phys. Rev. Lett.* **46**, 1351.
- Valentini, Antony, 1991, “Non-local correlations in quantum electrodynamics,” *Phys. Lett. A* **153**, 321–325.
- van de Kamp, Thomas W., Ryan J. Marshman, Sougato Bose, and Anupam Mazumdar, 2020, “Quantum gravity witness via entanglement of masses: Casimir screening,” *Phys. Rev. A* **102**, 062807.
- van der Stam, K. M. R., E. D. van Ooijen, R. Meppelink, J. M. Vogels, and P. van der Straten, 2007, “Large atom number Bose-Einstein condensate of sodium,” *Rev. Sci. Instrum.* **78**, 013102.
- Vanner, Michael R., Igor Pikovski, Garrett D. Cole, M. S. Kim, Č. Brukner, Klemens Hammerer, Gerard J. Milburn, and Markus Aspelmeyer, 2011, “Pulsed quantum optomechanics,” *Proc. Natl. Acad. Sci. U.S.A.* **108**, 16182–16187.
- van Zoest, T., *et al.*, 2010, “Bose-Einstein condensation in microgravity,” *Science* **328**, 1540–1543.
- Veneziano, Gabriele, 1986, “A stringy nature needs just two constants,” *Europhys. Lett.* **2**, 199.
- Verlinde, Erik P., and Kathryn M. Zurek, 2021, “Observational signatures of quantum gravity in interferometers,” *Phys. Lett. B* **822**, 136663.
- Vinante, A., Mohammad Bahrami, Angelo Bassi, O. Usenko, G. H. C. J. Wijts, and T. H. Oosterkamp, 2016, “Upper Bounds on Spontaneous Wave-Function Collapse Models Using Millikelvin-Cooled Nanocantilevers,” *Phys. Rev. Lett.* **116**, 090402.
- Vinante, Andrea, Giulio Gasbarri, Christopher Timberlake, M. Toroš, and Hendrik Ulbricht, 2020, “Testing dissipative collapse models with a levitated micromagnet,” *Phys. Rev. Res.* **2**, 043229.
- Viola, Lorenza, and Roberto Onofrio, 1997, “Testing the equivalence principle through freely falling quantum objects,” *Phys. Rev. D* **55**, 455.
- Visser, Matt, and Carmen Molina-París, 2010, “Acoustic geometry for general relativistic barotropic irrotational fluid flow,” *New J. Phys.* **12**, 095014.
- Vovrosh, Jamie, Muddassar Rashid, David Hempston, James Bateman, Mauro Paternostro, and Hendrik Ulbricht, 2017, “Parametric feedback cooling of levitated optomechanics in a parabolic mirror trap,” *J. Opt. Soc. Am. B* **34**, 1421–1428.
- Wagner, T. A., S. Schlamminger, J. H. Gundlach, and E. G. Adelberger, 2012, “Torsion-balance tests of the weak equivalence principle,” *Classical Quantum Gravity* **29**, 184002.
- Wan, C., M. Scala, G. W. Morley, A. T. M. A. Rahman, Hendrik Ulbricht, James Bateman, P. F. Barker, Sougato Bose, and M. S. Kim, 2016, “Free Nano-Object Ramsey Interferometry for Large Quantum Superpositions,” *Phys. Rev. Lett.* **117**, 143003.
- Wang, Charles H. T., Robert Bingham, and J. Tito Mendonça, 2006, “Quantum gravitational decoherence of matter waves,” *Classical Quantum Gravity* **23**, L59.
- Wang, Cheng, Louise Banniard, Laure Mercier de Lépinay, and Mika A. Sillanpää, 2023, “Fast Feedback Control of Mechanical Motion Using Circuit Optomechanics,” *Phys. Rev. Appl.* **19**, 054091.
- Wang, Hui, and Miles Blencowe, 2021, “Coherently amplifying photon production from vacuum with a dense cloud of accelerating photodetectors,” *Commun. Phys.* **4**, 128.
- Wanjura, Clara C., Matteo Brunelli, and Andreas Nunnenkamp, 2020, “Topological framework for directional amplification in driven-dissipative cavity arrays,” *Nat. Commun.* **11**, 3149.
- Wanjura, Clara C., Jesse J. Slim, Javier del Pino, Matteo Brunelli, Ewold Verhagen, and Andreas Nunnenkamp, 2023, “Quadrature nonreciprocity in bosonic networks without breaking time-reversal symmetry,” *Nat. Phys.* **19**, 1429–1436.
- Warren, Warren S., Herschel Rabitz, and Mohammed Dahleh, 1993, “Coherent control of quantum dynamics: The dream is alive,” *Science* **259**, 1581–1589.
- Weaver, Matthew J., Frank Buters, Fernando Luna, Hedwig Eerkens, Kier Heeck, Sven de Man, and Dirk Bouwmeester, 2017, “Coherent optomechanical state transfer between disparate mechanical resonators,” *Nat. Commun.* **8**, 824.
- Weber, J., 1969, “Evidence for Discovery of Gravitational Radiation,” *Phys. Rev. Lett.* **22**, 1320–1324.
- Weinberg, Steven, 1965, “Infrared photons and gravitons,” *Phys. Rev.* **140**, B516.
- Weinberg, Steven, 1980, “Ultraviolet divergences in quantum theories of gravitation,” in *General Relativity: An Einstein Centenary Survey*, edited by Stephen W. Hawking and W. Israel (Cambridge University Press, Cambridge, England), pp. 790–831.
- Weinberg, Steven, 1989a, “Precision Tests of Quantum Mechanics,” *Phys. Rev. Lett.* **62**, 485.
- Weinberg, Steven, 1989b, “Testing quantum mechanics,” *Ann. Phys. (N.Y.)* **194**, 336–386.
- Weiss, T., M. Roda-Llodes, E. Torrontegui, M. Aspelmeyer, and O. Romero-Isart, 2021, “Large Quantum Delocalization of a Levitated Nanoparticle Using Optimal Control: Applications for Force Sensing and Entangling via Weak Forces,” *Phys. Rev. Lett.* **127**, 023601.
- Werner, Michael, Philip K. Schwartz, Jan-Niclas Kirsten-Siemß, Naceur Gaaloul, Domenico Giulini, and Klemens Hammerer, 2024, “Atom interferometers in weakly curved spacetimes using Bragg diffraction and Bloch oscillations,” *Phys. Rev. D* **109**, 022008.

- Westphal, Tobias, Hans Hepach, Jeremias Pfaff, and Markus Aspelmeyer, 2021, “Measurement of gravitational coupling between millimetre-sized masses,” *Nature (London)* **591**, 225–228.
- Whittle, Chris, *et al.*, 2021, “Approaching the motional ground state of a 10-kg object,” *Science* **372**, 1333–1336.
- Wiener, Norbert, 1930, “Generalized harmonic analysis,” *Acta Math.* **55**, 117–258.
- Wilson, C. M., G. Johansson, A. Pourkabirian, M. Simoen, J. R. Johansson, T. Duty, F. Nori, and P. Delsing, 2011, “Observation of the dynamical Casimir effect in a superconducting circuit,” *Nature (London)* **479**, 376–379.
- Windey, Dominik, Carlos Gonzalez-Ballester, Patrick Maurer, Lukas Novotny, Oriol Romero-Isart, and René Reimann, 2019, “Cavity-Based 3D Cooling of a Levitated Nanoparticle via Coherent Scattering,” *Phys. Rev. Lett.* **122**, 123601.
- Winstone, George, Robert Bennett, Markus Rademacher, Muddassar Rashid, Stefan Buhmann, and Hendrik Ulbricht, 2018, “Direct measurement of the electrostatic image force of a levitated charged nanoparticle close to a surface,” *Phys. Rev. A* **98**, 053831.
- Wiseman, H. M., 1994, “Quantum theory of continuous feedback,” *Phys. Rev. A* **49**, 2133–2150.
- Wiseman, H. M., 1996, “Quantum trajectories and quantum measurement theory,” *Quantum Semiclassical Opt.* **8**, 205.
- Wiseman, H. M., and G. J. Milburn, 1993, “Quantum theory of field-quadrature measurements,” *Phys. Rev. A* **47**, 642–662.
- Wiseman, Howard M., and Gerard J. Milburn, 2009, *Quantum Measurement and Control* (Cambridge University Press, Cambridge, England).
- Wollack, E. Alex, Agnetta Y. Cleland, Rachel G. Gruenke, Zhaoyou Wang, Patricio Arrangoiz-Arriola, and Amir H. Safavi-Naeini, 2022, “Quantum state preparation and tomography of entangled mechanical resonators,” *Nature (London)* **604**, 463–467.
- Wollman, Emma Edwina, C. U. Lei, A. J. Weinstein, J. Suh, A. Kronwald, F. Marquardt, Aashish A. Clerk, and K. C. Schwab, 2015, “Quantum squeezing of motion in a mechanical resonator,” *Science* **349**, 952–955.
- Wood, B. D., S. Bose, and G. W. Morley, 2022, “Spin dynamical decoupling for generating macroscopic superpositions of a free-falling nanodiamond,” *Phys. Rev. A* **105**, 012824.
- Wood, B. D., *et al.*, 2022, “Long spin coherence times of nitrogen vacancy centers in milled nanodiamonds,” *Phys. Rev. B* **105**, 205401.
- Woolley, M. J., and A. A. Clerk, 2013, “Two-mode back-action-evading measurements in cavity optomechanics,” *Phys. Rev. A* **87**, 063846.
- Wu, Lian-Jie, Hao-Sheng Zeng, and Shu-Min Wu, 2023, “Quantum coherence of multi-partite fermionic fields in non-inertial frames beyond single-mode approximation,” *Quantum Inf. Process.* **22**, 377.
- Wu, Meng-Zhi, Marko Toroš, Sougato Bose, and Anupam Mazumdar, 2023, “Quantum gravitational sensor for space debris,” *Phys. Rev. D* **107**, 104053.
- Wu, Shu-Min, Dan-Dan Liu, Xiao-Wei Fan, Wen-Mei Li, Xiao-Li Huang, and Hao-Sheng Zeng, 2023, “Classifying quantum steering, entanglement, and discord for continuous variables in Schwarzschild spacetime,” *Quantum Inf. Process.* **22**, 372.
- Xia, Yi, Aman R. Agrawal, Christian M. Pluchar, Anthony J. Brady, Zhen Liu, Quntao Zhuang, Dalziel J. Wilson, and Zheshen Zhang, 2023, “Entanglement-enhanced optomechanical sensing,” *Nat. Photonics* **17**, 470–477.
- Xiong, Hao, and Ying Wu, 2018, “Fundamentals and applications of optomechanically induced transparency,” *Appl. Phys. Rev.* **5**, 031305.
- Xu, Qidong, and M. P. Blencowe, 2022, “Optomechanical Quantum Entanglement Mediated by Acoustic Phonon Fields,” *Phys. Rev. Lett.* **129**, 203604.
- Xu, Victoria, Matt Jaffe, Cristian D. Panda, Sofus L. Kristensen, Logan W. Clark, and Holger Müller, 2019, “Probing gravity by holding atoms for 20 seconds,” *Science* **366**, 745–749.
- Xu, Xunnong, and Jacob M. Taylor, 2014, “Squeezing in a coupled two-mode optomechanical system for force sensing below the standard quantum limit,” *Phys. Rev. A* **90**, 043848.
- Yang, Huan, Haixing Miao, Da-Shin Lee, Bassam Helou, and Yanbei Chen, 2013, “Macroscopic Quantum Mechanics in a Classical Spacetime,” *Phys. Rev. Lett.* **110**, 170401.
- Yeo, Inah, *et al.*, 2014, “Strain-mediated coupling in a quantum dot–mechanical oscillator hybrid system,” *Nat. Nanotechnol.* **9**, 106–110.
- Yi, Bin, Urbasi Sinha, Dipankar Home, Anupam Mazumdar, and Sougato Bose, 2021, “Massive spatial qubits for testing macroscopic nonclassicality and Casimir induced entanglement,” *arXiv:2106.11906*.
- Yi, Bin, Urbasi Sinha, Dipankar Home, Anupam Mazumdar, and Sougato Bose, 2022, “Spatial qubit entanglement witness for quantum natured gravity,” *arXiv:2211.03661*.
- Yin, Zhang-qi, Tongcang Li, Xiang Zhang, and L. M. Duan, 2013, “Large quantum superpositions of a levitated nanodiamond through spin-optomechanical coupling,” *Phys. Rev. A* **88**, 033614.
- Yoshida, Beni, 2019, “Firewalls vs. scrambling,” *J. High Energy Phys.* **10**, 132.
- Youssefi, Amir, Shingo Kono, Andrea Bancora, Mahdi Chegnizadeh, Jiahe Pan, Tatiana Vovk, and Tobias J. Kippenberg, 2022, “Topological lattices realized in superconducting circuit optomechanics,” *Nature (London)* **612**, 666–672.
- Youssefi, Amir, Shingo Kono, Mahdi Chegnizadeh, and Tobias J. Kippenberg, 2022, “A squeezed mechanical oscillator with millisecond quantum decoherence,” *arXiv:2208.13082*.
- Youssefi, Amir, Shingo Kono, Mahdi Chegnizadeh, and Tobias J. Kippenberg, 2023, “A squeezed mechanical oscillator with millisecond quantum decoherence,” *Nat. Phys.* **19**, 1697–1702.
- Zeldovich, Ya. B., 1959, “Storage of cold neutrons,” *Sov. Phys. JETP* **9**, 1389, <http://jetp.ras.ru/cgi-bin/e/index/e/9/6/p1389?a=list>.
- Zhang, Wei-Min, Ping-Yuan Lo, Heng-Na Xiong, Matisse Wei-Yuan Tu, and Franco Nori, 2012, “General Non-Markovian Dynamics of Open Quantum Systems,” *Phys. Rev. Lett.* **109**, 170402.
- Zhang, Wen-Zhao, Yan Han, Biao Xiong, and Ling Zhou, 2017, “Optomechanical force sensor in a non-Markovian regime,” *New J. Phys.* **19**, 083022.
- Zhang, Zheshen, and Quntao Zhuang, 2021, “Distributed quantum sensing,” *Quantum Sci. Technol.* **6**, 043001.
- Zhao, Wen, Sheng-Dian Zhang, Adam Miranowicz, and Hui Jing, 2020, “Weak-force sensing with squeezed optomechanics,” *Sci. China Phys. Mech. Astron.* **63**, 224211.
- Zhou, Run, Ryan J. Marshman, Sougato Bose, and Anupam Mazumdar, 2022, “Catapulting towards massive and large spatial quantum superposition,” *Phys. Rev. Res.* **4**, 043157.
- Zhou, Run, Ryan J. Marshman, Sougato Bose, and Anupam Mazumdar, 2023, “Mass-independent scheme for enhancing spatial quantum superpositions,” *Phys. Rev. A* **107**, 032212.

- Zhou, Sisi, Mengzhen Zhang, John Preskill, and Liang Jiang, 2018, “Achieving the Heisenberg limit in quantum metrology using quantum error correction,” *Nat. Commun.* **9**, 78.
- Zhuang, Quntao, Zheshen Zhang, and Jeffrey H. Shapiro, 2018, “Distributed quantum sensing using continuous-variable multipartite entanglement,” *Phys. Rev. A* **97**, 032329.
- Zimmer, Oliver, Florian M. Piegsa, and Sergey N. Ivanov, 2011, “Superthermal Source of Ultracold Neutrons for Fundamental Physics Experiments,” *Phys. Rev. Lett.* **107**, 134801.
- Zwierz, Marcin, Carlos A. Pérez-Delgado, and Pieter Kok, 2010, “General Optimality of the Heisenberg Limit for Quantum Metrology,” *Phys. Rev. Lett.* **105**, 180402.
- Zych, Magdalena, and Časlav Brukner, 2018, “Quantum formulation of the Einstein equivalence principle,” *Nat. Phys.* **14**, 1027–1031.
- Zych, Magdalena, Fabio Costa, Igor Pikovski, and Časlav Brukner, 2011, “Quantum interferometric visibility as a witness of general relativistic proper time,” *Nat. Commun.* **2**, 505.